Published in final edited form as:

Epilepsy Res. 2009 August ; 85(2-3): 150–161. doi:10.1016/j.epilepsyres.2009.03.006.

Postnatal neurogenesis as a therapeutic target in temporal lobe epilepsy

Helen E. Scharfman^{1,2} and Daniel P. McCloskey³

¹ Center for Dementia Research, The Nathan Kline Institute, 140 Old Orangeburg Rd., Bldg. 35, Orangeburg, NY 10962

² Departments of Child & Adolescent Psychiatry and Physiology & Neuroscience, New York University Langone Medical Center, 550 First Ave. New York, NY 10021

³ Department of Psychology and Program in Developmental Neuroscience, College of Staten Island-CUNY, Staten Island, NY 10314

Abstract

After it was first identified that seizures increase neurogenesis in the adult brain of laboratory animals, the idea that postnatal neurogenesis may be involved in epilepsy became a topic of widespread interest. Since that time, two perspectives have developed. They primarily address temporal lobe epilepsy (TLE), because the data have either been based on animal models of TLE or patients with intractable TLE. The first perspective is that postnatal neurogenesis contributes to the predisposition for seizures in TLE. This premise is founded in the observations showing that there is a dramatic rise in neurogenesis after many types of insults or injuries which ultimately lead to TLE. As a result of the increase in neurogenesis, several changes in the dentate gyrus occur, and the net effect appears to be an increase in excitability. One of the changes is the formation of a population of granule cells (GCs) that mismigrate, leading to ectopic granule cells in the hilus (hilar EGCs) that exhibit periodic bursts of action potentials, and contribute to recurrent excitatory circuitry. Atypical dendrites also form on a subset of GCs, and project into the hilus (hilar basal dendrites). Hilar basal dendrites appear to preferentially increase the glutamatergic input relative to GABAergic synapses, increasing excitability of the subset of GCs that form hilar basal dendrites. The alternate view is that postnatal neurogenesis is a homeostatic mechanism in epilepsy that maintains normal excitability. This idea is supported by studies showing that some of the new GCs that are born after seizures, and migrate into the correct location, have normal or reduced excitability.

Here we suggest that both perspectives may be important when considering a therapeutic strategy. It would seem advantageous to limit the numbers of mismigrating GCs and hilar basal dendrites, but maintain normal neurogenesis because it is potentially homeostatic. Maintaining normal neurogenesis is also important because it has been suggested that a decrease in dentate gyrus neurogenesis contributes to depression. It is challenging to design a strategy that would achieve these goals, and it is also difficult to propose how one could administer such a therapy prophylactically, that is, as an “antiepileptogenic” approach. Another issue to address is how a therapeutic intervention with these goals could be successful if it were administered after

Address for correspondence: The Nathan Kline Institute, 140 Old Orangeburg Rd., Bldg. 35, Orangeburg, NY 10962, Phone: (845) 398-5427, Fax: (845) 398-5422, E-mail: E-mail: hscharfman@nki.rfmh.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

chronic seizures develop, when most patients seek therapy. Although difficult, a number of approaches are possible, and technical advances suggest that there are more on the horizon.

I. BACKGROUND

A. Postnatal neurogenesis

It is now widely-accepted that neurogenesis occurs throughout the lifetime of mammals (Gage et al., 2008). The two places where this is best established include the dentate gyrus subgranular zone (SGZ; Figure 1) and the subventricular zone (SVZ), adjacent to the ventricular walls (Gage et al., 2008). Here we focus on the dentate gyrus primarily, because a substantial body of research about neurogenesis in epilepsy has focused on dentate gyrus neurogenesis. Temporal lobe epilepsy (TLE) is emphasized, because pathophysiology in the dentate gyrus is most relevant to this type of epilepsy. However, other types of epilepsy may also be important to consider, because it is possible that pathology in the dentate gyrus, or its effects on downstream targets (the hippocampus and elsewhere) contribute to other types of seizure syndromes. It is also important to bear in mind that the SVZ appears to generate cells that migrate into diverse parts of the brain, beyond the site of termination that is considered to be the main target, the olfactory bulb (Shapiro et al., 2009). Because of the potential for the SVZ to generate cells that migrate to diverse locations, it is possible that many types of epilepsy are influenced by the postnatal neurogenesis in the SVZ. Another consideration is that some types of neurons may be generated by cells outside the SGZ and SVZ. A recent report suggested that oligodendrocyte precursors that express NG2 (NG2 cells) in the piriform cortex may become cortical neurons (Rivers, 2008), which is potentially important in epilepsy, because the piriform cortex is an area which is epileptogenic, and sustains injury in animal models of epilepsy (Gale, 1992; Loscher and Ebert, 1996; McIntyre and Gilby, 2008).

In the postnatal dentate gyrus, the majority of progenitors reside in the SGZ, which is situated just below the granule cell layer (GCL), at the border with the hilus (Figure 2). Early progenitors are radial glia-like type 1 cells, which can be identified by expression of the astrocytic marker glial-fibrillary acid protein (GFAP), nestin or other markers (Figure 2). Although classification into Type 1a cells, Type 2a cells, Type 2b cells, and Type 3 cells, implies a sequential maturation in this order, it has not been proven definitively. However, it is clear that the progenitors typically commit to a neuronal fate and become GCs. Thus, under normal conditions, migrating GCs move into the GCL, where it is thought that they initially reside within the “inner” third of the layer, close to the hilus, before moving to the center and outer parts of the layer (Gage et al., 2008; Ge et al., 2008; Zhao et al., 2008). For the present discussion, it is notable that there may be less organization of the new GCs within the GCL in the epileptic brain and many GCs that migrate outside it (discussed further below).

Normal GCs develop dendrites that extend into the molecular layer and reach the hippocampal fissure. The normal inputs to the GCs that lie in the molecular layer, such as the perforant path, appear to innervate normal GCs in a similar manner as those that are born in adult life (van Praag et al., 2002). However, in TLE, this may not be true, because the perforant path fibers originate in layer II of the entorhinal cortex, which is an area of damage in patients with TLE (Scharfman, 2002). In rodent models of TLE, layer III is the area that is damaged primarily, so the issue is not easily studied.

The GC axon, also called a “mossy fiber,” projects to stratum lucidum of area CA3, and collateralizes in the hilus (Amaral et al., 2007; Ribak and Shapiro, 2007). GCs that are generated postnatally develop axonal projections which are very similar to the GCs born during development, i.e. the mossy fiber pathway appears the same (Faulkner et al., 2008; Gage et al., 2008; Ge et al., 2008; Toni et al., 2008; Zhao et al., 2008). In SE models of TLE, new neurons appear to develop a normal mossy fiber projection (Parent et al., 1999; Jessberger et al.,

2007), except for those that enter the hilus, which contribute to the plexus of mossy fibers that are in the inner molecular layer (Scharfman et al., 2000).

Although the cells that immediately arise as a result of progenitor division do not typically survive long-term, those that do survive and become GCs appear to persist (Kempermann et al., 2003). Several factors can alter the numbers of GCs that ultimately survive, such as age (Cuppini et al., 2006) and learning (Leuner et al., 2004). GCs that are born after adult SE appear to survive for the rest of the lifespan, similar to GCs that develop in the normal adult brain (Scharfman et al., 2000; McCloskey et al., 2006; Jessberger et al., 2007), although some studies suggest this is not true. The discrepancies are likely to be due to the fact that some experiments examined the cells that resulted from SE-induced proliferation, rather than SE-induced neurogenesis (Parent et al., 1997; Hattiangady et al., 2004).

Recordings from new neurons in the adult brain have shown that they develop characteristics similar to GCs born during development (Laplagne et al., 2006). However, when adult-born neurons are young, they demonstrate increased excitability. For example, new neurons appear to have T-type Ca^{2+} channels that facilitate Na^{+} -dependent action potential generation, and long-term potentiation (LTP) is easier to induce (Schmidt-Hieber et al., 2004). Other studies of postnatal neurogenesis show that it plays an important role in LTP of the perforant path input (Wojtowicz, 2008). Another characteristic of young GCs that are born in the adult is that they have a depolarized equilibrium potential for chloride, making them depolarize in response to GABA more than adult cells (Ge et al., 2006; Markwardt and Overstreet-Wadiche, 2008). There also are additional changes in GABAergic transmission that also increase in excitability of young neurons (Karten et al., 2006; Ge et al., 2007).

B. Postnatal neurogenesis as a contributing factor in epileptogenesis

1. Overview—Studies of dentate gyrus neurogenesis in the 1990's noted that postnatal neurogenesis is dynamic, regulated by many genes, and epigenetic factors. Activity is a major regulator of neurogenesis, so it might have been predicted that seizures would also. In 1997, it was shown that single after discharges increased neurogenesis, as did convulsants such as pilocarpine and kainic acid (Benzon et al., 1997; Parent et al., 1997). Because these convulsants are commonly used to initiate an epileptic state in laboratory animals, the new data suggested potential relevance of seizure-induced neurogenesis to epilepsy, and became a topic of widespread interest. It has now been shown that almost every method to induce seizures leads to an increase in the rate of dentate gyrus proliferation or neurogenesis (kainic acid i.c.v. (Gray and Sundstrom, 1998); kainic acid i.p. (Scharfman et al., 2000); kindling (Parent et al., 1998; Scott et al., 1998); electroconvulsive shock; (Scott et al., 2000)), and this occurs in both rats and mice (Jessberger et al., 2005; Overstreet-Wadiche et al., 2006). However, there are exceptions: experimental febrile seizures (Bender et al., 2003), and immature animals treated with convulsants (Porter, 2008), or several episodes of convulsant administration (Liu et al., 2003) do not appear to exhibit a change in the rate of neurogenesis thereafter, or the rate declines. Although other aspects of neurogenesis may be altered, such as maturation and survival of newly-generated neurons (Liu et al., 2003; Porter, 2008), it remains important to use caution in generalizing the data from animal models that evaluate adult animals using SE as the initiating mechanism to induce a state of epilepsy.

Although the early studies were intriguing, they did not clarify how a change in neurogenesis might contribute to epileptogenesis or chronic epilepsy. Several hypotheses developed based on anatomical and functional studies. Taken together, they support the concept that SE-induced neurogenesis leads to abnormal GCs, and these neurons contribute to increased excitability. They also suggest that some of the neurons born after SE are not normal, and these GCs may not contribute to an increase in excitability. Furthermore, the neurons that are abnormal appear to primarily develop in the weeks after SE, whereas long afterwards, the ability to generate

new neurons declines. The late decline in neurogenesis is potentially important, because it could contribute to cognitive impairment and depression in TLE, as discussed elsewhere (Hattiangady et al., 2004; Kuruba and Shetty, 2007; Scharfman and Gray, 2007). Therefore, there is a time-dependent and non-linear relationship between neurogenesis and TLE, diagrammed in Figure 3. Nevertheless, treatment strategies could be developed, theoretically (Figure 3 and below).

2. SE- induced changes in neurogenesis that increase excitability

a) Hilar EGCs: Recordings from animals that had experienced SE in adulthood, and demonstrated a dramatic rise in neurogenesis thereafter, led to the hypothesis that neurons which became abnormal in the epileptic brain may contribute to the increase in excitability. Evidence for this hypothesis was founded in early observations by Parent et al. (Parent et al., 1997), where it was shown that some of the neurons that were born in the dentate gyrus after pilocarpine-induced SE did not necessarily migrate correctly into the GCL. Instead, these cells appeared to migrate into the hilus, an area between GCL and CA3, where they formed ectopic clusters of GCs (hilar EGCs; Figure 4). Subsequent studies showed that hilar EGCs often exhibited dendrites that were not typical of GCs, although hilar EGC axons were extremely similar to axons of GCs. Thus, hilar EGCs generated mossy fibers that had the specializations and projection sites of normal GCs, including the inner molecular layer (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007), typical of the GC in SE models of TLE that exhibits mossy fiber “sprouting” (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007). Importantly, the hilar EGCs survived for long periods of time, so once they developed, their effects could persist (Scharfman et al., 2000; McCloskey et al., 2006). Other studies of hilar EGCs also demonstrate their long-term survival (Jessberger et al., 2007). Importantly, hilar EGCs could become a substantial fraction of hilar neurons (Scharfman et al., 2000; McCloskey et al., 2006).

In studies using hippocampal slices from epileptic rats that had pilocarpine-induced SE, many hilar EGCs exhibited spontaneous, intermittent bursts of action potentials, synchronized with CA3 pyramidal cells and other hilar neurons (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007). The burst discharges did not occur if rats were examined before spontaneous, recurrent seizures developed, or in animals without SE (Scharfman et al., 2000). Further studies, based on simultaneous recordings, suggested that bursts of action potentials in hilar EGCs are initiated by synaptic input from CA3 (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007). Thus, simultaneous recordings showed that the onset of burst discharges in CA3 pyramidal cells preceded the onset of bursts in hilar EGCs (Figure 5; (Scharfman et al., 2000; Scharfman et al., 2007)). Although these recordings were conducted *in vitro*, hilar EGCs were also active *in vivo*, and appeared to participate in seizures, because c-fos expression developed in hilar EGCs shortly after a spontaneous seizure in epileptic animals (Scharfman et al., 2002). Interestingly, the numbers of hilar EGCs were correlated with the frequency of stage 5 limbic seizures in the pilocarpine model (McCloskey et al., 2006). These data led to the hypothesis that hilar EGCs may increase dentate gyrus excitability in animal models of TLE, and possibly predispose the hippocampus to seizures (Parent and Lowenstein, 2002; Scharfman, 2004; Kuruba and Shetty, 2007; Scharfman and Gray, 2007; Parent and Murphy, 2008).

b) Hilar basal dendrites: Another abnormal type of GC that appears to develop in animal models of TLE is one that develops hilar basal dendrites. Numerous studies have been published that describe these cells in detail anatomically (Spigelman et al., 1998; Buckmaster, 1999; Ribak et al., 2000; Dashtipour et al., 2003; Jessberger et al., 2007; Walter et al., 2007; Thind et al., 2008), and they suggest that the hilar basal dendrites would lead to a greater net excitatory input to the cell, leading the cell to be closer to threshold. By extension, these cells

would increase the excitability of the network (Ribak and Dashtipour, 2002; Shapiro and Ribak, 2005; Danzer, 2008; Shapiro et al., 2008).

Why hilar basal dendrites form after SE is not completely clear at the present time, but it appears that the increase in immature neurons that occurs after SE may be one contributing factor, because immature GCs appear to be predisposed to develop hilar basal dendrites (Walter et al., 2007). Because the neurotrophin BDNF (brain-derived neurotrophic factor) increases after SE (Scharfman, 2005), and it facilitates the formation of hilar basal dendrites (Danzer, 2008), SE-induced increases in BDNF in the dentate gyrus may also contribute to the emergence hilar basal dendrites on GCs after SE.

Despite the striking changes in hilar basal dendrites in the literature, there have been few published recordings from GCs with hilar basal dendrites, so the functional influence of these neurons is not clear at this time. Nevertheless, computational modeling would suggest a potentially important role (Morgan and Soltesz, 2008).

3. Summary and caveats—Dentate gyrus neurogenesis changes dramatically after severe insults or injury in the adult rodent, and this leads to a dramatic rise in the rate of neurogenesis. The net effects include generation of abnormal GCs, which appear to increase excitability, as well as normal GCs, which do not necessarily increase excitability. The implications are that targeting the changes that cause increased excitability may be important in acquired TLE. In addition, one might want to preserve the aspects of neurogenesis that do not contribute to an increase in excitability, and long-term, one would want to preserve postnatal neurogenesis so that cognitive function and mood are maintained. We propose that the characteristics to target therapeutically include EGCs and hilar basal dendrites, while the adult-generated neurons that migrate normally and behave normally should be preserved, as well as the neurogenic niche (Figure 3).

It is important to emphasize that this strategy is based on a number of assumptions that have yet to be proved. For example, there has been no proof that increased neurogenesis, hilar EGCs or hilar basal dendrites are essential to epileptogenesis or chronic seizures. It was shown that hilar EGC number correlates with seizure frequency (McCloskey et al., 2006), but correlative data are not definitive. Hilar EGCs occur in patients with intractable TLE (Parent et al., 2006), which suggests that what is studied in the animal model has relevance to human TLE, but the relationship between EGCs and human TLE is not clear at this time. Two studies were conducted using the lithium-pilocarpine model of epilepsy by Jung and colleagues, and provided evidence that reducing neurogenesis and hilar EGCs was accompanied by a reduction in seizures (Jung et al., 2004; Jung et al., 2006). However, the procedures that were used to limit neurogenesis also had other effects besides those that reduced neurogenesis (Parent, 2005). One argument has been that SE is sometimes followed by seizures relatively soon thereafter, sooner than new GCs, and hilar basal dendrites, would be likely to mature and form strong interconnections. However, new neurons and process outgrowth develop quite rapidly after SE (Shapiro et al., 2007). Furthermore, the days after SE may be accompanied by changes in neurons that were almost mature at the time of SE, causing disruption to the end stages of their maturation. In addition, the earliest of seizures could be an acute response to the metabolic effects of SE, which can be severe, and not reflect the seizures which develop long-term. Below we consider potential therapeutic strategies based on the assumption that abnormal GCs contribute to epileptogenesis, and are an important target for intervention.

II. Therapeutic strategies

A. General strategies

1. Hilar EGCs—In light of the discussion above, a therapeutic strategy might include preventing hilar EGC formation. Although reasons why hilar EGCs form are not clear at this time, there appear to be two factors that contribute. One is the rapid increase in proliferation that occurs early after insult or injury. This is likely to lead to a surge in the production of new GCs, some of which will become hilar EGCs. Therefore, reducing the initial increase in proliferation that follows an initial insult could be an important strategy to reduce hilar EGCs. However, one of the problems with this idea is that some of the new neurons that would potentially be reduced might not be hilar EGCs, but neurons that would migrate into the GCL and potentially maintain excitability rather than increase it (Jakubs et al., 2006).

To target only those neurons that migrate incorrectly, one approach would be reducing stromal derived factor -1 (SDF-1), a “go” signal for migration (Bagri et al., 2002) that appears to be upregulated after seizures (Jung et al., 2008). Alternatively, it might be useful to preserve reelin, which is a “stop” signal for neuronal migration of GCs away from the GCL (Frotscher, 1997). Recent studies have demonstrated that hilar neurons which co-localize GABA and somatostatin also express reelin, and may prevent GCs from migrating into the hilus (Gong et al., 2007). These neurons are typically lost after SE because of their vulnerability, and the loss of the hilar reelin appears to cause hilar EGC formation (Gong et al., 2007). A third strategy is to reduce the abnormal activity that develops in hilar EGCs, i.e., the spontaneous rhythmic burst discharges. The periodic discharges appear to develop because CA3 pyramidal neurons project to hilar EGCs, and CA3 pyramidal cells develop intermittent burst discharges (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007). Methods to prevent the circuits that develop among hilar EGCs, CA3 and other neurons seem particularly important, because the ongoing network activity increases over time (Scharfman et al., 2000). Therefore, this circuitry could contribute to a progression of pathophysiology (Sutula and Hermann, 1999; Pitkanen and Sutula, 2002; Sutula, 2004). Even if hilar EGC formation could not be prevented, stopping the activity in these circuits might be valuable. Another strategy would be to reduce the survival of the hilar EGCs, when they form. Normalizing the lifespan of adult-generated neurons after SE could be extremely effective, because any increase in excitability that would be caused would self-terminate. In summary, it may be useful to target three characteristics that contribute to hilar EGCs and the presumed disruption of normal excitability they cause: the changes in expression of factors like SDF-1 and reelin that normally control migration, the formation of recurrent excitatory circuits that include hilar EGCs, and persistence of hilar EGCs.

2. Hilar basal dendrites—Preventing or reversing the formation of hilar basal dendrites seems a logical strategy given the evidence that they are likely to increase excitability (Shapiro et al., 2008). However, how to do so is difficult to propose, because the reasons why hilar basal dendrites form are not clear. It was shown that a large percentage of neurons with hilar basal dendrites are immature at the time of SE (Walter et al., 2007), suggesting that a cause of hilar basal dendrite formation is disruption of normal maturation of the dendritic tree. However, another study showed that many GCs with hilar basal dendrites form after SE (Jessberger et al., 2007). It has also been shown that SE disrupts the normal interactions between radial glia and immature GCs, which could play a role in the formation of hilar basal dendrites (Shapiro et al., 2005). The elevation in neurotrophins after SE may be important, because they stimulate process outgrowth. For example, brain-derived neurotrophic factor (BDNF) levels rise in the mossy fibers after SE, and remain elevated for long periods of time (Scharfman, 2005). It is not clear if that neurotrophin pool would be accessible to the extracellular milieu in sufficient concentrations to be trophic for hilar basal dendrites, but even if not, there are many other

examples of growth factors and cytokines that increase and decrease after SE in the hilus (Scharfman, 2005). Together, they may create a stimulus for hilar basal dendrite formation.

Thus, it is difficult to propose an intervention to prevent hilar basal dendrites from developing, or a strategy to reverse the process after the fact. However, even if the dendrites cannot be prevented from forming, or reversed, manipulations to increase the relative numbers of GABAergic synapses on hilar basal dendrites may be useful. This idea is suggested by quantitative studies of hilar basal dendrites, which showed that the majority of synapses are excitatory (Thind et al., 2008).

3. Seizures vs. co-morbidity—The main targets for treatment of seizures in TLE that are discussed above include preventing EGC formation and hilar basal dendrites, or the increase in excitability they cause. These could be addressed in a preventative (antiepileptogenic) manner or long after the problems develop and seizures become chronic, i.e., in an anticonvulsant manner (Figure 6 and see parts B–C, below). However, it is also important to consider the potential that a long-term reduction in proliferation rate occurs in the chronic disease, and leads to cognitive impairment and co-morbidities such as depression. Therefore, a strategy that would be potentially useful to reduce co-morbidities late in the course of the disease would be to increase the rate of neurogenesis. Presumably a stimulation of the rate of neurogenesis would be most useful if seizures were under control by AEDs, because increasing neurogenesis during a period of uncontrolled seizures could have adverse effects by disrupting normal maturation and migration.

B. Prevention (antiepileptogenic approaches)

1. Irradiation—Low doses of irradiation can reduce proliferating cells selectively compared to postmitotic neurons, because of the sensitivity of the dividing cell. Interestingly, it has been noted clinically that low dose irradiation to a seizure focus can be advantageous (Jenrow et al., 2006), although this effect may not be due to an influence of irradiation on neurogenesis.

With respect to the use of irradiation to modify the potential influence of neurogenesis in epilepsy, one of the first studies used this approach to ask whether new neurons born in SE models contributed to mossy fiber sprouting. Mossy fiber sprouting is an example of synaptic reorganization that is thought to contribute to epileptogenesis by increasing recurrent excitatory circuitry in the dentate gyrus (Dudek and Sutula, 2007; Sutula and Dudek, 2007). It was hypothesized that new neurons might be likely to contribute to the novel excitatory circuitry, and low-dose irradiation might preferentially block progenitor division without additional effects on mature cells (Parent et al., 1999). The reduction in neurogenesis that occurred after low-dose irradiation did not appear to impair mossy fiber sprouting (Parent et al., 1999), and it was not noted if the seizures in the animals were affected. Subsequent studies have attempted to address the influence of irradiation on seizures, using the kindling model. Mixed results have been obtained. In one study that used whole brain irradiation before the first kindling stimulus, afterdischarge threshold was reduced during kindling, but there was no net effect on the kindled state (Raedt et al., 2007). In another study, irradiation was aimed at the amygdala because it was the focus, i.e., the site that was chosen for kindling. Different effects were demonstrated, depending on when irradiation was conducted relative to stages during kindling acquisition (Jenrow et al., 2006). These studies suggest that irradiation can influence seizures, but it is not clear how, and whether it is due to reduction of neurogenesis by irradiation. Because neurogenesis is not the only effect of irradiation (Fike et al., 2007), other methods have been developed, such as genetic manipulations.

2. Genetic manipulations—Mice that have genetic manipulations allow deletion of the progenitor pool and are potentially powerful because they are more specific (Beech et al.,

2004; Saxe et al., 2006). They also have the potential to be tailored temporally. These inducible, conditional genetic manipulations could be a foundation for gene therapy, although it would be unlikely that gene therapy would be available in the near future. Nevertheless, gene therapy has substantial potential, as discussed elsewhere (Noe et al., 2006; Vezzani, 2007).

One strategy that has been used effectively in rodents to reduce neurogenesis selectively is to insert thymidine kinase in the progenitors that express glial-fibrillary acid protein (GFAP), the radial glia-like progenitors (Figure 2). Upon systemic exposure to the anti viral agent ganciclovir, thymidine kinase phosphorylates ganciclovir only in GFAP-expressing cells. Phosphorylated ganciclovir becomes a toxic thymidine analogue, which will lead to the death of any of the radial glial cells that begin to divide. After several weeks of exposure, most of the radial glia naturally divide, and therefore, most will have died. This is a powerful approach, but there is a caveat in using the mice to address consequences of SE. The reason is that injury leads to a division of mature astrocytes (which express GFAP) into reactive microglia (Faulkner et al., 2004). Because mature astrocytes express GFAP, there would be a loss of reactive microglia after SE, which could change the normal progression from SE to chronic seizures.

Another strategy is to target nestin-expressing cells. Nestin is expressed in progenitors like radial glia, as well as other types of progenitors (Figure 2). One caveat with this approach is that nestin is present in non-progenitors, but a new strategy circumvents the problem by targeting the nestin promoter that is specific for progenitors (Yu et al., 2008). The newer method is also advantageous in other ways, because some stem cells appear to remain after treatment with ganciclovir. Although normally dormant, it appears to be possible to induce division in this dormant population long after ganciclovir treatment is over (Yu et al., 2008). Therefore, such a strategy could potentially reduce neurogenesis as a preventative measure, but then restore it much later, allowing one to prevent seizures but maintain a neurogenic niche long-term.

There are a number of approaches that could potentially be used in the future. Increasing the expression of proteins that normally induce apoptosis of the newly-generated neurons, such as BAX or decreasing those that impair apoptosis (Bcl-2) could be advantageous. Another strategy would be to reduce levels of neurotrophins such as BDNF that increase neurogenesis, foster survival, process outgrowth, and plasticity (Scharfman, 2005; Binder, 2007). Both strategies would require selective targeting to avoid the adverse consequences of excess apoptosis, or neurotrophin depletion, making the desired goals seem difficult to achieve at the present time.

3. Pharmacological approaches—Currently-available anticonvulsants may be useful in the antepileptogenic context suggested here, because they might reduce excitability, and in so doing, prevent some of the stimulus for epileptogenesis. In other words, if one assumes that epileptogenesis is stimulated by a large increase in neuronal activity (i.e., SE), and is perpetuated by new GCs with increased excitability, drugs that reduce excitatory activity could be effective in blocking epileptogenesis. Bumetamide might be a logical choice, if the new GCs that accumulate in the dentate gyrus after SE have increased excitability due to depolarized responses to GABA (Overstreet-Wadiche et al., 2006), which are due the $\text{Na}^+\text{K}^+/\text{2Cl}^-$ transporter that bumetamide inhibits (Kahle and Staley, 2008). However, blocking the excitatory effects of GABA in normal adult-generated GCs lead to defects in their development (Ge et al., 2007). Other anticonvulsants, such as valproic acid, may be useful because valproic acid appears to normalize neurogenesis (Jessberger et al., 2007), although other agents that have been suggested to normalize neurogenesis, such as endoneuramidase-N, has not shown efficacy in reducing seizures in an animal model of TLE (Pekcec et al., 2008). Even if these drugs would not stop seizures, both valproic acid and endoneuramidase-N may be useful in preventing detrimental effects of chronic seizures on the neurogenic niche, and as a result,

stabilizing cognition or mood in patients with TLE (Jessberger et al., 2007; Scharfman and Gray, 2007; Pekcec et al., 2008).

C. Reversal (anticonvulsant approaches)

1. Futuristic approaches—The most valuable strategy for the future is probably the hardest - to attempt reversal or recovery of TLE, long after disruption of neurogenesis has occurred and potentially contributed to it. In other words, for the patient who already has recurrent seizures, what anticonvulsant treatment, or potential cure, could be offered that would reverse or reduce the effects of “aberrant” neurogenesis? Pharmacotherapy would be one strategy, using drugs that decrease excitability, which is presumed to be initiated by circuits of abnormal GCs with other dentate gyrus neurons. If a compound that preferentially acted on mossy fiber transmission were available, such a drug might be particularly effective, because the normal GCs and hilar EGCs both use these specialized synapses to release glutamate onto their targets and contribute to abnormal network activity. Of the drugs known to modulate mossy fiber transmission, some metabotropic receptor ligands such as DCG-IV are commonly used *in vitro*, but one would need to carefully tailor this approach so that the functions of the mossy fibers that are important to cognitive function would not be impaired, and the drug would not reach other sites in the brain where the receptors also are important to normal function.

Two new approaches are attractive, although in early stages of development at the present time. One of these new approaches is to selectively silence neurons of interest, so they do not influence their targets (Slimko et al., 2002; Lerchner et al., 2007). The manner in which silencing is achieved involves expression of an ion channel which will hyperpolarize the cell, such as a chloride or potassium channel (Slimko et al., 2002; Lerchner et al., 2007). In the present context, one might silence the hilar EGCs, because their burst discharges lead to greater activity in the dentate gyrus network than normal, facilitating seizure-like activity. Another possibility is to use this approach to silence CA3 pyramidal neurons, to reduce their ability to generate burst discharges that in turn activate hilar EGCs. It might be best to reduce the activity in a fraction of CA3 neurons rather than the entire population, however, because completely blocking the ability of CA3 pyramidal cells to discharge action potentials would presumably disturb normal hippocampal function.

Another approach is to use light-activated silencing (Zhang et al., 2007), where a miniaturized light source silences neurons that express channelrhodopsins, which are proteins that are sensitive to light. At the present time, this technique is not amenable for sites deep within the brain, but it has already been used *in vivo*, an important first step. Theoretically, if the light could be flashed intermittently to expose EGCs that express the channelrhodopsin protein, it might disrupt the periodic discharges of the EGCs and their associated circuitry. However, it is not clear that a specific protein is expressed in hilar EGCs that would allow one to target them selectively.

2. Surgical approaches—It already is clear that surgical removal of the hippocampus is effective in reducing seizures in TLE when the hippocampus is a defined and specific focal lesion (Engel et al., 1998; Schmidek and Roberts, 2005; Spencer and Huh, 2008). Although often not considered, it is possible that those patients which benefit the most are those with numerous hilar EGCs, and complex circuitry that develops between the dentate gyrus and residual CA3 neurons. If so, one surgical approach to consider is not only removal, but also transection or “disconnection.” In other words, one might transect the circuitry between the dentate gyrus and CA3 network to disrupt the output of the dentate gyrus, and reverberatory activity between CA3, hilar cells (such as EGCs) and GCs. This concept is analogous to subpial transection - instead of removal, transection could be better. However, transection would seem

very difficult, from the technical perspective, because of the difficulty in selectively transecting the junction between the dentate gyrus and CA3, which is not easily accessed.

3. Stem cells—Another potential approach for therapeutic use, and one that has already demonstrated preclinical success (Richardson et al., 2008), is transplantation. Recent studies have shown that transplanted neurons can restore neurogenesis (Kuruba et al., 2009), and GABAergic neurons can reduce seizures (Alvarez-Dolado et al., 2006). Moreover, one study has shown that a specific type of “stem cell,” transplanted into the dentate gyrus often matures into normal GCs, which could be used in a restorative manner if neurogenesis declines (Carpentino et al., 2008). Remarkably, these stem cells can lead to abnormal growths if a normal animal is used, but in an animal that has had seizures, the stem cells become GCs and tumors do not appear to develop (Carpentino et al., 2008). Infusion of neuropeptide Y may be a particularly effective strategy after such stem cell infusion, because it stimulates precursor division (Howell et al., 2003; Scharfman and Gray, 2006; Scharfman and Gray, 2007) and reduces seizures (Noe et al., 2006).

As intriguing as these possibilities are, there are many issues that need to be addressed before therapeutic use can truly be considered. For example, if some transplanted neurons become GCs, and they do not migrate correctly, would they increase the hilar EGC population and worsen the condition? If GABAergic neurons were used for a transplant procedure, and they were to project to other GABAergic neurons, seizures might increase due to disinhibition of principal cells. The remarkable plasticity of the dentate gyrus could be advantageous in allowing transplanted neurons to survive and develop synaptic connections, but be detrimental if inappropriate maturation and integration occur. There are many other issues to resolve, such as administration route, side effects, the type of TLE that should be examined, and whether age and comorbidity would complicate outcome. Still, there are a number of promising possibilities to consider that could improve a devastating condition, i.e., TLE.

Acknowledgments

Supported by NS 41490 and the Epilepsy Foundation.

References

- Alvarez-Dolado M, Calcagnotto M, Karkar K, Southwell D, Jones-Davis D, Estrada R, Rubenstein J, Alvarez-Buylla A, Baraban S. Cortical inhibition modified by embryonic neural precursors grafted into the postnatal brain. *J Neurosci* 2006;26:7380–7389. [PubMed: 16837585]
- Amaral DG, Scharfman HE, Lavenex P. The dentate gyrus: Fundamental neuroanatomical organization (dentate gyrus for dummies). *Prog Brain Res* 2007;163:3–22. [PubMed: 17765709]
- Bagri A, Gurney T, He X, Zou YR, Littman DR, Tessier-Lavigne M, Pleasure SJ. The chemokine sdf1 regulates migration of dentate granule cells. *Development* 2002;129:4249–4260. [PubMed: 12183377]
- Beech RD, Cleary MA, Treloar HB, Eisch AJ, Harriot AV, Zhong W, Greer CA, Duman RS, Picciotto MR. Nestin promoter/enhancer directs transgene expression to precursors of adult generated periglomerular neurons. *J Comp Neurol* 2004;475:128–141. [PubMed: 15176089]
- Bender RA, Dube C, Gonzalez-Vega R, Mina EW, Baram TZ. Mossy fiber plasticity and enhanced hippocampal excitability, without hippocampal cell loss or altered neurogenesis, in an animal model of prolonged febrile seizures. *Hippocampus* 2003;13:399–412. [PubMed: 12722980]
- Bengzon J, Kokaia Z, Elmer E, Nanobashvili A, Kokaia M, Lindvall O. Apoptosis and proliferation of dentate gyrus neurons after single and intermittent limbic seizures. *Proc Natl Acad Sci U S A* 1997;94:10432–10437. [PubMed: 9294228]
- Binder DK. Neurotrophins in the dentate gyrus. *Prog Brain Res* 2007;163:371–397. [PubMed: 17765730]
- Buckmaster PS. In vivo intracellular analysis of granule cell axon reorganization in epileptic rats. *J Neurophysiol* 1999;81:712–721. [PubMed: 10036272]

- Carpentino J, Hartman N, Grabel L, Naegele J. Region-specific differentiation of embryonic stem cell-derived neural progenitor transplants into the adult mouse hippocampus following seizures. *J Neurosci Res* 2008;86:512–524. [PubMed: 17918739]
- Carter DS, Harrison AJ, Falenski KW, Blair RE, DeLorenzo RJ. Long-term decrease in calbindin-d28k expression in the hippocampus of epileptic rats following pilocarpine-induced status epilepticus. *Epi Res* 2008;79:213–223.
- Cuppini R, Bucherelli C, Ambrogini P, Ciuffoli S, Orsini L, Ferri PEB. Age-related naturally occurring depression of hippocampal neurogenesis does not affect trace fear conditioning. 2006;16:141–148.
- Danzer S. Postnatal and adult neurogenesis in the development of human disease. *Neuroscientist* 2008;14:446–458. [PubMed: 18997123]
- Dashtipour K, Wong AM, Obenaus A, Spigelman I, Ribak CE. Temporal profile of hilar basal dendrite formation on dentate granule cells after status epilepticus. *Epilepsy Res* 2003;54:141–151. [PubMed: 12837565]
- Dudek FE, Sutula TP. Epileptogenesis in the dentate gyrus: a critical perspective. *Prog Brain Res* 2007;163:541–563. [PubMed: 17765737]
- Engel, J.; Cascino, G.; Shields, W. Surgically remediable syndromes. In: Engel, J.; Pedley, TA., editors. *Epilepsy: A comprehensive guide*. NY: Lippincott-Raven; 1998. p. 1687–1696.
- Faulkner JR, Herrmann JE, Woo MJ, Tansey KE, Doan NB, Sofroniew MV. Reactive astrocytes protect tissue and preserve function after spinal cord injury. *J Neurosci* 2004;24:2143 – 2155. [PubMed: 14999065]
- Faulkner RL, Jang MH, Liu XB, Duan X, Sailor KA, Kim JY, Ge S, Jones EG, Ming GL, Song H, Cheng HJ. Development of hippocampal mossy fiber synaptic outputs by new neurons in the adult brain. *Proc Natl Acad Sci U S A* 2008;105:14157–14162. [PubMed: 18780780]
- Fike JR, Rola R, Limoli CL. Radiation response of neural precursor cells. *Neurosurg Clin N Am* 2007;18:115–127. [PubMed: 17244559]
- Frotscher M. Dual role of cajal-retzius cells and reelin in cortical development. *Cell Tissue Res* 1997;290:315–322. [PubMed: 9321693]
- Gage, FH.; Kempermann, G.; Song, H. *Adult neurogenesis*. Cold Spring Harbor; Cold Spring Harbor Laboratory Press: 2008.
- Gale K. Subcortical structures and pathways involved in convulsive seizure generation. *J Clin Neurophysiol* 1992;9:264–277. [PubMed: 1350593]
- Ge S, Goh EL, Sailor KA, Kitabatake Y, Ming GL, Song H. GABA regulates synaptic integration of newly generated neurons in the adult brain. *Nature* 2006;439:589–593. [PubMed: 16341203]
- Ge S, Ming GL, Song H. GABA sets the tempo for activity-dependent adult neurogenesis. *Trends Neurosci* 2007;30:1–8. [PubMed: 17116335]
- Ge S, Sailor KA, Ming GL, Song H. Synaptic integration and plasticity of new neurons in the adult hippocampus. *J Physiol* 2008;586:3759–3765. [PubMed: 18499723]
- Gong W, Wang TW, Huang HS, Parent JM. Reelin regulates neuronal progenitor migration in intact and epileptic hippocampus. *J Neurosci* 2007;27:1803–1811. [PubMed: 17314278]
- Goodman JH, Wasterlain CG, Massarweh WF, Dean E, Sollas AL, Sloviter RS. Calbindin-D28k immunoreactivity and selective vulnerability to ischemia in the dentate gyrus of the developing rat. *Brain Res* 1993;606:309–314. [PubMed: 8490723]
- Gray WP, Sundstrom L. Kainic acid increases the proliferation of granule cell progenitors in the dentate gyrus of the adult rat. *Brain Res* 1998;790:52–59. [PubMed: 9593820]
- Hattiangady B, Rao MS, Shetty AK. Chronic temporal lobe epilepsy is associated with severely declined dentate neurogenesis in the adult hippocampus. *Neurobiol Dis* 2004;17:473–490. [PubMed: 15571983]
- Howell OW, Scharfman HE, Herzog H, Sundstrom LE, Beck-Sickinger A, Gray WP. Neuropeptide y is neuroproliferative for post-natal hippocampal precursor cells. *J Neurochem* 2003;86:646–659. [PubMed: 12859678]
- Jakubs K, Nanobashvili A, Bonde S, Ekdahl CT, Kokaia Z, Kokaia M, Lindvall O. Environment matters: synaptic properties of neurons born in the epileptic adult brain develop to reduce excitability. *Neuron* 2006;52:1047–1059. [PubMed: 17178407]

- Jenrow KA, Ratkewicz AE, Zalinski DN, Roszka KM, Lemke NW, Elisevich KV. Influence of ionizing radiation on the course of kindled epileptogenesis. *Brain Res* 2006;1094:207–216. [PubMed: 16762327]
- Jessberger S, Nakashima K, Clemenson GD, Mejia E, Mathews E, Ure K, Ogawa S, Sinton CM, Gage FH, Hsieh J. Epigenetic modulation of seizure-induced neurogenesis and cognitive decline. *J Neurosci* 2007;27:5967–5975. [PubMed: 17537967]
- Jessberger S, Romer B, Babu H, Kempermann G. Seizures induce proliferation and dispersion of doublecortin-positive hippocampal progenitor cells. *Exp Neurol* 2005;196:342–351. [PubMed: 16168988]
- Jessberger S, Zhao C, Toni N, Clemenson GD, Li Y, Gage FH. Seizure-associated, aberrant neurogenesis in adult rats characterized with retrovirus-mediated cell labeling. *J Neurosci* 2007;27:9400–9407. [PubMed: 17728453]
- Jung KH, Chu K, Kim M, Jeong SW, Song YM, Lee ST, Kim JY, Lee SK, Roh JK. Continuous cytosine-b-d-arabinofuranoside infusion reduces ectopic granule cells in adult hippocampus with attenuation of spontaneous recurrent seizures following pilocarpine-induced epileptics. *Eur J Neurosci* 2004;19:3219–3226. [PubMed: 15217378]
- Jung KH, Chu K, Lee ST, Kim J, Sinn DI, Kim JM, Park DK, Lee JJ, Kim SU, Kim M, Lee SK, Roh JK. Cyclooxygenase-2 inhibitor, celecoxib, inhibits the altered hippocampal neurogenesis with attenuation of spontaneous recurrent seizures following pilocarpine-induced status epilepticus. *Neurobiol Dis* 2006;23:237–246. [PubMed: 16806953]
- Jung KH, Chu K, Lee ST, Kim JH, Kang KM, Song EC, Kim SJ, Park HK, Kim M, Lee SK, Roh JK. Region-specific plasticity in the epileptic rat brain: a hippocampal and extrahippocampal analysis. *Epilepsia*. 2008[Epub ahead of print]
- Kahle KT, Staley KJ. The bumetanide-sensitive Na-K-2Cl cotransporter NKCC1 as a potential target of a novel mechanism-based treatment strategy for neonatal seizures. *Neurosurg Focus* 2008;25:E22. [PubMed: 18759624]
- Kempermann G. Adult neurogenesis. Oxford: Oxford University Press; 2006.
- Kempermann G, Gast D, Kronenberg G, Yamaguchi M, Gage FH. Early determination and long-term persistence of adult-generated new neurons in the hippocampus of mice. *Development* 2003;130:391–399. [PubMed: 12466205]
- Kempermann G, Jessberger S, Steiner B, Kronenberg G. Milestones of neuronal development in the adult hippocampus. *Trends Neurosci* 2004;27:447–452. [PubMed: 15271491]
- Kuruba R, Hattiangady B, Shetty AK. Hippocampal neurogenesis and neural stem cells in temporal lobe epilepsy. *Epilepsy Behav* 2009;14(S1):65–73. [PubMed: 18796338]
- Kuruba R, Shetty AK. Could hippocampal neurogenesis be a future drug target for treating temporal lobe epilepsy? *CNS Neurol Disord Drug Targets* 2007;6:342–357. [PubMed: 18045163]
- Laplagne DA, Espósito MS, Piatti VC, Morgenstern NA, Zhao C, van Praag H, Gage FH, Schinder AF. Functional convergence of neurons generated in the developing and adult hippocampus. *PLoS Biol* 2006;4:e409. [PubMed: 17121455]
- Lerchner W, Xiao C, Nashmi R, Slimko EM, van Trigt L, Lester HA, Anderson DJ. Reversible silencing of neuronal excitability in behaving mice by a genetically targeted, ivermectin-gated Cl^- channel. *Neuron* 2007;54:35–49. [PubMed: 17408576]
- Leuner B, Mendolia-Loffredo S, Kozorovitskiy Y, Samburg D, Gould E, Shors TJ. Learning enhances the survival of new neurons beyond the time when the hippocampus is required for memory. *J Neurosci* 2004;24:7477–7481. [PubMed: 15329394]
- Liu H, Kaur J, Dashtipour K, Kinyamu R, Ribak CE, Friedman LK. Suppression of hippocampal neurogenesis is associated with developmental stage, number of perinatal seizure episodes, and glucocorticosteroid level. *Exp Neurol* 2003;184:196–213. [PubMed: 14637092]
- Loscher W, Ebert U. The role of the piriform cortex in kindling. *Prog Neurobiol* 1996;50:427–481. [PubMed: 9015822]
- Markwardt S, Overstreet-Wadiche L. GABAergic signalling to adult-generated neurons. *J Physiol* 2008;586:3745–3749. [PubMed: 18511482]

- McCloskey DP, Hintz TM, Pierce JP, Scharfman HE. Stereological methods reveal the robust size and stability of ectopic hilar granule cells after pilocarpine-induced status epilepticus in the adult rat. *Eur J Neurosci* 2006;24:2203–2210. [PubMed: 17042797]
- McIntyre DC, Gilby KL. Mapping seizure pathways in the temporal lobe. *Epilepsia* 2008;49(S3):23–30. [PubMed: 18304253]
- Morgan RJ, Soltesz I. Nonrandom connectivity of the epileptic dentate gyrus predicts a major role for neuronal hubs in seizures. *Proc Natl Acad Sci USA* 2008;105:6179–6184. [PubMed: 18375756]
- Nagerl UV, Mody I, Jeub M, Lie AA, Elger CE, Beck H. Surviving granule cells of the sclerotic human hippocampus have reduced Ca^{2+} influx because of a loss of calbindin-D(28k) in temporal lobe epilepsy. *J Neurosci* 2000;20:1831–1836. [PubMed: 10684884]
- Noe F, Nissinen J, Pitkänen A, Gobbi M, Sperk G, During M, Vezzani A. Gene therapy in epilepsy: The focus on NPY. *Peptides* 2006;28:377–383. [PubMed: 17196301]
- Overstreet-Wadiche LS, Bromberg DA, Bensen AL, Westbrook GL. Seizures accelerate functional integration of adult-generated granule cells. *J Neurosci* 2006;26:4095–4103. [PubMed: 16611826]
- Parent JM. When newborn neurons stray. *Epilepsy Curr* 2005;5:231–233. [PubMed: 16372058]
- Parent JM, Elliott RC, Pleasure SJ, Barbaro NM, Lowenstein DH. Aberrant seizure-induced neurogenesis in experimental temporal lobe epilepsy. *Ann Neurol* 2006;59:81–91. [PubMed: 16261566]
- Parent JM, Janumpalli S, McNamara JO, Lowenstein DH. Increased dentate granule cell neurogenesis following amygdala kindling in the adult rat. *Neurosci Lett* 1998;247:9–12. [PubMed: 9637397]
- Parent JM, Lowenstein DH. Seizure-induced neurogenesis: Are more new neurons good for an adult brain? *Prog Brain Res* 2002;135:121–131. [PubMed: 12143334]
- Parent JM, Murphy GG. Mechanisms and functional significance of aberrant seizure-induced hippocampal neurogenesis. *Epilepsia* 2008;49:S19–S25.
- Parent JM, Tada E, Fike JR, Lowenstein DH. Inhibition of dentate granule cell neurogenesis with brain irradiation does not prevent seizure-induced mossy fiber synaptic reorganization in the rat. *J Neurosci* 1999;19:4508–4519. [PubMed: 10341251]
- Parent JM, Tada E, Fike JR, Lowenstein DH. Inhibition of dentate granule cell neurogenesis with brain irradiation does not prevent seizure-induced mossy fiber synaptic reorganization in the rat. *J Neurosci* 1999;18:4508–4519. [PubMed: 10341251]
- Parent JM, Yu TW, Leibowitz RT, Geschwind DH, Sloviter RS, Lowenstein DH. Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J Neurosci* 1997;17:3727–3738. [PubMed: 9133393]
- Pekcec A, Fuest C, Mühlenhoff M, Gerardy-Schahn R, Potschka H. Targeting epileptogenesis-associated induction of neurogenesis by enzymatic depolysialylation of ncam counteracts spatial learning dysfunction but fails to impact epilepsy development. *J Neurochem* 2008;105:389–400. [PubMed: 18194217]
- Pitkanen A, Sutula TP. Is epilepsy a progressive disorder? Prospects for new therapeutic approaches in temporal-lobe epilepsy. *Lancet Neurol* 2002;1:173–181. [PubMed: 12849486]
- Porter BE. Neurogenesis and epilepsy in the developing brain. *Epilepsia* 2008;49(S5):550–554.
- Raedt R, Boon P, Persson A, Alborn AM, Boterberg T, Van Dycke A, Linder B, De Smedt T, Wadman WJ, Ben-Menachem E, Eriksson PS. Radiation of the rat brain suppresses seizure-induced neurogenesis and transiently enhances excitability during kindling acquisition. *Epilepsia* 2007;48:1952–1963. [PubMed: 17555527]
- Ribak CE, Dashtipour K. Neuroplasticity in the damaged dentate gyrus of the epileptic brain. *Prog Brain Res* 2002;136:319–328. [PubMed: 12143392]
- Ribak CE, Shapiro LA. Ultrastructure and synaptic connectivity of cell types in the adult rat dentate gyrus. *Prog Brain Res* 2007;163:155–166. [PubMed: 17765717]
- Ribak CE, Tran PH, Spigelman I, Okazaki MM, Nadler JV. Status epilepticus-induced hilar basal dendrites on rodent granule cells contribute to recurrent excitatory circuitry. *J Comp Neurol* 2000;428:240–253. [PubMed: 11064364]
- Richardson RM, Barbaro NM, Alvarez-Bulley A, Baraban SC. Developing cell transplantation for temporal lobe epilepsy. *Neurosurg Focus* 2008;24:E17. [PubMed: 18341393]

- Rivers LE. Pdgfra/ng2 glia generate myelinating oligodendrocytes and piriform projection neurons in adult mice. *Nature Neurosci* 2008;11:1392–1401. [PubMed: 18849983]
- Saxe MD, Battaglia F, Wang J, Malleret G, David DJ, Monckton JE, Garcia AD, Sofroniew M, Kandel ER, Santarell L, Hen R, Drew MR. Ablation of hippocampal neurogenesis impairs contextual fear conditioning and synaptic plasticity in the dentate gyrus. *Proc Natl Acad Sci U S A* 2006;103:17501–17506. [PubMed: 17088541]
- Scharfman, HE. The parahippocampal region in temporal lobe epilepsy. In: Witter, MP.; Wouterlood, FG., editors. *The parahippocampal region: organization and role in cognitive functions*. Oxford University Press; 2002. p. 319–338.
- Scharfman HE. Functional implications of seizure-induced neurogenesis. *Adv Exp Med Biol* 2004;548:192–212. [PubMed: 15250595]
- Scharfman HE. Brain-derived neurotrophic factor and epilepsy—a missing link? *Epilepsy Curr* 2005;5:83–88. [PubMed: 16145610]
- Scharfman, HE. Seizure-induced neurogenesis and its dependence on growth factors and cytokines. In: Binder, DK.; Scharfman, HE., editors. *Growth factors and epilepsy*. New York: Nova Sciences; 2005.
- Scharfman HE, Goodman JH, McCloskey DP. Ectopic granule cells of the rat dentate gyrus. *Dev Neurosci* 2007;29:14–27. [PubMed: 17148946]
- Scharfman HE, Goodman JH, Sollas AL. Granule-like neurons at the hilar/ca3 border after status epilepticus and their synchrony with area ca3 pyramidal cells: Functional implications of seizure-induced neurogenesis. *J Neurosci* 2000;61:6144–6158. [PubMed: 10934264]
- Scharfman HE, Gray WP. Plasticity of neuropeptide y in the dentate gyrus after seizures, and its relevance to seizure-induced neurogenesis. *EXS* 2006;95:193–211. [PubMed: 16383008]
- Scharfman HE, Gray WP. Relevance of seizure-induced neurogenesis to the etiology of temporal lobe epilepsy. *Epilepsia* 2007;48(S2):33–41. [PubMed: 17571351]
- Scharfman HE, Sollas AL, Goodman JH. Spontaneous recurrent seizures after pilocarpine-induced status epilepticus activate calbindin-immunoreactive hilar cells of the rat dentate gyrus. *Neuroscience* 2002;111:71–81. [PubMed: 11955713]
- Schmidek, H.; Roberts, D. *Operative neurosurgical techniques*. New York: Elsevier; 2005.
- Schmidt-Hieber C, Jonas P, Bischofberger J. Enhanced synaptic plasticity in newly generated granule cells of the adult hippocampus. *Nature* 2004;429:184–187. [PubMed: 15107864]
- Scott BW, Wang S, Burnham WM, DeBoni U, Wojtowicz JM. Kindling-induced neurogenesis in the dentate gyrus of the rat. *Neurosci Lett* 1998;248:73–76. [PubMed: 9654345]
- Scott BW, Wojtowicz JM, Burnham WM. Neurogenesis in the dentate gyrus of the rat following electroconvulsive shock seizures. *Exp Neurol* 2000;165:231–236. [PubMed: 10993683]
- Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Bulley A. Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 2004;21:7153–7160. [PubMed: 11549726]
- Shapiro LA, Figueroa-Aragon S, Ribak CE. Newly generated granule cells show rapid neuroplastic changes in the adult rat dentate gyrus during the first five days following pilocarpine-induced seizures. *Eur J Neurosci* 2007;26:583–592. [PubMed: 17686039]
- Shapiro LA, Korn MJ, Ribak CE. Newly generated dentate granule cells from epileptic rats exhibit elongated hilar basal dendrites that align along gfap-immunolabeled processes. *Neuroscience* 2005;136:823–831. [PubMed: 16344154]
- Shapiro LA, Ng K, Zhou QY, Ribak CE. Subventricular zone-derived, newly generated neurons populate several olfactory and limbic forebrain regions. *Epilepsy Behav* 2009;14(S1):74–80. [PubMed: 18849007]
- Shapiro LA, Ribak CE. Integration of newly born dentate granule cells into adult brains: Hypotheses based on normal and epileptic rodents. *Brain Res Rev* 2005;48:43–56. [PubMed: 15708627]
- Shapiro LA, Ribak CE, Jessberger S. Structural changes for adult-born dentate granule cells after status epilepticus. *Epilepsia* 2008;49:S13–18.
- Slimko EM, McKinney S, Anderson DJ, Davidson N, Lester HA. Selective electrical silencing of mammalian neurons in vitro by the use of invertebrate ligand-gated chloride channels. *J Neurosci* 2002;22:7373–7379. [PubMed: 12196558]

- Spencer S, Huh L. Outcomes of epilepsy surgery in adults and children. *Lancet Neurol* 2008;7:525–537. [PubMed: 18485316]
- Spigelman I, Yan XX, Obenaus A, Lee EY, Wasterlain CG, Ribak CE. Dentate granule cells form novel basal dendrites in a rat model of temporal lobe epilepsy. *Neuroscience* 1998;86:109–120. [PubMed: 9692747]
- Sutula TP. Mechanisms of epilepsy progression: Current theories and perspectives from neuroplasticity in adulthood and development. *Epilepsy Res* 2004;60:161–171. [PubMed: 15380560]
- Sutula TP, Dudek FE. Unmasking recurrent excitation generated by mossy fiber sprouting in the epileptic dentate gyrus: An emergent property of a complex system. *Prog Brain Res* 2007;163:541–563. [PubMed: 17765737]
- Sutula TP, Hermann B. Progression in mesial temporal lobe epilepsy. *Ann Neurol* 1999;45:553–556. [PubMed: 10319875]
- Thind KK, Ribak CE, Buckmaster PS. Synaptic input to dentate granule cell basal dendrites in a rat model of temporal lobe epilepsy. *J Comp Neurol* 2008;509:190–202. [PubMed: 18461605]
- Toni N, Laplagne DA, Zhao C, Lombardi G, Ribak CE, Gage FH, Schinder AF. Neurons born in the adult dentate gyrus form functional synapses with target cells. *Nature Neurosci* 2008;11:901–907. [PubMed: 18622400]
- van Praag H, Schinder AF, Christie BR, Toni N, Palmer TD, Gage FH. Functional neurogenesis in the adult hippocampus. *Nature* 2002;415:1030–1034. [PubMed: 11875571]
- Vezzani A. The promise of gene therapy for the treatment of epilepsy. *Expert Rev Neurother* 2007;7:685–1692.
- Walter C, Murphy BL, Pun RY, Spieles-Engemann AL, Danzer SC. Pilocarpine-induced seizures cause selective time-dependent changes to adult-generated hippocampal dentate granule cells. *J Neurosci* 2007;27:7451–7452.
- Wojtowicz JM. Potential consequences of altered neurogenesis on learning and memory in the epileptic brain. *Epilepsia* 2008;49(S5):42–49. [PubMed: 18522599]
- Yu TS, Zhang GF, Liebl DJ, Kernie SG. Traumatic brain injury-induced hippocampal neurogenesis requires activation of early nestin-expressing progenitors. *J Neurosci* 2008;28:12901–12912. [PubMed: 19036984]
- Zhang F, Wang L, Brauner M, Liewald J, Kay K, Watzke N, Wood P, Bamberg E, Nagel G, Gottschalk A, Deisseroth K. Multimodal fast optical interrogation of neural circuitry. *Nature* 2007;446:633–639. [PubMed: 17410168]
- Zhao CS, Deng W, Gage FH. Mechanisms and functional implications of adult neurogenesis. *Cell* 2008;132:645–660. [PubMed: 18295581]

Figure 1. Postnatal neurogenesis in the hippocampus

A. A schematic of the hippocampus of the rodent in cross section showing the subfields (CA1, CA3, and dentate gyrus), and the major glutamatergic pathways (the trisynaptic circuit; blue arrows).

B. A schematic of the dentate gyrus illustrating major subtypes and the major output of GCs, the mossy fiber pathway. INT = GABAergic neuron; MC = mossy cell. The dotted line demarcates the border between the hilus and the CA3 subfield.

C. A schematic showing the location of the subgranular zone (SGZ; orange), where most progenitors are thought to reside in the adult dentate gyrus. MOL= molecular layer, GCL = granule cell layer; I= inner molecular layer; M= middle molecular layer; O = outer molecular layer.

Figure 2. Sequential maturation of progenitors in the normal adult dentate gyrus

TOP: Stages of progenitor maturation with patterns of expression of selective markers indicated in color.

BOTTOM: Schematic illustration of progenitor division in the subgranular zone (SGZ) or the dentate gyrus. In the SGZ, type 1 (radial glia-like progenitors, analogous to the Type B cells of the SVZ; (Seri et al., 2004)) become Type 2a, 2b, 3 cells. Most of the neurons become GCs, distinguished by calretinin immunoreactivity when they are immature, and calbindin immunoreactivity after maturity. GFAP, glial fibrillary acid protein. From (Kempermann et al., 2004; Kempermann, 2006).

Figure 3. Timeline of changes in postnatal neurogenesis in a SE model of TLE illustrates the changes that would be potential therapeutic targets

After SE (time 0), animals initially exhibit many changes in the dentate gyrus, including a dramatic increase in the rate of neurogenesis that lasts the first weeks after SE (Parent et al., 1997; Scharfman et al., 2000), and some neurons that migrate incorrectly appear to be epileptogenic (Scharfman et al., 2000; Scharfman, 2004; Scharfman et al., 2007). Others, that migrate normally, appear to have no such effect (Jakubs et al., 2006). Long term, after chronic seizures have developed, there is a reduction in neurogenesis (Hattiangady et al., 2004; Kuruba et al., 2009). The timing that these phases exhibit is only estimated in the schematic.

Figure 4. Hilar EGCs

A. Calbindin (D28K)-immunoreactivity in a control rat stains the cell bodies of GCs, GC dendrites in the molecular layer (MOL), as well as GC axons (mossy fibers) in the hilus (h). Calibration = 100 μ m.

B. Calbindin immunoreactivity in an epileptic rat illustrates a population of hilar EGCs near CA3 (arrows). Note that calbindin immunoreactivity is weak in the GCL in this section, because young GCs do not express calbindin (Goodman et al., 1993), and many mature GCs downregulate calbindin after seizures (Nagerl et al., 2000; Scharfman et al., 2002; Carter et al., 2008). **(B).** Prox1 immunoreactivity in a different pair of rats demonstrates numerous hilar EGCs in the epileptic rat (arrows).

Figure 5. Synchronization of CA3 pyramidal cells with hilar EGCs

Simultaneous intracellular recordings at a slow (left) and fast (right) time base show the synchronization of a hilar EGC with spontaneous bursts in a CA3 pyramidal cell. The arrow points to the capacitative artifact of the first CA3 pyramidal cell action potential in the EGC recording, illustrating that the onset of the burst in the pyramidal cell preceded the onset of the EGC burst. From (Scharfman et al., 2007).

Table 1

Therapeutic strategies to target abnormal postnatal neurogenesis in epilepsy

ANTIEPILEPTOGENIC Prevent hilar EGCs and hilar basal dendrites	ANTICONVULSANT Suppress dentate excitability
Control rate of neurogenesis	Anticonvulsants
Decrease migratory “go” signals	Targeted inactivation
Increase migratory “stop” signals	Surgical transection
Reduce survival	Stem cell therapy
Reduce process outgrowth	
Increase inhibitory innervation	

Legend: Antiepileptogenic (**Left**) and anticonvulsant (**Right**) strategies to address characteristics of postnatal neurogenesis that appear to contribute to the predisposition for seizures in adult SE models of TLE. For detailed description, see text.