Published in final edited form as:

J Org Chem. 2009 June 19; 74(12): 4565–4572. doi:10.1021/jo900180p.

Facile Pd(II)- and Ni(II)-Catalyzed Isomerization of Terminal Alkenes into 2-Alkenes

Hwan Jung Lim, Craig R. Smith, and T. V. RajanBabu *

Department of Chemistry, The Ohio State University, 100 W. 18th Avenue, Columbus, Ohio 43210

Abstract

Mono- and 2,2'-di-substituted terminal alkenes can be isomerized into the more stable internal *Z*- and *E*- alkenes by treating them with catalytic amounts of [(allyl)PdCl]₂ or [(allyl)NiBr]₂, a triarylphosphine and silver triflate at room temperature. The isomeric ratio (*E*:*Z*) depends on the alkenes, the (*E*)-isomer being the major one. The reaction is tolerant to a wide variety of functional groups including other reactive olefins. Unlike the more reactive Ir catalysts, monosubstituted alkenes give almost exclusively the 2-alkenes. Direct comparison to two of the best-known catalysts for this process, (Ir(PCy₃)₃)⁺ [BPh₄][−], and Grubbs Generation II metathesis catalyst) is also reported.

Introduction

Isomerization of allylic derivatives is a well known reaction¹ that has been used in unmasking allylic protecting groups,² and enantioselective synthesis of vinyl amines^{3a} and ethers.^{3b} Notable applications of the vinyl ether synthesis include the use of these intermediates for other C-C bond-forming reactions such as Claisen rearrangements.^{4a,b} The isomerization reaction has also been used for the stereoselective syntheses of silyl ethers of aldehyde enols.^{4c,d} Depending on the metal and the reaction conditions, varying selectivities for the isomerized *E*- and *Z*- alkenes have been observed. In general, while Ni(II)-catalysts deliver the best *Z*:*E* ratios,⁵ Ir-catalysts are the optimal reagents for the preparation of *E*-olefins (Scheme 1).^{4b–d} Low-yielding isomerizations of terminal alkenes catalyzed by Pt,^{7a} Rh^{7a} Pd^{7b} and Ni^{7c} have been known since 1964. The role of a bulky ligand and the intermediacy of the metal hydride implied in the [2-Me-C₆H₄-O)₃P]₃Ni/HCl^{7c} have been effectively used to design other protocols for this reaction.^{7d,e} Iridium (I) catalysts originally developed by Felkin^{6a} have been modified for the skeletal isomerization of functionalized terminal alkenes at room temperature.^{6a–e} The most recent entry into this group of catalysts capable of effecting the skeletal isomerization of *C*-allylic appendages in highly functionalized substrates is the Grubbs generation II metathesis catalyst that has been 'thermally modified' (eq 1).⁸

E-mail: rajanbabu.1@osu.edu.

SUPPORTING INFORMATION. Spectroscopic data (¹H and ¹³C NMR), gas chromatographic traces of starting materials, products and of crude isomerization products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(1)

During our studies on asymmetric hydrovinylation of 2,2'-disubstituted alkenes we noticed that some substrates consistently gave varying proportions of a minor component readily identified as the allylic rearrangement product as illustrated in eqs 2a and 2b.⁹ A similar rearrangement was also detected during cycloisomerization of dienes catalyzed by $\{[(\text{allyl})\text{PdCl}]_2/\text{AgOTf}/[(2\text{-Me-C}_6\text{H}_4)_3\text{P}]\}$ (eq 3).¹⁰

(2a)

(2b)

(3)

Since cationic metal hydrides have been implicated in these reactions, we wondered whether the hydrovinylation conditions could be modified to improve the yield of *selective isomerization* of a monosubstituted terminal alkene into a 2-alkene and of a 2,2'-disubstituted alkene into a trisubstituted alkene. We are especially interested in avoiding the complication from subsequent isomerization of the primary product, which often accompanies some of the isomerization reactions of terminal alkenes involving Ir^{6b,e} and Ru.^{8h} This isomerization is especially problematic when the homoallylic position of the carbon chain in the starting alkene is unsubstituted (eq 4). There are only limited reports of rearrangements of 2,2'-disubstituted alkenes to the corresponding internal trisubstituted isomer.^{8h} When coupled with a Wittig, Grignard or Keck radical allylation^{8f} reaction, to produce the ethylidene derivative, this transformation could be quite useful. In acyclic systems, the configuration of the newly formed carbon-carbon double bond would also be of some interest.¹¹ The results of these studies are reported here.

(4)

(5)

(6)

Results

Our initial studies concentrated on the little known^{8b} isomerization of the 2,2'-disubstituted alkenes. Results of isomerization of two prototypical substrates 1-methylenetetralin (**1**, eq 5) and the 1-(2-phthalimidoethyl)styrene (**3a**, eq 6) are shown in Table 1. When **1** was reacted under typical hydrovinylation conditions¹² using Ph_3P as a ligand, $\{[(\text{allyl})\text{NiBr}]_2, \text{Ph}_3\text{P}, \text{AgOTf}, 25^\circ\text{C}, 2\text{ d}\}$ in an atmosphere of ethylene, no reaction ensued (entry 1, column 4). In accordance with our previous observation that Pd(II) salts show enhanced reactivity under these conditions,^{12b} the use of $[(\text{allyl})\text{PdCl}]_2$ instead of the corresponding Ni salt leads to the expected product **2** in very good yield (entry 2, column 4).¹³ Typically the reaction involving ethylene is carried out as follows: appropriate amounts of the metal salt, the phosphine and AgOTf are mixed in CH_2Cl_2 in a drybox in a Schlenk tube, the mixture is taken out of the box and placed in an atmosphere of ethylene for 20 min. The alkene is introduced and the ethylene atmosphere is replaced by nitrogen, and the mixture is stirred for 2 days at rt. Unlike the hydrovinylation reaction, there is no need to remove the precipitated salts (AgX) for the isomerization reactions. While the use of ethylene has a slight advantage in the recovery of products, we find that an operationally simpler procedure is to use catalytic amounts of diallyl ether (entry 3). Presumably diallyl ether, which undergoes facile cycloisomerization to 3-methylene-4-methyltetrahydrofuran, readily produces^{10a,b} the requisite cationic metal hydride with minimum of contamination by other side products. Use of diallyl ether also helps the Ni-catalyzed reaction if a highly dissociated counter-anion, $[(3,5-(\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}^-]$, is used instead of OTf (entry 4, column 4).^{14,12b} Thus with 1 mol % of $[(\text{allyl})_2\text{NiBr}]_2$, and 2 mol % each of Ph_3P , $\text{Na}^+ [(3,5-(\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}^-]$ (BARF) and diallyl ether **1** gives very good yield of the isomerization product **2** (entry 4).

Isomerization of functionalized alkene **3a** (eq 6) is best carried out in the presence of ethylene or diallyl ether (entries 2 and 3, column 5) since other conditions gave virtually no reaction. The ratio of *E*:*Z* alkenes was found to depend on the additive, with ethylene giving the mixture in an *E*:*Z* ratio of 15:1 as determined by ^1H NMR. This reaction appears to be very sensitive to steric effects, since the *o*-substituted derivative **3b** gave no product.

Experimentally the use of commercially available $[(\text{allyl})\text{PdCl}]_2$ has a clear advantage over $[(\text{allyl})\text{NiBr}]_2$, which is a thermally sensitive solid that has to be kept at low temperature in a drybox.¹⁵ In subsequent experiments we discovered that for simple substrates such as **1** when the Pd-catalyzed reaction is carried out in the presence of (*o*-tolyl)₃phosphine, *there is no need to have the ethylene or diallyl ether present for a quantitative conversion to the product* (entry 5, column 4).

Entries 6^{6d} and 7^{8a} lists the use of two of the best catalysts capable of allylic isomerizations. Neither the Ir (I) nor the modified Grubbs generation II catalyst under the prescribed reaction conditions is useful for this transformation.

Further scope of the reaction is shown in Table 2, which lists other alkenes that undergo the isomerization under these conditions. The 1-indanone-derived alkene **5** gave an excellent yield

of the internal alkene at room temperature (entry 1). Methylene compounds prepared via Wittig reaction of aryl alkyl ketones are excellent substrates for this reaction. 2-Arylbutenes **6–8** gave excellent yields of the product, an internal alkene. While the 2-naphthyl derivative **7** gave good *E*-selectivity, the 1-naphthyl analog **8** gave only a 2:1 mixture of *E* and *Z* alkenes. Attempted isomerization of the furyl derivative **9** led to polymerization.

Monosubstituted terminal alkenes also undergo the isomerization with remarkable ease under the typical conditions outlined in the previous paragraph *giving exclusively the 2-alkene*. Several non-trivial examples of this transformation are shown in Table 3. 6-(*tert*-Butyldimethylsiloxy)hex-1-ene **13** undergoes exceptionally clean isomerization to the corresponding *Z*- and *E*-2-hexenes (**18**) under typical hydrovinylation conditions at -55°C (entry 1).¹³ The expected internal olefin **18** is formed in >95% yield in an *E*:*Z* ratio of 3.5:1.0 as judged by ^1H NMR spectroscopy and gas chromatography. No trace of any other alkenes is observed under these conditions. The reaction catalyzed by Pd(II) (entry 2) gives 80% yield of (*E*)- and (*Z*)-**18** in the same ratio. This reaction is less clean compared to the Ni-catalyzed reaction, the product being contaminated with up to 10% starting material, and 9% of another isomeric alkene.¹³ Isomerization of **13** using the metathesis catalyst (eq 1, 10 mol % [**Ru**], MeOH, 60°C , 12 h) yields <45% of the *E*/*Z* mixture (3.8:1.0) of **18** contaminated with other isomers. The Ir(I) catalyst $[(\text{Ir}(\text{COE})\text{Cl})_2, \text{PCy}_3, \text{NaBPh}_4, \text{CH}_2\text{Cl}_2/\text{acetone} (50:1)]^{4b}$ gave 94% yield of (*E*)-1-*t*-butyldimethylsiloxyhexene (Scheme 2).

The proportion of starting material in the equilibrium mixture of alkenes depends on the starting alkene, as indicated by the isomerization in 1-(but-3-enyloxy)-4-methoxybenzene (**14**) to the disubstituted alkene **19**. The *E*- and *Z*- isomers are obtained in >95% yield in a ratio of 5.9:1.0 with < 4% of the starting material (entry 3). The Ir(I)-catalyst gives a mixture of products arising from both 1,3- and 1,6-hydrogen shifts with the latter predominating (Scheme 2).¹³ The modified Grubbs catalyst gave a mixture of 14% starting material, 47% (*E*)-**19**, 12% (*Z*)-**19** along with a mixture of isomers.¹³ While 1,6-dienes readily undergo cycloisomerization followed by isomerization of the exocyclic alkylidene under Pd-catalysis (eq 3),^{10a} a substrate more resistant to cyclization, such as 1,7-octadiene (**15**), simply undergoes isomerization of the terminal alkenes to give a stereoisomeric mixture of 2,6-octadienes (**20**) [entry 4]. Surprisingly, the triene **16**, prone to isomerization of the endocyclic 1,4-diene, cleanly gives 90% yield of a mixture of *E*- and *Z*-3-butenyl derivatives (**21**) arising from selective isomerization of the side-chain (entry 5). Likewise, the enyne **17** undergoes just the isomerization to **22** *without the expected 1,6-cyclization of the enyne to a bisalkylidene*.

Finally, several sterically demanding substrates, **23–26**, failed to undergo isomerization even under more forcing conditions.

Discussion

Mechanistically the alkene isomerization is related to the olefin dimerization reaction since both start with a common reactive intermediate in the form of an active metal hydride. Such a hydride (**32**, Scheme 3) could be generated by initial formation of the [(allyl)metal (phosphine)]⁺ OTf[−] (**28**)¹⁴ from the metal halide followed by an insertion (**29** → **30**) and β -hydride elimination. The 1,4-diene **31** is formed as a byproduct.^{10b,17} Once the hydride is generated it adds to the terminal alkene to form an organometallic **33**, which undergoes yet

another β -hydride elimination to give the alkene isomers (**34**) in a thermodynamically controlled reaction. Note that almost invariably, the *E*-alkenes predominate in the isomerization reactions. The reluctance of the 1,1'-disubstituted alkene to undergo the reaction in the absence of a sterically unencumbered alkene, ethylene, is thus easily understood. A 1,1'-disubstituted alkene ($R' = \text{alkyl}$) could be slow to undergo the allylic insertion reaction (**29** $\rightarrow$ **30**) and ethylene is needed to generate the reactive metal hydride. The reluctance of the metal hydride to add to a 1,2-disubstituted alkene (**34**, $R' = \text{H}$) might also explain why the exhaustive isomerizations seen with other catalysts (Scheme 2), most notably Ir(I), is not seen in these reactions.

Conclusions

The cationic metal hydride-mediated selective skeletal isomerization of terminal alkenes provides a facile route to di- and trisubstituted alkenes. The reaction is sensitive to steric effects, yet tolerant to a variety of common functional groups and, in some cases, may provide synthetically useful levels of *E:Z* selectivity in the formation of the trisubstituted alkenes.

Experimental Section

General Information

See Supporting Information.

Attempted isomerization of 1-methylene-2,3,4-trihydronaphthalene **1** under hydrovinylation conditions. General procedure A (Table 1, entry 1, column 4)

In a flame-dried flask, [(allyl)NiBr]₂ or [(allyl)PdCl]₂, the phosphine ligand and the silver salt were mixed in that order in CH₂Cl₂ in a drybox. After the precipitated silver salt was removed by filtration through celite, the solution was taken out of the drybox and was connected to an ethylene line. The system was evacuated and refilled with ethylene three times. To the resulting complex was added **1** in CH₂Cl₂ dropwise at room temperature. After stirring for the prescribed time (Table 1), all volatile materials were removed and the product was purified by column chromatography. Gas chromatography and NMR showed complete recovery of starting materials. We have since found that for most isomerization reactions described below it is not necessary to remove the precipitated silver salt.

Use of [(allyl)PdCl]₂/phosphine/AgOTf activated by ethylene for the double bond migration. General procedure B (Table 1, entry 2, columns 4 and 5)

In a flame dried three-neck flask, [(allyl)PdCl]₂, PPh₃ and AgOTf were dissolved in CH₂Cl₂ in a glove box. After the reaction vessel was taken out of the box, an ethylene line was connected to the vessel and the line was evacuated, then ethylene was introduced. The process was repeated three times. The resulting monomeric allylic Pd complex was stirred for 20 min under one atmosphere of ethylene at rt. Then the exo-methylene substrate **1** or **3a** was added dropwise as a solution in CH₂Cl₂. The ethylene was exchanged with a N₂ atmosphere and the resulting mixture was stirred at ambient temperature for the prescribed time. All volatile materials were evaporated, and then the mixture was filtered using a short pad of silica gel using EtOAc/Hex solvent. The product was analyzed by GC and NMR.

Use of [(allyl)PdCl]₂/phosphine/AgOTf activated by diallyl ether for the double bond migration. General procedure C (Table 1, entry 3, columns 4 and 5)

To a premixed allyl-Pd complex in CH₂Cl₂ which was prepared in the same fashion as in the previous experiment (General Procedure B), an equivalent amount of diallylether was added instead of ethylene and then the mixture was stirred at rt for 20 min. The substrate dissolved in CH₂Cl₂ was added dropwise to the catalyst solution, and the resulting mixture was stirred

at ambient temperature for the indicated time. The product was purified and analyzed as indicated previously.

Use of [(allyl)PdCl]₂/phosphine/AgOTf without any additive for the double bond migration. General procedure D (Table 1, entry 5, columns 4 and 5)

To (allyl)Pd(phosphine)-complex in CH₂Cl₂ which was prepared in the same fashion as in the General Procedure B, the substrate dissolved in CH₂Cl₂ was added dropwise, and the resulting mixture was stirred at ambient temperature for the indicated time. The product was purified and analyzed as indicated previously.

Isomerization of 1 using [(allyl)NiBr]₂/Ph₃P/NaBARF and diallylether. (Table 1, entry 4, column 4)

The pre-catalyst was prepared as follows in a glovebox: To [(allyl)NiBr]₂ (10.8 mg, 0.03 mmol) in CH₂Cl₂ (1 mL) were added PPh₃ (15.7 mg, 0.06 mmol) and NaBARF (53.0 mg, 0.06 mmol). The catalyst solution prepared above was removed from the drybox, and diallylether (3.7 μL, 0.03 mmol) was added as a single portion under nitrogen and the mixture was allowed to stir for 2 min to form the active catalytic species. A solution of the substrate **1** (433 mg, 3.00 mmol) in 3 mL of CH₂Cl₂ was added dropwise over a period of one minute and the reaction was allowed to proceed for 2 d. The resulting product was filtered by flash column chromatography (eluted with pentane) to get the desired product **2** (0.35 g, 82 %) as a colorless oil, which was then used to acquire all analytical data without further purification. ¹H NMR (CDCl₃): δ 7.23-7.17 (m, 2 H), 7.15-7.10 (m, 2 H), 5.85-5.83 (m, 1 H), 2.75 (t, *J* = 8.0 Hz, 2 H), 2.26-2.20 (m, 2 H), 2.04 (dd, *J* = 3.2 Hz, *J* = 1.6 Hz, 3 H); ¹³C NMR (CDCl₃): δ 136.3, 135.9, 132.3, 127.3, 126.7, 126.3, 125.4, 122.8, 28.3, 23.2, 19.3.

Isomerization of 2-(3-(3-(benzyloxy)phenyl)but-3-enyl)isoindoline-1,3-dione, 3a (Table 1, entry 2, column 5). General procedure B

In a flame dried three-neck flask, [(allyl)PdCl]₂ (2.4 mg, 0.0061 mmol), PPh₃ (3.7 mg, 0.013 mmol) and AgOTf (3.4 mg, 0.013 mmol) were dissolved in distilled CH₂Cl₂ (3 mL) in a glove box. After the reaction vessel was taken out from a box, an ethylene line was connected to the vessel and the line was evacuated, and then ethylene was introduced. This process was repeated three times. The resulting Pd complex was stirred for 20 min under 1 atm of ethylene atmosphere at rt, and then the exomethylene substrate **3a** (29 mg, 0.076 mmol) in CH₂Cl₂ (1 mL) was added dropwise. Ethylene atmosphere was exchanged for a N₂ atmosphere, the resulting mixture was stirred at 35 °C for 2 d. All volatile materials were evaporated, and then the mixture was purified by flash column chromatography to get 28 mg (79%) of the product **4a** (E:Z = 15:1) contaminated with 21% of starting olefin, **3a**. ¹H NMR (CDCl₃): δ 7.90-7.69 (m, 4 H, phthalimidyl), 7.49-7.29 (m, 5 H, Ar), 7.19 (t, *J* = 6.4 Hz, 1 H, Ar), 7.05-6.85 (m, 2 H, Ar), 6.84-6.79 (m, 1 H, Ar), 5.85 (t, *J* = 7.0 Hz, 1 H, Ar(CH₃)C=CHCH₂N-phth), 5.03 (s, 2 H, OCH₂Ph), 4.50 (d, 7.5 Hz, 2 H, Ar(CH₃)C=CHCH₂N-phth, E), 4.22 (d, *J* = 6.5 Hz, 2 H, Ar(CH₃)C=CHCH₂N-phth, Z), 2.23 (s, 3 H, Ar(CH₃)C=CHCH₂NPhth, E), 2.01 (s, 3 H, Ar(CH₃)C=CHCH₂N-phth, Z).

The configuration of the major product was established by nOe studies. ^{13}C NMR (CDCl_3): δ 168.4, 144.5, 139.4, 137.3, 134.1, 132.5, 129.6, 128.1, 127.8, 123.4, 121.6, 119.1, 113.7, 113.1, 70.2, 36.5, 16.4. HRMS 406.1400 ($\text{M}+\text{Na}^+$; calcd for $\text{C}_{25}\text{H}_{21}\text{NNaO}_3$ 406.1419)

Isomerization of 1-methylene-1,2,3-trihydronaphthalene, **1** (Table 1, entry 2, column 4). General procedure D

Following the general procedure D using 10 mol% of preformed Pd-catalyst, the isomerization of **1** (0.145 g, 1.0 mmol) was checked. After the crude product was purified by column chromatography, the product (**2**, 0.145g, > 99 %) was analyzed by NMR.

Isomerization of 1-methylene-2,3-dihydro-1H-indene **5** (Table 2, entry 1). General procedure D

The isomerization reaction of **5** (0.131 g, 1.0 mmol) was carried out using $[(\text{allyl})\text{PdCl}]_2$ (5 mol%), (*o*-tol) $_3\text{P}$ (10 mol%) and AgOTf (10 mol%) in CH_2Cl_2 with no other additive at rt for 8 h. After the solvent was evaporated, the crude product was purified by column chromatography to yield the desired product **9** (94 %, isolated yield). ^1H NMR (CDCl_3): δ 7.47 (d, 1H, $J = 7.2$ Hz, Ar), 7.39-7.31 (m, 2H, Ar), 7.27-7.23 (m, 1H, Ar), 6.22 (s, 1H, vinyl), 3.33 (s, 2H, allylic CH_2), 2.19 (s, 3H, allylic CH_3); ^{13}C NMR (CDCl_3): δ 146.1, 144.3, 139.9, 128.7, 126.0, 124.4, 123.6, 118.8, 37.6, 13.0.

Isomerization of 1-(but-1-en-2-yl)-4-chlorobenzene **6** (Table 2, entry 2). General procedure D

The isomerization reaction of **6** (0.167 g, 1.0 mmol) was carried out using $[(\text{allyl})\text{PdCl}]_2$ (5 mol%), (*o*-tol) $_3\text{P}$ (10 mol%) and AgOTf (10 mol%) in CH_2Cl_2 with no other additive at rt for 16 h. After the solvent was evaporated, the crude product was purified by column chromatography to yield the desired product **10** (87 %, *E:Z* = 19:1). ^1H NMR (CDCl_3): δ 7.33-7.27 (m, 4H, Ar), 5.86 (q, 1H, $J = 6.50$ Hz, vinyl), 2.02 (s, 3H, allylic CH_3), 1.80 (d, 3H, $J = 6.50$ Hz, allylic CH_3 ; *E*), 1.59 (d, 3H, $J = 6.50$ Hz, allylic CH_3 ; *Z*); ^{13}C NMR (CDCl_3): δ 142.4, 134.5, 132.1, 128.2, 126.8, 123.0, 15.4, 14.3.

Isomerization of 2-(but-1-en-2-yl)naphthalene **7** (Table 2, entry 3). General procedure D

The isomerization reaction of **7** (0.182 g, 1.0 mmol) was carried out using $[(\text{allyl})\text{PdCl}]_2$ (5 mol%), (*o*-tol) $_3\text{P}$ (10 mol%) and AgOTf (10 mol%) in CH_2Cl_2 with no other additive at rt for 16 h. After the solvent was evaporated, the crude product was purified by column chromatography to yield the desired product **11** (97 %, *E:Z* = 9:1). ^1H NMR (CDCl_3): δ 7.83-7.81 (m, 2H, Ar), 7.79-7.76 (m, 2H, Ar), 7.58 (dd, 1H, $J = 9.0, 2.0$ Hz, Ar), 7.47-7.42 (m, 2H, Ar), 6.05 (q, 1H, $J = 7.0$ Hz, vinyl; *E*), 2.15 (s, 3H, allylic CH_3 ; *E*), 2.13 (s, 3H, allylic CH_3 ; *Z*), 1.87 (d, 3H, $J = 7.0$ Hz, allylic CH_3 ; *E*), 1.67 (d, 3H, $J = 7.0$ Hz, allylic CH_3 ; *Z*); ^{13}C NMR (CDCl_3): δ 141.2, 135.4, 133.5, 132.3, 128.0, 127.5, 127.4, 126.0, 125.3, 124.3, 123.8, 123.1, 15.5, 14.5.

Isomerization of 1-(but-1-en-2-yl)naphthalene 8 (Table 2, entry 4). General procedure D

The isomerization reaction of **8** (0.182 g, 1.0 mmol) was carried out using [(allyl)PdCl]₂ (5 mol%), (*o*-tol)₃P (10 mol%) and AgOTf (10 mol%) in CH₂Cl₂ at rt for 16 h. After the solvent was evaporated, the crude product was purified by column chromatography to yield the desired product **12** (99 %, *E*:*Z* = 2:1). ¹H NMR (CDCl₃): δ 8.03 (dd, 1H, *J* = 6.0, 2.5 Hz, Ar), 7.90 (dd, 1H, *J* = 8.0, 3.5 Hz, Ar), 7.79 (d, 1H, *J* = 8.0 Hz, Ar), 7.54–7.50 (m, 3H, Ar), 7.32 (d, 1H, *J* = 6.5 Hz, Ar), 5.87 (q, 1H, *J* = 7.0 Hz, vinyl; *Z*), 5.66 (q, 1H, *J* = 6.50 Hz, 1H; *E*), 2.16 (s, 3H), 1.94 (d, *J* = 6.50 Hz, 3H), 1.42 (d, *J* = 6.50 Hz; *Z*); ¹³C NMR (CDCl₃): δ *E* - 144.3, 135.7, 133.8, 131.4, 128.3, 126.7, 126.0, 125.6, 125.5, 125.4, 124.9, 18.7, 14.0. *Z* - 140.5, 135.8, 133.7, 130.8, 128.3, 125.8, 125.6, 125.2, 124.9, 123.3, 26.1, 14.8.

Isomerization of 1,7-octadiene (Table 3, entry 4). General procedure D

In a nitrogen filled drybox, a dry flask was charged with [(allyl)PdCl]₂ (0.016 g, 0.045 mmol) in 1 mL CH₂Cl₂. To this solution was added two equivalents of phosphine ligand (one equiv with respect to metal, 0.024 g, 0.09 mmol) followed by AgOTf (0.023 g, 0.09 mmol). The precipitate was filtered off using a celite-plugged pipette. After 30 min the reaction vessel was taken out from a box. To the resulting metal catalyst was added 1,7-octadiene (0.200 g, 1.81 mmol) in CH₂Cl₂ dropwise at ambient temperature. After the resulting mixture was stirred at ambient temperature for 2 h, most of the volatile materials were evaporated, and the crude product was analyzed by NMR and GC. Total absence of the starting olefinic peaks in the ¹H NMR due to the starting methylene compound (δ 5.70–5.90 m; 4.90–5.05 m), and appearance of new olefinic H at δ 5.25–5.40 (4 H) and vinyl-CH₃ signals at 1.50–1.51 (6 H) indicate isomerization of the double bonds without cyclization. Since the product is volatile the conversion was estimated to be >95 % by both NMR and GC. No further analysis was carried out.

Synthesis of 6-*tert*-butyldimethylsiloxy)hex-1-ene (13)

To a solution of 0.581 g (5.81 mmol) of 5-hexene-1-ol in 5 mL of DMF at 0 °C under nitrogen was added 1.19 g (17.4 mmol) of imidazole and 1.31 g (8.72 mmol) of *t*-butyldimethylsilyl chloride. The mixture was stirred at rt for 2 days and subsequently quenched with 10 mL of water. The aqueous layer was extracted with ether (3 × 10 mL). The combined organic layers were washed with 2 N aqueous NaOH solution then brine, and dried over MgSO₄ before concentration in vacuo. The residue was purified by flash chromatography on silica gel, eluting with hexane/ethyl acetate (98:2), to afford the silyl ether (**13**) as a clear oil (1.19 g, 96%). ¹H NMR (CDCl₃): δ 5.73–5.90 (m, 1 H), 4.91–5.06 (m, 2 H), 3.62 (t, *J* = 6.4 Hz, 2 H), 2.03–2.12 (m, 2 H), 1.38–1.60 (m, 4 H), 0.90 (s, 9 H), 0.05 (s, 6 H).

Isomerization of 6-*tert*-butyldimethylsiloxy)hex-1-ene. General procedure A (Table 3, entry 1)

To a solution of [(allyl)NiBr]₂ (5.2 mg, 0.14 mmol) in 1 mL of CH₂Cl₂ under nitrogen at room temperature was added a solution of Ph₃P (7.6 mg, 0.28 mmol) in 1 mL of CH₂Cl₂. The resulting brown solution was added to a mixture of AgOTf (10.3 mg, 0.40 mmol) in 1 mL of CH₂Cl₂. After stirring for 1.5 h at rt, the mixture was filtered through a small plug of celite, and the precipitate was rinsed with 2 mL of CH₂Cl₂. The filtrate was collected in a Schlenk flask, and was taken out of the drybox. The catalyst solution was cooled to –55 °C. Under an atmosphere of ethylene, 0.428 g (2.00 mmol) of alkene **13** was added dropwise to the catalyst solution. After stirring at –55 °C for 4 h, the mixture was quenched with saturated NH₄Cl solution and extracted CH₂Cl₂ (3 × 10 mL). The combined organic layers were dried over MgSO₄ and concentrated in vacuo. The crude product was analyzed by GC, which indicated that the alkene had completely isomerized to two new products in a ratio of 3.5:1.0. The volatile crude product (>95% estimated by GC) was purified by chromatography on silica, eluting

with ethyl acetate/hexane (98:2), to afford the product(s) **18** which was analyzed by GC and NMR. ^1H NMR (CDCl_3): δ 5.40–5.46 (m, 2 H), 3.57–3.65 (m, 2 H), 1.97–2.10 (m, 2 H), 1.54–1.67 (m, 5 H), 0.90 (s, 9 H), 0.05 (s, 6 H).

Palladium-catalyzed isomerization of 6-*tert*-butyldimethylsiloxy)hex-1-ene. General procedure D (Table 3, entry 2)

The isomerization reaction was carried out using $[(\text{allyl})\text{PdCl}]_2$ (0.50 mmol), (*o*-tol) $_3\text{P}$ (1 mol %) and AgOTf (1 mol%) in CD_2Cl_2 . The reaction was followed by ^1H NMR spectroscopy, which indicated maximum conversion to the products at 24 h. The products (*E*)- and (*Z*)-**18** (80 %, 3.7:1.0) were identified by comparison of gas chromatogram and spectral properties of the sample from the previous run. In addition to 10% starting material, ~9% of an unidentified product was also detected by gas chromatography (see GC traces).

Synthesis of 1-(but-3-enyloxy)-4-methoxybenzene (**14**)

To a stirred solution of 4-methoxyphenol (1 g, 8.06 mmol), 3-buten-1-ol (0.99 g, 10.48 mmol), and triphenylphosphine (2.75 g, 10.48 mmol) in anhydrous THF (20 mL) was added diisopropylazodicarboxylate (2.12 g, 10.48 mmol) slowly at 0 °C. After the reaction was stirred overnight at rt, the mixture was diluted with EtOAc (20 mL) and water (20 mL), and separated. The crude compound was extracted with EtOAc (3 $\times$ 20 mL) and the combined organic layers were dried over Na_2SO_4 . The resulting product was purified by chromatography to afford 1.5 g (93 %) of **14**. ^1H NMR (CDCl_3): δ 6.87–6.81 (m, 4 H, Ar), 5.94–5.86 (m, 1H, $\text{ROCH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 5.18–5.09 (q, 2H, $\text{ROCH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 3.97–3.95 (t, J = 7.0 Hz, 2H, $\text{ROCH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 2.53–2.49 (m, 2 H, $\text{ROCH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 3.75 (s, 3H, ArOCH_3), 1.75–1.70 (d, J = 5.0 Hz, 3H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$); ^{13}C NMR (CDCl_3): δ 154.0, 153.2, 134.8, 117.1, 115.8, 114.8, 68.1, 55.9, 33.9.

Isomerization of 1-(but-3-enyloxy)-4-methoxybenzene (**14**). General procedure D (Table 3, entry 3)

1-(But-3-enyloxy)-4-methoxybenzene (30 mg, 0.168 mmol) was reacted with $[(\text{allyl})\text{PdCl}]_2$ (1.6 mg, 0.0042 mmol), (*o*-tol) $_3\text{P}$ (2.6 mg, 0.0084 mmol), AgOTf (2.2 mg, 0.0084 mmol), and CH_2Cl_2 at rt for 1 d. The resulting product was purified by chromatography to afford 30 mg (> 99 %) of **19** as a mixture of *E* and *Z* isomers with the starting material. Based on ^1H NMR and GC, the conversion and *E/Z* ratio were estimated as 96 % and 5.9:1.0 respectively. ^1H NMR (CDCl_3): δ 6.86–6.80 (m, 4H, Ar), 5.84–5.80 (m, 1H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$, *E* and *Z*), 5.74–5.68 (m, 1 H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$, *E* and *Z*), 4.54–4.53 (d, J = 5.5 Hz, 2H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$, *Z*), 4.53–4.39 (m, 2H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$, *E*), 3.75 (s, 3H, ArOCH_3), 1.75–1.70 (d, J = 5.0 Hz, 3H, $\text{ROCH}_2\text{CH}=\text{CHCH}_3$); ^{13}C NMR δ (CDCl_3 , *E* only): 154.2, 153.3, 130.3, 126.8, 116.0, 69.7, 56.0, 17.9, 13.5.

The configuration of the major product was established by nOe studies.

Isomerization of methyl 1-(2-butenyl)-2,5-cyclohexadiene-1-carboxylate (16**). General procedure D (Table 3, entry 5)**

Substrate **16** (0.1 g, 0.5 mmol) was reacted with 5 mol% [(allyl)PdCl]₂ (10.4 mg, 0.026 mmol), 5 mol% of (*o*-CH₃-C₆H₄)₃P (16 mg, 0.05 mmol), AgOTf (13 mg, 0.05 mmol), and CH₂Cl₂ at rt for 1 d. The resulting product was purified by chromatography to yield 91 mg (91%) of **21** as a mixture E/Z isomers. Based on the ¹H NMR, the E/Z ratio was estimated as 3.6:1.0. ¹H NMR (CDCl₃): δ 5.88-5.73 (m, 4 H, -CH=CH- in the ring), 5.41-5.22 (m, 2 H, CH₂-CH=CH-CH₂-, E and Z), 3.69 (s, 3 H, OCH₃, Z), 3.67 (s, 3 H, OCH₃, E), 2.63 (m, 2 H, R=CH-CH₂-CH=R, E and Z), 2.43 (d, *J* = 5.2 Hz, 2 H, C-CH₂-CH=CHCH₃), 2.34 (d, *J* = 6.1 Hz, 2 H, C-CH₂-CH=CHCH₃), [δ 2.43 + δ 2.34] 2 H, 1.51-1.61 two sets of doublets (d, *J* = 6.7 Hz, 3 H, C=C-CH₃); ¹³C NMR (CDCl₃): δ 174.7, 128.6, 127.0, 126.7, 125.4, 125.3, 51.9, 47.9, 43.4, 37.2, 26.0, 17.9; HRMS 192.1132 (M⁺; calcd for C₁₂H₁₆O₂ 192.1150).

Isomerization of (*tert*-butyl[[2,2-dimethyl-1-(1-propynyl)-3-pentenyl]oxy] dimethylsilane (17**). General procedure D (Table 3, entry 6)**

Substrate **17**₁₆ (0.1 g, 0.8 mmol) was reacted with 5 mol% of [(allyl)PdCl]₂, 10 mol% of (2-Me-C₆H₄)₃P and 10 mol% AgOTf in CH₂Cl₂ at for 16 h. The reaction mixture was quenched with saturated ammonium chloride and the product was extracted with ether. The dried organic layer concentrated and the product was purified by flash column chromatography to get 95 mg (95%) of **22** as a thick oil. Based on the ¹H NMR, the E/Z ratio was estimated as 8.0:1.0. ¹H NMR (CDCl₃): δ 5.57-5.30 (m, 2H), 4.00 (m), 3.91 (q, *J* = 2.3 Hz) [together 1 H, R₃Si-O-CH], 1.79 (d, *J* = 2.3 Hz, 3H, R-CαC-CH₃), 1.66 (d, *J* = 5.6 Hz, 3 H, CH=CH-CH₃), 1.00 (2s, separated by ~ 2 Hz, 6 H, R(CH₃)₂), 0.10 (2s, 9H, *t*Bu in OTBS), 0.11 (s, 3H, Me in OTBS), 0.05 (s, 3H, Me in OTBS); ¹³C NMR (CDCl₃): δ 137.8, 120.1, 79.8, 79.3, 70.5, 40.9, 25.5, 25.4, 22.5, 22.1, 18.0, 2.9, -4.5; HRMS 266.2057 (M⁺; calcd for C₁₆H₃₀O₄Si 266.2066).

Isomerization of 6-*tert*-butyldimethylsiloxy)hex-1-ene (13**) using Grubbs second-generation catalyst.^{8a}**

Using the same procedure and catalyst (10 mol%) aforementioned, the isomerization of TBS-protected olefin **13** (11.2 mg, 0.052 mmol) was checked. The crude product was analyzed by GC. See Supporting Information for chromatograms.

Isomerization of 1-(but-3-enyloxy)-4-methoxybenzene (14**) using Grubbs second-generation catalyst shown in eq 1.^{8a}**

The starting olefin **14** (9.3 mg, 0.052 mmol) in anhydrous MeOH (0.8 mL) was treated with Grubbs second-generation catalyst (4.4 mg, 10 mol%). The resulting solution was stirred at 60

°C for 12 h. After the solvent was evaporated in vacuo, the residue was filtered on a silica pad and eluting with ether. The crude product was directly analyzed by GC. Supporting Information for chromatograms.

Isomerization of 1-methylene-2,3,4-trihydronaphthalene (1) using Grubbs second-generation catalyst

Using the same procedure and catalyst (10 mol%) aforementioned, the isomerization of olefin **1** (15.0 mg, 0.104 mmol) was checked. The crude product was purified by short column chromatography, and analyzed by ¹H NMR.

Isomerization of 2-(3-(3-(benzyloxy)phenyl)but-3-enyl)isoindoline-1,3-dione (3a) using Grubbs second-generation catalyst

Using the same procedure and catalyst (10 mol%) aforementioned, the isomerization of olefin **3a** (10.0 mg, 0.0261 mmol) was checked. The crude product was purified by short column chromatography, and analyzed by ¹H NMR.

Isomerization of 6-*tert*-butyldimethylsiloxy)hex-1-ene (13) using [Ir(COE)₂Cl]₂/PCy₃/NaBPh₄ in CH₂Cl₂/Acetone (Scheme 2).^{4b,6d}

Using the same procedure and catalyst (2 mol%) aforementioned, the isomerization of TBS-protected olefin **13** (25.5 mg, 0.119 mmol) was checked. After the resulting mixture was stirred at rt for 3 h, most of the volatile materials were evaporated, and the crude product was purified by column chromatography to get the isomerized silylenolether *tert*-butyl(hex-1-enyloxy)dimethylsilane (24.0 mg, 94 %). The product was analyzed by ¹H NMR and GC. ¹H NMR (CDCl₃): δ 6.21-6.18 (dt, 1H, J = 12.0, 1.5 Hz, nBuCH=CHOTBS), 4.97-4.95 (q, 1H, 3.8 Hz, nBuCH=CHOTBS), 1.89-1.82 (m, 2H, nBu), 1.30-1.26 (m, 4H, nBu), 0.90-0.84 (m, 12 H, tBuSi & nBu), 0.10 (s, 6H, SiMe₂).

Isomerization of 1-(but-3-enyloxy)-4-methoxybenzene (14) using [Ir(COE)₂Cl]₂/PCy₃/NaBPh₄ in CH₂Cl₂/Acetone.^{4b,6d}

In a nitrogen filled drybox, a dry flask was charged with [Ir(COE)Cl]₂ (1.1 mg, 0.0012 mmol), PCy₃ (1.9 mg, 0.0068 mmol), and NaBPh₄ (0.8 mg, 0.0029 mmol), and the mixture was dissolved in CH₂Cl₂/acetone (0.52 mL, 25/1). After 5 min stirring, the starting olefin **14** (21.2 mg, 0.119 mmol) in CH₂Cl₂ (0.5 mL) was added dropwise at rt. After the resulting mixture was stirred for 3 h at rt, most of the volatile materials were evaporated, and the crude product was purified by column chromatography to yield the mixture of the desired product (**19**) and further isomerized 1-(but-1-enyloxy)-4-methoxybenzene (2-OPMP) (20.4 mg (96 %), **19**:2-OPMP = 1.0:3.5 (based on ¹H NMR)). The product was analyzed by ¹H NMR and GC. ¹H NMR (CDCl₃): δ 6.92-6.88 (m, 2H, Ar), 6.84-6.79 (m, 2H, Ar), 6.36-6.33 (dt, 1H, J = 12.0, 1.5 Hz, ROCH=CH₂Et), 5.32-5.27 (m, 1H, ROCH=CH₂Et), 3.75 (s, 3H, OMe), 2.05-1.99 (m, 2H, RCH₂CH₃), 1.05-0.98 (t, 3H, J = 6.0 Hz, RCH₂CH₃).

Attempted isomerization of 1-methylene-2,3,4,5-trihydronaphthalene (1) using Ir(COE)₂Cl]₂/PCy₃/NaBPh₄ in CH₂Cl₂/Acetone.^{4b,6d}

Using the same procedure and catalyst (2 mol%) aforementioned, the isomerization of olefin **1** (17.1 mg, 0.119 mmol) was checked. The crude product was purified by short column chromatography, and analyzed by ¹H NMR.

Attempted isomerization of 2-(3-(3-(benzyloxy)phenyl)but-3-enyl)isoindoline-1,3-dione (**3a**) using $\text{Ir}(\text{COE})_2\text{Cl}]_2/\text{PCy}_3/\text{NaBPh}_4$ in $\text{CH}_2\text{Cl}_2/\text{Acetone}$.^{4b,6d}

Using the same procedure and catalyst (2 mol%) aforementioned, the isomerization of olefin **3a** (22.8 mg, 0.060 mmol) was checked. The crude product was purified by short column chromatography, and analyzed by ^1H NMR.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial assistance for this research by NSF (CHE- 0610349) and NIH (General Medical Sciences, R01 GM075107) is gratefully acknowledged. Craig Smith also thanks the ACS Organic Division for a graduate fellowship sponsored by The Schering Plough Corporation.

REFERENCES AND FOOTNOTES

- (a)Frauenrath H, Kropf H, Schaumann E. Houben-Weyl, E15/1 ThiemeStuttgart1995:1.(b) For a recent review describing applications in synthesis which appeared after the initial submission of this manuscript, see: (c)Donohue TJ, O’Riordan TJC, Rosa CP. Angew Chem Int Ed 2009;48:1014.
- Greene, TW.; Wuts, PGM. Protective Groups in Organic Synthesis. Vol. 2. Wiley; New York: 1991. Kocienski, PJ. Protecting Groups. Thieme Verlag; Stuttgart: 2000.
- (a) Otsuka S, Tani K. Synthesis 1991:665. (b) Frauenrath H, Brethauer D, Reim S, Maurer M, Raabe G. Angew Chem Int Ed 2001;40:177.
- (a)Reuter JM, Salomon RG. J Org Chem 1977;42:3360.(b)Nelson SG, Bungard CJ, Wang K. J Am Chem Soc 2003;125:13000. and references cited therein. [PubMed: 14570453](c)Stevens BD, Bungard CJ, Nelson SG. J Org Chem 2006;71:6397. [PubMed: 16901121](d)Ohmura T, Yamamoto Y, Miyaura N. Organometallics 1999;18:413.(e) A mineral acid-mediated isomerization of terminal methylene compounds is unlikely to be of broad value for reactions of sensitive substrates reported in this paper: (f)Buckle DR, Arch JRS, Edge C, Foster KA, Houge-Frydrych CSV, Pinto IL, Smith DG, Taylor JF, Taylor SG, Tedder JM, Webster RAB. J Med Chem 1991;34:919. [PubMed: 2002472]
- Wille A, Tomm S, Frauenrath H. Synthesis 1998:305.and references cited therein
- (a) Baudry D, Ephritikhine M, Felkin H. J Chem Soc, Chem Commun 1978:694. (b) Matsuda I, Kato T, Sato S, Izumi Y. Tetrahedron Lett 1986;27:5747. (c) Baxendale IR, Lee AL, Ley SV. J Chem Soc, Perkin Trans 1 2002:1850. (d) Shen X, Wasmuth AS, Zhao J, Zhu C, Nelson SG. J Am Chem Soc 2006;128:7438. [PubMed: 16756287] (e) Patnam R, Juárez-Ruiz JM, Roy R. Org Lett 2006;8:2691. [PubMed: 16774233]
- (a)Harrod JF, Chalk AJ. J Am Chem Soc 1964;86:1776.(b)Davies NR. Aust J Chem 1964;17:212.(c) Gosser LW, Parshall GW. Tetrahedron Lett 1971:2555.(d)Lochow CF, Miller RG. J Org Chem 1976;41:3020.(e)Malanga C, Urso A, Lardicci L. Tetrahedron Lett 1995;36:1133.(f) For reports on the use of isomerization of C-allyl compounds in the context of complex molecule synthesis, see: via Rh-intermediates: (g)Warmerdam E, Tranoy I, Renoux B, Gesson JP. Tetrahedron Lett 1998;39:8077. (h)Nakamura H, Arata K, Wakamatsu T, Ban Y, Shibasaki M. Chem Pharm Bull 1990;38:2435.Pd: (i)Whang K, Cooke RJ, Okay G, Cha JK. J Am Chem Soc 1990;112:8985.Use of a ruthenium hydride: (j)Curran DP, Jacobs PB, Elliott RL, Kim BH. J Am Chem Soc 1987;109:5280.
- (a)Hanessian S, Giroux S, Larsson A. Org Lett 2006;8:5481. [PubMed: 17107052](b)Dinger MB, Mol JC. Eur J Inorg Chem 2003:2827.See, also: (c)Arisawa M, Terada Y, Nakagawa M, Nishida A. Angew Chem Int Ed 2002;41:4732. and references cited therein.(d)Yue CJ, Liu Y, He R. J Mol Catal A: Chem 2006;259:17.(e)Böhrsch V, Blechert S. Chem Commun 2006:1968.(f)Wipf P, Spencer SR. J Am Chem Soc 2005;127:225. [PubMed: 15631472](g)Schmidt B. Eur J Org Chem 2004:1865.For a bifunctional Ru-catalyst which brings about isomerization of double bonds over 30 positions, see: (h)Grotjahn DB, Larsen CR, Gustafson JL, Nair R, Sharma A. J Am Chem Soc 2007;129:9592. [PubMed: 17636924] See also: (i)Cadot C, Dalko PI, Cossy J. Tetrahedron Lett 2002;43:1839.(j)Alcaide B, Almendros P,

- Alonso JM, Aly MF. *Org Lett* 2001;3:3781. [PubMed: 11700137](k)Donohue TJ, Rosa CP. *Org Lett* 2007;9:5509. [PubMed: 18004864]
9. (a)Zhang A, RajanBabu TV. *J Am Chem Soc* 2006;128:5620. [PubMed: 16637613](b)Smith CR, RajanBabu TV. *Org Lett* 2008;10:1657. [PubMed: 18351772](c)Smith CR, Lim HJ, RajanBabu TV. *Synthesis*. 2009 (in press).For a recent reviews of hydrovinylation see: (d)Jolly PW, Wilke G. Cornils B, Herrmann WA. *Hydrovinylation. Applied Homogeneous Catalysis with Organometallic Compounds VCH*New York1996;2:1024–1048.1048(e)RajanBabu TV. *Chem Rev* 2003;103:2845. [PubMed: 12914483](f)RajanBabu TV. *Synlett* 2009:853. [PubMed: 19606231]
10. (a)Radetich B, RajanBabu TV. *J Am Chem Soc* 1998;120:8007.For other related observations see: (b)Lloyd-Jones GC. *Org Biomol Chem* 2003:215. [PubMed: 12929414](c)Kisanga P, Widenhoefer RA. *J Am Chem Soc* 2000;122:10017.(d)Yamamoto Y, Ohkoshi N, Kameda M, Itoh K. *J Org Chem* 1999;64:2178.(e)Heumann A, Moukhliiss M. *Synlett* 1998:1211.(f)Grigg R, Malone JF, Mitchell TRB, Ramasubbu A, Scott RM. *J Chem Soc, Perkin Trans 1* 1984:1745.(g)Behr A, Freudenberg U, Keim W. *J Mol Catal* 1986;35:9.(h)Bogdanovic B. *Adv Organomet Chem* 1979;17:105.
11. Importance of tri-substituted alkenes, (a)Chatterjee AK, Grubbs RH. *Org Lett* 1999;1:1751. [PubMed: 10836036]A recent report on E/Z isomerization of arylalkenes. see: (b)Yu J, Gaunt MJ, Spencer JB. *J Org Chem* 2002;67:4627. [PubMed: 12076172]
12. (a) Nomura N, Jin J, Park H, RajanBabu TV. *J Am Chem Soc* 1998;120:459. (b) RajanBabu TV, Nomura N, Jin J, Nandi M, Park H, Sun X. *J Org Chem* 2003;68:8431. [PubMed: 14575468]
13. See Supporting Information for ¹H and ¹³C NMR spectra of key starting materials and products, and gas chromatograms of products from various reactions.
14. Effect of counteranions in related reactions, see: Nandi M, Jin J, RajanBabu TV. *J Am Chem Soc* 1999;121:9899.
15. (a)Wilke G, Bogdonovic B, Hardt P, Heimbach P, Keim W, Kröner M, Oberkirck W, Tanaka K, Steinrück E, Walter D, Zimmermann H. *Angew Chem Int Ed Engl* 1966;5:151.(b)Herrmann WA, Salzer A. *Synthetic Methods of Organometallic and Inorganic Chemistry Georg Thieme Verlag*Stuttgart1996;1:156.Details of the preparation of [(allyl)NiBr]₂: (c)Smith CR, Zhang A, Mans D, RajanBabu TV. *Org Synth* 2008;85:248.
16. (a) Trost BM, Lautens M. *J Am Chem Soc* 1985;107:1781. (b) Trost BM. *Acc Chem Res* 1990;23:34.
17. DiRenzo, GM. For isolation of such an insertion product in styrene dimerization. Ph D Thesis. University of North Carolina; 1997. *Mechanistic Studies of Catalytic Olefin Dimerization Reactions Using Electrophilic η^3 -allyl- Palladium(II) Complexes*.
18. (a) Kobayashi H, Sonoda A, Iwamoto H, Yoshimura M. *Chem Lett* 1981;10:579. (b) Brookhart M, Grant B, Volpe AF Jr. *Organometallics* 1992;11:3920–3922.
20. Exon C, Magnus P. *J Am Chem Soc* 1983;105:2477.

Figure 1.
Terminal alkene substrates for isomerization listed in Table 3

Scheme 1.
Isomerization of Allyl Ethers into Vinyl Ethers

Scheme 2. Isomerization of a Terminal Alkene using an Ir (I) Catalyst^{4b,6d}

Scheme 3.
A Possible Mechanism for the Alkene Isomerization

Table 1
Isomerization of 1-Methylenetetralin (**1**) and 1-(2-Phthalimidoethyl)styrene **3a**^a

entry	catalyst (mol%), additives (mol%)	conditions	yield (%)	
			2	4 (E:Z)
1.	[(allyl)NiBr] ₂ (5), Ph ₃ P (10)AgOTf (5), ethylene	ethylene (1 atm), 25 °C, 2 d	0	< 5%
2.	[(allyl)PdCl] ₂ (5), Ph ₃ P (5)AgOTf (5), ethylene	ethylene (1 atm, 20 min), 25 °C, 2 d	90	79 (15:1) ^b
3.	[(allyl)PdCl] ₂ (1), Ph ₃ P (5)AgOTf (1), diallyl ether (1)	25 °C, 1.5 d	90	79 (8:1) ^b
4.	[(allyl)NiBr] ₂ (1), Ph ₃ P (2)NaBARF(1), diallyl ether (1)	25 °C, 2 d	82	--
5.	[(allyl)PdCl] ₂ (5), (<i>o</i> -tol) ₃ P (10)AgOTf (10), CH ₂ Cl ₂	25 °C, 2 d	>99	~10 ^c
6. ^d	(Ir(COE) ₂ Cl) ₂ , PCy ₃ , NaBPh ₄ , CH ₂ Cl ₂ /acetone (50:1)	25 °C, 3 h	< 2	<2
7. ^e	[Cl ₂ RuL ₃] Grubbs 2 nd Gen. (10)MeOH	60 °C, 1 d	~ 7%	~ 7%

^a See eq 5,6 and experimental section for details.

^b 16 mol% cat., 35 °C,

^c 10 mol% cat.

^d ref. 6d, Scheme 1.

^e ref. 8a, see eq 1.

Table 2Isomerization of 2,2'-Disubstituted Alkenes^a

entry	alkene	time (h)
1.	5	8
2.	6	16
3.	7	20

entry	alkene	time (h)
-------	--------	----------

4.

12

5.

^a [(allyl)PdCl]₂ (5 mol%), (*o*-tol)₃P (10 mol%), AgOTf (10 mol%), CH₂Cl₂ (0.05M).

^b determined by ¹H NMR analysis.

Table 3

Selective Isomerization of C- and O-Allyl Derivatives

entry	substrate	catalyst (mol%), additives (mol%)	conditions	product(s)	yield (%) (E/Z) ^a
1.	13	[allyl] NiBr ₂ (0.7), (C ₆ H ₅) ₃ P (1.4), AgOTf (1.4), ethylene	-55 °C, 4 h		>95 (3.5:1.0)
2.	13	[allyl] PdCl ₂ (0.5), (o-Me) C ₆ H ₄) ₃ P (1.0), AgOTf (1.0), CD ₂ Cl ₂	rt, 24 h	18	80 (3.7:1.0) ^b

entry	substrate	catalyst (mol%), additives (mol%)	conditions	product(s)	yield (%) (E/Z) ^a
3.	14	$[(\text{allyl})\text{PdCl}]_2$ (2.5), <i>o</i> -Me $\text{C}_6\text{H}_4\text{P}$ (5.0), AgOTf (5.0), CD_2Cl_2 ,	rt, 24 h		96 (5.9:1.0) ^c
4.	15	$[(\text{allyl})\text{PdCl}]_2$ (2.5), $(\text{C}_6\text{H}_5)_3\text{P}$ (5), AgOTf (5)	23 °C, 2 h		90 (–) ^d

entry	substrate	catalyst (mol%), additives (mol%)	conditions	product(s)	yield (%) (E/Z) ^a
5.	16	[allyl] PdCl ₂ (5) (<i>o</i> -Me- C ₆ H ₄) ₃ P (10), AgOTf (10)	23 °C, 1 d		90 (3.6:1.0)
6.	17	[allyl] PdCl ₂ (5), (<i>o</i> -Me- C ₆ H ₄) ₃ P (10), AgOTf (10)	23 °C, 1 d		92 (8:1)

^aDetermined by NMR or GC.

^b9% unidentified material and 10% starting material, rest starting material.

^cmixture of E/Z, configuration not determined.