A survey on head lice infestation in Korea (2001) and the therapeutic efficacy of oral trimethoprim/sulfamethoxazole adding to lindane shampoo

Seobo SIM¹⁾, In-Yong LEE¹⁾, Kyu-Jae LEE²⁾, Jang-Hoon SEO³⁾, Kyung-Il IM¹⁾,
Myeong Heon SHIN¹⁾ and Tai-Soon YONG^{1)*}

¹⁾Department of Parasitology and Institute of Tropical Medicine, Yonsei University College of Medicine, Seoul 120-752,

²⁾Department of Parasitology, Yonsei University Wonju College of Medicine, Wonju 220-701,

³⁾Department of Clinical Pathology, Shinheung College, Uijeongbu 480-701, Korea

Abstract: Total of 7,495 children including 3,908 boys and 3,587 girls from a kindergarten and 15 primary schools were examined for head lice infestation (HLI). The overall prevalence of HLI in this study was found to be 5.8%. Head lice were much more commonly detected in girls than in boys with prevalence of 11.2% and 0.9%, respectively. Sixty-nine children with HLI were treated with 1% lindane shampoo alone (group 1), and 45 children with HLI were treated with 1% lindane shampoo and oral trimethoprim/sulfamethoxazole (group 2), and follow-up visits were conducted 2 and 4 weeks later. The children who still had HLI 2 weeks after the primary treatment were treated again. At the 2-week follow-up visit, the treatment success rates of groups 1 and 2 were 76.8% and 86.7%, respectively, and at the 4-week follow-up visit, the rates were 91.3% and 97.8%, respectively. No statistically significant synergistic effect was observed for the combination of a 1% lindane shampoo and oral trimethoprim/sulfamethoxazole.

Key words: *Pediculus capitis*, lindane, trimethoprim-sulfamethoxazole combination, Korea

INTRODUCTION

Head lice infestation (HLI) caused by *Pediculus capitis*, which has existed for more than 10,000 years, is a common, worldwide problem (Gratz et al., 1997; Chouela et al., 1999; Araujo et al., 2000; Ha et al., 2000). The prevalence of infestation is ubiquitous, particularly among children, however, it is also

frequent among adults (Gratz et al., 1997). Therefore, up-to-date the information on the extent of lice infestation is important in order to determine the priority that should be given to control it (Gratz et al., 1997). The head louse is an obligate human parasite, which is transmitted directly by close contact with an infected individual's head or indirectly via fomites such as hair, hats, or combs (Speare, 2000; Bartels et al., 2001). This infestation readily spreads to co-dwelling adults and children, and therefore, presents a public health problem that is difficult to solve. Classmates, playmates, and

• Received 12 November 2002, accepted after revision 27 November 2002.

*Corresponding author (e-mail: tsyong212@yumc.yonsei.ac.kr)

family members of infested children should be inspected for head lice, and efforts should also be directed to the control of fomites and nits (Elston DM, 1999). Head lice feed by sucking blood and cause pruritus to host due to sensitization to louse saliva, and subsequent skin excoriation may lead to secondary bacterial infections. In addition, chronic irritation and secondary infection may disrupt behavior and school performance (Bailey and Prociv, 2000), and HLI may cause psychological distress (Dodd, 2001; Mumcuoglu, 1999), however, is not considered to be a serious health problem. Nevertheless, HLI consume considerable time and attention of the public and private health communities.

Mechanical removal and pediculocidal agents are often used to treat HLI (Kuwahara and Rasberry, 2001). However, resistance to pediculocidal agents has been reported (Schachner, 1997), and the high prevalence of head lice and consequent heavy use of insecticidal products may explain the rapid increase of resistant strains. Sublethal doses of residual pediculicides may also predispose to resistance (Bailey and Prociv, 2000; Meinking et al., 2001). Patterns of resistance will almost certainly vary geographically, therefore, it is not possible to make an overall estimate of the level of resistance to the various pediculicides in different areas. The choice of treatment depends on local resistance patterns, and whenever treatment failed this should be assessed on a case-by-case basis. It has been known that combination therapy of 1% permethrin with trimethoprim/sulfamethoxazole is effective in cases of lice-related resistance to therapy (Hipolito et al., 2001).

Lindane is a prescription product that kills lice by continuously stimulating the nervous system, but it is also neurotoxic to humans (Gratz et al., 1997; Bartels et al., 2001). On the other hand, trimethoprim/sulfamethoxazole has been reported as an effective treatment for HLI (Hipolito et al., 2001), and it kills lice by destroying symbiotic bacteria which synthesize B group vitamins in the lice gut (Burns, 1987).

Until better agents are available, clinicians may find that rotational or combination treatments with different agents may help to slow the emergence of

resistance (Elston, 1999).

This study was undertaken to survey the recent status of HLI in Korean children and evaluate data on the therapeutic efficacy of oral administration of trimethoprim/sulfamethoxazole adding to lindane shampoo.

MATERIALS AND METHODS

Study population

Total of 7,495 children including 3,908 boys and 3,587 girls from kindergarten and 15 primary schools throughout the country were examined for head lice from April 2001 to October 2001. Two schools were in urban areas (one in Seoul and the other in Gyeonggi-do) and the remainder in rural areas (two in Gyeonggi-do, four in Gangwon-do, two in Chungcheongbuk-do, two in Chungcheongnam-do, one in Gyeongsangnam-do, and two in Jeollanam-do). HLI was identified by examining the scalp and hairs for the presence of adult lice, nymphs or viable nits. The viability of nits was confirmed by stereomicroscopic inspection. Viable nits were plump, shiny, and tan to coffee colored with an intact operculum and frequently a pigmented eye-spot. Nits that were empty, shriveled, misshapen, or indented, and those that contained air pockets were considered non-viable (Meinking TL et al., 1986).

Study design

Group 1

A total of 69 children with HLI from 3 schools (one in urban area, Seoul; two in rural areas, Gyeonggi-do) were treated with 1% lindane shampoo (Sinsin Pharm. Co., Seoul, Korea). A quantity sufficient to saturate the hair was applied directly to dry hair and a small amount of water was added to form a good lather. The hair was rinsed with clean water after 10 minutes and towel dried.

Group 2

A total of 45 children with HLI from the same school were treated with a combination of 1% lindane shampoo as described above and trimethoprim/sulfamethoxazole orally administered at a dosage of 8 mg/kg/day based on trimethoprim

for 12 days in 2 divided doses.

Follow-up observations and subsequent treatment

Follow-up examinations were carried out 2 and 4 weeks after the initial treatment. Children who still had HLI 2 weeks after the primary treatment were treated again as described above and reexamined at 4 weeks later.

Questionnaire

In 8 of 15 primary schools, children were asked whether or not they were living with both parents. Children were also asked about academic qualifications and jobs of their parents.

Statistical analysis

Statistical analysis of results was performed using the chi-square test. Statistical significance was assumed at a P -value < 0.05 .

RESULTS

The overall prevalence of HLI was 5.8%, and the

rate was higher in rural areas (10.6%) than that in urban areas (3.2%). Head lice were much more commonly detected in girls than in boys with prevalence of 11.2% and 0.9%, respectively (Table 1). The prevalence of HLI by province in urban areas was 2.3% and 4.2% for Seoul and Gyeonggi-do, respectively, and that in rural areas was 10.1%, 9.4%, 5.6%, 14.2%, 5.6%, and 20.2% for Gyeonggi-do, Gangwon-do, Chungcheongbuk-do, Chungcheongnam-do, Gyeongsangnam-do, and Jeollanam-do, respectively (Table 2).

The prevalence of HLI by school grade was 8.0%, 2.9%, 5.4%, 6.4%, 7.6%, 7.9%, and 4.2% for kindergarten, first, second, third, fourth, fifth, and sixth grades, respectively (Table 3).

At 2 weeks after the primary treatment, the success rates of treatment in the groups 1 and 2 were 76.8% and 86.7%, respectively. The success rates of secondary treatment at the 4-week follow-up in the groups 1 and 2 were 91.3% and 97.8%, respectively (Table 4). No statistically significant difference was found between the two groups ($P > 0.1$). The success rate of the secondary treatment at the 4-week follow-up for each group was higher than that of the

Table 1. Prevalence of head lice infestation by area and sex

Area	Male		Female		Total	
	No. examined	No. positive (%)	No. examined	No. positive (%)	No. examined	No. positive (%)
Urban	2,512	14 (0.6)	2,304	140 (6.1)	4,816	154 (3.2)
Rural	1,396	22 (1.6)	1,283	261 (20.3)	2,679	283 (10.6)
Total	3,908	36 (0.9)	3,587	401 (11.2)	7,495	437 (5.8)

Table 2. Prevalence of head lice infestation by province

Area	Province (No. school)	Male		Female		Total	
		No. examined	No. positive (%)	No. examined	No. positive (%)	No. examined	No. positive (%)
Urban	Seoul (1)	1,293	5 (0.4)	1,172	51 (4.4)	2,465	56 (2.3)
	Gyeonggi-do (1)	1,219	9 (0.7)	1,132	89 (7.9)	2,351	98 (4.2)
Rural	Gyeonggi-do (2)	362	7 (1.9)	320	62 (19.4)	682	69 (10.1)
	Gangwon-do (4)	247	6 (2.4)	241	40 (16.6)	488	46 (9.4)
	Chungcheongbuk-do (2)	108	1 (0.9)	87	10 (11.5)	195	11 (5.6)
	Chungcheongnam-do (2)	192	2 (1.0)	224	57 (25.4)	416	59 (14.2)
	Gyeongsangnam-do (1)	314	4 (1.3)	253	28 (11.1)	567	32 (5.6)
	Jeollanam-do (2)	173	3 (1.7)	158	64 (40.5)	331	67 (20.2)
Total	(15)	3,908	36 (0.9)	3,587	401 (11.2)	7,495	437 (5.8)

Table 3. Prevalence of head lice infestation by school grade and sex

Grade	Male		Female		Total	
	No. examined	No. positive (%)	No. examined	No. positive (%)	No. examined	No. positive (%)
K ^{a)}	104	1 (1.0)	120	17 (14.2)	224	18 (8.0)
1	653	9 (1.4)	540	26 (4.8)	1,193	35 (2.9)
2	659	5 (0.8)	594	63 (10.6)	1,253	68 (5.4)
3	679	10 (1.5)	631	74 (11.7)	1,310	84 (6.4)
4	638	4 (0.6)	586	89 (15.2)	1,224	93 (7.6)
5	576	6 (1.0)	575	85 (14.8)	1,151	91 (7.9)
6	599	1 (0.2)	541	47 (8.7)	1,140	48 (4.2)
Total	3,908	36 (0.9)	3,587	401 (11.2)	7,495	437 (5.8)

^{a)}kindergarten.

Table 4. Comparison of therapeutic efficacies between 1% lindane shampoo alone (group 1) and a combination of 1% lindane shampoo and oral administration of trimethoprim/sulfamethoxazole (group 2) at 2 weeks and at 4 weeks

	No. positive at 2 weeks (%)	No. positive at 4 weeks (%)	Total No. children
Group 1	16 (23.2)	6 (8.7)	69
Group 2	6 (13.3)	1 (2.2)	45

Table 5. Status of head lice infestation among children living with both parents and children not living with both parents

	No. examined	No. positive (%)
Children living with both parents	3,789	233 (6.1)
Children not living with both parents	172	25 (14.5)
Total	3,961	258 (6.5)

primary treatment at the 2-week follow-up ($P < 0.05$).

The prevalences of HLI in children living with both parents and those not living with both parents were 6.1% (233/3789) and 14.5% (25/172), respectively (Table 5). Other factors such as parents' academic qualifications and jobs had no statistically significant influence on the prevalence of HLI ($P > 0.1$) (data not shown).

DISCUSSION

The present study shows that the overall prevalence (5.8%) of HLI in Korea has decreased. In Korea, it was previously reported that the prevalence of pediculosis in urban areas was 5.0% to 40.5% and that in rural areas was 12.8% to 43.6% over the past 10 years (Ree et al., 1992; Huh et al., 1993; Hong et al., 1995; Ha et al., 2000). Improved hygiene, decreased birth rate which reduces the frequency of close contact in the classroom and home, increased economic status, and increased parental concern about children might have resulted in this decrease of HLI. Urban areas had lower prevalence of HLI than rural areas, which is attributed to better hygiene, less time to closely contact friends after school, and frequent change of hairstyles. Girls had a higher infestation rate than boys. Difference in behavior patterns between boys and girls might have affect transmission rates and susceptibility to HLI: These include frequency of personal grooming, close contact with friends and family, frequency of hairstyle changes, shared sleeping environment, shared hair accessories, and hair length.

The infestation rate of kindergarten children was higher than that of any other grade. Kindergarten children are organized into small learning groups and often clustered around a table, where cooperation and sharing are encouraged. This might increase the transmission chance, possibly explaining the higher prevalence among kindergarten children.

The prevalence of HLI among children living with both parents was lower than that among children not

living with both parents ($P < 0.001$), which suggested that childcare was an important factor in the prevalence of HLI.

Of 114 children who received the treatment for HLI, 6 (5.3%) had side effect due to the treatment, one had a mild scalp irritation after application of 1% lindane shampoo, and 5 children had transient abdominal discomfort, diarrhea and/or dizziness due to oral trimethoprim/sulfamethoxazole. However, no major side effect was observed such as CNS toxicity, Stevens-Johnson syndrome, or aplastic anemia.

Pediculus capitis has an incubation period of 6 to 10 days to hatch out, and treatment with closely spaced applications of pediculicides appears to kill adult lice, but does not deal with the possibility of subsequent hatching (Mathias et al., 1984). The 1% lindane shampoo has incomplete ovicidal activity (Brandenburg et al., 1986; Meinking et al., 1986), therefore, we expected to observe a synergistic effect by adding oral trimethoprim/sulfamethoxazole. However, no such statistically significant synergistic effect was observed. In the both groups, the secondary treatment 2 weeks after the primary treatment had marked effect on HLI eradication and although these two applications are expensive, they may significantly reduce the recurrence of HLI.

Educational campaigns by community nurses, public health doctors, and teachers are expected to be helpful for head lice control. It is essential that school authorities and public health centers cooperate to successfully control HLI in primary schools.

REFERENCES

- Araujo A, Ferreira LF, Guidon N, et al. (2000) Ten thousand years of head lice infection. *Parasitol Today* **16**: 269. 10.
- Bartels CL, Peterson KE, Taylor KL. (2001) Head lice resistance: itching that just won't stop. *Ann Pharmacother* **35**: 109-112.
- Bailey AM, Prociv P. (2000) Persistent head lice following multiple treatments: evidence for insecticide resistance in *Pediculus humanus capitis*. *Australas J Dermatol* **41**: 250-254.
- Bartels CL, Peterson KE, Taylor KL. (2001) Head lice resistance: itching that just won't stop. *Ann Pharmacother* **35**: 109-112.
- Brandenburg K, Deinard AS, DiNapoli J, et al. (1986) 1% permethrin cream rinse vs 1% lindane shampoo in treating pediculosis capitis. *Am J Dis Child* **140**: 894-896.
- Burns DA. (1987) Action of cotrimoxazole on head lice. *Br J Dermatol* **117**: 399-400.
- Chouela E, Abeldano A, Cirigliano M, et al. (1999) Head louse infestations: epidemiologic survey and treatment evaluation in Argentinian schoolchildren. *Int J Dermatol* **36**: 819-825.
- Dodd C. (2001) Treatment of head lice. *BMJ* **323**: 1084.
- Elston DM. (1999) What's eating you? *Pediculus humanus* (Head louse and body louse). *Cutis* **63**: 259-264.
- Gratz NG. Human Lice, their Prevalence, Control and Resistance to Insecticides, a Review, 1985-1997. Geneva, Switzerland: World Health Organization, Division of Control of Tropical Diseases, WHO Pesticide Evaluation Scheme; 1997. Report WHO/CTD/WHOPES/97.8.
- Ha YC, Heo JM, Kim HJ, et al. (2000) Infestation status of head louse and treatment with lindane shampoo in children of primary school and kindergarten in Chinju-shi, Kyongsangnam-do, Korea. *Korean J Parasitol* **38**: 41-43.
- Hipolito RB, Mallorca FG, Zuniga-Macaraig ZO, et al. (2001) Head lice infestation: single drug versus combination therapy with one percent permethrin and trimethoprim/sulfamethoxazole. *Pediatrics* **107**: E30.
- Hong HK, Kim CM, Lee JS, et al. (1995) Infestation rate of head lice in primary school children in Incheon, Korea. *Korean J Parasitol* **33**: 243-244.
- Huh S, Pai KS, Lee SJ, et al. (1993) Prevalence of head louse infestation in primary school children in Kangwon-do, Korea. *Korean J Parasitol* **31**: 67-69.
- Kuwahara RT, Rasberry RD. (2001) Inexpensive head lice comb. *Pediatr Dermatol* **18**: 85.
- Mathias RG, Huggins DR, Leroux SJ, Proctor EM. (1984) Comparative trial of treatment with Prioderm lotion and Kwellada shampoo in children with head lice. *Can Med Assoc J* **130**: 407-409.
- Meinking TL, Entzel P, Villar ME, et al. (2001) Comparative efficacy of treatments for pediculosis capitis infestations. *Arch Dermatol* **137**: 287-292.
- Meinking TL, Taplin D, Kalter DC, Eberle MW. (1986) Comparative efficacy of treatments for pediculosis capitis infestations. *Arch Dermatol* **122**: 267-271.
- Mumcuoglu KY. (1999) Prevention and treatment of head lice in children. *Paediatr Drugs* **1**: 211-218.
- Ree HI, Yong TS, Shin HJ, et al. (1992) Mass treatment of head louse infestation with Sumithrin powder in primary schools in Korea. *Korean J Parasitol* **30**: 349-354.
- Schachner LA. (1997) Treatment resistant head lice: alternative therapeutic approaches. *Pediatr Dermatol* **14**: 409-410.
- Speare R. (2000) Hot water kills head lice experimentally placed on brushes. *Int J Dermatol* **39**: 954-956.