Effects of anti-allergic drugs on intestinal mastocytosis and worm expulsion of rats infected with *Neodiplostomum seoulense*

Eun-Hee SHIN¹⁾, Tae-Heung KIM²⁾, Sung-Jong HONG³⁾, Jae-Hwan PARK¹⁾,
Sang-Mee GUK¹⁾ and Jong-Yil CHAI^{1)*}

¹⁾Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799,

²⁾Department of Dermatology, College of Medicine, Gyeongsang National University, Jinju 660-280,

³⁾Department of Parasitology, Chung-Ang University College of Medicine, Seoul 156-756, Korea

Abstract: The effects of anti-allergic drugs on intestinal mastocytosis and the expulsion of *Neodiplostomum seoulense* were observed in Sprague-Dawley rats, after oral infection with 500 metacercariae. The drugs used were hydroxyzine (a histamine receptor H₁ blocker), cimetidine (a H₂ blocker), cyclosporin-A (a helper T-cell suppressant), and prednisolone (a T- and B-cell suppressant). Infected, but untreated controls, and uninfected controls, were prepared. Worm recovery rate and intestinal mastocytosis were measured on weeks 1, 2, 3, 5, and 7 post-infection. Compared with the infected controls, worm expulsion was significantly ($P < 0.05$) delayed in hydroxyzine- and cimetidine-treated rats, despite mastocytosis being equally marked in the duodenum of all three groups. In the cyclosporin-A- and prednisolone-treated groups, mastocytosis was suppressed, but worm expulsion was only slightly delayed, without statistical significance. Our results suggest that binding of histamine to its receptors on intestinal smooth muscles is more important in terms of the expulsion of *N. seoulense* from rats than the levels of histamine alone, or mastocytosis.

Key words: *Neodiplostomum seoulense*, expulsion of worm, mast cell (mucosal), goblet cell, hydroxyzine, cimetidine, cyclosporin, prednisolone

INTRODUCTION

Neodiplostomum seoulense (Digenea: Neodiplostomidae) is an intestinal fluke of humans and rodents in the Republic of Korea (Seo, 1990). The grass snake, *Rhabdophis tigrina*, a paratenic host, is an important

source of human infection (Hong et al., 1982; Seo, 1990). The adult flukes parasitize the duodenum of rodents and, if worm burdens are high, they extend to the jejunum and ileum (Hong, 1982). In experimentally infected rodents, villous atrophy (including a decreased villus-crypt ratio, blunting and fusion of the villi), and crypt hyperplasia (with stromal inflammations) were found to be the major histopathological changes in the small intestine (Lee et al., 1985). Infection with this trematode can be lethal to mice (Huh et al., 1988; Kook et al., 1998; Chai et al.,

• Received 2 April 2003, accepted after revision 12 May 2003.

• This study was supported by a grant from the Seoul National University College of Medicine Research Fund (2000).

*Corresponding author (e-mail: cji@plaza.snu.ac.kr)

2000), and infected people may suffer from symptoms, which include diarrhea, abdominal pain, and fever (Seo, 1990).

In rats infected with *N. seoulense*, the worm recovery rate (WRR) decreases from four weeks post-infection (PI) (Hong et al., 1983). Mucosal mast cells (MMC)s, which increased markedly in the duodenum at around week 3 PI, were proposed to be an important effector of this spontaneous worm expulsion (Kho et al., 1990). In mice, however, MMCs were found to play only a minor role in the expulsion of *N. seoulense* (Chai et al., 1998). Therefore, the mechanisms of *N. seoulense* expulsion may be different in rats and mice, and the role of MMCs in rats requires verification.

In this study, we investigated the effects of four types of anti-allergic drugs on mastocytosis and on the worm expulsion of *N. seoulense* from the rat small intestines. These drugs examined were, prednisolone (a T- and B-cell suppressant), cyclosporin-A (a helper T-cell suppressant), hydroxyzine (a histamine receptor H1 blocker), and cimetidine (a H₂ blocker).

MATERIALS AND METHODS

Collection of the metacercariae and the experimental infection of rats

The metacercariae of *N. seoulense* were collected from 50 grass snakes (*Rhabdophis tigrina*), which were purchased from a local snake collector in Jinju, Gyeongsangnam-do, using an artificial digestion technique. A total of 185 Sprague-Dawley rats 3-week-old were purchased from the Laboratory Animal Center, at Seoul National University. The animals were treated twice with mebendazole (10 mg/kg) and praziquantel (25 mg/kg), and confirmed to be free of helminth eggs by fecal examinations before being used in the experiment.

The rats were divided into 4 drug-treated and 2 control groups; i.e., hydroxyzine-, cimetidine-, cyclosporin-A- and prednisolone-treated, and uninfected and infected but untreated controls. Each group was composed of 30 rats, but 5 rats were added to the prednisolone-treated group because prednisolone treatment was expected to affect rat survival during the experiment. Each rat in the

infected groups was fed 500 metacercariae of *N. seoulense* orally using a gavage needle. The uninfected controls were given only water through the same needle type.

Drug administration

Hydroxyzine, at a daily dose of 1 mg/kg, and cimetidine, at a daily dose of 20 mg/kg, were diluted in water and administered orally from the day of infection until sacrifice. Cyclosporin-A, which is insoluble in water, was diluted initially in a small amount of absolute ethanol; this solution was then diluted with water just prior to use, and administered orally at a daily dose of 7.5 mg/kg, until sacrifice. Prednisolone (10 mg/kg) was injected intramuscularly every other day into the inner thighs of the rats, until sacrifice.

Histochemistry for MMCs and worm recovery

Rats were sacrificed on weeks 1, 2, 3, 5 and 7 PI by cervical dislocation. Abdomens were incised, and intestinal segments, about 1.5 cm in length, were taken from the middle portion of the duodenum, jejunum, and ileum. The segments were washed with saline 2-3 times, and fixed in Carnoy's fixative for 6 to 12 hr. The fixed tissues were then embedded in paraffin, and sectioned at 4 μ m. The tissue sections obtained were stained with 1% alcian blue (Janssen Pharmaceutica, Belgium) (pH = 0.3), and counter-stained with safranin O (pH = 1.0) (Merck, USA), as described by Strobel et al. (1981). The number of MMCs in a defined region of 10 villus-crypt units (VCU) was counted for each stained sample, and data are expressed as means and SD of the number of MMCs per VCU.

Adult flukes of *N. seoulense* were recovered from the segments of the small intestines, using Baermann's apparatus (Chai et al., 1998). Worms were collected from the bottom of a test tube, and counted under a stereomicroscope to determine the WRRs. The WRR was calculated for each rat by dividing the number of recovered worms by the number of metacercariae given.

Statistical test

Statistical significance between the groups was

determined by using the paired *t*-test. Values of $P < 0.05$ were considered significant.

RESULTS

Worm recovery

In the infected controls, the WRRs did not change significantly over weeks 1-3 PI, ranging from $33.6 \pm 6.2\%$ to $38.4 \pm 5.5\%$, but they decreased significantly ($P < 0.05$) to $23.6 \pm 5.2\%$ and $19.9 \pm 8.0\%$, at weeks 5 and 7 PI, respectively (Fig. 1). The four drug treatment groups had similar chronological patterns. However, throughout the observation period, the WRRs were significantly ($P < 0.05$) higher in hydroxyzine- and in cimetidine-treated rats, than in the infected controls (P

< 0.05 except at week 2 PI; Fig. 1). The WRRs of the cyclosporin-A- and prednisolone-treated groups during weeks 1 and 7 PI were slightly higher than in the infected controls; however, this difference was only statistically significant ($P < 0.05$) at week 5 PI for the cyclosporin-A-treated animals (Fig. 1).

Changes in the number of MMCs in the rat intestine

In the duodenum of uninfected control rats, the number of MMCs per VCU ranged from 11.5 ± 3.3 (at week 2 PI) to 15.6 ± 4.0 (at week 5 PI), and showed little variation (Fig. 2). In the jejunum and ileum, MMC numbers were almost identical to those of the duodenum of the uninfected controls. In *N. seoulense*-

Fig. 1. Worm recovery of *Neodiplostomum seoulense* in rats treated with anti-allergic drugs versus infected controls. Each rat was infected with 500 metacercariae, and sacrificed at weeks 1-7 post-infection (PI). The worm recovery rates (WRRs) were significantly higher in hydroxyzine-, and cimetidine-treated rats compared with the controls ($P < 0.05$; except at week 2 PI). The WRRs in cyclosporin-A- and prednisolone-treated groups were slightly higher than in the infected controls; however, statistical significance was observed only at week 5 PI between the cyclosporin-A-treated animals and the infected control group ($P < 0.05$).

Fig. 2. Mucosal mast cell (MMC) responses in the small intestines of rats, after an infection with *Neodiplostomum seoulense* and treatment with anti-allergic drugs. Data are expressed as the numbers of MMCs per villus-crypt unit (VCU*). In *N. seoulense*-infected rats, the number of duodenal MMCs peaked at week 3 PI, and then decreased gradually. Jejunal and ileal MMCs increased only slightly through the infection period. In hydroxyzine- and cimetidine-treated rats, mastocytosis levels in the duodenum were similar to that of the infected controls. By comparison, in rats treated with cyclosporin-A or prednisolone, mastocytosis was lower than in the infected controls. In the jejunum and ileum, mastocytosis was unremarkable in all drug-treated animals.

infected rats, however, the numbers of duodenal MMCs per VCU were greater, 18.9 ± 5.5 at week 1 PI, and this increased rapidly and remarkably to 41.6 ± 11.4 and 68.9 ± 21.7 at weeks 2 and 3 PI, respectively (Fig. 2). After peaking at week 3 PI, the number of duodenal MMCs decreased gradually to 53.7 ± 10.9 and 42.3 ± 10.4 , at weeks 5 and 7 PI, respectively (Fig. 2). On the other hand, in both the jejunum and ileum of rats infected with *N. seoulense*, MMC numbers increased only slightly, but significantly ($P < 0.05$), throughout the infection period (Fig. 2).

In the duodenums of hydroxyzine- and cimetidine-treated rats, the chronological patterns of mastocytosis were similar to those of the infected controls (Fig. 2). MMC numbers were slightly greater in the drug-treated groups than in the untreated controls; however, this was not statistically significant ($P > 0.05$). The chronological patterns of MMCs in the jejunum and ileum were similar for hydroxyzine- and cimetidine-treated animals and infected controls (Fig. 2).

Different chronological patterns of MMCs were observed in the cyclosporin-A- and prednisolone-treated groups (Fig. 2). In the duodenum of the cyclosporin-A-treated group, the number of MMCs increased gradually and peaked at week 5 PI, and then decreased from week 7 PI (Fig. 2). However, compared with the infected controls, the number of MMCs during weeks 1-3 PI was significantly ($P < 0.05$) lower in the cyclosporin-A-treated group (Fig. 2). In the duodenum of prednisolone-treated rats, the number of MMCs increased gradually and peaked at week 3 PI, and then decreased at weeks 5 and 7 PI (Fig. 2). However, the peak level of mastocytosis in the prednisolone-treated group was only a half of that of the infected controls and of the hydroxyzine- and cimetidine-treated groups; MMC numbers through weeks 1-7 PI were significantly ($P < 0.05$) lower than in the infected controls (Fig. 2). In the jejunum and ileum of cyclosporin-A- and prednisolone-treated rats, mastocytosis was unremarkable through weeks 1-7 PI (Fig. 2).

DISCUSSION

Host-parasite relationships related to the expulsion

mechanisms of helminths from the rodent intestine have been studied using a variety of nematode and trematode models (Chai et al., 1993; Fujino et al., 1998; Koyama and Ito, 2000). Reviews of these studies have led to the conclusion that intestinal MMCs and goblet cells (GCs) are the two major effector cell types (Else and Finkelman, 1998; Onah and Nawa, 2000).

The relative importances of the effector cell are, however, different and depend on parasite species. For example, MMCs, but not GCs, play a vital role in the expulsion of *Strongyloides ratti*, whereas GCs, but not MMCs, play an essential role in the expulsion of *N. brasiliensis* and *Echinostoma trivolvis* from rodents (Nawa et al., 1994; Fujino et al., 1998; Onah and Nawa, 2000). It has also been shown that MMCs are not essential for the protection of mice against infection with the nematode, *Trichuris muris* (Koyama and Ito, 2000), or for protecting rats against infection with the cestode, *Hymenolepis diminuta* (Starke and Oaks, 2001). Moreover, mucosal mastocytosis was suppressed by unknown reasons in a nematode infection with *Nematospiroides dubius* in mice (Dehlawi et al., 1987).

In the case of *N. seoulense* infection, spontaneous worm expulsion, in association with intestinal mastocytosis (notably in the duodenum), has been observed in rats (Kho et al., 1990), and in mice (Chai et al., 1998). The kinetics and extent of mastocytosis, however, were different between the two animal hosts. In rats, the number of MMCs peaked at week 3 PI, and then decreased gradually up to week 7 PI (Kho et al., 1990); but in mice, it peaked at week 1 PI, and quickly decreased to the normal level at week 2 PI (Chai et al., 1998). The peak MMC number in rats was about 80 per VCU (Kho et al., 1990), 4~5-fold higher than the peak value in BALB/c mice (Chai et al., 1998). Therefore, the role of MMCs in the expulsion of *N. seoulense* may be different for mice and rats. Consequently, in mice, neither MMCs nor GCs were shown to play a major role in worm expulsion by BALB/c and C3H mice (Chai et al., 1998), whereas the role of MMCs remains to be elucidated in rats (Kho et al., 1990). In this study, rats were used as the experimental host, and worm expulsion with intestinal mastocytosis was observed; nevertheless, the importance of MMCs in worm expulsion is difficult to determine.

In relation to the actions of MMCs, the enzymes produced by these cells, which include amines like histamine and serotonin, as well as proteases and peroxidases, are candidate biochemical mediators responsible for intestinal nematode expulsion (Woodbury et al., 1984). In particular, histamine is known to enhance intestinal smooth muscle contraction, which could facilitate worm expulsion, by binding to histamine receptors (Mary et al., 2000). The present study supports for the importance of histamine in worm expulsion, because treatments with hydroxyzine or cimetidine, antagonists of the histamine receptors, significantly delayed worm expulsion from rats. Hydroxyzine is known to block the H_1 receptor (Mary et al., 2000), and thus inhibits the action of histamine and worm expulsion, regardless of the level of mastocytosis. Similarly, mepyramine, another H_1 antagonist, inhibits the expulsion of *Trichostrongylus colubriformis* from guinea pigs (Rothwell et al., 1978). Cimetidine has the ability to inhibit histamine-induced smooth muscle contraction (Scott and Maric, 1993), by blocking the histamine receptor H_2 , and thus can inhibit worm expulsion regardless of the level of mastocytosis.

In this study, however, the effects of cyclosporin-A and prednisolone on the expulsion of *N. seoulense* are unclear. Cyclosporin-A has a reversible immunosuppressive effect on helper T-cells (McLauchlan et al., 1999), and prednisolone reduces mast cell numbers and mast cell protease levels dose-dependently (King et al., 1985; Goldsmith et al., 1990); both inhibit mastocytosis in the host intestinal mucosa. Unexpectedly, however, the WRRs in cyclosporin-A- and prednisolone-treated rats were only slightly higher than that in infected controls; this difference was not statistically significant. Mastocytosis was markedly depressed in both drug-treated groups. This discrepancy strongly suggests the possible existence of effectors other than MMCs and histamine for the expulsion of *N. seoulense* from rats. Intestinal GC is a candidate effector, as has been reported in other trematode species, such as *E. trivolvis* (Weinstein and Fried, 1991) and *E. caproni* (Fujino et al., 1996). In mice, GCs have been shown to play only minor roles in BALB/c and C3H strains in the expulsion of *N. seoulense* (Chai et al., 1998),

however, their role in rats needs to be clarified.

In this connection, it is of considerable interest to note that histamine can stimulate airway GC secretion in guinea pigs, and that cimetidine, an antagonist of histamine H_2 receptor, can abolish this mucus response (Takeyama et al., 1996; Tamaoki et al., 1997). We speculate, therefore, that cimetidine, and possibly hydroxyzine affect the histamine receptors on GCs in the intestinal tracts of rats, and suppress the action of GCs. Information upon the effects of histamine on intestinal GCs, however, is lacking. Studies on the effects of antihistamines on the intestinal GC responses of rats infected with *N. seoulense* are required.

REFERENCES

- Chai JY, Kim TH, Kho WG, et al. (1993) Mucosal mast cell responses to experimental *Metagonimus yokogawai* infection in rats. *Korean J Parasitol* **31**: 129-134.
- Chai JY, Kim TK, Cho WH, et al. (1998) Intestinal mastocytosis and goblet cell hyperplasia in BALB/c and C3H mice infected with *Neodiplostomum seoulense*. *Korean J Parasitol* **36**: 109-119.
- Chai JY, Shin EH, Han ET, Guk SM, Choi MH, Lee SH (2000) Genetic difference in susceptibility and fatality of three strains of mice experimentally infected with *Neodiplostomum seoulense*. *J Parasitol* **86**: 1140-1144.
- Dehlawi MS, Wakelin D, Behnke JM (1987) Suppression of mucosal mastocytosis by infection with the intestinal nematode *Nematospiroides dubius*. *Parasite Immunol* **9**: 187-194.
- Else KJ, Finkelman FD (1998) Intestinal nematode parasites, cytokines and effector mechanisms. *Int J Parasitol* **28**: 1145-1158.
- Fujino T, Fried B, Ichikawa H, Tada I (1996) Rapid expulsion of the intestinal trematodes *Echinostoma trivolvis* and *E. caproni* from C3H mice by trapping with increased goblet cell mucins. *Int J Parasitol* **26**: 319-324.
- Fujino T, Ichikawa H, Fukuda K, Fried B (1998) The expulsion of *Echinostoma trivolvis* caused by goblet cell hyperplasia in severe combined immunodeficient (SCID) mice. *Parasite* **5**: 219-222.
- Goldsmith P, McGarity B, Walls AF, et al. (1990) Corticosteroid treatment reduces mast cell numbers in inflammatory bowel disease. *Dig Dis Sci* **35**: 1409-1413.
- Hong SJ, Lee SH, Seo BS, et al. (1983) Studies on intestinal trematodes in Korea IX. Recovery rate and development of *Fibricola seoulensis* in experimental animals. *Korean J*

- Parasitol* **21**: 224-233.
- Hong ST, Hong SJ, Lee SH, et al. (1982) Studies on intestinal trematodes in Korea VI. On the metacercaria and the second intermediate host of *Fibricola seoulensis*. *Korean J Parasitol* **20**: 101-111.
- Hong ST (1982) Studies on intestinal trematodes in Korea VII. Growth, development and recovery of *Fibricola seoulensis* from experimentally infected rats and mice. *Korean J Parasitol* **20**: 112-121.
- Huh S, Chai JY, Hong ST, et al. (1988) Clinical and histopathologic findings in mice heavily infected with *Fibricola seoulensis*. *Korean J Parasitol* **26**: 45-53.
- King SJ, Miller HR, Newlands GF, Woodbury RG (1985) Depletion of mucosal mast cell protease by corticosteroids: effect on intestinal anaphylaxis in the rat. *Proc Natl Acad Sci USA* **82**: 1214-1218.
- Kho WG, Chai JY, Chun CH, et al. (1990) Mucosal mast cell response to experimental *Fibricola seoulensis* infection in rats. *Seoul J Med* **31**: 191-199.
- Kook J, Nawa Y, Lee SH, Chai JY (1998) Pathogenicity and lethality of a minute intestinal fluke, *Neodiplostomum seoulense*, to various strains of mice. *J Parasitol* **84**: 1178-1183.
- Koyama K, Ito Y (2000) Mucosal mast cell responses are not required for protection against infection with the murine nematode parasite *Trichuris muris*. *Parasite Immunol* **22**: 13-20.
- Lee SH, Yoo BH, Hong ST, et al. (1985) A Histopathological study on the intestine of mice and rats infected by *Fibricola seoulensis*. *Korean J Parasitol* **23**: 58-72.
- Mary JM, Richard AH, Pamela CC (2000) Antihistamines; Lippincott's illustrated reviews. In Pharmacology. pp 420-421, Lippincott Williams & Wilkins, USA.
- McLauchlan PE, Roberts HC, Loxton NJ, Wastling JM, Newlands GF, Chappell LH (1999) Mucosal mast cell responses and release of mast cell protease-I in infections of mice with *Hymenolepis diminuta* and *H. microstoma*: modulation by cyclosporin A. *Parasite Immunol* **21**: 151-61.
- Nawa Y, Ishikawa N, Tsuchiya K, et al. (1994) Selective effector mechanisms for the expulsion of intestinal helminths. *Parasite Immunol* **16**: 333-338.
- Onah DN, Nawa Y (2000) Mucosal immunity against parasitic gastrointestinal nematodes. *Korean J Parasitol* **38**: 209-236.
- Rothwell TL, Love RJ, Evans DP (1978) Studies on the role of histamine and 5-hydroxytryptamine in immunity against the nematode *Trichostrongylus colubriformis*. IV. Inhibition of the expulsion of worms transplanted into the duodenum of immune guinea pigs. *Int Arch Allergy Appl Immunol* **56**: 457-462.
- Scott RB, Maric M (1993) Mediation of anaphylaxis-induced jejunal circular smooth muscle contraction in rats. *Dig Dis Sci* **38**: 396-402.
- Seo BS (1990) *Fibricola seoulensis* Seo, Rim and Lee, 1964 (Trematoda) and fibricoliasis in man. *Seoul J Med* **31**: 61-96.
- Starke WA, Oaks JA (2001) Ileal mucosal mast cell, eosinophil, and goblet cell populations during *Hymenolepis diminuta* infection of the rat. *J Parasitol* **87**: 1222-1225.
- Strobel S, Miller HR, Ferguson A (1981) Human intestinal mucosal mast cells: evaluation of fixation and staining techniques. *J Clin Pathol* **34**: 851-858.
- Takeyama K, Tamaoki J, Nakata J, Konno K (1996) Effect of oxitropium bromide on histamine-induced airway goblet cell secretion. *Am J Respir Crit Care Med* **154**: 231-236.
- Tamaoki J, Nakata J, Takeyama K, Chiyotani A, Konno K (1997) Histamine H₂ receptor-mediated airway goblet cell secretion and its modulation by histamine-degrading enzymes. *J Allergy Clin Immunol* **99**: 233-238.
- Weinstein MS, Fried B (1991) The expulsion of *Echinostoma trivolvis* and retention of *Echinostoma caproni* in the ICR mouse: pathological effects. *Int J Parasitol* **21**: 255-257.
- Woodbury RG, Miller HRP, Huntly JF, et al. (1984) Mucosal mast cells are functionally active during spontaneous expulsion of intestinal nematode infections in rat. *Nature* **312**: 450-452.