Published in final edited form as:

Biochem Biophys Res Commun. 2007 November 9; 363(1): 194–196. doi:10.1016/j.bbrc.2007.08.131.

Two-Pore K⁺ channels, NO and Metabolic Inhibition

Zhongju Lu^{1,#}, Junyuan Gao¹, Joan Zuckerman¹, Richard T. Mathias¹, Glenn Gaudette², Irvin Krukenkamp^{1,3}, and Ira S. Cohen¹

¹Department of Physiology & Biophysics, Institute for Molecular Cardiology, State University of New York at Stony Brook, NY 11794-8661, USA

²Department of Biomedical Engineering, Worcester Polytechnic Institute, Worcester, MA 01609, USA

³Division of Cardiothoracic Surgery, State University of New York at Stony Brook, NY 11794, USA

Abstract

Ischemic preconditioning is a potent endogenous mechanism protecting many organs from the devastating effects of prolonged ischemia. In the heart, NO is one mediator of this myoprotective response thought to involve activation of the K_{ATP} channel. Ischemic preconditioning is known to be induced by metabolic inhibition using sodium cyanide (NaCN) in single cardiomyocytes. In the present study we show for the first time that the end effector channel activated by NaCN has been incorrectly identified. The channel activated is not K_{ATP} but instead belongs to the relatively new family of two-pore domain potassium channels (K2P). Further when activated by metabolic ischemia, the amplitude of K2P current is directly modulated by activators and inhibitors of the NO pathway.

Keywords

Ischemic Preconditioning; Ion Channel; NO

Since the initial report by Murry et al [1], ischemic preconditioning has been recognized as a potent endogenous mechanism of myoprotection. In an attempt to investigate the mechanism of this protection, Liu et al. [2] developed a single cell model of metabolic ischemia. Upon exposure to NaCN in glucose free solution an outward current is induced (Figure 1A) in isolated ventricular myocytes (*Materials and Methods*). The amplitude of this current increases and the time to appearance of this current shortens when the preparation is first exposed to preconditioning agents [3]. In the initial study and subsequently [2;4], the current activated by metabolic ischemia was identified as I_{K_{ATP}} because it declined when glibenclamide, a blocker of the K_{ATP} channel, was applied at the peak of the response. However, even in the absence of this channel blocker, the current declines on its own (Figure 1A). Further when glibenclamide is included throughout the exposure to NaCN there is no effect on the amplitude of the response (inset of Figure 2A). This result led us to question whether the channel activated in preconditioning had been correctly identified. In 1996 a new class of ion channels was identified [5]. These channels were K⁺ specific, also insensitive to classic K⁺ channel blockers like Ba²⁺ and Cs⁺ but instead were blocked by either Zn²⁺ or Quinidine [6]. Figure 2A

Corresponding author: Zhongju Lu, MD, PhD, Department of Physiology and Biophysics, State University of New York at Stony Brook, Stony Brook, NY 11794-8661, USA. Telephone: (631)-444 3613, Fax: (631)-444 3432, Email: E-mail: zlu@notes.cc.sunysb.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

demonstrates that the current activated by metabolic ischemia is not blocked by glibenclamide but is blocked by Zn^{2+} . Since glibenclamide is a poor blocker of K_{ATP} channels in acidotic conditions we asked whether the current activated by metabolic ischemia might still be K_{ATP} and that this channel might also be blocked by Zn^{2+} or Quinidine. Figures 2B and 2D show our results. IK_{ATP} was activated by pinacidil together with a low concentration of intracellular ATP (0.1mM). This current is identified as IK_{ATP} by its sensitivity to glibenclamide [7]. It is however unaffected by either Zn^{2+} or quinidine. There are at least 15 members in the K2P family, all of which are K^+ specific [6;8;9]. The current activated by NaCN is K^+ specific (Figure 1C-E). A number of these channels such as TALK-1 and TALK-2 are directly activated by nitric oxide (NO) [10]. Given the importance of NO to preconditioning, we examined the effects of an activator (L-Arginine, 400 μM) and an inhibitor (L-NAME, 200 μM) of the NO pathway on the NaCN induced current. The results are provided in Figure 2C. Activating the NO pathway increases the NaCN induced current while inhibiting the pathway reduces the current. These results (summarized in Figure 2D) suggest that there is constitutive NO production in guinea pig ventricular myocytes and that the NaCN induced current is a K2P channel that is modulated by NO.

In summary, the current initiated by metabolic ischemia is not mediated by the K_{ATP} channel. Instead, an NO sensitive member of the K2P family is implicated. It is well known that the K2P channels help to regulate cell volume [11]. It is possible that their role in volume regulation plays a key role in the protection they afford from prolonged ischemia where cell swelling can induce apoptosis [9]. Finally, with the identification of a novel sarcolemmal channel involved in preconditioning, there is a new therapeutic target to investigate. Other activators of the K2P channels should have the potential to induce preconditioning.

Materials and Methods

Single ventricular cells were enzymatically isolated from adult male guinea pig hearts as described in Gao et al [12]. An Axopatch 1D amplifier (Axon Instruments, Inc) and the classical whole-cell patch clamp technique were employed to observe cell membrane current [13]. The pipette solution contained (in mM): K-Aspartic Acid 125; KCl 15; KOH 10; MgCl_2 1; HEPES 10; EGTA 11; Mg-ATP 1 (pH 7.2). The external Tyrode solution contained (in mM): NaCl 137.7; NaOH 2.3; KCl 4; MgCl_2 1; HEPES 5; CaCl_2 1; CdCl_2 1; Glucose 10 (pH 7.4). Sodium cyanide (NaCN) was dissolved in Tyrode solution without glucose and prepared at the target concentration to simulate metabolic ischemia [3]. All patch clamp data were digitized by the data acquisition program pClamp8 (Axon Instruments, Inc) for later analysis. Cell capacitance was obtained for each cell and currents were normalized to cell capacitances. NaCN, glibenclamide, collagenase (type II) and other reagents were obtained from Sigma Chemical (St. Louis, MO).

Acknowledgments

This work was supported by grant HL70161 to IBK, grants HL67101 and HL28958 to ISC and grant HL85221 to RTM

References

1. Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation* 1986;74:1124–1136. [PubMed: 3769170]
2. Liu Y, Gao WD, O'Rourke B, Marban E. Priming effect of adenosine on K_{ATP} currents in intact ventricular myocytes: implications for preconditioning. *Am J Physiol* 1997;273:H1637–43. [PubMed: 9362225]

3. Irie H, Gao J, Gaudette GR, Cohen IS, Mathias RT, Saltman AE, Krukenkamp IB. Both metabolic inhibition and mitochondrial K_{ATP} channel opening are myoprotective and initiate a compensatory sarcolemmal outward membrane current. *Circulation* 2003;108:II341–II347. [PubMed: 12970257]
4. Liu Y, Gao WD, O'Rourke B, Marban E. Synergistic modulation of ATP-sensitive K^+ currents by protein kinase C and adenosine. implications for ischemic preconditioning. *Circ Res* 1996;78:443–454. [PubMed: 8593703]
5. Fink M, Duprat F, Lesage F, Reyes R, Romey G, Heurteaux C, Lazdunski M. Cloning, functional expression and brain localization of a novel unconventional outward rectifier K^+ channel. *EMBO J* 1996;15:6854–6862. [PubMed: 9003761]
6. Kim D. Physiology and pharmacology of two-pore domain potassium channels. *Curr Pharm Des* 2005;11:2717–2736. [PubMed: 16101451]
7. Song Y, Belardinelli L. Electrophysiological and functional effects of adenosine on ventricular myocytes of various mammalian species. *Am J Physiol* 1996;271:C1233–43. [PubMed: 8897829]
8. Lesage F, Lazdunski M. Molecular and functional properties of two-pore-domain potassium channels. *Am J Physiol Renal Physiol* 2000;279:F793–F801. [PubMed: 11053038]
9. Patel AJ, Lazdunski M. The 2P-domain K^+ channels: role in apoptosis and tumorigenesis. *Pflugers Arch* 2004;448:261–273. [PubMed: 15133669]
10. Duprat F, Girard C, Jarretou G, Lazdunski M. Pancreatic two P domain K^+ channels TALK-1 and TALK-2 are activated by nitric oxide and reactive oxygen species. *J Physiol* 2005;562:235–244. [PubMed: 15513946]
11. Trimarchi JR, Liu L, Smith PJ, Keefe DL. Apoptosis recruits two-pore domain potassium channels used for homeostatic volume regulation. *Am J Physiol Cell Physiol* 2002;282:C588–C594. [PubMed: 11832344]
12. Gao J, Mathias RT, Cohen IS, Baldo GJ. Isoprenaline, Ca^{2+} and the Na^+-K^+ pump in guinea-pig ventricular myocytes. *J Physiol (Lond)* 1992;449:689–704. [PubMed: 1326051]
13. Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch* 1981;391:85–100. [PubMed: 6270629]

Figure 1. Characterization of the K^+ selective outward induced by NaCN from isolated guinea pig ventricular myocytes

A: Sample trace of a large outward current induced by extracellular application of sodium cyanide (NaCN, 2mM). The $[K^+]_o$ was 5.4mM and the patch pipette $[K^+]$ was 150mM. Note a relatively slow onset phase and sustained phase in the NaCN induced current, and that the current can decay on its own. Similar results were observed in all of the cells ($n=18$) studied in the same conditions. The cells were held at 0 mV and experiments were performed at 22° C. **B:** The outward current induced by NaCN did not appear when both external and pipette K^+ were absent. The external K^+ was absent without substitute and the pipette K^+ was substituted with L-Aspartic Acid, and the pH was adjusted with Trizma Base. Similar results

were observed in all of the (n=6) cells studied in the same conditions. **C**: A typical recording of NaCN induced outward current measured in which voltage ramps were applied to obtain the current-voltage relationship. The $[K^+]_o$ was 3mM and the pipette $[K^+]$ was 150mM. The cells were held at 0mV and were subject to hyperpolarizing ramps from +50mV to -100mV (2s duration) with a frequency of 0.1Hz (*Upper panel*). The corresponding I/V relationship of NaCN induced current (*Middle Panel*), which is constructed by subtraction of the I/V curve measured at the base (**a**) from that measured at the peak (**b**) (*Lower Panel*). **D**. The I/V curves of the NaCN induced current in three different $[K^+]_o$ were plotted using the same protocol shown in **C**. **E**. The linear relationship between averaged reversal potentials and equilibrium potential of K^+ (E_K) calculated with Nernst equation at different $[K^+]_o$, suggested that the current induced by NaCN induced is K^+ selective. Note that our measured reversal potentials are not the same as E_K (possibly due to the large intracellular K^+ loss induced by the NaCN induced-current). The numbers in the parentheses indicate the cells studied.

Figure 2. The NaCN induced current from guinea pig ventricular myocytes is not $I_{K_{ATP}}$ nor I_{K_1}

A: The outward current induced by NaCN can be reversibly abolished by Zn²⁺ (3mM). Inset of **A**: Glibenclamide (200μM) cannot prevent the appearance of the NaCN induced current.

B: The $I_{K_{ATP}}$ activated by pinacidil (100μM) together with low intracellular ATP (0.1mM) cannot be blocked by Zn²⁺ (3mM), but is abolished by glibenclamide. **C:** NO-regulation of the NaCN induced current. Note the NaCN induced current is reduced by the NOS inhibitor, L-NAME (200μM), and the NaCN induced current is reactivated after washout of Zn²⁺. Inset of **C**: A typical current trace showing that L-arginine (400μM) can additionally activate an outward current which is sensitive to Quinidine (1mM).

D: In summary, the average results suggest that the NaCN induced current is not blocked by either the K_{ATP} channel blocker (glibenclamide) or classical K⁺ channel blockers (1mM Ba²⁺ or 1mM Cs⁺), it is sensitive to typical K2P channel blockers (both Zn²⁺ and Quinidine) and modulated by NO. In contrast, classical $I_{K_{ATP}}$ is completely blocked by glibenclamide and unaffected by typical K2P channel blockers. All the cells were held at 0 mV. $P < 0.05$ was considered statistically significant by unpaired student's T-test.