Published in final edited form as:

Org Lett. 2009 July 16; 11(14): 3128–3131. doi:10.1021/ol9010008.

Synthesis of the Tetracyclic Core of the Neomangicols Using a Late-Stage Indene Alkylation

Jessica L. Wood, Brian G. Pujanauski, and Richmond Sarpong*

Department of Chemistry, University of California, Berkeley, California 94720

Abstract

A general approach to the tetracyclic core of the neomangicol natural products via a late-stage indene alkylation reaction is presented. This strategy sets the stage for access to the neomangicol family and, in addition, provides a potential biogenetically inspired entry to the mangicol natural products.

Historically, the carbon framework of terpenoid natural products has proven to be a synthetic challenge due to the lack of functional groups on these molecules that can direct C-C bond formation.¹ As a result, synthetic chemists must design strategies for terpenoid syntheses that maximize carbon-carbon bond formation. Enolate methodology has been featured prominently in a wide range of C-C bond forming cyclization tactics (e.g., intramolecular aldol and Dieckmann reactions) en route to terpene natural products, which often necessitate the removal of the enolate carbonyl oxygen at a late stage. In the context of [5–6] bicyclic ring-containing terpene natural products, we envisioned that indenenes, which possess acidities comparable to carbonyl compounds (e.g., ketones), could serve as enolate equivalents in late-stage bond-forming events. In this communication, we present the application of this tactic to the synthesis of the tetracyclic core of the natural product neomangicol C (**2**, Figure 1).

Neomangicol C is a member of a group of rearranged sesterterpenoids isolated from a marine fungus of the genus *Fusarium*, which possess varied bioactivity.² For example, neomangicols A and B (**1a** and **1b**, Figure 1) have shown in vitro cytotoxicity against human colon carcinoma.

Furthermore, neomangicol B (**1b**) has shown potency against the Gram-positive bacterium *Bacillus subtilis* similar to that of the aminoglycoside gentamycin and may prove to be of general utility as an antibiotic.² Preliminary studies have not identified significant bioactivity for neomangicol C (**2**), and there is evidence that suggests it may be an isolation artifact arising from the net loss of HCl or HBr from **1a** or **1b**, respectively.²

To date, there have been no reports of synthetic work toward the neomangicols. However, studies by Uemura³ and Paquette⁴ have recently begun to address the synthesis of the related

E-mail: rsarpong@berkeley.edu.

Supporting Information Available. Experimental details and characterization data for all new compounds are available free of charge via the Internet at <http://pubs.acs.org>.

mangicols⁵ (e.g., mangicol A (**3**) and B (**4**), Figure 1). Mangicol A is of interest as a unique structural motif that possesses anti-inflammatory activity.

A series of feeding studies has demonstrated that the neomangicols may arise from a rearrangement of the mangicol skeleton (see **B** → **A**, Figure 1).⁵ We envisioned that our model studies on the neomangicols would offer an opportunity to test the reverse of the biosynthetic proposal by employing a semi-pinacol/Wagner-Meerwein⁶ ring contraction (see **A** → **B**, Figure 1) to construct the mangicol skeleton.

Despite their relatively small size, the neomangicols and mangicols possess several challenging features from a synthetic standpoint. For example, these natural product families possess nine and eleven stereocenters, respectively. Additionally, the vinyl halide moiety present in neomangicols A and B is highly unusual in sesterterpene natural products. Our initial studies have focused on the neomangicols and specifically the tetracyclic core of neomangicol C. This presents an opportunity to investigate the potential application of indene alkylation chemistry to the synthesis of these highly complex rearranged sesterterpenoids. Importantly, the core of neomangicol C will serve as a starting point for the synthesis of neomangicols A and B, as well as the mangicols.

Our retrosynthetic analysis of **2** (Scheme 1) features a late-stage disconnection to indene **5**, which offers a potential solution to the construction of terpenoid frameworks that incorporate a fused [5–6] bicyclic ring system. An appealing aspect of our approach was that the indene cyclization precursor **5** could be readily constructed in a convergent manner from vinyl triflate **6** and boronic ester **7**.

Our synthetic studies commenced with the preparation of boronic ester **7** as illustrated in Scheme 2. The sequence began with Knoevenagel condensation of 2-bromo-5-methoxybenzaldehyde (**8**)⁷ and the sodium salt of Meldrum's acid (**9**), which provided adduct **10** in 85% yield.⁸ Of note, numerous attempts to effect the Knoevenagel condensation using the conditions of Fillion (cat. pyrrolidinium acetate),⁹ which work well for electron-rich aryl aldehydes, resulted in lower yields. A conjugate reduction of alkylidene **10** was effected with sodium triacetoxyborohydride (STAB) and the resulting Meldrum's acid derivative was methylated under standard conditions (K₂CO₃, MeI). At this stage, a formal Friedel-Crafts acylation using polyphosphoric acid (PPA) proceeded with concomitant loss of acetone and carbon dioxide to yield indanone **11** in 83% yield over the three steps.¹⁰

Reduction of indanone **11** upon treatment with DIBAL-H gave an inconsequential diastereomeric mixture (3:1 dr) of indanol products,¹¹ which was immediately protected to afford MOM ether **12**. Installation of the boronic ester moiety was accomplished via halogen-metal exchange with *t*-BuLi at –78 °C followed by a quench of the resulting aryl anion with dioxaborolane **13** to give **7** in 93% yield. The overall sequence for the preparation of boronic ester **7** can be routinely performed on gram scale.

Having established reliable access to **7**, we turned our attention to the preparation of vinyl triflate **6** (Eq. 1). This compound was available in short order by enolate formation and triflation starting from easily obtained β-ketoester **13**.¹² Of note, **13** can be prepared in highly enantioenriched form,¹³ which presents an avenue for the enantioselective synthesis of these natural products.

(1)

Subsequent screening revealed an optimal set of conditions for the Suzuki cross-coupling of **6** with boronic ester **7** (Scheme 3; 10 mol % $\text{PdCl}_2(\text{PPh}_3)_2$, 1:1 *i*-PrOH/2 M aq. Na_2CO_3), which afforded **14** in 80% yield.¹⁴ At this juncture, two-stage reduction of the ethyl ester moiety followed by elimination of the MOM ether using pyridinium *p*-toluenesulfonate (PPTS) gave indene conjugate **15**. Attempted removal of the MOM ether group using other protic conditions (e.g., KHSO_4 in PhMe at $> 60^\circ\text{C}$) resulted in formation of a tetrasubstituted double bond (see **18** $\rightarrow$ **19**, Scheme 4), presumably via the mechanism as shown. Oxidation of the primary hydroxyl group of **15** (Scheme 3) set the stage for the key indenide anion-mediated C-C bond formation to forge the tetracyclic core of neomangicol C.

Deprotonation of indenenes and subsequent reactions of the resulting anions have been studied in great detail.¹⁵ In line with literature precedent, we investigated a range of bases such as *t*-BuLi, KO*t*-Bu and the amide base LiTMP (see Table 1, entries 1–3, respectively) to effect deprotonation of indene **5**. The use of *t*-BuLi was ineffective and returned starting material along with significant decomposition (entry 1). Both KO*t*-Bu and LiTMP led to isomerization of the indene double bond (entries 2–3), suggesting that deprotonation was occurring; however, none of the desired tetracycle was observed.

On the basis of the pioneering studies of Sprinzak, we were drawn to the use of hydroxide bases such as Triton B® to effect indene alkylations.¹⁶ Gratifyingly, subjecting indenyl aldehyde **5** to Triton B® (8 equiv) in DMSO at rt over 2 min provided a 50% yield of alcohol **16** as a 2:1 mixture of epimers, which existed as an inseparable mixture of double-bond positional isomers, along with several unidentified byproducts (entry 4).

Encouraged by this initial result, we embarked on optimization studies of the Triton B®-mediated cyclization of **5** by varying the solvent, temperature and reaction times. Ultimately, we identified a set of optimal conditions which utilized 1 equiv of Triton B® in DMF at -60°C (entry 6).

Because the allylic alcohol product (**16**) was relatively unstable, it was oxidized immediately following work-up to the more stable enone **17**. This compound possesses functional handles in the A and B rings that make it an attractive precursor to the neomangicol natural products.¹⁷

In summary, we report the first synthesis of the tetracyclic core of the neomangicols via an efficient indene alkylation strategy. This design provides a starting point for the syntheses of the biosynthetically related neomangicol and mangicol natural products by utilizing a powerful alternative to enolate chemistry in the synthesis of this subset of terpenes. Current efforts are focused on advancing **17** to neomangicols A–C.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

The authors are grateful to UC Berkeley, the NIH (NIGMS RO1 GM84906-01), and the American Cancer Society (RSG-09-017-01-CDD) for generous financial support. BGP thanks the CA•TRDRP for a Thesis Fellowship Award. The authors are indebted to Eric Simmons (UC Berkeley) for discussions and suggestions. RS is a 2009 Alfred P. Sloan Foundation Fellow and a 2009 Camille Dreyfus Teacher-Scholar.

References

1. For a recent discussion on terpene synthesis, see Hudlicky T, Reed J. *The Way of Synthesis: Evolution of Design and Methods for Natural Products Synthesis* (1st ed.) 2007 1st ed. Wiley-VCH Weinheim: 207–215.215
2. Renner MK, Jensen PR, Fenical W. *J. Org. Chem* 1998;63:8346–8354.
3. Araki K, Saito K, Arimoto H, Uemura D. *Angew. Chem. Int. Ed* 2004;43:81–84.
4. (a) Pichlmair S, de Lera Ruiz M, Basu K, Paquette LA. *Tetrahedron* 2006;62:5178–5194. (b) Pichlmair S, de Lera Ruiz M, Vilotijevic I, Paquette LA. *Tetrahedron* 2006;62:5791–5802.
5. Renner MK, Jensen PR, Fenical W. *J. Org. Chem* 2000;65:4843–4852. [PubMed: 10956462]
6. For a review, see Hanson JR, Trost BM, Fleming I. *Wagner-Meerwein Rearrangements. Comp. Org. Synth* 1991;Vol. 3 Oxford Pergamon:705–719.719
7. Tietze LF, Brasche G, Grube A, Böhnke N, Stadler C. *Chem. Eur. J* 2007;13:8543–8563.
8. Margaretha P, Polansky OE. *Tetrahedron Lett* 1969;57:4983–4986.
9. Dumas AM, Seed A, Zorzitto AK, Fillion E. *Tetrahedron Lett* 2007;48:7072–7074.
10. The success of the Friedel-Crafts reaction using PPA was significant, given that earlier attempts with Lewis acids such as Sc(OTf)₃ returned carboxylic acid **I** whereas AlCl₃ did effect cyclization, but also led to the cleavage of the methyl ether to yield **ii**.
11. Reduction also proceeded efficiently with Li(*Or*-Bu)₃AlH, NaBH₄ and LiEt₃BH with equal but opposite diastereocontrol.
12. Barco A, Beretti S, Pollini GP. *Synthesis* 1973:316.
13. (a) Saigo K, Koda H, Nohira H. *Bull. Chem. Soc. Jpn* 1979;52:3119–3120. (b) Ramachandran PV, Chen GM, Brown HC. *J. Org. Chem* 1996;61:95–99.
14. Initial attempts to prepare adducts related to **14** using a Grignard reagent generated from **12** and addition to β-ketoester **13** gave low returns of the desired product.
15. (a) Cedheim L, Eberson L. *Synthesis* 1973;3:159. (b) Makosza M. *Tetrahedron Lett* 1966;38:4621–4624.4624 For applications in organometallic chemistry, see (c) Bringtzinger HH, Fischer D, Mülhaupt R, Rieger B, Waymouth RM. *Angew. Chem. Int. Ed* 1995;34:1143–1170.1170 For enantioenriched indenide anions, see (d) Hoppe I, Marsch M, Harms K, Buchi G, Hoppe D. *Angew. Chem. Int. Ed* 1995;34:2158–2160.2160
16. (a) Ghera E, Sprinzak Y. *J. Am. Chem. Soc* 1960;82:4945–4952. (b) Avramoff M, Sprinzak Y. *J. Am. Chem. Soc* 1960;82:4953–4955.
17. Initial attempts to prepare tetracycle **17** from indene ester **iii** by employing the optimized conditions for the cyclization of **5**, or the conditions of Birman,¹⁸ resulted in the recovery of **iii** or the corresponding indene double-bond isomer.
18. Birman VB, Zhao Z, Guo L. *Org. Lett* 2007;9:1223–1225. [PubMed: 17328553]

products

Neomangicol A (**1a** R = Cl)
Neomangicol B (**1b** R = Br)

Neomangicol C (**2**)

Mangicol A (**3**)

Mangicol B (**4**)

Figure 1.
Selected neomangicol and mangicol natural products

Scheme 1.
Retrosynthetic analysis of neomangicol C

Scheme 2.
Synthesis of indene precursor 7

Scheme 3.
Preparation of the neomangicol tetracycle

Scheme 4.
Proposed deformylation pathway

Table 1.Optimization of the tetracyclization step^a

entry	conditions	time	result
1	<i>t</i> -BuLi (16 equiv) THF, -78 °C	15 min	SM + decomp.
2	KOr-Bu (6 equiv) MeOH, 23 °C to 50 °C	12 h	Indene isomerization + decomp.
3	LiTMP (3 equiv) THF, -78 °C to 23 °C	5 h	Indene isomerization
4	Triton B (8 equiv) DMSO, 23 °C	2 min	50% yield of 16 + byproducts
5 ^b	Triton B (1 equiv) DMF, -42 °C	10 min	50% yield of 16 + byproducts
6 ^{b,c}	Triton B (1 equiv) DMF, -60 °C	40 min	92% yield of 17

^aStandard protocol involved addition a solution of Triton B® in methanol to a solution of **5** in the indicated solvent (0.04 M).

^bA reverse addition of the substrate (**5**) to a solution of the Triton B® was used.

^cYield was determined following oxidation of the crude product (**16**) with Dess-Martin periodinane.