Molecular cloning and characterization of an antigenic protein with a repeating region from *Clonorchis sinensis*

Tae Yun KIM¹⁾, Shin-Yong KANG¹⁾, Il-Young Ahn¹⁾,
Seung-Yull CHO²⁾ and Sung-Jong HONG^{1)*}

Department of Parasitology¹⁾, Chung-Ang University Faculty of Medicine, Seoul 156-756,
Section of Molecular Parasitology²⁾, Department of Molecular Cell Biology,
Sungkyunkwan University School of Medicine, Suwon 440-746, Korea

Abstract: In the course of immunoscreening of *Clonorchis sinensis* cDNA library, a cDNA CsRP12 containing a tandem repeat was isolated. The cDNA CsRP12 encodes two putative peptides of open reading frames (ORFs) 1 and 2 (CsRP12-1 and -2). The repetitive region is composed of 15 repeats of 10 amino acids. Of the two putative peptides, CsRP12-1 was proline-rich and found to have homologues in several organisms. Recombinant proteins of the putative peptides were bacterially produced and purified by an affinity chromatography. Recombinant CsRP12-1 protein was recognized by sera of clonorchiasis patients and experimental rabbits, but recombinant CsRP12-2 was not. One of the putative peptide, CsRP12-1, is designated CsPRA, proline-rich antigen of *C. sinensis*. Both the C-termini of CsRP12-1 and -2 were bacterially produced and analysed to show no antigenicity. Recombinant CsPRA protein showed high sensitivity and specificity. In experimental rabbits, IgG antibodies to CsPRA was produced between 4 and 8 weeks after the infection and decreased thereafter over one year. These results indicate that CsPRA is equivalent to a natural protein and a useful antigenic protein for serodiagnosis of human clonorchiasis.

Key words: *Clonorchis sinensis*, repetitive peptide, antigenic, proline-rich, CsPRA

INTRODUCTION

About 30-40% of eukaryotic genome is composed of repetitive sequences. The repetitive sequences have been understood as functional components of eukaryotic genome representing genome plasticity and adaptability and contributing to genomic evolution (Brosius and Tiedge, 1996; Kidwell and Lisch, 1997; Levin, 1997). Although the specific

function of the repeats remains to be elucidated, they have been applied to determine phylogenetic relationships (Hamilton et al., 1990; Minnick et al., 1992), and to provide information about the organization of genomes (Porter, 1994; Connolly et al., 1995).

However, the genomic structures of parasites have not been intensively studied compared to those of other eukaryotes and prokaryotes. The genome of *Schistosoma mansoni* contains short repetitive DNA elements designated SM α elements (Spotila et al., 1989). The elements homologous to RNA polymerase III promoter are thought to disperse throughout the genome. Tandemly repeated DNA elements were cloned and

• Received 6 February 2001, accepted after revision 26 February 2001

*Corresponding author (e-mail: hongsj@cau.ac.kr)

sequenced from *Echinococcus granulosus*, which is clustered in the genome (Rosenzvit et al., 1997). The repetitive elements showing sex-specific polymorphism were identified from *S. mansoni* (Quack et al., 1998).

In diagnosis of parasitic human infections, subtelomeric tandem repeat sequence specific to *Leishmania braziliensis* showed usefulness in diagnosis for the parasitic infection by PCR and endonuclease digestion (Fu, 1998). From *Clonorchis sinensis*, the gene containing a repetitive sequence, Cs44, was identified and its gene product was evaluated as a potent antigenic protein for serodiagnosis of human clonorchiasis (Yong et al., 1998).

The diagnosis of clonorchiasis mainly depends on the detection of eggs in the stool. However, the difficulty of collecting stools from patients in the field have urged the researchers to recruit another diagnostic tests such as serodiagnosis. Among them, the enzyme-linked immunosorbent assay (ELISA) has been employed as an effective serodiagnostic technique due to its specificity and sensitivity (Hahm et al., 1984; Yong et al., 1996). Studies on appropriate immunodiagnostic techniques should be paralleled to search specific and sensitive antigenic proteins. The specific and sensitive antigenic proteins of *C. sinensis* have been studied for several decades but antigenic molecules superior to the crude worm extract have not been identified yet.

In this study, to identify genes encoding antigenic proteins, an expression cDNA library of adult *C. sinensis* was screened with a clonorchiasis patient serum. A clone with repetitive sequences was purified and its gene product produced bacterially was isolated and characterized. It was sensitive and specific for serodiagnosis of human clonorchiasis.

MATERIALS AND METHODS

Cloning a repetitive cDNA

A cDNA expression library of adult *C. sinensis* constructed previously (Hong et al., 2000) was used. The cDNA library, 6×10^6 pfu, was mixed with *Escherichia coli* XL1-Blue and cultured on a LB-agar plate. The plate was overlaid with a nitrocellulose membrane

(BioTrace™ NT; Gelman Science, Ann Arbor, MI, USA) soaked previously in 10 mM isopropyl-D-thiogalactoside (IPTG), and incubated further at 37°C for 4 hr. The membrane alone was then incubated for 3 hr within a *C. sinensis*-infected human serum diluted by 1:100 at room temperature, followed by incubation with secondary antibody, alkaline phosphatase-conjugated anti-human IgG diluted by 1:2,000. A color signal was developed with the addition of 5-bromo-4-chloro-3-indolyl phosphate and nitroblue tetrazolium (BCIP/NBT; Sigma Chemical Co., St. Louis, MO, USA) solution. Positive clones were purified through secondary and tertiary screenings using the same serum. The purified clones were in vivo excised to single-stranded phagemids by employing a helper phage (ExAssist™, Stratagene, La Jolla, CA, USA) and converted to double-stranded plasmids by following manufacturer's instruction.

cDNA sequencing and data base search

The cDNA sequences were determined by using BigDye™ Terminator Cycle sequencing kit (Applied Biosystems, Foster City, CA, USA) at Korea Basic Science Institute (Taejeon, Korea). Briefly, plasmid DNA and primers were added to the BigDye mixture, and thermal cycling was carried out by 25 cycles at 96°C for 30 sec, at 50°C for 15 sec and at 60°C for 4 min. After the removal of unincorporated nucleotides, the products were denatured by heating at 90°C for 2 min and electrophoresed on an automatic DNA sequencer (Model 377, Applied Biosystems). To get overlapping and both strand sequences, specific internal oligonucleotides (Fig. 1) were synthesized at Bioneer Co. (Chungwon-gun, Korea) and used in the cycle DNA sequencing. Putative peptide sequence of the cDNA was analysed using DNASIS program (Hitachi Software Engineering Co., Yokohama, Japan). Sequence homology of the cDNA clone was searched using GenBank database by BLAST program (Altschul et al., 1997). Multiple alignments of peptide sequences were generated by CLUSTAL W program (Thompson et al., 1994) and were optimized manually in part.

Production of recombinant CsRP12-1 and CsRP12-2 proteins

To produce recombinant proteins corresponding to the open reading frames (ORFs) 1 and 2 of the CsRP12 cDNA, coding region of CsRP12 was enzymatically cut and ligated into an expression vector pRSET™ A or B (Invitrogen, Carlsbad, CA, USA) with respect to the frame set of each respective ORF. CsRP12 cDNA was digested with *EcoR* I endonuclease and run on 1% agarose gel. The cDNA fragment corresponding to the coding region was excised and recovered from the gel slice by gene cleaning method. pRSET A and B were also digested and recovered as aforementioned. The recovered *EcoR* I-fragment of CsRP12 was ligated into pRSET A or B by using T4 DNA ligase. Bacterial host cells, *E. coli* XL-1 Blue, were transformed and in frame connection of CsRP12 ORF 1 and 2 to the tag peptide of the respective vector was confirmed by DNA sequencing. The plasmids, pRSET A and B constructs, were then transformed into *E. coli* BLR(DE3)pLysS (Novagen, Madison, WI, USA) by heat shock method. Positive colonies were grown to $A_{600} = 0.6$ and expression of the recombinant fusion proteins were induced by adding IPTG at 1 mM final concentration and further incubation at 37°C for 4 hr. The bacterial pellet harvested by spinning the culture medium was resuspended in mouse tonicity phosphate-buffered saline (15 mM NaCl, 160 mM NaH_2PO_4 , 40 mM Na_2HPO_4 , 0.5 mM phenylmethylsulfonyl fluoride), and electrophoresed in reducing condition on 13.5% SDS-polyacrylamide gel. Localization of the expressed fusion proteins were checked by Coomassie Brilliant blue staining and by immunoblotting using anti-Xpress™ mouse IgG (Invitrogen) as a primary antibody that specifically recognize the tag peptide of the pRSET vectors.

Since the expressed protein is localized in cytosolic fractions as an insoluble form, the recombinant fusion proteins were purified by an affinity chromatography employing nickel-nitrotri-acetic acid (Ni-NTA) resin (Qiagen GmbH, Hilden, Germany) under denaturation condition according to the manufacturer's instruction. The bacterial pellets were

resuspended in buffer A (6 M Guanidine hydrochloride, 0.1 M NaH_2PO_4 , 10 mM β -mercaptoethanol, 0.01 M Tris, pH 8.0). The suspension was spun, and supernatant was removed and loaded onto Ni-NTA column. The column was sequentially washed out with buffers B and C (8 M urea, 0.1 M NaH_2PO_4 , 0.01 M Tris, pH 8.0 and 6.3, respectively). The bound fusion protein was eluted with buffers D and E (8 M urea, 0.1 M NaH_2PO_4 , 0.01 M Tris, pH 5.9 and 4.5, respectively) containing 250 mM imidazole. The fractions collected were dialyzed exhaustively 3 times for 3 hr each against distilled water and deployed by SDS-PAGE. The fusion proteins were detected by immunoblotting with anti-Xpress IgG (Invitrogen).

Flukes and sera

Clonorchis sinensis metacercariae were collected from *Pseudorasbora parva* caught in Chinju, Kyeongsangnam-do and administered orally, 500 metacercariae each, to four New Zealand White rabbits. Adult *C. sinensis* were recovered from bile ducts of the experimental rabbits 6 months after the metacercarial infection. The recovered flukes were used immediately, or stored at -70°C.

Sera were collected from the experimental rabbits at 1-6 week interval over one year, and stored at -20°C until used. After one year, the rabbits were sacrificed and adult worms were recovered.

Sera was collected from 35 clonorchiasis and 5 paragonimiasis patients proven parasitologically infected with the respective flukes. Cysticercosis sera were obtained from 5 patients diagnosed by CT/MRI imaging. Sparganosis sera were collected from 5 patients proven by surgical removing of the worm(s).

Antigenicity of recombinant CsRP12-1 and CsRP12-2 proteins

The purified recombinant proteins (CsRP12-1 and -2) were separated on 13.5% SDS-PAGE and electrotransferred onto nitrocellulose membranes (Hybond™-C, Amersham Pharmacia Biotech, Uppsala, Sweden). The membranes were immunoblotted with sera of helminth-infected humans and experimental

As both recombinant CsRP12-1 and -2 proteins were localized in cytosolic insoluble fraction in bacterial host BLR(DE3)pLysS, both of the recombinant fusion proteins were purified by Ni-NTA affinity chromatography under denaturation condition. The recombinant fusion proteins CsRP12-1 and -2 were efficiently eluted as 37 and 30 kDa molecules,

Fig. 3. Purification of recombinant CsRP12-1 and -2 proteins. Recombinant proteins bacterially produced were charged in a denaturation condition onto Ni-NTA column and eluted with a buffer containing 250 mM imidazole. The recombinant proteins purified and deployed in SDS-polyacrylamide gel were detected by Coomassie blue staining (A) and immunoblotting with anti-Xpress antibody (B). Lane 1, induced bacterial lysate; lane 2, flow-through; lane 3, eluate.

1. All of the clonorchiasis patient sera reacted to recombinant CsRP12-1 protein (Fig. 4). Cross-reactivity of the recombinant CsRP12-1 appeared to be 12-21% against the tissue invading helminth-infected patient sera. However, the signals in immunoblottings were marginal for positive. The recombinant CsRP12-2 protein did not react to clonorchiasis patient sera nor to any helminth-infected patient sera (Table 1).

Table 1. Seroreactivity of recombinant CsRP12 proteins to various helminth-infected patient sera.

	CsRP12-1		CsRP12-2	
	No. of sera	Positive sera (%)	No. of sera	Positive sera (%)
Clonorchiasis	35	35 (100%)	5	0 (0%)
Paragonimiasis	19	4 (21%)	5	0 (0%)
Sparganosis	17	2 (12%)	5	0 (0%)
Cysticercosis	17	3 (18%)	5	0 (0%)
Normal control	2	0 (0%)	2	0 (0%)

Antibodies reactive to CsRP12-1 in *C. sinensis*-infected rabbits

As shown in Fig. 5, IgG antibodies reacting to the recombinant CsRP12-1 were produced between week 4 and 8 after the metacercarial infection. The IgG antibody level decreased steadily from week 8 to week 52 after the infection.

Antigenic property of recombinant CsRP12 C-terminal peptides.

Both of the recombinant C-terminus peptides, CsRP12ctm1 and CsRP12ctm2, were found in insoluble forms in *E. coli* lysate and thereafter purified by Ni-NTA affinity chromatography under denaturation condition (Fig. 6A). The immunoreactivities were analyzed by immunoblotting with *C. sinensis*-infected patient sera and anti-RGS His antibody. Of seven *C. sinensis*-infected sera, only one reacted with the recombinant CsRP12ctm1 protein. Recombinant CsRP12ctm2 protein did not produce any positive signal against the sera (Fig. 6 B).

DISCUSSION

A cDNA, CsRP12, was cloned by immunoscreening the adult *C. sinensis* cDNA library. Of the two putative peptides deduced from the CsRP12, CsRP12-1 corresponding to ORF-1 showed the highest homology to a glycine-rich *C. sinensis* protein (GRCSF; Yang et al., 2000) and a wide range of sequential identities with various proteins of invertebrate animals (Fig. 2). The homologous proteins

Fig. 4. Immunoblots of purified recombinant CsRP12-1 to helminth-infected patients' sera. A. Immunoblot against clonorchiasis patients' sera. B. Other helminth-infected patients' sera. Pw, paragonimiasis; Sp, sparganosis; Cy, cysticercosis; N, normal human sera; C, control by anti-Xpress antibody.

Fig. 5. An immunoblot of recombinant CsRP12-1 protein to *Clonorchis sinensis*-infected rabbit sera.

contain the repetitive peptides rendering sequential identities to CsRP12-1. Of the two bacterially produced recombinant proteins of CsRP12, CsRP12-1 was recognized by the sera of *C. sinensis*-infected patients and experimental rabbits, but CsRP12-2 was not. These results indicate that the peptide corresponding to CsRP12-1 is the natural protein occurring in *C. sinensis*. With superior antigenicity and proline-richness in the repeating unit, the CsRP12-1 protein designates CsPRA (proline-rich antigen of *C. sinensis*) in this report.

The peptide structure of CsPRA is different from GRCSP in amino acid composition of the

repeating units, proline-rich versus glycine-rich and number of repetition, 15-times versus 23 times (Yang et al., 2000). However, CsPRA and GRCSP share high sequential identities in their N- and C-terminus regions. The homologous proteins to CsPRA are identified to be surface-associated and antigenic. The proteins homologous to CsPRA were identified to be the surface proteins of the respective organisms. In this respect, it is assumed that CsPRA may be associated with surface membrane of *C. sinensis*.

The recombinant CsPRA protein, although its N-terminus was truncated, was a good antigen showing high sensitivity and specificity toward clonorchiasis patient sera. A counterpart, GRCSP, which shows the highest homology to CsPRA, showed a high antigenic specificity and moderate sensitivity for serodiagnosis of clonorchiasis (Yong et al., 1998). In protozoan and helminthic parasites, proteins containing repetitive peptides have been identified to be highly antigenic with acceptable sensitivities and specificities for serological diagnosis of the respective infections (Imboden et al., 1995; Charest and Matlashewski, 1994; Burns et al., 1993; Marin et al., 1992). Both of the recombinant C-terminal peptides of CsPRA, CsRP12ctm1 and CsRP12ctm2, were not antigenic toward *C.*

Fig. 6. Purification and antigenicity of bacterial recombinant C-terminus peptides of CsRP12. A. Purification of the recombinant proteins by Ni-NTA affinity chromatography. Lane 1, induced lysate; lane 2, pass-through fraction; lanes 3 and 4, buffers B and C, respectively; lanes 5, 6, 7 and 8, eluates with buffers C, D, E and F containing 250 mM imidazole, respectively. B. Immunoblots against *C. sinensis*-infected patient sera. Lanes 1 to 7, clonorchiasis patients sera; lane 8, anti-RGS His antibody.

sinensis-infected patient sera (Fig. 6) indicating that antigenic epitope of CsPRA protein is most probably in the repetitive region. In many antigenic proteins, the repetitive regions are found to contain the major antigenic determinant(s) (Kulane et al., 1999; Banic et al., 1998; Burns et al., 1993) when compared to less antigenic N- and C-terminus regions (Lopez et al., 1996; Sharma et al., 1996). The repetitive amino acid sequences are variable in length and the number of repeats and show low sequence similarities with each other. The variability of the repetitive peptides has been recognized to render antigenic specificities of the protein (Kulane et al., 1999; Imboden et al., 1995).

The CsPRA-specific antibodies were produced in the experimental rabbits as early as 5-7 weeks after the *C. sinensis* infection. In cattle, serum IgG antibodies to a *Fasciola hepatica* antigenic protein bearing repetitive

sequences appeared as early as 4-5 weeks after the infection, increased to a peak at 11-13 weeks, and decreased slowly thereafter (Marin et al., 1992), as observed in *C. sinensis* infection. A repetitive peptide was expressed stage-specifically in plerocercoids of *Spirometra erinaceieuropaei* (Liu et al., 1996). It is, therefore, tempting to note that CsPRA is expressed in the juvenile worms, irrespective of its biological role, and presented as an antigen to the host animals including humans in the early stage of infection.

Interestingly, when kept at room temperature, the purified CsPRA was readily degraded showing a ladder-like feature in SDS-polyacrylamide gel (data not shown). The ladder pattern was highly ordered by 15 fragments equal to the number of repeats. The PEST sequences (proline, glutamate, serine and threonine-rich sequence) involved in protein degradation (Rogers et al., 1986)

creating structural domains recognized by proteolytic enzymes were not found in the CsRP12 protein sequence. Moreover, since proteinase inhibitor cocktail was added immediately after the purification, the degradation of recombinant CsPRA is assumed to be arisen from auto-proteolysis. This may support the smearing under the CsPRA major band in immunoblotting reacted with clonorchiasis patients sera (Fig. 3).

ACKNOWLEDGEMENT

This work was supported by the grant from the Health Technology Planing and Evaluation (HMP-96-M-2-1059), the Ministry of Health and Social Affairs, the Republic of Korea.

REFERENCES

- Altschul SF, Madden TL, Schaffer AA, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389-3402.
- Banic DM, de Oliveira-Ferreira J, Pratt-Riccio LR, et al. (1998) Immune response and lack of immune response to *Plasmodium falciparum* P126 antigen and its amino-terminal repeat in malaria-infected humans. *Am J Trop Med Hyg* **58**: 768-774.
- Brosius J, Tiedge H (1995) Reverse transcriptase: mediator of genomic plasticity. *Virus Genes* **11**: 163-179.
- Brown H, Kemp DJ, Barzaga N, Brown GV, Anders RF, Coppel RL (1987) Sequence variation in S-antigen genes of *Plasmodium falciparum*. *Mol Biol Med* **4**: 365-376.
- Burns JM, Shreffler WG, Benson DR, Ghalib, HW, Badaro R, Reed SG (1993) Molecular characterization of a kinesin-related antigen of *Leishmania chagasi* that detects specific antibody in African and American visceral leishmaniasis. *Proc Natl Acad Sci USA* **90**: 775-779.
- Buschiazio A, Campetella OE, Marcina RA, et al (1992) Sequence of the gene for a *Trypanosoma cruzi* protein antigenic during the chronic phase of human Chagas disease. *Mol Biochem Parasitol* **54**: 125-128.
- Charest H, Matlashewski G (1994) Developmental gene expression in *Leishmania donovani*: differential cloning and analysis of an amastigote-stage-specific gene. *Mol Cell Biol* **14**: 2975-2984.
- Connolly B, Ingram LJ, Smith DF (1995) *Trichinella spiralis*: cloning and characterization of two repetitive DNA sequences. *Exp Parasitol* **80**: 488-498.
- Fu G, Perona-Wright G, Barker DC (1998) *Leishmania braziliensis*: characterization of a complex specific subtelomeric repeat sequence and its use in the detection of parasites. *Exp Parasitol* **90**: 236-243.
- Hahm WH, Lee JS, Rim HJ (1984) Comparative study on the indirect immunofluorescent antibody test, complement test and ELISA in diagnosis of human clonorchiasis. *Korea Univ Med J* **21**: 177-184.
- Hamilton MJ, Honeycutt RL, Baker RJ (1990) Intragenomic movement, sequence amplification and concerted evolution in satellite DNA in harvest mice, *Reithrodontomys*: evidence from in situ hybridization. *Chromosoma* **99**: 321-329.
- Hauser F, Roeben C, Hoffmann W (1992) xP2, a new member of the P-domain peptide family of potential growth factors, is synthesized in *Xenopus laevis* skin. *J Biol Chem* **267**: 14451-14455.
- Hong SJ, Seong KY, Sohn WM, Song KY (2000) Molecular cloning and immunological characterizations of phosphoglycerate kinase from *Clonorchis sinensis*. *Mol Biochem Parasitol* **108**: 207-216.
- Hong ST, Park KH, Seo M, Choi BI, Chai JY, Lee SH (1994) Correlation of sonographic findings with histopathological changes of the bile ducts in rabbits infected with *Clonorchis sinensis*. *Korean J Parasitol* **32**: 223-230.
- Imboden M, Muller N, Hemphill A, Mattioli R, Seebeck T (1995) Repetitive proteins from the flagellar cytoskeleton of African trypanosomes are diagnostically useful antigens. *Parasitology* **110**: 249-258.
- Kidwell MG, Lisch D (1997) Transposable elements as sources of variation in animals and plants. *Proc Natl Acad Sci USA* **94**: 7704-7711.
- Kulane A, Siddique AB, Sarthou JL, et al. (1999) Human immune responses to the highly repetitive *Plasmodium falciparum* antigen Pf332. *Am J Trop Med Hyg* **61**: 141-148.
- Levin HL (1997) It's prime time for reverse transcriptase. *Cell* **88**: 5-8.
- Liu DW, Kato H, Kazuo S (1996) Isolation of a cDNA encoding an antigenic polypeptide containing repeating units of *Spirometra*

- erinaceieuropaei*. *Jpn J Parasitol* **45**: 324-327.
- Lopez JA, Roggero MA, Duombo O, et al. (1996) Recognition of synthetic 104-mer and 102-mer peptides corresponding to N- and C-terminal nonrepeat regions of the *Plasmodium falciparum* circumsporozoite protein by sera from human donors. *Am J Trop Med Hyg* **55**: 424-429.
- Marin MS, Prieto M, Martin JM, Casais R, Boga JA, Parra F (1992) Identification and expression of a *Fasciola hepatica* gene encoding a gut antigen protein bearing repetitive sequences. *Mol Biochem Parasitol* **55**: 155-165.
- Minnick MF, Stillwell LC, Heineman JM, Stiegler GL (1992) A highly repetitive DNA sequence possibly unique to canids. *Gene* **110**: 235-238.
- Peterson DS, Wrightsman RA, Manning JF (1986) Cloning of a major surface-antigen gene of *Trypanosoma cruzi* and identification of a nonapeptide repeat. *Nature* **322**: 566-568.
- Porter CA (1994) Organization and chromosomal location of repetitive DNA sequences in three species of squamate reptiles. *Chromosome Res* **2**: 263-273.
- Quack T, Doenhoff M, Kunz W, Grevelding CG (1998) *Schistosoma mansoni*: the varying occurrence of repetitive elements in different strains shows sex-specific polymorphisms. *Exp Parasitol* **89**: 222-227.
- Rosenzvit MC, Canova SG, Kamenetzky L, Ledesma BA, Guarnera EA (1997) *Echinococcus granulosus*: cloning and characterization of a tandemly repeated DNA element. *Exp Parasitol* **87**: 65-68.
- Rogers S, Wells R, Rechsteiner M (1986) Amino acid sequences common to rapidly degraded proteins: the PEST hypothesis. *Science* **234**: 364-368.
- Sharma S, Goswami A, Singh NJ, Kabilan L, Deodhar SS (1996) Immunogenicity of the nonrepetitive regions of the circumsporozoite protein of *Plasmodium knowlesi*. *Am J Trop Med Hyg* **55**: 635-641.
- Spotila LD, Hirai H, Rekosh DM, Lo Verde PT (1989) A retroposon-like short repetitive DNA element in the genome of the human blood fluke, *Schistosoma mansoni*. *Chromosoma* **97**: 421-428.
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673-4680.
- Wilson R, Ainscough R, Anderson K, et al. (1994) 2.2 Mb of contiguous nucleotide sequence from chromosome III of *C. elegans*. *Nature* **368**: 32-38.
- Yang HJ, Park SJ, Im KI, Yong TS (2000) Identification of a *Clonorchis sinensis* gene encoding a vitellaria antigenic protein containing repetitive sequences. *Mol Biochem Parasitol* **111**: 213-216.
- Yong TS, Lee JS, Cho SN, Seo JH, Park H (1996) A carbohydrate antigen of *Clonorchis sinensis* recognized by a species-specific monoclonal antibody. *Korean J Parasitol* **34**: 279-281.
- Yong TS, Yang HJ, Park SJ, Kim YK, Lee DH, Lee SM (1998) Immunodiagnosis of clonorchiasis using a recombinant antigen. *Korean J Parasitol* **36**: 183-190.