Determination of antigenic domain in GST fused major surface protein (Nc-p43) of *Neospora caninum*

Eui-Sun SON¹⁾, Hye-Jin AHN¹⁾, Jae-Hoon KIM²⁾,
Dae-Yong KIM³⁾ and Ho-Woo NAM^{1)*}

Department of Parasitology and Catholic Institute of Parasitic Diseases¹⁾,
Catholic University of Korea, College of Medicine, Seoul 137-701, National Veterinary Research and
Quarantine Service²⁾, Anyang City 430-016, and Department of Veterinary Pathology³⁾,
College of Veterinary Medicine and School of Agricultural Biotechnology, Seoul National University,
Suwon City 441-744, Korea

Abstract: The antigenic domain of the major surface protein (Nc-p43) of *Neospora caninum* was examined by polymerase chain reaction of its gene fragments and recombinant expression as GST fusion proteins. The fragments of Nc-p43 were as follow: a total open reading frame (OFR), T; OFR without signal sequence and C-terminal hydrophobic sequence, S; N-terminal 2/3 parts of S, A; C-terminal 2/3 parts, P; N-terminal 1/3 part, X; middle 1/3 part, Y; and C-terminal 1/3 part, Z, respectively. The DNA fragments were cloned into pGEX-4T vector. Recombinant plasmids transformed into *Escherichia coli* of BL21 pLysS (DE3) strain were induced to express GST or GST fused fragments of Nc-p43 such as 69 kDa protein for T, 66 kDa for S, 52 kDa for A, 53 kDa for P, and 40 kDa proteins for X, Y, and Z, respectively in SDS-PAGE. The Nc-p43 fragments of T, S, and P reacted with a bovine serum of neosporosis while those of A, X, Y, and Z together with GST did not in the western blot. These findings suggest that the antigenic domain of Nc-p43 of *N. caninum* may be localized in the C-terminal 2/3 parts. Together with A19 clone in SAG1 of *Toxoplasma gondii* (Nam et al., 1996), the P fragment of Nc-p43 could be used as efficient antigens to diagnose and differentiate those infections with both species.

Key words: *Neospora caninum*, Nc-p43, expression, western blot, antigenic domain

INTRODUCTION

Neospora caninum is an apicomplexan parasite that was originally identified as an aetiological agent of neurological disease in dogs

and has been known to be associated with abortions in cattle (Dubey et al., 1988; Dubey and Lindsay, 1993). Both tachyzoites and cyst-forming bradyzoites have been characterized in the asexual phase of *N. caninum* (Lindsay et al., 1993). And recently, it was described that dogs are definitive hosts of this parasite (McAllister et al., 1998).

The surface proteins of apicomplexan parasites are often immunodominant and seem to be of particular interest as a diagnostic tool and/or as vaccine antigens (Bulow and Boothroyd, 1991; Lunden et al.,

Received 27 July 2001, accepted after revision 22 August 2001.

This work was financially supported by the Ministry of Agriculture and Forestry (399002-3) of the Republic of Korea.

*Corresponding author (e-mail: howoo@cmc.cuk.ac.kr)

1997). Many surface proteins of *N. caninum* have been identified including Nc-p43 (Hemphill and Gottstein, 1996), p29 and p35 (Howe et al., 1998), and p38 (Schaes et al., 2000). Their functions have not been elucidated, however, there is an indirect evidence that at least one of these antigens, Nc-p43, is involved in the attachment of the parasite to host cells (Hemphill, 1996). A complete Nc-p43 gene has been cloned for the DNA vaccine trials (Nishikawa et al., 2000, 2001). Therefore, the essential antigenic determinants should be investigated for cloning in order to produce a large quantity of antigens in recombinant forms for the diagnostic purpose. And for the protein or DNA vaccine, cloning of crucial epitope should be performed to eliminate the unnecessary interaction between processed antigen of non-essential and host immune machinery. In this study, fragments of Nc-p43 were expressed as GST fusion proteins to examine the antigenic domains.

MATERIALS AND METHODS

Parasite

N. caninum tachyzoites of the Nc-1 strain and the Korean isolate (KBA-2, Kim et al., 2000) were maintained in Vero cells (CRL 6318, ATCC, Rockville, MD) in DMEM supplemented with 10% FBS (Gibco BRL Co.,

Rockville, MD). Pure *N. caninum* tachyzoites were obtained from the supernatant of 3 to 4 days confluent culture.

Reverse transcriptase-polymerase chain reaction (RT-PCR) and amplification of Nc-p43 gene fragments

Total RNA was purified from the tachyzoite extracts with Tri reagent (Sigma Chem. Co., St. Louis, MO) and used as templates. RT-PCR was performed with a primer set of T in Table 1, which amplified the open reading frame of the major surface membrane antigen (Nc-p43) (GeneBank accession number U93870). Amplified DNAs were cloned into a pGEM-T Easy vector (Promega Corp., Madison, WI) and sequenced with T7 and SP6 primers. Oligonucleotide primers were synthesized according to the coding sequence of GeneBank

Fig. 1. Design of fragmentation of Nc-p43 gene into hydrophilic and hydrophobic moieties.

Table 1. Primers designed for the amplification of NC-p43 gene fragments

Name of primer			Sequence	Size of DNA Amplified (bp)	Estimated GST-protein (kDa)
T	Sense		5'-ATG GCG ACG CAT GCT TGT-3'	1,203	69
	Antisense		5'-GTA CGC AAA GAT TGC CGT-3'		
S	Sense		5'-AAG GCG GAT GCT GCT TGC-3'	1,104	66
	Antisense		5'-TGC ATC TCC TCT TAA CAC-3'		
A	Sense		5'-AAG GCG GAT GCT GCT TGC-3'	738	53
	Antisense		5'-CCA AAA ACT ACT GGA GAA-3'		
P	Sense		5'-AAA GAG TGG GTG ACT GGA-3'	732	53
	Antisense		5'-TGC ATC TCC TCT TAA CAC-3'		
X	Sense		5'-AAG GCG GAT GCT GCT TGC-3'	372	40
	Antisense		5'-TTT GGC CCC CGG AAG CAA-3'		
Y	Sense		5'-AAA GAG TGG GTG ACT GGA-3'	366	40
	Antisense		5'-CCA AAA ACT ACT GGA GAA-3'		
Z	Sense		5'-ACT GGA GAA GCT TCC GGC-3'	366	40
	Antisense		5'-TGC ATC TCC TCT TAA CAC-3'		

accession number U93870 as designed in Fig. 1. The DNA sequences of the synthesized primers are listed in Table 1. PCR was done on a subcloned pGEM-T vector with Nc-p43 gene from KBA-2 strain.

Construction of recombinant plasmids and expression of fusion proteins

The amplified DNAs were inserted into pGEM-T Easy vector, and then subcloned into pGEX-4T vector (Amersham Pharmacia Biotech., Uppsala, Sweden) with *Eco* RI (Gibco BRL) digestion. After confirming the correct orientation of an insert, plasmids were transformed into *Escherichia coli* of BL21 pLysS (DE3) strain (Invitrogen, Carolsbad, CA). *E. coli* cells of log phase were treated with 0.5 mM isopropyl β -D-thiogalactoside (IPTG) for 3 hr at 30°C to induce the expression of the fusion proteins.

Western blot

Western blot was performed by the method of Towbin et al. (1979). The cell extracts were separated on 12% SDS-PAGE gels and transferred onto nitrocellulose sheets (NC, Schleicher and Shuell, Keene, NH). The fusion proteins were confirmed with GST detection kit (Amersham Pharmacia Biotech). For the determination of the antigenic domain, NC paper was blocked by 10% rabbit serum in PBS/0.05% Tween-20 overnight. NC paper was then incubated with 1:500 diluted bovine serum of neosporosis (Bae et al., 2000) followed by incubation with 1:5,000 diluted HRP-conjugated goat anti-bovine IgG antibody (Sigma). They were soaked in enhanced chemiluminescence (ECL) solution (Intron, Daejeon, Korea) for 1 min and exposed to an X-ray film (Konica, Tokyo, Japan).

RESULTS

With a set of primers used to amplify the ORF of Nc-p43 gene, a 1,203 bp DNA fragment was obtained by RT-PCR from each Nc-1 and KBA-2 strain of *N. caninum* (Fig. 2). Each DNA was subcloned into a pGEM-T easy vector for sequencing. The DNA sequencing data of the Nc-p43 gene of the KBA-2 strain revealed a point mutation at the 625th position where

Fig. 2. PCR amplified Nc-p43 coding genes of Nc-1 (N) and KBA-2 (K) strains. Numerals indicate bp.

Fig. 3. Sequence variation in Nc-p43 gene and deduced amino acid of Nc-1 and KBA-2 strains. Point mutation in DNA sequence and deduced amino acid sequence are italicized.

cytosine was in place of guanine in the sequence of Nc-1. This affected the deduced amino acid sequence of the 209th residue, which was changed to glutamine (Q) from glutamic acid (E) as shown in Fig. 3. When using the Nc-p43 (T) as a template, a number of Nc-p43 fragments was obtained with sets of primers listed in Table 1. The sizes of these fragments were 1,104 bp for S, 738 bp for A, 732 bp for P, 372 bp for X, 366 bp for Y, and 366 bp for Z as designated in Fig. 1 (Fig. 4). Each DNA was inserted into a pGEM-T Easy vector and then subcloned into a pGEX-4T with *Eco* RI digestion (data not shown).

pGEX-4T and recombinant plasmids transformed into *E. coli* of BL21 pLysS (DE3) strain were induced to express GST or GST fused fragments of Nc-p43 with the addition of IPTG. As shown in Fig. 5, GST was expressed

Fig. 4. PCR-amplified Nc-p43 coding gene fragments of KBA-2 strain separated in 1.2% agarose gel. Numerals indicate bp.

Fig. 5. SDS-PAGE of expressed GST-Nc-p43 fusion proteins. Expressed GST and GST fused proteins are indicated by asterisks.

as a 26 kDa protein. GST fusion proteins were expressed as 69 kDa protein for T, 66 kDa for S, 52 kDa for A, 53 kDa for P, and 40 kDa proteins for X, Y, and Z, respectively, when visualized on SDS-PAGE. Each fusion protein was confirmed with GST detection kit (Fig. 6A) as comparable molecular weight with that of SDS-PAGE. In the western blot, Nc-p43 fragments of T, S, and P reacted with a bovine serum of neosporosis while those of A, X, Y, and Z together with GST did not as shown in Fig. 6B.

Fig. 6. Western blot detection of GST fusion proteins of Nc-p43 fragments (A) and with bovine serum confirmed to be infected with *Neospora caninum*. Asterisks indicate GST and GST fused proteins as in Fig. 5.

DISCUSSION

In the western blot, bovine serum of neosporosis detected a P fragment without binding to the subfragments of it, Y nor Z. This suggests that the antigenic domain of Nc-p43 of *N. caninum* may be localized in the C-terminal 2/3 parts, of which polypeptides of the second 1/3 with the presence of the third 1/3 or the third 1/3 with the presence of the second 1/3 or in the oligopeptides between the margins of the second and the third 1/3 parts. This is comparable to the antigenic domain of the major surface antigen of *Toxoplasma gondii*, SAG1, in the N-terminal 1/3 with the presence of middle 1/3 (Nam et al., 1996). Although oligopeptides of about 20 amino acids are sufficient to induce B- and T-cell immune responses (Godard et al., 1994), a conformational aid by the other parts seems to be essential to express antigenicity in western blot.

The Korean isolate KBA-2 has a point mutation in the coding gene of Nc-p43, which affects the deduced amino acid at 209th residue from glutamic acid of Nc-1 strain to glutamine. A substitution of glutamic acid, which has an acidic side chain, to an uncharged glutamine residue may contribute

to the change in conformation of the antigen. This change in amino acids occurs in the antigenic domain of C-terminal 2/3 of Nc-p43 protein. The extracts of KBA-2 strain was more suitable as a diagnostic antigen than those of Nc-1 for *N. caninum* infection in Korea (Bae et al., 2000), though there were no differences in morphology of the parasites and pathologic findings in the hosts by the infections (Kim et al., 2000). This partially reflects the prevalent strain of KBA-2 in Korea.

Nc-p43 has been known to play an important role during the initial physical interaction between the parasite and the host cell surface (Hemphill and Gottstein, 1996). And the affinity-purified anti-Nc-p43 antibodies inhibited adhesion and invasion of *N. caninum* tachyzoites into the host cells (Hemphill, 1996). But the inhibitory effect may result from the aggregation of tachyzoites themselves by the antibody, which merely delay the entry of tachyzoites. Controversially, it is also possible for multiple entry of the tachyzoites of *N. caninum* into host cells through the crosslinking by the antibody as in the case of *Plasmodium falciparum* (Ramasamy et al., 2001). And the surface antigens are inserted into the membrane by glycosylphosphatidylinositol (GPI)-anchor (Howe et al., 1998) to evade the host immune by shaking off the attack of the antibody.

Prior to 1988, *N. caninum* was misdiagnosed as *T. gondii* due to its close structural similarity (Dubey, 1992). However, due to the ultrastructural differences (Lindsay et al., 1993) and difference in genetic level (Marsh et al., 1995), it has been possible to distinguish these two species from each other. Although the gene encoding Nc-p43 revealed a sequence homology to the SRS-2 gene of *T. gondii* (Hemphill et al., 1997; Howe et al., 1998), cross reactivity of this antigen with *T. gondii* infection seem to be negligible as determined by immunoblotting (Bjerkas et al., 1994; Pare et al., 1997; Bae et al., 2000). Together with A19 in SAG1 of *T. gondii* (Nam et al., 1996), the P fragment of Nc-p43 could be used as antigens for the diagnosis of *N. caninum* infection, and furthermore, the antigens could be used to efficiently differentiate *N. caninum* from *T. gondii*.

REFERENCES

- Bae JS, Kim DY, Hwang WS, Kim JH, Lee NS, Nam HW (2000) Detection of IgG antibody against *Neospora caninum* in cattle in Korea. *Korean J Parasitol* **38**: 245-249.
- Bjerkas I, Jenkins MC, Dubey JP (1994) Identification and characterization of *Neospora caninum* tachyzoite antigens useful for diagnosis of neosporosis. *Clin Diagn Lab Immunol* **1**: 214-221.
- Bulow R, Boothroyd JC (1991) Protection of mice from fatal *Toxoplasma* infection by immunization with p30 antigen in liposomes. *J Immunol* **141**: 3584-3591.
- Dubey JP, Carpenter JL, Speer A, Topper MJ, Uggla A (1988) Newly recognized fatal protozoan disease of dogs. *J Am Vet Med Assoc* **198**: 1269-1285.
- Dubey JP (1992) A review of *Neospora caninum* and *Neospora*-like infections in animals. *J Protozool Res* **2**: 40-54.
- Dubey JP, Lindsay DS (1993) Neosporosis. *Parasitol Today* **9**: 452-458.
- Godard I, Estaquier J, Zenner L, et al. (1994) Antigenicity and immunogenicity of p30-derived peptides in experimental models of toxoplasmosis. *Mol Immunol* **31**: 1353-1363.
- Hemphill A (1996) Subcellular localization and functional characterization of Nc-p43, a major *Neospora caninum* tachyzoite surface protein. *Infect Immun* **64**: 4279-4287.
- Hemphill A, Gottstein B (1996) Identification of a major surface protein on *Neospora caninum* tachyzoites. *Parasitol Res* **82**: 497-504.
- Hemphill A, Felleisen R, Connolly B, Gottstein B, Hentrich B, Muller N (1997) Characterization of a cDNA-clone encoding Nc-p43, a major *Neospora caninum* tachyzoite surface protein. *Parasitology* **115**: 581-590.
- Howe DK, Crawford AC, Lindsay D, Sibley LD (1998) The p29 and p35 immunodominant antigens of *Neospora caninum* tachyzoites are homologous to the family of surface antigens of *Toxoplasma gondii*. *Infect Immun* **65**: 5322-5328.
- Kim JH, Sohn HJ, Hwang WS, et al. (2000) In vitro isolation and characterization of bovine *Neospora caninum* in Korea. *Vet Parasitol* **90**: 147-154.
- Lindsay DS, Speer CA, Toivio-Kinnucan MA, Dubey JP, Blagburn BL (1993) Use of infected cultured cells to compare ultrastructural features of *Neospora caninum* from dogs and

- Toxoplasma gondii*. *Am J Vet Res* **54**: 103-106.
- Lunden A, Parmley SF, Bengtsson KL, Araujo FG (1997) Use of a recombinant antigen, SAG2, expressed as a glutathione-S-transferase fusion protein to immunize mice against *Toxoplasma gondii*. *Parasitol Res* **83**: 6-9.
- Marsh AE, Barr BC, Sverlow K, Ho M, Dubey JP, Conrad PA (1995) Sequence analysis and comparison of ribosomal DNA from bovine *Neospora* to similar coccidial parasites. *J Parasitol* **81**: 530-535.
- McAllister MM, Dubey JP, Lindsay DS, Jolley WR, Wills RA, McGuire AM (1998) Dogs are definitive hosts of *Neospora caninum*. *Int J Parasitol* **28**: 1473-1478.
- Nam HW, Im KS, Baek EJ, Choi WY, Cho SY (1996) Analysis of antigenic domain of GST fused major surface protein (p30) fragments of *Toxoplasma gondii*. *Korean J Parasitol* **34**: 135-141.
- Nishikawa Y, Kousaka Y, Fukumoto S, et al. (2000) Delivery of *Neospora caninum* surface protein, NcSRS2 (Nc-p43), to mouse using recombinant vaccinia virus. *Parasitol Res* **86**: 934-939.
- Nishikawa Y, Xuan X, Nagasawa H, et al. (2001) Prevention of vertical transmission of *Neospora caninum* in BALB/c mice by recombinant vaccinia virus carrying NcSRS2 gene. *Vaccine* **19**: 1710-1716.
- Pare J, Hietala SK, Thurmond MC (1995) An enzyme-linked immunosorbent assay (ELISA) for serological diagnosis of *Neospora* sp. Infection in cattle. *J Vet Diagn Invest* **7**: 353-359.
- Ramasamy R, Ramasamy M, Yasawardena S (2001) Antibodies and *Plasmodium falciparum* merozoites. *Trends Parasitol* **17**: 194-197.
- Schares G, Rauser M, Sondgen P, et al. (2000) Use of purified tachyzoite surface antigen p38 in an ELISA to diagnose bovine neosporosis. *Int J Parasitol* **30**: 1123-1130.
- Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* **76**: 4350-4354.