Case Report

Nosocomial submandibular infections with dipterous fly larvae

Chong Yoon JOO^{1)*} and Jong Bae KIM²⁾

Department of Parasitology¹⁾ and Department of Dentistry²⁾ Keimyung University
School of Medicine, Taegu, Korea

Abstract: In September 1998, a case of nosocomial cutaneous myiasis caused by *Lucilia sericata* (Meigen, 1826) in a 77-year-old male was found. The patient had been receiving partial maxillectomy due to the presence of malignant tumor on premaxilla. This is the first verified case involving *Lucilia sericata* in Taegu, Korea. In the present paper, the salient morphological features of the third instar larvae involved have been studied.

Key words: myiasis, nosocomial infection, *Lucilia sericata*, third instar larva

INTRODUCTION

Myiasis is well recognized as causing infections in humans and vertebrate animals with dipterous larvae which, at least for a certain period of time, feed on the host's dead or living tissue, liquid body substances, or ingested food (Zumpt, 1965). As noted by Beaver et al. (1984), these Diptera are medically classified into three groups according to the site of the lesions: specific, semispecific and accidental myiasis producing flies. Nosocomial myiasis, although rare, is sometimes reported in debilitated patients, some of which were diabetic (Mielke and Schlote, 1980; Smith and Clevenger, 1986; Burgess or Davies, 1991).

Some of the contributing factors include disturbed consciousness or hypoesthesia that prevent the patient's sensation of fly contact, and paralysis or immobility that prevent from fending off the fly detected (Lettau, 1991;

Daniel et al., 1994; Amitay et al., 1998). Recent findings from the United States, Puerto Rico, Canada, Honduras, Israel and others indicate that nosocomial myiasis is probably under-reported.

It is likely that hospital-acquired myiasis is under-reported for a number of reasons and one of the reasons may be related to the institutional risk management decisions. It usually occurs during summer months when fly populations are most dense, and most of the larvae are identified as *Phaenicia sericata* (Greenberg, 1984), *Lucilia sericata* (Amitay et al., 1998), *Cochliomyia macellaria* (Fox and Rodriguez-Torrens, 1972; Smith and Clevenger, 1986; Josephson and Krajden, 1993), and *Cochliomyia hominivorax* (de Kaminsky RG, 1993), all of which belong to the family Calliphoridae.

We are not aware of any previous reports with respect to opportunistic cutaneous myiasis following radiotherapy for a squamous cell carcinoma on the submandibular area. This article describes a recent case and discusses nosocomial myiasis.

Received 26 January 2001, accepted after revision 22 May 2001.

*Corresponding author (e-mail: jyjoo@dsm.or.kr)

CASE HISTORY

A 77-year-old man with partial maxillectomy due to the presence of metastatic tumor on the premaxilla was admitted to the Dongsan Hospital, Taegu, Korea, from 3 August to 18 September, 1998. After 7 weeks of radiotherapy for a 7 cm-diameter squamous cell carcinoma on the submandibular area with 7000 cGy (Siemens linear accelerator-6MeV photon), he was discharged from the hospital on 18 September, 1998.

The patient reported increasing pain from the lesion 3 days prior to presentation, and ivory-colored larvae were seen in the central portion of the lesion when he was admitted to the Dongsan Hospital in Taegu, Korea, on 20 of September, 1998.

Examination showed a 7 cm-diameter necrotic ulcer, with heaped up borders, in the submandibular area. The lesion was filled by serosanguinous fluid containing 20 fly larvae. He was treated intravenous injection with benzyl penicillin. All the necrotic tissue and the infesting larvae were removed. The larvae were 8 to 11 mm long (Fig. 1). The wound was dressed with 10% povidone iodine and paraffin gauze. A few organisms were fixed in 10% formalin and used for identification. The larva was identified by Dr. Hiromu Kurahashi at Department of Medical Entomology, National Institute of Infectious Diseases, Japan, as a third instar larva of *Lucilia sericata* (Meigen, 1826).

For scanning electron microscopic studies, specimens were washed three times in buffer solution, fixed with 3% glutaraldehyde for 2 hours in 0.1 M phosphate buffer solution, pH 7.4, and post-fixed with 1% osmium tetroxide. They were further dehydrated with a graded series of ethylalcohol and iso-amyl acetate, and critical pointed dried. The specimens were coated with gold, using Hitachi E-1030 ion sputtering apparatus, and then examined with a S-4200 scanning electron microscope at an accelerating voltage of 15 kv.

The scanning electron microscopic inspection of larvae is rewarding. The pointed end of the larva is anterior, and the broadly truncate end is posterior. The larva holds on with two sickle-shaped oral hooks on its

bulbous head (Fig. 2 and Fig. 3), and it also has rows of concentric, backward-facing, spines and hooks (Fig. 3) which facilitate burrowing and rasping of the host tissues. This explains the sensation of pain as the larva moves around (Gordon et al., 1995). The larva breathes through the posterior spiracles (Fig. 4, Fig. 5 and Fig. 6) which have three slit-like opening. Posterior spiracles with button area well chitinized and ring (peritreme) complete, button area a part of the ring; slits nearly straight; Spiracular area surrounded by 10 tubercles (Fig. 6).

DISCUSSION

Nosocomial infection is defined as the infection that is caused during or after the hospitalization which was neither present nor incubating at the time of admission (Wenzel, 1987; Barrett-Connor et al., 1978). Of the various types of myiasis, only the secondary (accidental, facultative) myiasis is potentially nosocomial. As noted by Greenburg (1984), some of the factors that frequently produce myiasis include (1) helpless and debilitated individuals — helpless individuals can't be a factor causing myiasis ..., (2) blood or odors of decomposition, (3) neglect of nursing or custodial personnel, and (4) summer season. Seventeen reported cases from various countries showed that of nosocomial myiasis were caused by maggots of facultative and obligatory parasitic species of flies, and their characteristics are summarized in Table 1.

In practice, there is no indication of 'true' myiasis in both community-acquired cases as well as nosocomial infestations, because hospital-acquired myiasis is probably under-reported for a number of reasons and some of the cases are squelched by hospital administrators, risk managers, and public relations departments for obvious medicolegal and political reasons. Such considerations were also recognized by Smith and Clevenger (1986), Lettau (1991), Josephson and Krajden (1993), and Daniel et al. (1994). The under-reporting of nosocomial myiasis is further supported by a prospective multihospital study in Brisbane, Australia, that found 14 infestations, of which six were nosocomial,

Fig. 1. Twenty larvae of *Lucilia sericata* removed from the wound. **Figs. 2-6.** Scanning electron microscopy of the larva. Anterior end with oral hooks (X 50). **3.** Oral hooks (X 180). **4.** Caudal end with posterior spiracles. Spiracular area surrounded by 10 tubercles (X 25). **5.** Posterior spiracles which have three slit-like opening (X 50). **6.** Posterior spiracles (X 180).

Table 1. The reported cases of hospital-acquired myiasis

Author	Country	Age/Sex	Diagnosis	Mental status ^{a)}	Site of larvae
Mielke & Schlote (1980)	Germany	73/NS ^{b)}	Diabetic foot infection	Alert	Foot wound
Jacobson et al. (1980)	Germany	66/NS	Diabetic foot infection	Alert	Foot wound
Magnarelli & Andreadis (1981)	United States	73/F	Heart surgery	General anesthesia	Sternal wound
Gupta et al. (1983)	United States	67/M	Heart surgery	General anesthesia	Nose
Rawlins & Barnett (1983)	United States	8/M	Encephalopathy	Comatose	Nose
Greenberg (1984)	India	25/M	Multiple stab wounds	Sedated	Urethra
	Jamaica	65/F	Cerebrovascular accident	Comatose	Nasopharynx
	United States	NS/M	Terminal illness	NS	Tracheostomy wound
	United States	NS/F	Renal failure	NS	Nose
Smith & Clevenger (1986)	United States	64/M	Diabetic hyperosmolality	Comatose	Nose
Bosniak & Schiller (1990)	United States	63/M	Diabetic	Alert	Eyelid reconstruction
Burgess & Davies (1991)	England	57/F	Diabetic	Alert	Leg ulcer
Burgess & Spraggs (1992)	England	79/M	Peripheral vascular disease	Alert	Leg ulcer
Josephson & Krajden (1993)	Canada	82/F	Myocardial infarct	Unconscious	Tracheostomy wound
de Kaminsky (1993)	Honduras	19/F	Vulvar condyloma	NS	Vulvar wound
Philips & Marsden (1993)	Honduras	2/M	Retropharyngeal abscess	NS	Nose
Chung et al. (1996)	England	73/M	Squamous cell carcinoma	NS	Left temple wound
Amitay et al. (1998)	Korea	71/M	Right facial palsy	Comatose	Oral cavity
	Israel	Infant/F	Premature	Sedated	Vaginal orifice

^{a)}At time of egg deposition by flies

^{b)}Not stated

over a period of 17 months with clustering of cases during hot weather (Lukin, 1989). He also recorded that most patients were elderly, with either diabetes or peripheral vascular disease, and had myiasis in foot or ankle wounds. The predisposing conditions included chronic ulcerations of the ankle, burns, gangrene, traumatic laceration, and surgery, including skin grafting. Chigusa et al. (1996) also indicated that patients with psychiatric disorders such as schizophrenia, as well elderly and debilitated persons, should be protected from flies, because of their autism and/or diminished sensitivity, which may make it easy for flies to deposit eggs or larvae on the patient's body surface or orifices.

The reported cases, as shown in Table 1, had similar predisposing features. All occurred during summer months and were hospital-acquired. We are not aware of any previous reports of cutaneous myiasis complicating radiotherapy in Korea. Although opportunistic cutaneous myiasis of the head is rare, a case has been reported complicating the excision of a basal cell carcinoma of the eyelid by Bosniak and Schiller (1990) in the United States, and a case of opportunistic cutaneous myiasis following radiotherapy for a squamous cell carcinoma of the left temple by Phillips and Marsden (1993) in the United Kingdom. On the basis of previously reported data and our own findings, we found, as expected, that all three cases occurred during the summer. This is when the fly population is in its greatest density. Although two previously reported cases involved *Calliphorid* larvae, our case was due to *Lucilia* larvae.

As indicated by Lettau (1991), the prevention measures of nosocomial myiasis is directly related to the flies. Sanitary and efficient waste disposals supplemented by some insecticide sprays should minimize fly density. Screens or sealed windows will provide physical barriers. Wound care and dressing, as well as attention to the hygiene of the patients will also decrease the attractions of flies to the patients. These considerations were also made by Josephson and Krajden (1993).

REFERENCES

- Amitay M, Efrat M, McGarry JW, Shinwell ES (1998) Nosocomial myiasis in an extremely premature infant caused by the sheep blowfly *Lucilia sericata*. *Pediatr Infect Dis J* **17**: 1056-1057.
- Barrett-Conner E, Brandt SL, Simon HJ, Dechairo DC (1978) Epidemiology for the infection control nurse. The CV Mosby Co., St. Louis.
- Beaver PC, Jung RC, Cupp EW (1984) Clinical parasitology. 9th ed. Lea & Febiger, Philadelphia.
- Bosniak SL, Schiller JD (1990) Ophthalmomyiasis in an eyelid reconstruction. *Am J Ophthalmol* **109**: 101-102.
- Burgess I, Davies EA (1991) Cutaneous myiasis caused by the housefly, *Musca domestica*. *Br J Dermatol* **125**: 377-379.
- Burgess I, Spraggs PD (1992) Myiasis due to *Parasarcophaga argyrostoma* — first recorded case in Britain. *Clin Exp Dermatol* **17**: 261-263.
- Chigusa Y, Kirinoki M, Yokoi H, et al. (1996) Two cases of wound myiasis due to *Lucilia sericata* and *L. illustris* (Diptera: Calliphoridae). *Med Entomol Zool* **47**: 73-76.
- Chodosh J, Clarridge J (1992) Ophthalmomyiasis: a review with special reference to *Cochliomyia hominivorax*. *Clin Infect Dis* **14**: 444-449.
- Chung PR, Jung YH, Kim KS, Cho SK, Jeong S, Ree HI (1996) A human case of internal myiasis in Korea. *Korean J Parasitol* **34**: 151-154 (in Korean).
- Daniel M, Sramova H, Zalabska E (1994) *Lucilia sericata* (Diptera: Calliphoridae) causing hospital-acquired myiasis of a traumatic wound. *J Hosp Infect* **28**: 149-152.
- de Kaminsky RG (1993) Nosocomial myiasis by *Cochliomyia hominivorax* in Honduras. *Trans Roy Soc Trop Med Hyg* **87**: 199-200.
- Fox I, Rodriquez-Torrens R (1972) Human macellaria myiasis in Puerto Rico. *Bol Assoc Med P R* **64**: 53-55.
- Gordon PM, Hepburn NC, Williams AE, Bunney MH (1995) Cutaneous myiasis due to *Dermatobia hominis*: a report of six cases. *Br J Dermatol* **132**: 811-814.
- Greenberg B (1984) Two cases of human myiasis caused by *Phaenicia sericata* (Diptera: Calliphoridae) in Chicago area hospitals. *J Med Entomol* **21**: 615.

- Gupta SC, Kumar S, Srivastava A (1983) Urethral myiasis. *Trop Geogr Med* **35**: 73-74.
- Jacobson JA, Kolts RL, Conti M, Burke JP (1980) Hospital-acquired myiasis. *Infect Control* **1**: 319-320.
- Josephson RL, Krajden S (1993) An unusual nosocomial infection: Nasotracheal myiasis. *J Otolaryngol* **22**: 46-47.
- Lettau LA (1991) Nosocomial transmission and infection control aspects of parasitic and ectoparasitic diseases. Part III. Ectoparasites/Summary and conclusions. *Infect Control Hosp Epidemiol* **12**: 179-185.
- Lukin LG (1989) Human cutaneous myiasis in Brisbane: a prospective study. *Med J Aust* **150**: 237-240.
- Magnarelli LA, Andreadis TG (1981) Human cases of furuncular, traumatic, and nasal myiasis in Connecticut. *Am J Trop Med Hyg* **30**: 984-896.
- Mielke U, Schlote A (1980) Hospital infection resulting from maggots. *Z Arztl Fortbild* **74**: 556-558.
- Phillips WG, Marsden JR (1993) Opportunistic cutaneous myiasis following radiotherapy for squamous cell carcinoma of the left temple. *Br J Dermatol* **129**: 502-503.
- Rawlins SC, Barnett DB (1983) Internal human myiasis. *West India Med J* **32**: 184-186.
- Smith DR, Clevenger RR (1986) Nosocomial nasal myiasis. *Arch Pathol Lab Med* **110**: 439-440.
- Spigel GT (1988) Opportunistic cutaneous myiasis. *Arch Dermatol* **124**: 1014-1015.
- Wenzel RP (1987) Prevention and control of nosocomial infections. Williams & Wilkins Co., Blatimore.
- Zumpt F (1965) Myiasis in man and animals in the Old World. A textbook for physicians, veterinarians and zoologists. Butterworth Co., London.