Published in final edited form as:

Bioorg Med Chem. 2009 July 15; 17(14): 5146–5152. doi:10.1016/j.bmc.2009.05.061.

Inhibition of Human Vascular NADPH Oxidase by Apocynin Derived Oligophenols

Mauricio Mora-Pale¹, Michel Weïwer², Jingjing Yu³, Robert J. Linhardt^{1,2,3}, and Jonathan S. Dordick^{1,3}

¹ Department of Chemical and Biological Engineering, Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy NY, 12180, USA

² Department of Chemistry and Chemical Biology, Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy NY, 12180, USA

³ Department of Biology, Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute, Troy NY, 12180, USA

Abstract

Enzymatic oxidation of apocynin, which may mimic *in vivo* metabolism, affords a large number of oligomers (apocynin oxidation products, AOP) that inhibit vascular NADPH oxidase. *In vitro* studies of NADPH oxidase activity were performed to identify active inhibitors, resulting in a trimer hydroxylated quinone (IIIHyQ) that inhibited NADPH oxidase with an $IC_{50} = 31$ nM. Apocynin itself possessed minimal inhibitory activity. NADPH oxidase is believed to be inhibited through prevention of the interaction between two NADPH oxidase subunits, p47^{phox} and p22^{phox}. To that end, while apocynin was unable to block the interaction of his-tagged p47^{phox} with a surface immobilized biotinylated p22^{phox} peptide, the IIIHyQ product strongly interfered with this interaction (apparent $IC_{50} = 1.6$ μ M). These results provide evidence that peroxidase-catalyzed AOP, which consist of oligomeric phenols and quinones, inhibit critical interactions that are involved in the assembly and activation of human vascular NADPH oxidase.

1. Introduction

Recent years have seen substantial improvement in our understanding of the role of superoxide anion ($\bullet O_2^-$) in eliciting oxidative stress and vascular diseases [1–5]. The production of $\bullet O_2^-$ is catalyzed by a variety of enzymes, including xanthine oxidase, cytochromes P450, lipoxygenase, enzymes in the mitochondrial respiratory chain, and NADPH oxidases [6]. The latter, in particular, have been identified as the major source of $\bullet O_2^-$ in vascular endothelial cells (VECs). Excessive production of $\bullet O_2^-$ in VECs leads to increased oxidative stress and endothelial dysfunction. This in turn can result in a diverse array of cardiovascular diseases, including atherosclerosis, hypertension, diabetes, heart failure, stroke, and restenosis [1,7–11].

The most well-studied NADPH oxidase in humans is found in neutrophils. In the resting state, the neutrophil enzyme consists of at least six partially dissociated components [12]. Tight

Correspondence authors: dordick@rpi.edu, linhar@rpi.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

regulation of NADPH oxidase activity is achieved by at least two mechanisms; the association of the cytosolic subunits and the modulation of reversible protein-protein and protein-membrane interactions [13]. Despite near 100% homology with the neutrophil NADPH oxidase [14,15], the exact assembly and activation of VEC NADPH oxidase is poorly understood [16]. Nevertheless, current evidence suggests that phosphorylation of key serine residues in p47^{phox} facilitates interaction of a Src homology 3 (SH3) domain of this protein with a proline-rich region (PRR) of p22^{phox}, thereby forming the active membrane-associated NADPH oxidase complex [17,18] (Scheme 1).

The catalytic subunit of NADPH oxidase (Nox) has several isoforms and, at least three of them are expressed in VEC (Nox1, Nox2, and Nox4). Their precise activation mechanisms and cellular regulation remain unclear [19,20]. Nox1 appears to be of only minor importance in the generation of VEC reactive oxygen species (ROS). Nox4, however, is abundantly expressed in endothelial cells, more than Nox2 [21]. Nevertheless, Li et. al. [21] showed that, despite low expression relative to Nox4, under starvation conditions Nox2 was upregulated ~ 8-fold and, as a consequence of this nutrient deprivation-induced oxidative stress, the production of •O₂⁻ increased ~2.3-fold. Importantly, Nox2 requires assembly of p47^{phox} with other cytosolic subunits prior to translocation to the membrane to form the active NADPH oxidase complex [22,23].

Due to the key role VEC NADPH oxidase appears to play in vascular diseases, identification of selective inhibitors is of great interest. Along these lines, several inhibitors have been identified, including nitrovasodilators [24], the flavonoid derivative 6,8-diallyl-5,7-dihydroxy-2-(2-allyl-3-hydroxy-4-methoxyphenyl)-1-H-benzo-[b]-pyran-4-one [25], and peptides such as the antibiotic PR-39 [26]. Interestingly, PRR regions are also known to bind polyphenols (such as flavonoids) [27]. It is not surprising, therefore, that phenolics have been found to have NADPH oxidase inhibitory activity.

Apocynin (4'-hydroxy-3'-methoxyacetophenone) [28,29] is a particularly interesting phenol that has been used as inhibitor of NADPH oxidase. While apocynin itself was found to have low activity *in vitro* [29,30] metabolism *in vivo* converts the phenol into active metabolites that inhibit the enzyme [30–34]. This may be due to peroxidase catalysis [29,30,35,36] leading to disruption of the p47^{phox}-p22^{phox} interaction, which is required for translocation of the cytosolic enzyme components to the membrane leading to activation of the enzyme complex. In the current work, we demonstrate that several oligomeric apocynin oxidation products generated by peroxidases are extremely potent inhibitors of VEC NADPH oxidase *in vitro*. Moreover, a strong correlation exists between the inhibition of VEC NADPH oxidase in endothelial cell-based assays and disruption of the interaction of EC p47^{phox}-p22^{phox} in cell-free assays. These results provide additional mechanistic insight into the nature and function of active metabolites of apocynin.

2. Results and discussion

Under conditions of oxidative stress, overactive NADPH oxidase in the vasculature generates •O₂⁻, thereby leading to increased levels of H₂O₂. In the presence of peroxidases in the blood, e.g., myeloperoxidase, and reducing substrates of peroxidases, such as phenols, peroxidatic reactions can occur. To mimic this scenario, we used a simple commercially available peroxidase from soybean (SBP) to catalyze the oxidation of apocynin in the presence of H₂O₂ following our earlier published procedure [37]. Such reaction would then be expected to mimic peroxidatic metabolism in the vasculature.

Following the enzymatic oxidation of apocynin, a water-soluble fraction was extracted into ethyl acetate to give fraction AOP-1 and a chloroform soluble precipitate was fractionated by

silica chromatography to yield nine fractions (AOP-2 to AOP-10). Each fraction was analyzed by LC-MS to qualitatively identify the AOP in each mixture, giving rise to the identification of oligomers in their demethylated, hydroxylated, or quinone forms (Table 1). The total conversion of the enzymatic reaction was ~50%; the precipitate representing 87% of the total products while 13% remained in the aqueous phase. The ability of each AOP fraction to inhibit VEC NADPH oxidase was then assessed at a dose range from 0 to 1000 μM (based on apocynin monomer unit mass), as determined by cytochrome c reduction for extracellular superoxide detection and by dihydroethidium (DHE) staining for intracellular superoxide detection.

NADPH oxidase activity - Cytochrome c reduction

The NADPH oxidase inhibitory activity of apocynin and AOP fractions was assessed in an endothelial cell-based assay by measuring the generation of $\bullet\text{O}_2^-$ via cytochrome c reduction. Apocynin itself possesses minimal inhibitory activity ($\text{IC}_{50} > 1 \text{ mM}$; Figure 1), which is consistent with reports in the literature [29,30]. However, the extracted water-soluble phase (AOP-1) exhibited an apparent IC_{50} value of 155 nM (Figure 1), despite analysis of this mixture by NMR, TLC, and LC-MS, which indicated that the major component was unreacted apocynin (~90%). Thus, the inhibition of NADPH oxidase must result from the presence of at least one very strong NADPH oxidase inhibitor present in the remaining 10% of the water-soluble fraction. Following purification via silica chromatography, a trimer hydroxylated quinone (IIIHyQ, 508 m/z, Figure 1) was identified as the major active compound in AOP-1, with strong inhibitory activity against VEC NADPH oxidase ($\text{IC}_{50} = 31 \text{ nM}$).

A similar study was performed for the chloroform-soluble enzyme reaction precipitate, which following chromatographic separation, resulted in nine distinct fractions (AOP-2 through AOP-10, IC_{50} values summarized in Table 2). Fractions AOP-2 – AOP-4 showed substantial NADPH oxidase inhibitory activity ($< 1.0 \mu\text{M}$). Fractions AOP-1 and AOP-2 consisted of IIIHyQ along with other oligomeric species. AOP-4, however, consisted of other trimeric compounds, including trimeric quinones (IIIQ, IIIHy-MeQ, and III3HyQ) that also inhibited NADPH oxidase. Higher oligomeric species (e.g., tetrameric to heptameric) showed relatively low inhibitory activity. These results demonstrate that some of the AOP possess strong VEC NADPH oxidase inhibitory activity, as reflected in the cell-based enzyme assay.

Intracellular superoxide detection – DHE staining

To establish the ability of AOP to suppress intracellular $\bullet\text{O}_2^-$ formation by NADPH oxidases, we used DHE staining of whole EC following incubation with apocynin, IIIHyQ, and phenylarsine oxide (PAO, an inhibitor of NADPH oxidase [39] and serving as a positive control). DHE is a cell-permeable reagent that reacts with $\bullet\text{O}_2^-$ to form oxyethidium [38], which in turn interacts with nucleic acids to emit a bright red color detectable qualitatively by fluorescence microscopy. Figure 2 shows images of DHE stained EC after incubation with the aforementioned compounds. In the absence of compound, the EC fluoresce as a result of strong and expected $\bullet\text{O}_2^-$ production. Apocynin, even up to 1 mM, did not appreciably reduce this fluorescence (Figure 2A). In contrast, IIIHyQ strongly reduced the $\bullet\text{O}_2^-$ induced fluorescence (Figure 2B), which was consistent with the effect of the positive control compound (PAO, Figure 2C). These results suggest that the IIIHyQ inhibits NADPH oxidase intracellularly. In addition, the inability of very high concentrations of apocynin to inhibit NADPH oxidase (as reflected in the lack of inhibition of $\bullet\text{O}_2^-$ production) also indicates that apocynin is unable to scavenge $\bullet\text{O}_2^-$.

Interaction between his-p47^{phox} protein and biotin-p22 peptide

The p47^{phox} and p22^{phox} protein subunits play a significant role in the activation of NADPH oxidase [13]. Translocation of p47^{phox} from the cytosol to bind to membrane-associated p22^{phox} is a key event in the mechanism of NADPH oxidase activation and is presumably

driven by the interaction of an SH3 domain on the p47^{phox} with a proline-rich region on the p22^{phox} [17,18]. To ascertain whether the mechanism of AOP inhibition of NADPH oxidase is due to disruption of the interaction of p47^{phox} with p22^{phox}, we performed a well-plate ELISA assay. Isolated IIIHyQ and the AOP-1 mixture were potent inhibitors of this interaction (Figure 3; Table 2) with IC₅₀ values of 1.60 μ M and 0.546 μ M, respectively. Apocynin had no effect on protein-peptide interaction.

We then evaluated the remaining AOP fractions for their ability to disrupt the interaction of p47^{phox} with p22^{phox} (Table 2). ELISA results followed a similar pattern to that observed in the cell based studies, with low activity in fractions consisting of higher oligomeric species. A plot of the log(IC₅₀) values from the cell-based assay and from ELISA (Figure 4) shows good linear correlation ($R^2 = 0.87$), suggesting that VEC NADPH oxidase inhibition is likely explained by the ability of the AOP to disrupt the interaction of p47^{phox} with p22^{phox}.

Proposed mechanism of AOP-mediated inhibition of NADPH oxidase

Apocynin was a poor inhibitor of NADPH oxidase in VEC-based assays, suggesting that previous results describing the effectiveness of apocynin as an NADPH oxidase inhibitor were a consequence of its conversion into active metabolites, likely catalyzed by peroxidases. In the present study, we produced the oligomers *in vitro*, which allowed the structural characterization of AOP. As a result, a trimer hydroxylated quinone (IIIHyQ) was identified as a strong inhibitor from AOP-1 and its structure was characterized by high resolution mass spectrometry (HRMS) and nuclear magnetic resonance (¹H-NMR and ¹³C-NMR); detailed information on HRMS and NMR is provided in the Supporting Information. However, not all the oligomers were strong inhibitors. Dimeric and trimeric AOP were more effective inhibitors of VEC NADPH oxidase than higher order oligomers identified in fractions AOP-6 to AOP-10. A possible explanation is that large oligomers are poorly soluble in aqueous media and are, therefore less capable of interacting with key subunits of VEC NADPH to inhibit assembly.

One possible mechanism of inhibition is the ability of reactive metabolites (e.g., quinones such as IIIHyQ) to form Michael adducts with the cysteine residues of p47^{phox} [40,41]. This would be consistent with the known neutrophils NADPH oxidase inhibitor, phenylarsine oxide, which binds covalently with thiol groups in the enzyme, thereby preventing assembly of protein subunits [42]. Human p47^{phox} contains four cysteine residues at positions 98, 111, 196 and 378. Indeed, Cys-196 is located in one of the two SH3 domains of p47^{phox} (SH3-N, amino acids 156–215). As indicated above, the interaction of the p47^{phox} (in complex with the other cytosolic NADPH oxidase subunits) with the membrane-associated p22^{phox} occurs through this SH3 site on p47^{phox} with a PRR on p22^{phox} [43]. Modification of Cys-196 by quinone-containing may disrupt this critical interaction. However, one cannot rule out the modification of the other three cysteine residues, which may lead to additional structural changes of p47^{phox} that could disrupt binding of the p47^{phox} to the membrane or interaction of the p47^{phox} with PRR of the p67^{phox} subunit. This is a key part of the formation of the cytosolic complex [13,44,45] and is required for translocation of p67^{phox} to the membrane [46,47].

To determine the importance of Cys on the interaction between p22^{phox} and p47^{phox} we examined the interaction of immobilized biotin-p22 (2 μ M) with his-p47^{phox} (0 – 1.0 μ M) in the presence of iodoacetamide (IAA) (Figure 5), which reacts with thiol groups on cysteine residues. The presence of 1 μ M IAA results in a > 50% drop in the amount of bound p47 to the immobilized biotin-p22^{phox}; hence, thiol reactivity of quinone-based AOPs may be a critical mechanism of inhibition of VEC NADPH oxidase assembly.

In summary, we have demonstrated that peroxidase-generated apocynin metabolites serve as strong inhibitors of VEC NADPH oxidase. The IIIHyQ was a particularly strong (nanomolar) inhibitor of the enzyme as determined through EC-based assays. This compound was also

effective at disrupting the p47^{phox}-p22^{phox} interaction *in vitro*. This suggests that the mechanism of apocynin inhibition of NADPH oxidase is a result of peroxidase metabolism to yield reactive quinones that bind to Cys residues in p47^{phox} and disrupt translocation to the membrane through SH3-PRR association with the p22^{phox}. Since peroxidases are common enzymes in the vasculature (e.g., myeloperoxidases) one may anticipate that similar metabolites are generated *in vivo*. Further work is needed to determine whether these metabolites provide a means to the generation of potential therapeutic molecules.

3. Experimental procedures

Apocynin, SBP, solvents, H₂O₂, LDL, superoxide dismutase (SOD), low density lipoprotein (LDL), cytochrome c, Tween 20, 3,3',5,5'-tetramethyl-benzidine (TMB), sodium caseinate, fetal bovine serum, heparin, and endothelial growth supplement were purchased from Sigma-Aldrich. Endothelial cells and medium were purchased from ATCC. *E. coli* BL21 (DE3), IPTG and Ni-affinity column (Probond system) were purchased from Invitrogen. Antibodies were purchased from Upstate. High-affinity streptavidin-coated-96 well plates were purchased from Pierce. LC-MS analyses were performed in a Shimadzu LCMS-2010A. Samples for LC-MS were separated in an Agilent Zorbax 300SB-C₁₈ column (5 μ m, 2.1 $\times$ 150 mm). Silica gel 230–400 mesh was purchased from Natland International Corporation. Thin layer chromatography (TLC) plates were purchased from Merck. Microplate reader analyses were performed in a Perkin-Elmer, HTS 7000, Bio Assay Reader.

Enzymatic production of Apocynin Oxidation Products (AOP)

AOP were synthesized by soybean peroxidase (SBP)-catalyzed oxidation of apocynin, as described by Antoniotti et. al. [37]. Briefly, apocynin (6 mmol) was dissolved in 5 mL of dimethylformamide (DMF) and transferred to 490 mL phosphate buffer (50 mM, pH 7). SBP (5 mL of a 1mg/mL solution) was added and the reaction was initiated by using a syringe pump to introduce H₂O₂ (30% w/v) at 0.1 mL/min for 12 min to afford 12 mmol H₂O₂. Finally, the reaction was stopped after 2 h. Soluble and precipitated phases were separated by centrifugation and ethyl acetate was added to supernatant to extract organic compounds (AOP-1). Both precipitated and extracted supernatant fractions were dried and stored at –20°C under argon.

Fractionation of non-soluble phase

The precipitated fraction (60 mg) was dissolved in the minimum amount of chloroform and loaded onto a silica gel column (flash chromatography, 4 g silica gel 230–400 mesh, Natland International Corporation) and eluted with a gradient of petroleum ether:ethyl acetate 2:1 to 0:1. Nine fractions (AOP-2 to AOP-10) were collected and analyzed by LC-MS (Shimadzu LCMS-2010A) using an Agilent Zorbax 300SB-C₁₈ column, 5 μ m, 2.1 $\times$ 150 mm with isocratic elution (MeCN:H₂O, 3:7; 0.2 mL/min).

Purification of the trimer hydroxylated quinone (IIIHyQ)

AOP-1 (290 mg) was dissolved in chloroform and loaded on a silica gel column (15 g) and eluted with a gradient of petroleum ether:ethyl acetate (2:1 to 0:1). Unmodified apocynin was recovered in the early fractions (210 mg, R_f 0.62 with petroleum ether:ethyl acetate, 1:1) and further elution with pure ethyl acetate furnished the IIIHyQ as a white powder (14 mg, R_f 0.34 with petroleum ether:ethyl acetate, 1:1). **HRMS** *m/z*, calculated for C₂₇H₂₅O₁₀ [M+H]⁺ 509.1442, found 509.1442. Calculated for C₂₇H₂₄O₁₀Na [M+Na]⁺ 531.1258, found 531.1261. Calculated for C₂₇H₂₄O₁₀K [M+K]⁺ 547.1001, found 547.0997. **¹H-NMR** (500 MHz, CDCl₃) 7.65 (1H, d, *J* = 1.5 Hz), 7.57 (1H, d, *J* = 1.8 Hz), 7.40 (1H, d, *J* = 1.5 Hz), 7.20 (1H, d, *J* = 1.5 Hz), 6.07 (1H, s), 4.05 (1H, s), 3.99 (3H, s), 3.98 (3H, s), 3.69 (3H, s), 2.57 (3H, s), 2.44 (3H, s), 2.18 (3H, s). **¹³C-NMR** (125 MHz, CDCl₃) 201.22, 196.02, 195.63, 195.46, 153.08, 148.88, 145.82, 144.94, 133.14, 132.85, 123.70, 121.53, 120.41, 119.30, 113.34,

111.62, 98.31, 90.18, 63.13, 62.59, 62.16, 56.40, 56.23, 53.21, 26.48, 26.29, 23.65. ^1H -NMR and ^{13}C -NMR spectra were recorded at room temperature, in CDCl_3 (Varian 500 MHz or Bruker 600 MHz and 800 MHz). Chemical shifts (δ) are indicated in ppm and coupling constants (J) in Hz. Flash chromatography was performed using silica gel 230–400 mesh (Natland International Corporation). Thin-layer chromatography (TLC) was carried out using Merck plates of silica gel 60 with fluorescent indicator and revealed with UV light (254 nm) and 5% H_2SO_4 in EtOH.

Endothelial cell culture

Human umbilical vascular endothelial cells (HUVEC; ATCC) were cultured in F-12K medium supplemented with 10% (v/v) fetal bovine serum, heparin (0.1 mg/mL), and endothelial growth supplement (0.04 mg/mL). The medium was replenished every two days until confluence was achieved. The cells were propagated by detaching them with 0.25% (w/v) trypsin - 0.53 mM EDTA solution, adding 8 mL of F-12K medium, and centrifuging, and then the cells were sub-cultured in new culture vessels until the desired number of cells was obtained.

NADPH oxidase activity - Cytochrome c reduction

The inhibitory effect of AOP on NADPH oxidase was assessed by the inhibition of $\bullet\text{O}_2^-$ generated by VEC and measured by reduction of cytochrome c [16]. Cells were resuspended in DMEM without phenol red and incubated in 96-well flat bottom culture plates (10^5 cells/mL) for 10 min at 37°C in a humidified CO_2 incubator. Low density lipoprotein (100 $\mu\text{g/mL}$) was used to induce activation of NADPH oxidase. AOP were incubated at concentrations ranging from 0 to 1000 μM in the presence of 100 μM NADPH, with or without superoxide dismutase (SOD, 200 $\mu\text{g/mL}$), and in the presence of cytochrome c (250 μM) for 30 min at room temperature. Cytochrome c reduction was measured by reading absorbance at 550 nm in a microplate reader. Inhibition of NADPH oxidase was calculated from the difference between the absorbance of sample with or without SOD and the extinction coefficient for the change of oxidized cytochrome c to reduced cytochrome c ($18.7 \text{ cm}^{-1}\text{mM}^{-1}$); experiments were performed in triplicate.

Intracellular superoxide production – DHE staining

Endothelial cells were incubated in black, clear-bottom 96-well cell binding surface plates and incubated with apocynin, IIIHyQ, or PAO at concentration ranging from 0–1 mM for 2 h in DMEM in the absence of phenol red. DMEM was removed and the cells were washed twice with PBS and then resuspended in fresh DMEM. NADPH oxidase was activated with phorbol myristate acetate (PMA, 1 μM) for 30 min and the cells were then incubated with DHE (3 μM) for 30 min and then NADPH (100 μM) to generate $\bullet\text{O}_2^-$; the experiment was performed in the dark. Cell images were captured after 30 min with a Zeiss LSM 510 laser scanning confocal microscope at excitation and emission wavelengths of 520 and 610 nm, respectively.

Production and purification of his-p47^{phox} and biotin-p22

A proline-rich p22^{phox} peptide N'-151PPSNPPPRPPAEARK165-C', which was biotinylated at the N-terminus and amidated at the C-terminus was obtained from Genemed Synthesis Inc. (South San Francisco, CA). The biotin group was attached through a 4-residue spacer consisting of SGSG. The purity of the peptide was 99.99%. Endothelial cell derived p47^{phox} DNA (6 His-tagged) was obtained from SUNY Albany and Stratton VA Medical Center and confirmed by DNA sequence analysis (U. of Maine). His-p47^{phox} protein was expressed in BL21 (DE3) cells for 9 h using 0.5 mM isopropyl- β -D-thiogalactopyranoside (IPTG) at 35°C . The protein was purified using a Ni-affinity column (ProBond System) and confirmed by western blot analysis with anti p47^{phox} antibody and the purity (80%) was calculated with the

Image J software (NIH, USA, public domain) based on the intensity of each protein band on the electrophoresis gel.

Biotin-p22 and his-p47^{phox} interaction

Interaction of p47^{phox} with the p22^{phox} peptide was studied using ELISA, which was modified from the technique reported by Dahan et al. [48]. Experiments were performed in high-affinity streptavidin-coated-96 well plates. To block non-specific binding sites, each well was re-blocked with 300 μ L of PBS supplemented with 0.1% (v/v) Tween 20 and 1% sodium caseinate. To each well, 100 μ L of 2 μ M biotin-p22^{phox} peptide solution were added and incubated at room temperature for 1 h. After washing each well four-times with 300 μ L PBS-Tween solution, 100 μ L of 0.30 μ M his-p47^{phox} (in PBS-Tween solution containing 1% sodium caseinate) and AOP (0 – 1000 μ M) were added to each well and incubated at room temperature for 1 h. Unbound components were removed by washing four times with 300 μ L/well PBS-Tween solution. The amounts of bound his-p47^{phox} were quantified by adding 100 μ L/well of polyclonal goat anti-p47^{phox} (diluted 1:2000 in PBS-Tween solution containing 1% sodium caseinate) and incubating at room temperature for 1 h. Each well was washed four times with 300 μ L PBS-Tween solution and incubated with 100 μ L/well of HRP-conjugated rabbit anti-goat IgG secondary antibody (diluted 1:10000 in PBS-Tween solution containing 1% sodium caseinate) at room temperature for 1 h. The plate was finally washed four times with 300 μ L/well of PBS-Tween solution and two additional washes with 300 μ L/well of PBS. Detection of peroxidase activity was performed with a ready-to-use TMB liquid substrate by adding 200 μ L/well and incubating at room temperature for 30 min. The reaction was terminated by adding 100 μ L/well of 0.5 M H₂SO₄ solution and the absorbance was read at 450 nm in the microplate reader. All experiments were performed in triplicate and results were quantified from a standard curve of the interaction between biotin-p22^{phox} (2 μ M) and his-p47^{phox} (0 to 0.40 μ M).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to NIH (AT002115) for the financial support of this project and to Dr. Christopher Bjornsson, Director of Microscopy and Imaging Core Facility (Center for Biotechnology and Interdisciplinary Studies, Rensselaer Polytechnic Institute), for his help with the confocal fluorescent microscope analysis.

References

1. Li J, Mullen A, Shah A. J Mol Cell Cardiol 2001;33:1119–1131. [PubMed: 11444917]
2. Touyz RM. Hypertension 2004;44:248–252. [PubMed: 15262903]
3. Li Q, Subbulakshmi V, Fields AP, Murray NR, Cathcart MK. J Biol Chem 1999;274(6):3764–3771. [PubMed: 9920929]
4. Walch L, Massade L, Dufilho M, Brunet A, Rendu F. Atherosclerosis 2006;187:285–291. [PubMed: 16249002]
5. Holland JA, Ziegler LM, Meyer JW. J Cell Physiol 1996;166:144–151. [PubMed: 8557764]
6. Lassegue B, Clemens RE. Am J Physiol Regul Integr Comp Physiol 2003;285:277–297.
7. Babior BM. Blood 1999;93(5):1464–1476. [PubMed: 10029572]
8. Finkel T. Curr Opin Cell Biol 2003;15:247–254. [PubMed: 12648682]
9. Meng TC, Fukada T, Tonks NK. Mol Cell 2002;9:387–399. [PubMed: 11864611]
10. Taniyama Y, Griendling KK. Hypertension 2003;42:1075–1081. [PubMed: 14581295]
11. Meyer JW, Schmitt ME. FEBS Lett 2000;472:1–4. [PubMed: 10781793]
12. Seguchi H, Kobayashi T. J Electron Microsc 2002;51(2):87–91.

13. Groemping Y, Rittinger K. *Biochem J* 2005;386:401–416. [PubMed: 15588255]
14. Gorlach A, Brandes RP, Nguyen K, Amidi M, Dehghani F, Busse R. *Circ Res* 2000;87:26–32. [PubMed: 10884368]
15. Li J, Shah AM. *Cardiovasc Res* 2001;52:477–486. [PubMed: 11738065]
16. Li JM, Shah AM. *J Biol Chem* 2002;277(22):19952–19960. [PubMed: 11893732]
17. Hiroaki H, Ago T, Ito T, Sumimoto H, Kohda D. *Nature Struct Biol* 2001;8(6):526–530. [PubMed: 11373621]
18. Sumimoto H, Kage Y, Nunoi H, Sasaki H, Nose T, Fukumari Y, Ohno M, Minakami S, Takeshige K. *Proc Natl Acad Sci* 1994;91:5345–5349. [PubMed: 8202490]
19. Martyn KD, Frederick LM, von Loehneysen K, Dinauer MC, Knaus UG. *Cell Signalling* 2006;18:69–82. [PubMed: 15927447]
20. Kuroda J, Nakagawa K, Yamasaki T, Nakamura K, Takeya R, Kuribayashi F, Imajoh-Ohmi S, Igarashi K, Shibata Y, Sueishi K, Sumimoto H. *Genes to Cells* 2005;10:1139–1151. [PubMed: 16324151]
21. Li J, Fan LM, George VT, Brooks G. *Free Radical Biol* 2007;43:976–986.
22. Taura M, Miyano K, Miakaru R, Kamabura S, Takeya R, Sumimoto H. *Biochem J*. 2009Epub ahead of print
23. Frey RS, Ushio-Fukai M, Malik A. *Antioxid Redox Signal*. 2008Ahead of print
24. Selemidis S, Dusting GJ, Peshavariya H, Kemp-Harper BK, Drummond GR. *Cardiovasc Res* 2007;75:349–358. [PubMed: 17568572]
25. Cayatte AJ, Rupin A, Oliver-Krasinski J, Maitland K, Sansilvestri-Morel P, Boussard M, Wierzbicki M, Verbeuren TJ, Cohen RA. *Arterioscler Thromb Vasc Biol* 2001;21:1577–1584.
26. Shi J, Ross CR, Leto TL, Blecha F. *Proc Natl Acad Sci* 1996;93:6014–6018. [PubMed: 8650211]
27. Kanegae MPP, da Fonseca LM, Brunetti IL, de Oliveira Silva S, Ximenes VF. *Biochem Pharmacol* 2007;74:457–464. [PubMed: 17544376]
28. Yu J, Weiwer M, Linhardt R, Dordick JS. *Curr Vasc Pharmacol* 2008;6:1–14. [PubMed: 18220934]
29. Johnson DK, Schillinger KJ, Kwait DM, Hughes CV, McNamara EJ, Ishmael F, O'Donnell RW, Chang M, Hogg MG, Dordick JS, Santhanam L, Ziegler LM, Holland JA. *Endothelium* 2002;9:191–203. [PubMed: 12380644]
30. Heumüller S, Wind S, Barbosa-Sicard E, Schmidt HHHW, Busse R, Schröder K, Brandes RP. *Hypertension* 2008;51:211–217. [PubMed: 18086956]
31. Vejraska M, Micek R, Stipek S. *Biochim Biophys Acta* 2005;1722:143–147. [PubMed: 15716123]
32. Dodd OJM, Pearse DB. *Am J Physiol Heart Circ Physiol* 2000;279:303–312.
33. Muijers RBR, van den Worm E, Folkerts G, Beukelman CJ, Koster AS, Postma DS, Nijkamp FP. *Brit J Pharmacol* 2000;130:932–936. [PubMed: 10864902]
34. Chan EC, Datla SR, Dilley R, Hickey H, Drummond GR, Dusting GJ. *Cardiovasc Res* 2007;75:710–718. [PubMed: 17659266]
35. Ximenes VF, Kanegae MPP, Rissato SR, Galhiane MS. *Arch Biochem Biophys* 2007;457:134–141. [PubMed: 17166480]
36. Stolk J, Hiltermann TJ, Dijkman JH, Verhoeven AJ. *Am J Respir Cell Mol Biol* 1994;11(1):95–102. [PubMed: 8018341]
37. Antoniotti S, Santhanam L, Ahuja D, Hogg M, Dordick JS. *Org Lett* 2004;6(12):1975–1978. [PubMed: 15176797]
38. Bendall JK, Rinze R, Adlam D, Tatham AL, de Bono J, Channon KM. *Circ Res* 2007;100:1016–1025. [PubMed: 17363703]
39. Doussiere J, Poinas A, Blais C, Vignais P. *Eur J Biochem* 1998;251:649–658. [PubMed: 9490037]
40. 't Hart BA, Simons JM, Rijkers GT, Hoogvliet JC, Van Dijk H, Labadie RP. *Free Radical Bio Med* 1990;8:241–249. [PubMed: 2341054]
41. Simons JM, 't Hart BA, Vai Ching Tvan Dijk H, Labadie RP. *Free Rad Biol Med* 1990;8(3):251–258. [PubMed: 2160411]
42. Le Cabec V, Maridonneau-Parini I. *J Biol Chem* 1995;270(5):2067–2073. [PubMed: 7530716]
43. Leto TL, Adams AG, De Mendez I. *Proc Natl Acad Sci* 1994;91:10650–10654. [PubMed: 7938008]

44. Takeya R, Ueno N, Kami K, Taura M, Kohjima M, Izaki T, Nunoi H, Sumimoto H. *J Biol Chem* 2003;278(27):25234–25246. [PubMed: 12716910]
45. Lapouge K, Smith SJM, Groemping Y, Rittinger K. *J Biol Chem* 2002;277(12):10121–10128. [PubMed: 11796733]
46. Morozov I, Lotan O, Joseph G, Gorzalczany Y, Pick E. *J Biol Chem* 1998;273(25):15435–15444. [PubMed: 9624128]
47. Park J. *Biochem Biophys Res Commun* 1996;220:31–35. [PubMed: 8602852]
48. Dahan I, Issaeva I, Gorzalczany Y, Sigal N, Hirshberg M. *J Biol Chem* 2002;277(10):8421–8432. [PubMed: 11733522]

Trimer Quinone Hydroxylated IIIHyQ
Chemical Formula: $C_{27}H_{24}O_{10}$
Exact Mass: 508.1369

Apocynin

Chemical Formula: $C_9H_{10}O_3$
Exact Mass: 166.0630
Molecular Weight: 166.1739

Figure 1.

Effect of apocynin, IIIHyQ, and AOP-1 on NADPH oxidase activity. Experiments were performed by reduction of cytochrome c using EC in a 96-well plate with AOP concentrations of 0 – 1 mM.

Figure 2. Intracellular detection of $\bullet\text{O}_2^-$ by DHE staining in EC. Cells were incubated with (A) Apocynin (B) IIIHyQ and (C) PAO. Dose range: 0 – 1mM. Images were taken by confocal microscopy.

Figure 3. Effect of (A) isolated IIIHyQ, (B) AOP-2 and (C) apocynin, on biotin-p22^{phox}-his-p47^{phox} interaction by ELISA. Biotin- p22^{phox} = 2.0 μM, his- p47^{phox} = 0.3 μM.

Figure 4. Correlation of experimental log (IC₅₀) values of cell-based and ELISA assays.

Figure 5. Interaction between biotin-p22^{phox} (2.0 μM) and his-p47^{phox} (0 – 1.0 μM) in the absence and presence of IAA (1.0 μM).

Scheme 1.

(A) Active complex of NADPH oxidase. Cytosolic subunits (p47^{phox}, p67^{phox}, p40^{phox} and Rac 1 or 2 translocate to the membrane to bind p22^{phox} and catalytic gp91^{phox}. (B) Protein complexes (in the cytosol and membrane) occur by SH3 domain interactions with PRR [7], [13].

Table 1

AOP present in soluble and precipitate fractions produced from the peroxidatic oxidation of apocynin based on LC-MS analysis. Nomenclature: A (apocynin), I (monomer), II (dimer), III (trimer), IV (tetramer), V (pentamer), VI (hexamer), VII (heptamer), Q (quinone), Hy (hydroxylated), -Me (demethylated)

Fraction	Components
AOP-1	A (166m/z); II or IIHy-MeQ (330m/z); IIIHyQ (508m/z); III2Hy or III3HyQ (526m/z); IV-3MeQ (614m/z); IV or IVHy-MeQ (658m/z)
AOP-2	A (166m/z); III3OH (542m/z); II or IIHy-MeQ (330m/z); III4Hy-Me (530m/z); IIIHyQ (508m/z); III-Me (478m/z); IIIHy (480m/z); III-3Me (450m/z); IV-Me (644m/z); III-2MeQ (464m/z); III-3Me (500m/z); IV-Me (692m/z)
AOP-3	II or IIHy-MeQ (330m/z); III or IIIHy-MeQ (494m/z); III-Me or IIIHy-2MeQ (480m/z); IIIHyQ (508m/z); IVHy (674m/z); IV3Hy (706m/z); IV3Hy-2Me (678m/z); IV-2MeQ (628m/z)
AOP-4	A (166m/z); IIIHy (510m/z); II or IIHy-MeQ (330m/z); III3Hy (542m/z); IVHy (674m/z); III2Hy or III3HyQ (526m/z); IIIQ (492m/z); III or IIIHy-MeQ (494m/z)
AOP-5	A (166m/z); II or IIHy-MeQ (330m/z); IV (658m/z); III3Hy (542m/z); III-2Me (466m/z); III2Hy or III3HyQ (526m/z); IV4Hy-3Me (680m/z); III or IIIHy-MeQ (494m/z); IV or IVHy-MeQ (658m/z); IIIQ (492m/z); III-3Me (500m/z); VHy-MeQ (822m/z)
AOP-6	II or IIHy-MeQ (330m/z); III or IIIHy-MeQ (494m/z); IV2Hy (690m/z); III-3Me (452m/z); IV4Hy-3Me (680m/z); IV or IVHy-MeQ (658m/z); IV-3Me (616m/z); V4Hy-3Me (844m/z); VII4Hy-2Me (1186m/z) V or VHy-MeQ (822m/z)
AOP-7	III3Hy (500m/z); II or IIHy-MeQ (330m/z); III or IIIHy-MeQ (494m/z); VI5Hy-4Me (1010m/z); IIQ (328m/z); IV2Hy (690m/z); IV4Hy-3Me (680m/z); VI4Hy-3Me (1008m/z); IV or IVHy-MeQ (658m/z);
AOP-8	II or IIHy-MeQ (330m/z); IV5Hy-4Me (682m/z); III3Hy-2Me (514m/z); IVHy (674m/z); IV or IVHy-MeQ (658m/z); VI4Hy-4Me (994m/z)
AOP-9	II or IIHy-MeQ (330m/z); IV5Hy-4Me (682m/z); IVHy-Me (660m/z); IVHy-4Me, IVHyQ (672m/z); VII3Hy-2Me (1170m/z); IV2Hy-3Me (648m/z); III2Hy (526m/z); IV2Hy or IV3Hy-MeQ (690m/z); III or IIIHy-MeQ (494m/z); IV4Hy-3Me (680m/z); IV or IVHy-MeQ (658m/z); VHy (838m/z)
AOP-10	II or IIHy-MeQ (330m/z); IV5Hy-4Me (682m/z); IVHyQ (672m/z); IVHy-4Me (618m/z); V-2Me (794m/z); IV2Hy-2Me (662m/z); VHy (838m/z); IV2Hy or IV3Hy-MeQ (690m/z); III or IIIHy-MeQ (494m/z); IV4Hy-3Me (680m/z); III-Me or IIIHy-2MeQ (480m/z); IV-Me or IVHy-2MeQ (644m/z); IHy-Me (168m/z); IV4Hy-Me (694m/z)

Table 2

Dose response IC₅₀ values of AOP fractions following peroxidase-catalyzed oxidation of apocynin and silica gel chromatography.

Fraction	Log (IC ₅₀) (IC ₅₀ in parentheses, μ M) Cell-based assay	Log (IC ₅₀) (IC ₅₀ in parentheses, μ M) ELISA assay
Apocynin	3.04 $\pm$ 0.49 (1090)	4.78 $\pm$ 6.65 (61,000)
IIIHyQ	-1.51 $\pm$ 0.05 (0.03)	0.20 $\pm$ 0.15 (1.60)
AOP-2	-1.40 $\pm$ 0.05 (0.04)	-0.27 $\pm$ 0.15 (0.55)
AOP-3	-0.61 $\pm$ 0.08 (0.25)	1.70 $\pm$ 0.15 (50.0)
AOP-4	-0.67 $\pm$ 0.10 (0.22)	0.47 $\pm$ 0.18 (2.96)
AOP-5	0.34 $\pm$ 0.11 (2.20)	0.44 $\pm$ 0.90 (7.96)
AOP-6	-0.09 $\pm$ 0.09 (0.82)	2.01 $\pm$ 0.16 (103)
AOP-7	0.001 $\pm$ 0.06 (1.00)	2.66 $\pm$ 456 (453)
AOP-8	0.03 $\pm$ 0.22 (1.08)	2.38 $\pm$ 0.08 (239)
AOP-9	-0.04 $\pm$ 0.13 (0.92)	1.54 $\pm$ 0.08 (34.8)
AOP-10	-0.11 $\pm$ 0.12 (0.78)	2.03 $\pm$ 0.08 (107)