Published in final edited form as:

J Org Chem. 2009 March 20; 74(6): 2613–2615. doi:10.1021/jo802632v.

Steric Modulation of Chiral Biaryl Diamines via Pd-Catalyzed Directed C–H Arylation

Christopher C. Scarborough, Richard I. McDonald, Caroline Hartmann, Graham T. Sazama, Ana Bergant, and Shannon S. Stahl

Department of Chemistry, University of Wisconsin-Madison, Madison, Wisconsin 53706

Abstract

Palladium-catalyzed directed arylation of 2,2'-diacetamidobiaryls with aryl iodides provides efficient access to chiral *ortho*-substituted biaryl diamines. Aryl iodides with *para*- and *meta*-substituents are tolerated. Deprotection of the acetyl groups under basic conditions furnishes the free diamines, which should find broad utility in asymmetric catalysis.

Chiral biaryl diamines (Chart 1)¹ find widespread use in asymmetric catalysis, ranging from oxidations^{2–5} and reductions^{6–8} to addition reactions.^{9–19} Derivatives of these diamines have been employed as ligands for transition-metal catalysts and directly as non-metal-based catalysts, for example, as chiral nucleophilic catalysts. In several cases, enantioselectivities of reactions with DABN and H₈-DABN-based catalysts (cf. Chart 1) can be increased substantially by introducing substituents in the 3- and 3'-positions of the aromatic rings (i.e., *ortho* to the amino groups).^{6a,14b,17–19} Here, we describe a Pd-catalyzed C–H arylation method that provides efficient access to *ortho*-functionalized biaryl diamine derivatives.

Recent studies in our lab have employed biphenyl diamines as a scaffold for the development of axially-chiral, seven-membered *N*-heterocyclic carbenes (**A**).²⁰ In the course of this work, we sought access to derivatives with bulky substituents in the 3- and 3'-positions of biphenyl diamine derivative **1** (Chart 1). Existing methods for the introduction of substituents in these positions are based on Suzuki-Miyaura coupling and require preparation of the corresponding halogenated biaryl diamines.^{6a,14b,17–19} The synthetic inefficiency and low selectivity that we encountered with the halogenation procedures prompted us to consider alternative strategies for *ortho*-functionalization of biphenyl diamines.

Correspondence to: Shannon S. Stahl.
stahl@chem.wisc.edu.

Several methods have been reported recently for arylation of aromatic C–H bonds, directed by an adjacent functional group.²¹ Daugulis and Zaitsev, for example, have shown that anilides undergo arylation selectively in the *ortho* position(s) with aryl iodides in the presence of stoichiometric AgOAc and catalytic Pd(OAc)₂ in trifluoroacetic acid.^{21a,22} These methods are attractive because they employ aryl iodides, which are more-readily available than the diaryliodonium salts^{21c} and trialkoxylarylsilanes^{21d} employed in other C–H arylation methods. We first attempted a Daugulis-Zaitsev (DZ) coupling reaction with iodobenzene and 2,2'-diacetylamino-biphenyl (**2**).²³ This reaction was successful and afforded the quaterphenyldiamine **3** in 70% isolated yield (eq 1). Extension of this reactivity to *chiral* biaryldiamines was also successful. The *N,N'*-diacetyl derivative of diamine **1** is a highly effective substrate for this reaction (Scheme 1); the C–H arylation product **5a** is obtained in essentially quantitative yield from the DZ coupling reaction (Scheme 1). Removal of the acetyl directing groups under basic conditions (1:1 EtOH:50% KOH_(aq)) in a stainless steel pressure vessel, 150 °C, 48h), affords the quaterphenyl diamine **6** in very good overall yield.

(1)

In order to probe the scope of the DZ coupling conditions for the synthesis of *ortho*-arylated biphenyldiamines, we examined other iodoarene substrates (Table 1). A number of the reactions proceed in good-to-excellent yields (entries 1–3, 5, 6). Limitations include strongly electron-withdrawing groups and the presence of *ortho*-substitution in the aryl iodide (entries 4 and 7). The presence of *meta*-substituents are tolerated, although somewhat more-forcing conditions were required in these reactions (20% Pd loading, 3.0 equiv AgOAc, 14h, entries 5 & 6).

Both the DZ coupling and amide hydrolysis conditions require rather high temperatures, and it was unclear if these conditions could lead to epimerization of the atropisomeric biaryls. Enantiomerically resolved (*S*)-**1** (>99% e.e.) was subjected to the diacetylation, DZ coupling and hydrolysis conditions in Scheme 1. The expected quaterphenyl diamine (*S*)-**5a** was obtained in 94% overall yield, and chiral HPLC analysis demonstrated that the product retained >99% e.e. (eq 2). Therefore, these conditions are well suited for the preparation of enantiomerically resolved diamines.

(2)

We have briefly investigated whether the methods described here can be applied to other common biaryl diamines. DABN is an attractive biaryl diamine substrate because the resolved enantiomers are commercially available. Pd-catalyzed ortho arylation of *N,N'*-diacetyl-DABN with PhI was successful, and the desired diamine **7** was obtained in 86% yield, following removal of the acetyl groups (eq 3).²⁴ The partially hydrogenated DABN derivative H₈-DABN²⁵ also undergoes effective arylation: DZ coupling of the *N,N'*-diacetyl derivative of H₈-DABN with PhI, followed by amide hydrolysis, afforded quaterphenyldiamine **8** in nearly quantitative yield (eq 4).^{17,18,19}

(3)

(4)

In conclusion, we have developed a simple and efficient three-step procedure for the installation of *ortho*-aryl groups onto 2,2'-diaminobiaryls. This procedure takes advantage of the Daugulis-Zaitsev coupling method for amide-directed arylation of aromatic C–H bonds. The incorporation of bulky *ortho*-substituents onto BINOL has had dramatic beneficial effects in asymmetric catalysis.²⁶ We anticipate the methods described here will provide the basis for similar advances in asymmetric reactions with biaryl diamine-derived catalysts.

Experimental Section

General procedure for the Daugulis-Zaitsev coupling

A representative example is given for the synthesis of **5a**. Bisamide **4** (296 mg, 1.0 mmol), Pd(OAc)₂ (34 mg, 0.15 mmol), AgOAc (384 mg, 2.3 mmol), and PhI (560 μ L, 5.0 mmol) were dissolved in 3 mL trifluoroacetic acid. A reflux condenser was attached, and the reaction was heated to a vigorous reflux for 6 h. After cooling to room temperature, 10 mL toluene was added to the dark mixture, which was then filtered over Celite, and the solvent removed under vacuum. The mixture was purified by flash chromatography with silica gel (gradient, 1:1 Et₂O/toluene to pure Et₂O, *R_f* = 0.29 in 1:1 Et₂O/toluene) to afford 446 mg off-white powder, 99.6% yield. ¹H NMR (300 MHz, CDCl₃, δ) 1.60 (s, 6H), 1.94 (s, 6H), 7.26–7.40 (m, 14H). ¹³C NMR (75 MHz, CDCl₃, δ) 20.1, 23.0, 127.3, 128.4, 128.8, 129.4, 129.8, 132.8, 138.8, 140.2. HRMS: *m/z* (ESI) calculated [MH]⁺ = 449.2224, measured 449.2242 (Δ = 4.0 ppm).

General procedure for the deacetylation of 2,2'-diacetamidobiaryls

A representative example is given for the synthesis of **6. 5a** (1.19g, 2.65 mmol) was added to 100 mL of a 1:1 mixture of EtOH and 50% KOH_(aq) in a stainless steel Parr bomb. The bomb was sealed and heated to 150 °C (based on an internal temperature probe) for 48 h. After cooling to room temperature, 100 mL saturated NH₄Cl_(aq) was added to the suspension, which was then extracted 2 $\times$ with CH₂Cl₂. The combined CH₂Cl₂ solutions were dried over MgSO₄, filtered, and the solvent was removed in vacuo to afford 905 mg pure **6** (94% yield) without further purification. ¹H NMR (300 MHz, CDCl₃, δ) 2.04 (s, 6H), 3.67 (br. s, 4H), 6.79 (dd, *J* = 7.5, 0.3 Hz, 2H), 7.07 (d, *J* = 7.8 Hz, 2H), 7.30 (tt, *J* = 7.5, 1.5 Hz, 2H), 7.39–7.44 (m, 4H), 7.46–7.50 (m, 4H). ¹³C NMR (75 MHz, CDCl₃, δ) 19.9, 120.1, 122.7, 125.5, 127.1, 128.9, 129.4, 129.7, 137.3, 140.2, 141.5. (**S**)-**6**: [α]_D²⁵ –81.4 (c 8.9, CH₂Cl₂). HRMS: *m/z* (ESI) calculated [MH]⁺ = 365.2013, measured 365.2027 (Δ = 3.8 ppm).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We thank Sigma-Aldrich and Amgen for resolving ($\pm$)-2,2'-diamino-6,6'-dimethylbiphenyl (**1**). This work was supported by the NIH (R01 GM67163) and a UW-Madison Draper Technology Innovation Fund Grant. CCS is grateful to the ACS Division of Organic Chemistry for a graduate research fellowship.

References

1. 2,2'-Diamino-1,1'-binaphthyl (DABN) is commercially available. For information regarding the synthesis of diamines in Chart 1, see: (a) Takasaki M, Motoyama Y, Yoon SH, Mochida I, Nagashima H. *J. Org. Chem* 2007;72:10291–10293.10293 [PubMed: 18044926] (b) Korostylev A, Tararov VI, Fischer C, Monsees A, Borner A. *J. Org. Chem* 2004;69:3220–3221.3221 [PubMed: 15104470] (c) Guo H, Ding KL. *Tetrahedron Lett* 2000;41:10061–10064.10064 (d) Gillespie KM, Sanders CJ, O'Shaughnessy P, Westmoreland I, Thickitt CP, Scott P. *J. Org. Chem* 2002;67:3450–3458.3458 [PubMed: 12003558] (e) Chen YX, Li YM, Lam KH, Chan ASC. *Chin. J. Chem* 2001;19:794–799.799
2. Pd-catalyzed aerobic kinetic resolution of secondary alcohols: Chen T, Jiang JJ, Xu Q, Shi M. *Org. Lett* 2007;9:865–868.868 [PubMed: 17261005]
3. Cu-catalyzed aziridination: (a) ref. 1d. (b) Shi M, Wang CJ. *Chirality* 2002;14:412–416.416 [PubMed: 11984757] (c) Gillespie KM, Crust EJ, Deeth RJ, Scott P. *Chem. Commun* 2001:785–786.786 (d) Sanders CJ, Gillespie KM, Bell D, Scott P. *J. Am. Chem. Soc* 2000;122:7132–7133.7133
4. Cu-catalyzed aerobic oxidative coupling of 3-hydroxy-2-naphthoate: Kim KH, Lee DW, Lee YS, Ko DH, Ha DC. *Tetrahedron* 2004;60:9037–9042.9042
5. Co-catalyzed epoxidation of styrene: Nishinaga A, Yamato H, Abe T, Maruyama K, Matsuura T. *Tetrahedron Lett* 1988;29:6309–6312.6312
6. Rh- and Ru-catalyzed hydrogenation: (a) Huang HM, Okuno T, Tsuda K, Yoshimura M, Kitamura M. *J. Am. Chem. Soc* 2006;128:8716–8717.8717 [PubMed: 16819850] (b) Chen YX, Li YM, Lam KH, Chan ASC. *Chin. J. Chem* 2003;21:66–70.70 (c) Perez C, Perez S, Fuentes GA, Corma A. *J. Mol. Catal. A: Chem* 2003;197:275–281.281 (d) Chen YX, Li YM, Lam KH, Chan ASC. *Chin. J. Chem* 2002;20:606–609.609 (e) Zhang FY, Kwok WH, Chan ASC. *Tetrahedron: Asymmetry* 2001;12:2337–2342.2342 (f) Zhang FY, Pai CC, Chan ASC. *J. Am. Chem. Soc* 1998;120:5808–5809.5809
7. Rh- and Y-catalyzed hydrosilylation: (a) Xu Q, Gu XX, Liu SJ, Dou QY, Shi M. *J. Org. Chem* 2007;72:2240–2242.2242 [PubMed: 17311460] (b) Gountchev TI, Tilley TD. *Organometallics* 1999;18:5661–5667.5667
8. Reduction of ketones by LiAlH_4 : Suda H, Motoi M, Fujii M, Kanoh S, Yoshida H. *Tetrahedron Lett* 1979:4565–4568.4568
9. Pd-catalyzed allylic alkylation: Chen XH, Guo RW, Li YM, Chen G, Yeung CH, Chan ASC. *Tetrahedron: Asymmetry* 2004;15:213–217.217
10. Cu-catalyzed alkynylation of aldimines: Hatano M, Asai T, Ishihara K. *Tetrahedron Lett* 2008;49:379–382.382
11. Cu- and Ru-catalyzed cyclopropanation: (a) Sanders CJ, Gillespie KM, Scott P. *Tetrahedron: Asymmetry* 2001;12:1055–1061.1061 (b) Munslow IJ, Gillespie KM, Deeth RJ, Scott P. *Chem. Commun* 2001:1638–1639.1639 (c) Bachmann S, Furler M, Mezzetti A. *Organometallics* 2001;20:2102–2108.2108
12. Zr-, Y-, Sm-, and La-catalyzed intramolecular hydroamination of alkenes: (a) Wood MC, Leitch DC, Yeung CS, Kozak JA, Schafer LL. *Angew. Chem. Int. Ed* 2007;46:354–358.358 (b) Gott AL, Clarke AJ, Clarkson GJ, Scott P. *Organometallics* 2007;26:1729–1737.1737 (c) Knight PD, Munslow I, O'Shaughnessy PN, Scott P. *Chem. Commun* 2004:894–895.895 (d) O'Shaughnessy PN, Gillespie KM, Knight PD, Munslow IJ, Scott P. *Dalton Trans* 2004:2251–2256.2256 [PubMed: 15278115] (e) O'Shaughnessy PN, Knight PD, Morton C, Gillespie KM, Scott P. *Chem. Commun* 2003:1770–1771.1771 (f) O'Shaughnessy PN, Scott P. *Tetrahedron: Asymmetry* 2003;14:1979–1983.1983

13. Dialkylzinc to cyclic enones: Hatano M, Asai T, Ishihara K. *Tetrahedron Lett* 2007;48:8590–8594.8594
14. Diels-Alder cycladditions: (a) Sakakura A, Suzuki K, Ishihara K. *Adv. Synth. Catal* 2006;348:2457–2465.2465 (b) Kano T, Tanaka Y, Maruoka K. *Org. Lett* 2006;8:2687–2689.2689 [PubMed: 16774232]
15. Glyoxylate-ene reactions: Mikami K, Aikawa K. *Org. Lett* 2002;4:99–101.101 [PubMed: 11772100]
16. Silylcyanation of aldehydes: Zhou XG, Huang JS, Ko PH, Cheung KK, Che CM. *J. Chem. Soc.; Dalton Trans* 1999:3303–3309.3309
17. Morita-Baylis-Hillman reactions: Shi M, Liu XG. *Org. Lett* 2008;10:1043–1046.1046 [PubMed: 18284244]
18. Oxy-Michael addition reactions: Kano T, Tanaka Y, Maruoka K. *Tetrahedron* 2007;63:8658–8664.8664
19. Henry reactions: Liu XG, Jiang JJ, Shi M. *Tetrahedron: Asymmetry* 2007;18:2773–2781.2781
20. Scarborough CC, Popp BV, Guzei IA, Stahl SS. *J. Organomet. Chem* 2005;690:6143–6155. Scarborough CC, Grady MJW, Guzei IA, Gandhi BA, Bunel EE, Stahl SS. *Angew. Chem. Int. Ed* 2005;44:5269–5272.
21. See, for example: (a) Daugulis O, Zaitsev VG. *Angew. Chem. Int. Ed* 2005;44:4046–4048.4048 (b) Shabashov D, Daugulis O. *J. Org. Chem* 2007;72:7720–7725.7725 [PubMed: 17824657] (c) Kalyani D, Deprez NR, Desai LV, Sanford MS. *J. Am. Chem. Soc* 2005;127:7330–7331.7331 [PubMed: 15898779] (d) Yang SD, Li BJ, Wan XB, Shi ZJ. *J. Am. Chem. Soc* 2007;129:6066–6067.6067 [PubMed: 17455938]
22. For earlier demonstration of related reactivity with alkyl iodides and stoichiometric Pd, see: Tremont SJ, Rahman HU. *J. Am Chem. Soc* 1984;106:5759–5760.5760
23. Zhang SJ, Zhang DW, Liebeskind LS. *J. Org. Chem* 1997;62:2312–2313. [PubMed: 11671553]
24. The DABN substrate appears to be more reactive and susceptible to side reactions. Efforts to perform reactions with aryl iodides other than PhI typically resulted in lower yields. For example, DZ coupling of N,N'-diacetyl-DABN with 1-iodo-3,5-dimethylbenzene proceeded in 46% yield with complete consumption of starting material (cf. entry 5, Table 1).
25. Partial hydrogenation of DABN to form Hg-DABN can be achieved without loss of enantiopurity.
26. (a) Akiyama T, Itoh J, Fuchibe K. *Adv. Synth. Catal* 2006;348:999–1010. (b) Pihko PM. *Lett. Org. Chem* 2005;2:398–403. (c) Chen Y, Yekta S, Yudin AK. *Chem. Rev* 2003;103:3155–3212. [PubMed: 12914495]

SCHEME 1.
Synthetic Sequence for Bis-*Ortho*-Arylation of Biphenyl Diamine **1**.

CHART 1.

Common Biaryl Diamine Structures (DABN = 2,2'-diamino-1,1'-binaphthyl).

Table 1

Scope of *ortho*-Arylation of **4**.^a

entry	Ar	product
1		5a
2		5b
3		5c
4		--

entry

Art

produc

5

5d

entry

Art

product

6

5e

entry

Art

produc

7

--

^a Substrate (1.0 mmol), ArI (5.0 mmol), Pd(OAc)₂ (0.15 mmol), AgOAc (2.2 mmol), 3 mL CF₃CO₂H, reflux, 6h.

^b 20% Pd(OAc)₂, 3.0 equiv AgOAc, 14h.