Published in final edited form as:

J Organomet Chem. 2007 October 1; 692(21): 4618–4629. doi:10.1016/j.jorgchem.2007.05.035.

Synthesis of Functionalized Isoindolinones: Addition of *In Situ* Generated Organoalanes to Acyliminium Ions[#]

Joshua G. Pierce, David L. Waller, and Peter Wipf^{*}

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, U.S.A.

Abstract

Addition of *in situ* generated di- or trisubstituted alkenylalanes to *N*-acyliminium ions provides rapid access to functionalized isoindolinones. Subsequent ring closing metathesis leads to tricyclic products. These transformations proceed under mild conditions and allow for the convergent synthesis of biologically significant scaffolds from readily available starting materials.

Keywords

Isoindolinone; aluminum; zirconium; hydrozirconation; carbometallation; *N*-acyliminium ion

[#]Dedicated to Prof. Gerhard Erker in celebration of his 60th birthday and in recognition of his pioneering contributions to transition metal chemistry.

e-mail: pwipf@pitt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Isoindolinones constitute the core structures of numerous naturally occurring biologically active compounds such as magallanesine **1**ⁱ and lennoxamine **2**ⁱⁱ as well as many drug candidates such as pagoclone **3**ⁱⁱⁱ (Figure 1). Isoindolinones demonstrate a remarkably wide array of biological activities, including anti-inflammatory,^{iv} antihypertensive,^v antipsychotic,^{vi} vasodilatory^{vii} and antileukemic^{viii} effects, which renders the development of new synthetic routes toward these heterocycles particularly attractive. While several methods have been reported for the preparation of isoindolinones,^{ix} some leading to the synthesis of natural products^x and compound libraries,^{xi} many approaches suffer from a lack of generality or functional group compatibility.

In view of the potent and diverse biological spectrum of isoindolinones, we initiated studies directed at their preparation as part of our program for the synthesis of nitrogen containing heterocycles.^{xii} We initially proposed combining our water-accelerated carboalumination^{xiii} or hydrozirconation^{xiv} with an intramolecular *N*-acyliminium ion addition in a cascade process that would generate medium-ring containing isoindolinones (Scheme 1). *N*-acyliminium ions are versatile synthetic intermediates^{xv} and have previously been employed in the formation of simple isoindolinones;^{xvi} however, with the exception of allylsilanes, the addition of functionalized organometallic reagents to these systems has not yet been exploited. Moreover, the intramolecular addition would potentially provide an entry toward strained medium rings containing (*E*)-alkenes.

2. Results and Discussion

Our investigations began with the synthesis of methoxy-**8** and phenoxy lactam **9** (Scheme 2). Alcohol **5** was obtained in 85% yield by the isomerization of the commercially available hept-3-yne-1-ol with KAPA reagent in 85% yield.^{xvii} Imide **6** could then be constructed by a Mitsunobu alkylation with alkynol **5**, which proceeded in 82% yield. Reduction with sodium borohydride was followed by a methanol exchange to give methoxylactam **8** in 40% overall yield (Scheme 2). With the key substrate in hand, we submitted **8** to our water accelerated carboalumination conditions (AlMe₃ (3 eq), H₂O (1 eq), zirconocene dichloride (10 mol%)) which only returned starting material with no evidence for elimination of methanol or carboalumination of the alkyne moiety (Scheme 3).

We anticipated the need for a better leaving group and hence prepared phenoxy lactam **9**, which was generated by treatment of **7** with SOCl₂ and catalytic DMF. The resulting unstable chloride intermediate was immediately treated with phenol in the presence of triethylamine to yield **9** in 34% yield.^{xviii} When **9** was subjected to the carboalumination conditions, only alkylated lactam **10** (10%), starting material (52%), and decomposition products were observed by NMR analysis (Scheme 3).

These experiments suggested that the functionality present in these lactam acetals was inhibiting the carboalumination step. We postulated that the lactam group was responsible for the observed inhibition, as highly Lewis basic functionality has been noted to decrease the reactivity of aluminoxanes.^{xix} Therefore, we performed a series of GC experiments to test this hypothesis with 1-heptyne, which is known to undergo rapid water-accelerated carboalumination. A control experiment demonstrated that the carboalumination of heptyne with AlMe₃ (3 eq), H₂O (1 eq), and Cp₂ZrCl₂ (10 mol%) proceeded to completion in less than 5 min at low temperatures (−78 °C to −25 °C) to furnish a 97:3 mixture of regioisomers by GC analysis (Scheme 4). Conducting the identical experiment in the presence of 1 equivalent of methoxylactam **8** dramatically inhibited carboalumination, with only 4.4% of heptyne undergoing conversion after 1 h at ambient temperature. Most likely, this inhibition is due to

the coordination of Lewis basic functional groups to AlMe_3 oligomers, thus preventing alkyne carboalumination.

Due to the lack of reactivity in the carboalumination approach, we turned to hydrozirconation, known to be quite tolerant of diverse functionality.¹⁴ When lactam **8** was treated with Cp_2ZrHCl , only the corresponding alkene was recovered (Scheme 5). Treatment of the intermediate alkenylzirconocene with AgClO_4 or trimethylaluminum also provided no desired cyclization products. Our attempts to enhance leaving group ability by using pivaloate **11** also failed to promote cyclization under any of the above conditions. Highly concentrated reaction mixtures or prolonged heating resulted only in slow methyl group addition. Possibly, the ring strain present in the desired products is too high to allow the reaction to proceed under these experimental conditions.

Because the intramolecular cyclizations proved elusive, we turned our attention toward an alternative construction of these systems starting with an intermolecular addition process. Alkenylation of *N*-acyliminium ions can often require harsh reaction conditions^{xx} or unusual anomeric leaving groups;^{xxi} a noteworthy exception is the mild intermolecular addition of alkenyl boronates reported by Batey et al.^{xxii} We first explored the hydrozirconation of terminal alkynes as a method to generate alkenyl nucleophiles (Table 1). Hydrozirconation of 1-hexyne, followed by treatment with silver perchlorate^{xxiii} and lactam **13** yielded only a trace of addition product after 24 h at rt (entry 1, Table 1). Transmetalation to dimethylzinc also proved unsuccessful (entry 2, Table 1). Gratifyingly, hydrozirconation-transmetalation to trimethylaluminum^{xxiv} generated an alkenylalane that reacted with lactam **13** in a moderate 43% yield (entry 3, Table 1). We further increased the leaving group ability by using acetate **14**; however, product **17** was only formed in 30% yield while the remaining starting material was consumed (entry 4, Table 1). Attempts to perform this reaction in THF or toluene led to recovered starting material. In an effort to prevent decomposition pathways while retaining good leaving group ability, pivaloate **15** was synthesized and treated with the *in situ* generated alane to provide hexenyl isoindolinone **17** in 81% yield without any observed isomerization of the alkene moiety (entry 5, Table 1). Attempts to achieve this transformation under cationic conditions with silver perchlorate²⁰ led to complex mixtures, seemingly arising from alkene isomerization. The mild conditions under which the addition proceeds and the ability to utilize readily available alkynes and pivaloate iminium ion precursors encouraged us to explore this process further.

Among the numerous methods known for the synthesis of substituted isoindolinones, only a few examples of Heck-type cyclizations install an alkenyl moiety at the 3-position.^{xxv} The modularity of our approach warranted further investigation, and therefore we evaluated the substrate scope of this process. As shown in Table 2, silyl ether, carbamate, and sulfonamide functionalities are well tolerated, generating functionalized isoindolinones in a high yielding, one-pot procedure.

To further probe the scope of this reaction, we synthesized succinimide-derived pivaloate **25** from 1-benzyl-3,3-dimethylpyrrolidine-2,5-dione (**24**) and subjected it to our optimized conditions (Scheme 6). Addition product **26** was isolated in 83% yield. This success bodes well for the extension of the cyclic iminium ion alkenylation methodology toward a broad class of heterocyclic electrophiles.

We also explored the use of trisubstituted alkenes in this transformation. Although the lactam functionality had prevented an intramolecular addition, pre-forming the alkenylalane under our water-accelerated conditions, followed by addition of **15**, provided isoindolinones **27** and **29** in 77% yield after only 15 minutes (Table 3). The flexibility and mild nature of these

transformations are well suited for their application in the synthesis of biologically interesting target molecules and libraries.

Since this strategy provided a rapid access toward alkenyl-functionalized phthalimides, we sought to revisit our initial goal of generating medium rings. As a model system, we synthesized alkyne **30** and allyl-pivaloate **31** and subjected them to our hydrozirconation-transmetallation conditions to generate **32** in 55% yield accompanied by considerable amounts of methyl addition side product (Scheme 7). It should be noted that carbometallation of **30** failed and the rate of addition for the alkenylalane generated via the transmetallation reaction was retarded due to the presence of the ether functionality. With **32** in hand, we screened conditions to form the desired 12-membered macrocycle. Although ring closing metathesis methodology²³ has been demonstrated to perform well for the synthesis of a variety of macrocycles,^{xxvi} rapid formation of the undesired 5-membered ring **33** was observed (Scheme 7).^{xxvii} Increasing dilution, change of metathesis catalyst, addition of $\text{Ti}(\text{O}^i\text{Pr})_4$,^{xxviii} or varying solvents did not influence the reaction pathway.

Due to the ease of formation of pyrrolizidine **33**, we explored the synthesis of this compound via addition of hexyne to **31** and subsequent ring closing metathesis to provide tricycle **33** in 75% yield (Scheme 8). Extension of this approach to the synthesis of fused azepines was achieved from indolinone **37**, prepared from 2-(pent-4-enyl)isoindoline-1,3-dione (**35**) via pivaloate **36** and 1-hexyne in 72% yield. Ring closing metathesis of **37** proved to be problematic in CH_2Cl_2 with several ruthenium catalysts, yielding only starting material or decomposition products. Addition of $\text{Ti}(\text{O}^i\text{Pr})_4$ in toluene at room temperature resolved these issues and yielded 67% of **38**. Possible deactivation of the metathesis intermediate by the neighboring amide carbonyl could explain the difficulty in this transformation.^{xxix} While not fully optimized, these reactions provide proof of concept for the conversion of our phthalimide substrates to structurally diverse tricyclic products.

3. Conclusions

We have demonstrated that *in situ* generated alkenylalanes represent versatile nucleophiles for additions to *N*-acyliminium ions. While direct intramolecular cyclization strategies currently suffer from inhibition of the carboalumination reaction by Lewis basic functional groups, preforming alkenylalanes via hydrozirconation-transmetallation or carboalumination and subsequent addition to the lactam acetal substrates yield functionalized isoindolinones. This method applies easily prepared or commercially available starting materials that provide opportunities for diversification at numerous points and yields synthetically useful heterocyclic products in a one-pot transformation. Further elaboration of the alkenyl heterocycles through the use of ring closing metathesis leads to tricyclic products that are common motifs in natural products and drug-like molecules.

4. Experimental Section

4.1 General Details

All reactions were performed under an N_2 atmosphere and all glassware was dried in an oven at 140 °C for 2 h prior to use. THF and Et_2O was distilled over sodium/benzophenone ketyl, Et_3N was distilled from CaH_2 , and CH_2Cl_2 and toluene were purified using an alumina filtration system. Cp_2ZrHCl ,^{xxx} **18**,^{xxxi} **20**,^{xxxii} and **22**²⁴ were prepared according to literature procedures and all other compounds were purchased and used as received.

Reactions were monitored by TLC analysis (EM Science pre-coated silica gel 60 F₂₅₄ plates, 250 mm layer thickness) and visualization was accomplished with a 254 nm UV light and by staining with a PMA solution (5 g of phosphomolybdic acid in 100 mL of 95% EtOH), *p*-

anisaldehyde solution (2.5 mL of *p*-anisaldehyde, 2 mL of AcOH, and 3.5 mL of conc. H₂SO₄ in 100 mL of 95% EtOH), Vaughn's reagent (4.8 g of (NH₄)₆Mo₇O₂₄•4 H₂O and 0.2 g of Ce(SO₄)₂ in 100 mL of a 3.5 N H₂SO₄ solution) or a KMnO₄ solution (1.5 g of KMnO₄ and 1.5 g of K₂CO₃ in 100 mL of a 0.1% NaOH solution). Flash chromatography on SiO₂ was used to purify the crude reaction mixtures.

Melting points were determined using a Laboratory Devices Mel-Temp II. Infrared spectra were determined on a Nicolet Avatar 360 FT-IR spectrometer. ¹H and ¹³C NMR spectra were obtained on a Bruker Avance 300 instrument in CDCl₃ unless otherwise noted. Chemical shifts were reported in parts per million with the residual solvent peak used as an internal standard. ¹H NMR spectra were run at 300 MHz and are tabulated as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, m = multiplet), number of protons, and coupling constant(s). ¹³C NMR spectra were run at 76 MHz using a proton-decoupled pulse sequence with a d₁ of 3 sec, and are tabulated by observed peak. Mass spectra were obtained on a Micromass Autospec double focusing instrument.

2-(Hept-6-ynyl)isoindoline-1,3-dione (6)—A solution of hept-6-yn-1-ol (**5**) (228 mg, 2.04 mmol), phthalimide **4** (302 mg, 2.04 mmol), and PPh₃ (540 mg, 2.04 mmol) in THF (20 mL) was cooled to 0 °C and treated with DIAD (403 μL, 2.04 mmol) over 5 min. The reaction mixture was warmed to rt and stirred for 6 h. The solvent was evaporated, the residue was dissolved in EtOAc/hexanes (1:1, 10 mL) and the solids were filtered off. The filtrate was concentrated and chromatographed on SiO₂ (EtOAc:hexanes, 1:5) to yield imide **6** (407 mg, 1.69 mmol, 82%) as a colorless oil: IR (neat) 3466, 3283, 2941, 2862, 2115, 1772, 1709, 1615, 1397, 1046, 720 cm⁻¹; ¹H NMR δ 7.84-7.79 (m, 2 H), 7.72-7.67 (m, 2 H), 3.67 (t, 2 H, *J* = 7.1 Hz), 2.17 (td, 2 H, *J* = 6.8, 2.6 Hz), 1.91 (t, 1 H, *J* = 2.6 Hz), 1.75-1.61 (m, 2 H), 1.58-1.51 (m, 2 H), 1.49-1.38 (m, 2 H); ¹³C NMR δ 168.3 (2 C), 133.8 (2 C), 132.1 (2 C), 123.1 (2 C), 84.2, 68.4, 37.7, 28.0, 27.9, 25.8, 18.2; MS (EI) *m/z* (rel intensity) 241 ([M]⁺, 11), 186 (6), 173 (10), 160 (100), 148 (31), 130 (29), 104 (27); HRMS (EI) *m/z* calcd for C₁₅H₁₅NO₂ 241.1103, found 241.1107.

2-(Hept-6-ynyl)-3-methoxyisoindolin-1-one (8)—To a 0 °C solution of imide **6** (40.0 mg, 0.166 mmol) in MeOH (4 mL) was added NaBH₄ (5.00 mg, 0.125 mmol). The reaction mixture was stirred at rt for 2.5 h, quenched with H₂O (1 mL) and the MeOH was removed *in vacuo*. The residue was extracted with CH₂Cl₂ (5 × 3 mL) and the combined organic layers were washed with brine, dried (Na₂SO₄), and concentrated to provide crude hydroxylactam **7** which was immediately dissolved in MeOH (4 mL), treated with *d,l*-camphorsulfonic acid (3.85 mg, 0.0166 mmol) and stirred for 16 h. The solvent was removed *in vacuo* and the residue was partitioned between H₂O (5 mL) and CH₂Cl₂ (5 mL). The aqueous layer was extracted with CH₂Cl₂ (3 × 5 mL). The combined organic layers were washed with brine, dried (Na₂SO₄), concentrated and chromatographed on SiO₂ (EtOAc:hexanes, 1:4) to yield methoxy lactam **8** (17.0 mg, 0.0661 mmol, 40% over two steps) as a colorless oil: IR (neat) 3298, 2925, 2854, 2115, 1702, 1466, 1412, 1059, 746 cm⁻¹; ¹H NMR δ 7.84-7.80 (m, 1 H), 7.61-7.49 (m, 3 H), 5.88 (s, 1 H), 3.79 (ddd, 1 H, *J* = 13.7, 8.1, 7.3 Hz), 3.24 (ddd, 1 H, *J* = 14.0, 8.0, 6.3 Hz), 2.87 (s, 3 H), 2.19 (td, 2 H, *J* = 6.7, 2.6 Hz), 1.92 (t, 1 H, *J* = 2.6 Hz), 1.75-1.63 (m, 2 H), 1.62-1.43 (m, 4 H); ¹³C NMR δ 167.6, 140.3, 133.2, 131.9, 129.9, 123.4 (2 C), 86.2, 84.3, 68.4, 49.1, 39.3, 28.0, 27.6, 26.0, 18.3; MS (EI) *m/z* (rel intensity) 257 ([M]⁺, 18), 242 (17), 226 (20), 176 (66), 146 (100), 132 (39), 117 (21); HRMS (EI) *m/z* calcd for C₁₆H₁₉NO₂ 257.1416, found 257.1419.

2-(Hept-6-ynyl)-3-phenoxyisoindolin-1-one (9)—A 0 °C solution of imide **6** (307 mg, 1.27 mmol) in MeOH (20 mL) was treated with NaBH₄ (36.3 mg, 0.953 mmol), warmed to rt and stirred for 1 h. The reaction mixture was quenched with H₂O (10 mL) and the MeOH was removed *in vacuo*. The residue was extracted with CH₂Cl₂ (6 × 10 mL) and the combined

organic layers were washed with brine, dried (Na_2SO_4) and concentrated to yield crude hydroxylactam **7** as a colorless oil. A portion of the crude hydroxylactam (101 mg, 0.416 mmol) was immediately dissolved in THF (5 mL) at rt and treated with SOCl_2 (30.3 μL , 0.416 mmol) and DMF (1 drop) and stirred for 14 h. The volatiles were removed under reduced pressure and the residue was dried *in vacuo* for 12 h. The unstable crude chloride was obtained as a colorless oil and immediately dissolved in THF (4.1 mL) at ambient temperature. This solution was treated with phenol (58.0 mg, 0.620 mmol) and Et_3N (288 μL , 2.07 mmol) and stirred for 2 h. The reaction mixture was quenched with H_2O (1 mL) and the THF was removed *in vacuo*. The residue was extracted with CH_2Cl_2 (4×5 mL) and the combined organic layers were washed with brine, dried (Na_2SO_4), and concentrated. The residue was chromatographed on SiO_2 (EtOAc:hexanes, 1:3) to yield phenoxylactam **9** (45.0 mg, 1.41 mmol, 34% over two steps) as a colorless oil: IR (neat) 3296, 2936, 2115, 1707, 1588, 1491, 1415, 1222, 992, 780 cm^{-1} ; ^1H NMR δ 7.85-7.79 (m, 1 H), 7.53-7.46 (m, 3 H), 7.33-7.27 (m, 2 H), 7.08-6.98 (m, 3 H), 6.42 (s, 1 H), 3.80 (dt, 1 H, $J = 14.2$, 7.4 Hz), 3.44 (ddd, 1 H, $J = 14.0$, 7.8, 6.4 Hz), 2.16 (td, 2 H, $J = 6.7$, 2.6 Hz), 1.91 (t, 1 H, $J = 2.6$ Hz), 1.78-1.61 (m, 2 H), 1.57-1.39 (m, 4 H); ^{13}C NMR δ 167.4, 156.3, 141.3, 132.3, 132.0, 130.0, 129.7 (2 C), 123.5, 123.3, 123.1, 118.1 (2 C), 86.8, 84.2, 68.4, 40.0, 27.9, 27.6, 25.9, 18.2; MS (EI) m/z (rel intensity) 319 ($[\text{M}]^+$, 70), 265 (28), 226 (100), 198 (8), 146 (52), 132 (65); HRMS (EI) m/z calcd for $\text{C}_{21}\text{H}_{21}\text{NO}_2$ 319.1572, found 319.1573.

2-Benzyl-3-oxoisindolin-1-yl pivaloate (15)—To a solution of benzyl phthalimide (8.89 g, 37.5 mmol) in MeOH (130 mL) was added sodium borohydride (1.42 g, 37.5 mmol) at 0 °C. The resulting reaction mixture was stirred at 0 °C for 2 h and carefully quenched with H_2O . The aqueous layer was extracted with EtOAc (2 $\times$) and the organic layer was separated, dried (MgSO_4) and concentrated. The residue was dried to yield a white solid which was carried on to the next step without further purification. To a solution of the crude reduced phthalimide (1.00 g, 4.20 mmol) in THF (30 mL) was added Et_3N (1.17 mL, 8.40 mmol) and pivaloyl chloride (621 μL , 5.04 mmol) at 0 °C and the reaction mixture was allowed to warm to rt and stirred at this temperature for 4 h. The mixture was quenched with sat. aq. NaHCO_3 , extracted with EtOAc (2 $\times$), dried (MgSO_4) and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc:hexanes, 3:7) to yield 1.09 g (80% over 2 steps) of **15** as a colorless solid: mp 88.1 – 89.0 °C (CH_2Cl_2); IR (neat) 3419, 3062, 3031, 2973, 2934, 2872, 1717, 1408, 1276, 1122, 959, 751 cm^{-1} ; ^1H NMR δ 7.92-7.84 (m, 1 H), 7.61-7.53 (m, 2 H), 7.50-7.43 (m, 1 H), 7.38-7.22 (m, 5 H), 6.89 (s, 1 H), 5.01 (d, 1 H, $J = 15$ Hz), 4.44 (d, 1 H, $J = 15$ Hz), 1.13 (s, 9 H); ^{13}C NMR δ 178.4, 167.9, 141.3, 136.8, 132.5, 131.9, 130.2, 128.7, 128.1, 127.7, 123.7, 123.7, 81.0, 44.2, 39.0, 26.8; MS (EI) m/z (rel intensity) 323 ($[\text{M}]^+$, 35), 221 (100), 133 (65), 91 (100); HRMS (EI) m/z calcd for $\text{C}_{21}\text{H}_{23}\text{NO}$ 323.1521, found 323.1515.

(E)-2-Benzyl-3-(hex-1-enyl)isindolin-1-one (17). General Protocol A—To a solution of 1-hexyne (**16**) (60.5 μL , 0.526 mmol) in CH_2Cl_2 (1.5 mL) was added zirconocene hydrochloride (156 mg, 0.605 mmol) and the resulting suspension was stirred at rt for 10 min. The resulting yellow solution was cooled to 0 °C and Me_3Al (1.0 M in CH_2Cl_2 , 0.605 mL, 0.605 mmol) and **15** (85.0 mg, 0.263 mmol) were added. The mixture was warmed to rt and stirred at this temperature for 1 h, quenched with sat. aq. NH_4Cl , and extracted with CH_2Cl_2 . The organic layers were separated, dried (MgSO_4) and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc:hexanes, 3:7) to yield 65.1 mg (81%) of **17** as a colorless oil: IR (neat) 3479, 3031, 2927, 2857, 1694, 1615, 1400, 972 cm^{-1} ; ^1H NMR δ 7.91-7.86 (m, 1 H), 7.56-7.42 (m, 2 H), 7.36-7.21 (m, 6 H), 5.90 (dt, 1 H, $J = 15.0$, 6.8 Hz), 5.30 (d, 1 H, $J = 14.8$ Hz), 5.09 (dd, 1 H, $J = 15.2$, 9.2 Hz), 4.70 (d, 1 H, $J = 9.2$ Hz), 4.17 (d, 1 H, $J = 14.9$ Hz), 2.13 (app q, 2 H, $J = 6.7$ Hz), 1.50-1.30 (m, 4 H), 0.94 (t, 3 H, $J = 7.2$ Hz); ^{13}C NMR δ 167.9, 145.1, 138.3, 137.5, 131.8, 131.5, 128.6, 128.3, 127.3, 126.1, 123.6, 123.0, 62.7, 43.8,

31.8, 31.1, 22.1, 13.8; MS (EI) m/z (rel intensity) 305 ($[M]^+$, 100), 248 (40), 237 (70), 214 (40); HRMS (EI) m/z calcd for $C_{21}H_{23}NO$ 305.1780, found 305.1785.

(E)-2-Benzyl-3-(2-cyclohexylvinyl)isoindolin-1-one (19)—According to general protocol A, alkyne **18** (66.9 mg, 0.618 mmol), CH_2Cl_2 (1.5 mL), zirconocene hydrochloride (183 mg, 0.711 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 0.711 mL, 0.711 mmol) and **15** (100 mg, 0.309 mmol) afforded 80.8 mg (79%) of **19** as a colorless oil after purification on SiO_2 (EtOAc:hexanes, 3:7): IR (neat) 3375, 3030, 2921, 2851, 2243, 1220, 1200, 1097, 844 cm^{-1} ; 1H NMR δ 7.92–7.86 (m, 1 H), 7.54–7.41 (m, 2 H), 7.35–7.20 (m, 6 H), 5.85 (dd, 1 H, $J = 15.3, 6.6$ Hz), 5.28 (d, 1 H, $J = 14.8$ Hz), 5.02 (ddd, 1 H, $J = 15.3, 9.2, 1.2$ Hz), 4.67 (d, 1 H, $J = 9.2$ Hz), 4.18 (d, 1 H, $J = 14.8$ Hz), 2.14–1.96 (m, 1 H), 1.85–1.60 (m, 4 H), 1.42–1.00 (m, 6 H); ^{13}C NMR δ 167.9, 145.1, 144.1, 137.4, 131.8, 131.4, 128.5, 128.3, 128.2, 127.3, 123.6, 122.9, 62.8, 43.8, 40.3, 32.6, 26.0, 25.8; MS (EI) m/z (rel intensity) 331 ($[M]^+$, 86), 248 (40), 237 (100); HRMS (EI) m/z calcd for $C_{23}H_{25}NO$ 331.1936, found 331.1951.

(E)-2-Benzyl-3-(4-(tert-butyl)phenylsilyloxy)but-1-enylisoindolin-1-one (21)—According to general protocol A, alkyne **20** (94.4 mg, 0.306 mmol), CH_2Cl_2 (1 mL), zirconocene hydrochloride (78.9 mg, 0.306 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 0.306 mL, 0.306 mmol) and **15** (50.0 mg, 0.153) afforded 57.6 mg (71%) of **21** as a colorless oil after purification on SiO_2 (EtOAc:hexanes, 2:8): IR (neat) 3450, 2930, 2857, 1692, 1428, 1111, 735, 701 cm^{-1} ; 1H NMR δ 7.92–7.86 (m, 1 H), 7.72–7.65 (m, 4 H), 7.53–7.35 (m, 9 H), 7.32–7.21 (m, 5 H), 5.96 (dt, 1 H, $J = 15.3, 6.8$ Hz), 5.27 (d, 1 H, $J = 14.9$ Hz), 5.16 (dddd, 1 H, $J = 15.3, 9.1, 1.2, 1.2$ Hz), 4.71 (d, 1 H, $J = 9.1$ Hz), 4.16 (d, 1 H, $J = 14.9$ Hz), 3.77 (dt, 2 H, $J = 6.3, 1.2$ Hz), 2.37 (app q, 2 H, $J = 6.6$ Hz), 1.08 (s, 9 H); ^{13}C NMR δ 168.0, 144.9, 137.5, 135.6, 134.8, 133.8, 131.9, 131.5, 129.7, 128.6, 128.4, 128.2, 127.7, 127.4, 123.6, 123.1, 63.2, 62.6, 43.9, 35.6, 26.9, 19.2; MS (EI) m/z (rel intensity) 532 ($[M]^+$, 35), 488 (100), 474 (45), 306 (65), 252 (75); HRMS (EI) m/z calcd for $C_{35}H_{37}NO_2Si$ 532.2664, found 532.2664.

(E)-Methyl 4-(2-benzyl-3-oxoisoindolin-1-yl)but-3-enyl(tosyl)carbamate (23)—According to general protocol A, alkyne **22** (174 mg, 0.618 mmol), CH_2Cl_2 (1.5 mL), zirconocene hydrochloride (183 mg, 0.711 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 0.711 mL, 0.711 mmol) and **15** (100 mg, 0.309) afforded 95.1 mg (62%) of **23** as a colorless oil after purification on SiO_2 (acetone: CH_2Cl_2 , 0.3:9.7): IR (neat) 3467, 3032, 2957, 2245, 1735, 1686, 1359, 1168, 733 cm^{-1} ; 1H NMR δ 7.92–7.78 (m, 3 H), 7.56–7.41 (m, 2 H), 7.39–7.18 (m, 9 H), 5.93 (dt, 1 H, $J = 15.2, 7.0$ Hz), 5.26 (d, 1 H, $J = 14.9$ Hz), 5.21 (dd, 1 H, $J = 15.1, 9.1$ Hz), 4.73 (d, 1 H, $J = 9.0$), 4.18 (d, 1 H, $J = 14.9$ Hz), 3.95 (t, 2 H, $J = 7.0$ Hz), 3.69 (s, 3 H), 2.58 (app q, 2 H, $J = 7.0$ Hz), 2.43 (s, 3 H); ^{13}C NMR δ 168.0, 152.8, 144.8, 144.6, 137.4, 136.5, 133.0, 131.7, 131.6, 129.7, 129.4, 128.6, 128.3, 128.3, 127.4, 123.6, 123.2, 62.3, 53.8, 46.5, 43.8, 32.9, 21.6; MS (EI) m/z (rel intensity) 504 ($[M]^+$, 50), 262 (40), 155 (100); HRMS (EI) m/z calcd for $C_{28}H_{28}N_2O_5S$ 504.1719, found 504.1724.

1-Benzyl-3,3-dimethylpyrrolidine-2,5-dione (24)—A mixture of 2,2-dimethylsuccinic anhydride (710 mg, 5.55 mmol) and benzyl amine (712 mg, 6.66 mmol, 1.2 eq) was heated over a bunsen burner for ~ 1 min. The cooled mixture was purified by chromatography on SiO_2 (EtOAc:hexanes, 1:1) to furnish 1.14 g (95%) of **24** as a colorless oil: IR (neat) 2969, 2933, 1777, 1702, 1344, 1142, 709 cm^{-1} ; 1H NMR δ 7.29–7.21 (m, 5 H), 4.57 (s, 2 H), 2.47 (s, 2 H), 1.22 (s, 6 H); ^{13}C NMR δ 182.5, 175.1, 135.8, 128.3, 128.1, 127.5, 43.2, 42.0, 39.7, 25.1; MS (EI) m/z (rel intensity) 217 ($[M]^+$, 100), 174 (33), 133 (22); HRMS (EI) m/z calcd for $C_{13}H_{15}NO_2$ 217.1103, found 217.1104.

1-Benzyl-4,4-dimethyl-5-oxopyrrolidin-2-yl pivaloate (25)—A solution of imide **24** (1.14 g, 5.25 mmol) in MeOH (53 mL) was treated with $NaBH_4$ (200 mg, 2.62 mmol) at ambient

temperature. After 6 h, the reaction mixture was concentrated, the residue dissolved in CH_2Cl_2 (50 mL) and treated with sat. aq. NH_4Cl . The aqueous layer was separated and washed with CH_2Cl_2 (2 $\times$) and the combined organic layers were dried (MgSO_4), filtered, concentrated and the resulting oil used without further purification. A solution of the crude oil in CH_2Cl_2 (50 mL) was treated sequentially with Et_3N (2.19 mL, 15.7 mmol, 3 eq), DMAP (128 mg, 1.05 mmol, 20 mol %) and pivaloyl chloride (1.29 mL, 10.5 mmol, 2 eq) at ambient temperature. After 6 h, the reaction mixture was quenched with 3 M HCl and the aqueous layer washed with CH_2Cl_2 (3 $\times$). The combined organic layers were washed with H_2O , dried (MgSO_4), filtered, and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc :hexanes, 1:2) to yield 398 mg (25%) of **25** as a colorless oil: IR (neat) 2971, 1710, 1419, 1125, 707 cm^{-1} ; ^1H NMR δ 7.33-7.21 (m, 5 H), 5.99 (d, 1 H, J = 6.3 Hz), 4.74 (d, 1 H, J = 14.7 Hz), 4.15 (d, 1 H, J = 14.7 Hz), 2.14 (dd, 1 H, J = 14.1, 6.3 Hz), 1.87 (d, 1 H, J = 14.4 Hz), 1.32 (s, 3 H), 1.22 (s, 3 H), 1.10 (s, 9 H); ^{13}C NMR δ 180.6, 177.9, 136.6, 128.7, 128.2, 127.6, 82.1, 44.8, 41.5, 39.3, 38.7, 26.8, 26.5, 25.6; MS (EI) m/z (rel intensity) 303 ($[\text{M}]^+$, 36), 275 (15), 202 (85), 158 (46); HRMS (EI) m/z calcd for $\text{C}_{18}\text{H}_{25}\text{NO}_3$ 303.1834, found 303.1827.

(E)-1-Benzyl-5-(hex-1-enyl)-3,3-dimethylpyrrolidin-2-one (26)—According to general protocol A, hexyne (**16**) (9.77 mg, 0.119 mmol), CH_2Cl_2 (1 mL), zirconocene hydrochloride (30.7 mg, 0.119 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 0.119 mL, 0.119 mmol) and **25** (18.0 mg, 0.0593 mmol) afforded 14.0 mg (83%) of **26** as a colorless oil after purification on SiO_2 (EtOAc :hexanes, 2.5:7.5): IR (neat) 2958, 2928, 2868, 1692, 1411, 1262, 972, 752 cm^{-1} ; ^1H NMR δ 7.35-7.23 (m, 3 H), 7.21-7.13 (m, 2 H), 5.53 (dt, 1 H, J = 15.0, 6.6 Hz), 5.16 (dd, 1 H, J = 15.3, 9.0 Hz), 4.94 (d, 1 H, J = 14.4 Hz), 3.88 (d, 1 H, J = 14.4 Hz), 3.73 (app q, 1 H, J = 7.8 Hz), 2.10-1.98 (m, 2 H), 1.99 (dd, 1 H, J = 12.9, 7.2 Hz), 1.58 (dd, 1 H, J = 12.9, 8.1 Hz), 1.42-1.28 (m, 4 H), 1.24 (s, 3 H), 1.11 (s, 3 H), 0.92 (t, 3 H, J = 6.9 Hz); ^{13}C NMR δ 179.6, 137.2, 135.8, 129.8, 128.4, 128.3, 127.2, 56.9, 44.2, 42.0, 40.3, 31.8, 31.2, 25.5, 24.7, 22.2, 13.9; MS (EI) m/z (rel intensity) 285 ($[\text{M}]^+$, 20), 228 (20), 175 (30), 91 (100); HRMS (EI) m/z calcd for $\text{C}_{19}\text{H}_{27}\text{NO}$ 285.2093, found 285.2090.

(E)-2-Benzyl-3-(2-methylhex-1-enyl)isoindolin-1-one (27)

General Protocol B: To a $-30\text{ }^\circ\text{C}$ solution of AlMe_3 (89.0 mg, 1.24 mmol) and Cp_2ZrCl_2 (9.00 mg, 0.0309 mmol) in CH_2Cl_2 (1.5 mL) was added H_2O (11.1 mg, 0.619 mmol) dropwise. The reaction mixture was warmed to ambient temperature, cooled to $0\text{ }^\circ\text{C}$, treated with 1-hexyne (**16**) (71.0 μL , 0.619 mmol), stirred for 30 min and treated with **15** (100 mg, 0.309 mmol). The reaction mixture was warmed to rt and stirred at this temperature for 1 h, quenched with sat. aq. NH_4Cl , and extracted with CH_2Cl_2 . The organic layers were separated, dried (MgSO_4) and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc :hexanes, 3:7) to yield 76.0 mg (77%) of **27** as a colorless oil: IR (neat) 2956, 2929, 2858, 1694, 1468, 1401, 749, 703 cm^{-1} ; ^1H NMR δ 7.89-7.50 (m, 1 H), 7.53-7.40 (m, 2 H), 7.35-7.20 (m, 6 H), 5.31 (d, 1 H, J = 14.9 Hz), 5.06 (d, 1 H, J = 9.8), 4.81 (dq, 1 H, J = 9.8, 1.2 Hz), 4.09 (d, 1 H, J = 14.9 Hz), 2.07 (t, 2 H, J = 7.1 Hz), 1.67 (d, 3 H, J = 1.3 Hz), 1.50-1.22 (m, 4 H), 0.91 (t, 3 H, J = 7.2 Hz); ^{13}C NMR δ 168.1, 145.7, 143.4, 137.5, 132.0, 131.4, 128.5, 128.2, 128.0, 127.3, 123.6, 122.8, 120.6, 57.9, 43.9, 39.3, 29.8, 22.2, 16.5, 13.8; MS (EI) m/z (rel intensity) 319 ($[\text{M}]^+$, 100), 221 (40); HRMS (EI) m/z calcd for $\text{C}_{22}\text{H}_{25}\text{NO}$ 319.1936, found 319.1921.

(E)-2-Benzyl-3-(2-phenylprop-1-enyl)isoindolin-1-one (29)—According to general protocol B, alkyne **28** (68.0 μL , 0.619 mmol), AlMe_3 (89.0 mg, 1.24 mmol), Cp_2ZrCl_2 (9.00 mg, 0.0309 mmol), CH_2Cl_2 (1.5 mL), H_2O (11.1 mg, 0.619 mmol) and **15** (100 mg, 0.309 mmol) afforded 81.0 mg (77%) of **29** as a colorless oil and a 95:5 mixture of regioisomers after purification on SiO_2 (EtOAc :hexanes, 1:3): Major isomer: IR (neat) 3030, 2918, 1693, 1602, 1432, 1400, 1250, 749 cm^{-1} ; ^1H NMR δ 7.93 (dd, 1 H, J = 6.3, 2.1 Hz), 7.54-7.46 (m, 2 H),

7.35-7.26 (m, 11 H), 5.41 (dd, 1 H, $J = 9.6, 0.9$ Hz), 5.35 (d, 1 H, $J = 15.0$ Hz), 5.27 (d, 1 H, $J = 9.9$ Hz), 4.21 (d, 1 H, $J = 15.0$ Hz), 2.12 (d, 3 H, $J = 0.9$ Hz); ^{13}C NMR δ 168.1, 145.0, 142.0, 141.3, 137.4, 132.0, 131.6, 128.6, 128.3, 127.8, 127.5, 125.8, 123.8, 123.7, 122.9, 58.2, 44.3, 16.3; MS (EI) m/z (rel intensity) 339 ($[\text{M}]^+$, 41), 248 (47), 234 (77), 91 (100); HRMS (EI) m/z calcd for $\text{C}_{24}\text{H}_{21}\text{NO}$ 339.1623, found 339.1631. Minor isomer (characteristic peaks): ^1H NMR δ 1.44 (d, 3 H, $J = 6.9$ Hz); ^{13}C NMR δ 128.0, 121.9, 18.0;

(E)-(3-(Pent-4-ynyloxy)prop-1-enyl)benzene (30)—To a solution of 4-pentyn-1-ol (4.03 mL, 43.3 mmol) in hexanes (75 mL) was added 50% NaOH (75 mL), TBAI (801 mg, 2.17 mmol) and cinnamyl bromide (8.97 g, 45.5 mmol). The reaction mixture was rapidly stirred for 10 h. The organic layer was separated, dried (MgSO_4) and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc:hexanes, 1:9) to yield 9.00 g (69%) of **30** as a colorless oil: IR (neat) 3298, 3027, 2951, 2854, 2117, 1478, 1365, 1111, 967 cm^{-1} ; ^1H NMR δ 7.44-7.37 (m, 2 H), 7.37-7.27 (m, 2 H), 7.27-7.21 (m, 1 H), 6.62 (d, 1 H, $J = 15.9$ Hz), 6.30 (dt, 1 H, $J = 15.9, 6.0$), 4.15 (dd, 2 H, $J = 6.0, 0.9$ Hz), 3.60 (t, 2 H, $J = 6.3$ Hz), 2.34 (dt, 2 H, $J = 7.2, 2.7$ Hz), 1.96 (t, 1 H, $J = 2.7$ Hz), 1.85 (app p, 2 H, $J = 6.6$ Hz); ^{13}C NMR δ 136.5, 131.8, 128.3, 127.4, 126.2, 126.0, 83.7, 71.2, 68.5, 68.3, 28.5, 15.1; MS (EI) m/z (rel intensity) 199 ($[\text{M}]^+$, 100), 186 (35), 173 (35), 131 (100); HRMS (EI) m/z calcd for $\text{C}_{14}\text{H}_{16}\text{O}$ 199.1123, found 199.1126.

2-Allyl-3-oxoisindolin-1-yl pivaloate (31)—To a solution of 2-allylisindoline-1,3-dione (3.64 g, 19.4 mmol) in MeOH (100 mL) was added sodium borohydride (734 mg, 19.4 mmol) at 0 °C. The resulting reaction mixture was stirred at 0 °C for 2 h and carefully quenched with H_2O . The aqueous layer was extracted with EtOAc (2 $\times$) and the organic layer was separated, dried (MgSO_4) and concentrated. The residue was dried to yield a white solid which was carried on to the next step without further purification. To a solution of the crude reduced phthalamide (3.60 g, 19.0 mmol) in THF (100 mL) was added Et_3N (7.90 mL, 57.0 mmol) and pivaloyl chloride (2.81 mL, 22.8 mmol) at 0 °C and the reaction mixture was allowed to warm to rt and stirred at this temperature for 4 h. The mixture was quenched with sat. aq. NaHCO_3 , extracted with EtOAc (2 $\times$), dried (MgSO_4) and concentrated. The residue was purified by chromatography on SiO_2 (EtOAc:hexanes, 2.5:7.5) to yield 4.40 g (83% over 2 steps) of **31** as a colorless oil: IR (neat) 2976, 1716, 1405, 1135, 753; ^1H NMR δ 7.88-7.80 (m, 1 H), 7.62-7.46 (m, 3 H), 6.99 (s, 1 H), 5.94-5.77 (m, 1 H), 5.29-5.15 (m, 2 H), 4.44 (dd, 1 H, $J = 15.9, 5.1$ Hz), 3.88 (dd, 1 H, $J = 15.3, 6.6$ Hz), 1.23 (s, 9 H) cm^{-1} ; ^{13}C NMR δ 178.3, 167.5, 141.2, 132.4, 132.4, 131.8, 130.0, 123.6, 123.5, 118.0, 80.9, 42.8, 39.0, 26.9; MS (EI) m/z (rel intensity) 273 ($[\text{M}]^+$, 60), 172 (100); HRMS (EI) m/z calcd for $\text{C}_{16}\text{H}_{19}\text{NO}_3$ 273.1365, found 273.1358.

2-Allyl-3-((E)-5-(cinnamyloxy)pent-1-enyl)isoindolin-1-one (32)—According to general protocol A, alkyne **30** (733 mg, 3.66 mmol), CH_2Cl_2 (10 mL), zirconocene hydrochloride (944 mg, 3.66 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 3.66 mL, 3.66 mmol) and **31** (500 mg, 1.83 mmol) (reaction time increased to 12 h) afforded 376 mg (55%) of **32** as a colorless oil after purification on SiO_2 (Acetone: CH_2Cl_2 , 0.7:9.3): IR (neat) 2924, 2853, 1694, 1468, 1289, 1098, 968, 746 cm^{-1} ; ^1H NMR δ 7.85 (d, 1 H, $J = 6.6$ Hz), 7.57-7.20 (m, 8 H), 6.61 (d, 1 H, $J = 15.9$ Hz), 6.29 (dt, 1 H, $J = 15.9, 6.0$ Hz), 6.02 (dt, 1 H, $J = 13.8, 6.9$ Hz), 5.90-5.73 (m, 1 H), 5.25-5.07 (m, 3 H), 4.87 (d, 1 H, $J = 9.0$ Hz), 4.60 (ddd, 1 H, $J = 15.6, 4.5, 2.7$ Hz), 4.15 (dd, 2 H, $J = 6.0, 1.2$ Hz), 3.70 (dd, 1 H, $J = 15.3, 7.2$ Hz), 3.53 (t, 2 H, $J = 6.3$ Hz), 2.25 (app q, 2 H, $J = 6.9$ Hz), 1.77 (app p, 2 H, $J = 7.8$ Hz); ^{13}C NMR δ 167.7, 144.9, 137.2, 136.6, 133.2, 132.3, 131.8, 131.4, 128.5, 128.3, 127.6, 126.7, 126.4, 126.1, 123.5, 122.9, 117.5, 71.5, 69.4, 62.9, 42.5, 29.2, 28.9; MS (ESI) m/z (rel intensity) 396 ($[\text{M}+\text{Na}]^+$, 100), 307 (20), 297 (20); HRMS (ESI) m/z calcd for $\text{C}_{25}\text{H}_{27}\text{NO}_2\text{Na}$ 396.1939, found 396.1928.

3H-Pyrrolo[2,1-a]isoindol-5(9bH)-one (33)—To a solution of **32** (41.7 mg, 0.112 mmol) in CH₂Cl₂ (6 mL) was added Grubbs 2nd generation catalyst (4.75 mg, 5.60 μmol) and the red solution was heated at 50 °C for 1 h, cooled to rt and concentrated. The residue was purified by chromatography on SiO₂ (Acetone:CH₂Cl₂, 0.4:9.6) to yield 11.7 mg (61%) of **33** as a colorless oil: IR (neat) 2872, 1614, 1468, 1395, 1366, 1080, 746 cm⁻¹; ¹H NMR (acetone-d₆) δ 7.74-7.66 (m, 2 H), 7.66-7.58 (m, 1 H), 7.54-7.45 (m, 1 H), 6.28-6.20 (m, 1 H), 6.08-5.99 (m, 1 H), 5.53 (app d, 1 H, *J* = 1.8 Hz), 4.57-4.43 (m, 1 H), 3.98-3.83 (m, 1 H); ¹³C NMR (acetone-d₆) δ 175.4, 148.4, 133.6, 133.1, 131.9, 129.4, 129.2, 124.6, 124.0, 71.1, 52.0; MS (EI) *m/z* (rel intensity) 171 ([M]⁺, 60), 160 (70), 130 (55), 105 (60), 83 (100); HRMS (EI) *m/z* calcd for C₁₁H₉NO 171.0684, found 171.0676.

(E)-2-Allyl-3-(hex-1-enyl)isoindolin-1-one (34)—According to general protocol A, 1-hexyne (**16**) (420 μL, 3.66 mmol), CH₂Cl₂ (10 mL), zirconocene hydrochloride (944 mg, 3.66 mmol), Me₃Al (1.0 M in CH₂Cl₂, 3.66 mL, 3.66 mmol) and **31** (500 mg, 1.83 mmol) afforded 341 mg (73%) of **34** as a colorless oil after purification on SiO₂ (EtOAc:hexanes, 3:7): IR (neat) 2957, 2926, 1697, 1468, 1396, 971, 748 cm⁻¹; ¹H NMR δ 7.83 (d, 1 H, *J* = 7.2 Hz), 7.57-7.38 (m, 2 H), 7.33 (d, 1 H, *J* = 7.5 Hz), 5.97 (dt, 1 H, *J* = 15.0, 6.9 Hz), 5.88-5.71 (m, 1 H), 5.22-4.97 (m, 3 H), 4.85 (d, 1 H, *J* = 9.3 Hz), 4.59 (dd, 1 H, *J* = 15.6, 4.5 Hz), 3.69 (dd, 1 H, *J* = 15.3, 7.2 Hz), 2.12 (app q, 2 H, *J* = 6.6 Hz), 1.48-1.28 (m, 4 H), 0.91 (t, 3 H, *J* = 6.9 Hz); ¹³C NMR δ 167.7, 145.0, 138.2, 133.1, 131.7, 131.4, 128.2, 126.0, 123.4, 122.9, 117.4, 62.9, 42.5, 31.8, 31.1, 27.0, 22.0, 13.8; MS (EI) *m/z* (rel intensity) 255 ([M]⁺, 30), 198 (100), 172 (55); HRMS (EI) *m/z* calcd for C₁₇H₂₁NO 255.1623, found 255.1627.

2-(Pent-4-enyl)isoindoline-1,3-dione (35)—A solution of 4-penten-1-ol (3.51 mL, 34.0 mmol), phthalimide **4** (5.00 mg, 34.0 mmol), and PPh₃ (8.92 g, 34.0 mmol) in THF (220 mL) was cooled to 0 °C and treated with DIAD (6.69 mL, 34.0 mmol) over 5 min. The reaction mixture was warmed to rt and stirred for 6 h. The solvent was evaporated, the residue was dissolved in EtOAc/hexanes (1:1, 100 mL) and the solids were filtered off. The filtrate was concentrated and chromatographed on SiO₂ (EtOAc:hexanes, 1:5) to yield 6.44 g (88%) of imide **35** as a colorless oil: IR (neat) 3466, 3077, 2939, 1773, 1641, 1397, 995, 720 cm⁻¹; ¹H NMR δ 7.88-7.81 (m, 2 H), 7.76-7.67 (m, 2 H), 5.90-5.73 (m, 1 H), 5.12-4.94 (m, 2 H), 3.70 (dt, 2 H, *J* = 7.5, 4.5 Hz), 2.20-2.07 (m, 2 H), 1.87-1.73 (m, 2 H); ¹³C NMR δ 168.1, 137.1, 133.7, 131.9, 122.9, 115.1, 37.3, 30.8, 27.4; MS (EI) *m/z* (rel intensity) 215 ([M]⁺, 45), 173 (70), 160 (100), 148 (80), 130 (80), 104 (90); HRMS (EI) *m/z* calcd for C₁₃H₁₃NO₂ 215.0946, found 215.0946.

3-Oxo-2-(pent-4-enyl)isoindolin-1-yl pivalate (36)—To a solution of imide **35** (4.42 g, 20.5 mmol) in MeOH (100 mL) was added sodium borohydride (776 mg, 20.5 mmol) at 0 °C. The resulting reaction mixture was stirred at 0 °C for 2 h and carefully quenched with H₂O. The aqueous layer was extracted with EtOAc (2×) and the organic layer was separated, dried (MgSO₄) and concentrated. The residue was dried to yield a white solid which was carried on to the next step without further purification. To a solution of the crude reduced phthalimide (4.23 g, 19.5 mmol) in THF (100 mL) was added Et₃N (8.15 mL, 58.5 mmol), pivaloyl chloride (3.61 mL, 29.3 mmol) and DMAP (119 mg, 0.975 mmol) at 0 °C and the reaction mixture was allowed to warm to rt and stirred at this temperature for 8 h. The mixture was quenched with sat. aq. NaHCO₃, extracted with EtOAc (2×), dried (MgSO₄) and concentrated. The residue was purified by chromatography on SiO₂ (EtOAc:hexanes, 1:5) to yield 4.70 g (76% over 2 steps) of **36** as a colorless oil: IR (neat) 3077, 2974, 2873, 1739, 1641, 1618, 1369, 1208, 1141, 959, 751 cm⁻¹; ¹H NMR δ 7.67 (d, 1 H, *J* = 6.0 Hz), 7.49-7.32 (m, 3 H), 6.89 (s, 1 H), 5.79-5.58 (m, 1 H), 4.91 (d, 1 H, *J* = 17.1 Hz), 4.84 (d, 1 H, *J* = 10.5 Hz), 3.73-3.55 (m, 1 H), 3.27-3.09 (m, 1 H), 2.07-1.93 (m, 2 H), 1.77-1.50 (m, 2 H), 1.11 (s, 9 H); ¹³C NMR δ 178.2, 167.4, 140.9, 137.1, 132.0, 131.8, 129.8, 123.2, 123.1, 114.9, 80.8, 39.5, 38.7, 30.7, 27.1, 26.6; MS (EI) *m/z*

z (rel intensity) 301 ($[M]^+$, 15), 246 (45), 216 (65), 200 (85), 146 (85), 133 (65); HRMS (EI) m/z calcd for $C_{18}H_{23}NO_3$ 301.1678, found 301.1681.

(E)-3-(Hex-1-enyl)-2-(pent-4-enyl)isoindolin-1-one (37)—According to general protocol A, hexyne (**16**) (420 μ L, 3.66 mmol), CH_2Cl_2 (10 mL), zirconocene hydrochloride (944 mg, 3.66 mmol), Me_3Al (1.0 M in CH_2Cl_2 , 3.66 mL, 3.66 mmol) and **36** (552 mg, 1.83 mmol) afforded 368 mg (72%) of **37** as a colorless oil after purification on SiO_2 (EtOAc:hexanes, 3:7): IR (neat) 3076, 2967, 2928, 2860, 1693, 1468, 1404, 972, 749 cm^{-1} ; 1H NMR δ 7.72 (d, 1 H, $J = 7.5$ Hz), 7.44–7.28 (m, 2 H), 7.23 (d, 1 H, $J = 7.2$ Hz), 5.94 (dt, 1 H, $J = 14.4, 6.6$ Hz), 5.80–5.63 (m, 1 H), 5.03–4.82 (m, 3 H), 4.75 (d, 1 H, $J = 9.0$ Hz), 3.83–3.65 (m, 1 H), 3.23–3.07 (m, 1 H), 2.13–1.93 (m, 4 H), 1.75–1.51 (m, 2 H), 1.42–1.19 (m, 4 H), 0.83 (t, 3 H, $J = 6.9$ Hz); ^{13}C NMR δ 167.6, 144.6, 137.5, 137.3, 131.7, 131.0, 127.9, 126.2, 122.9, 122.6, 114.7, 63.2, 39.4, 31.5, 30.8, 30.7, 27.3, 26.9, 21.8, 13.5; MS (EI) m/z (rel intensity) 283 ($[M]^+$, 20), 228 (70), 160 (100), 146 (50), 76 (50); HRMS (EI) m/z calcd for $C_{19}H_{25}NO$ 283.1936, found 283.1934.

(Z)-7,8,9,11a-Tetrahydro-5H-azepino[2,1-a]isoindol-5-one (38)—To a solution of **37** (150 mg, 0.529 mmol) in toluene (100 mL) was added $Ti(O^iPr)_4$ (157 μ L, 0.529 mmol) and Grubbs 2nd generation catalyst (22.5 mg, 0.0265 mmol) and the red solution was stirred at rt for 12 h and concentrated. The residue was purified by chromatography on SiO_2 (Acetone: CH_2Cl_2 , 0.4:9.6) to yield 70.7 mg (67%) of **38** as a colorless oil: IR (neat) 3024, 2926, 1680, 1469, 1419, 1298, 938, 709 cm^{-1} ; 1H NMR (CD_2Cl_2) δ 7.76 (dd, 1 H, $J = 6.9, 1.2$ Hz), 7.58–7.51 (m, 1 H), 7.46 (app t, 2 H, 7.2 Hz), 6.13–6.02 (m, 1 H), 6.02–5.91 (m, 1 H), 4.49 (dd, 1 H, $J = 11.4, 2.1$ Hz), 4.26 (ddd, 1 H, $J = 13.5, 6.6, 3.0$ Hz), 3.22 (ddd, 1 H, $J = 12.6, 9.6, 2.7$ Hz), 2.81 (app ddd, 1 H, $J = 15.9, 7.8, 2.4$ Hz), 2.53–2.26 (m, 2 H), 2.25–2.09 (m, 1 H); ^{13}C NMR (CD_2Cl_2) δ 167.5, 145.9, 133.0, 132.7, 131.6, 128.8, 128.5, 123.5, 122.4, 60.6, 41.6, 35.2, 28.2; MS (EI) m/z (rel intensity) 199 ($[M]^+$, 45), 145 (100), 117 (40), 90 (35); HRMS (EI) m/z calcd for $C_{13}H_{13}NO$ 199.0997, found 199.0991.

Acknowledgments

This work has been supported by the National Institutes of Health - NIGMS CMLD Program (GM067082) and Merck Research Laboratories. JGP thanks the ACS Division of Organic Chemistry for a Graduate Fellowship sponsored by Wyeth.

References

- i. Valencia E, Fajardo V, Freyer AJ, Shamma M. *Tetrahedron Lett* 1985;26:993.
- ii. Valencia E, Freyer AJ, Shamma M, Fajardo V. *Tetrahedron Lett* 1984;25:599.
- iii. Sorbera LA, Leeson PA, Silvestre J, Castaner J. *Drugs Fut* 2001;26:651.
- iv. Li S, Wang X, Guo H, Chen L. Yiyano Gongye 1985;16:543. *Chem Abstr* 1985;105:3788.
- v. Ferland JM, Demerson CA, Humber LG. *Can J Chem* 1985;63:361.
- vi. (a) Linden M, Hadler D, Hofmann S. *Hum Psychopharmacol* 1997;12:445. (b) Zhuang ZP, Kung MP, Mu M, Kung HF. *J Med Chem* 1998;41:157. [PubMed: 9457239] (c) Norman MH, Minick DJ, Rigdon GC. *J Med Chem* 1996;39:149. [PubMed: 8568802]
- vii. Achinami K, Ashizawa N, Kobayasui F. *Chem Abstr* 1991;115:255977j. Japanese Patent 03,133,955
- viii. Taylor EC, Zhou P, Jennings LD, Mao Z, Hu B, Jun JG. *Tetrahedron Lett* 1997;38:521.
- ix. For recent racemic syntheses, see: Yao T, Larock RC. *J Org Chem* 2005;70:1432. [PubMed: 15704980] Shen L, Hsung RP. *Org Lett* 2005;7:775. and references cited therein. [PubMed: 15727438] For recent asymmetric syntheses, see: Comins DL, Schilling S, Zhang Y. *Org Lett* 2005;7:95. and references cited therein. [PubMed: 15624986]

- x. For recent examples, see: (a)Comins DL, Schilling S, Zhang Y. *Org Lett* 2005;7:95. [PubMed: 15624986](b)Heaney H, Shuhaibar KF. *Synlett* 1995:47.(c)Othman M, Decroix B. *Synth Commun* 1996;26:2803.(d)Othman M, Pigeon P, Decroix B. *Tetrahedron* 1997;53:2495.
- xi. Boger DL, Lee JK, Goldberg J, Jin Q. *J Org Chem* 2000;65:1467. [PubMed: 10814111]
- xii. For recent examples, see: (a)Wipf P, Graham TH. *Org Biomol Chem* 2005;3:31–35.35 [PubMed: 15602595](b)Wipf P, Fletcher JM, Scarone L. *Tetrahedron Lett* 2005;46:5463. and references cited therein.
- xiii. (a) Wipf P, Lim S. *Angew Chem Int Ed Engl* 1993;32:1068. (b) Wipf P, Nunes RL, Ribe S. *Helv Chim Acta* 2002;85:3478. (c) Negishi EI. *Bull Chem Soc Jpn* 2007;80:233.
- xiv. For reviews, see: (a)Wipf P, Jahn H. *Tetrahedron* 1996;52:12853.(b)Wipf P, Nunes RL. *Tetrahedron* 2004;60:1269.(c)Wipf P, Kendall C. *Top Organomet Chem* 2005;8:1.
- xv. Hiemstra, H.; Speckamp, WN. *Comprehensive Organic Synthesis*. Trost, BM.; Fleming, I., editors. Vol. 2. Oxford; Pergamon: 1991. Chapter 4.5 (b) Maryanoff BE, Zhang HC, Cohen JH, Turchi IJ, Maryanoff CA. *Chem Rev* 2004;104:1431. [PubMed: 15008627]and references cited therein.
- xvi. (a) Nikitin KV, Andryukhova NP. *Can J Chem* 2000;78:1285. (b) Tranchant MJ, Moine C, Ben Othman R, Bousquet T, Othman M, Dalla V. *Tetrahedron Lett* 2006;47:4477. (c) Ben Othman R, Bousquet T, Fousse A, Othman M, Dalla V. *Org Lett* 2005;7:2825. [PubMed: 15987146] (d) Allin SM, Northfield CJ, Page MI, Slawin AMZ. *Tetrahedron Lett* 1999;40:141. (e) Ukaji Y, Tsukamoto K, Nasada Y, Shimizu M, Fujisawa T. *Chem Lett* 1993;2:221.
- xvii. Abrams SR. *Can J Chem* 1984;62:1333.
- xviii. Kitching MS, Clegg W, Elsegood MRJ, Griffin RJ, Golding BT. *Synlett* 1999;1:997.
- xix. Wipf P, Waller DL, Reeves JT. *J Org Chem* 2005;70:8096. [PubMed: 16277333]
- xx. Coulton S, Francois L, Southgate R. *Tetrahedron Lett* 1990;31:6923.
- xxi. (a) Washburn DG, Heidbrecht RW Jr, Martin SF. *Org Lett* 2003;5:3523. [PubMed: 12967315] (b) Neipp CE, Martin SF. *J Org Chem* 2003;68:8867. [PubMed: 14604356]
- xxii. (a) Batey RA, MacKay DB, Santhakumar V. *J Am Chem Soc* 1999;121:5075. (b) Batey RA, MacKay DB. *Tetrahedron Lett* 2000;41:9935.
- xxiii. For recent applications of cationic zirconocenes, see: Wipf P, Pierce JG, Zhuang N. *Org Lett* 2005;7:483. and references cited therein. [PubMed: 15673270]
- xxiv. Wipf P, Pierce JG. *Org Lett* 2006;8:3375. [PubMed: 16836409]
- xxv. (a) Cho CS, Wu X, Jiang L-H, Shim S-C, Choi H-J, Kim Tae J. *J Het Chem* 1998;35:265. (b) Burns B, Grigg R, Santhakumar V, Sridharan V, Stevenson P, Worakun T. *Tetrahedron* 1992;48:7297.
- xxvi. For a review, see: Fürstner A. *Angew Chem Int Ed Engl* 2000;39:3012. [PubMed: 11028025]
- xxvii. For related examples, see: (a)Ahn JB, Yun CS, Kim KH, Ha DC. *J Org Chem* 2000;65:9249. [PubMed: 11149882](b)Kim IS, Zee OP, Jung YH. *Org Lett* 2006;8:4101. [PubMed: 16928084]
- xxviii. Fürstner A, Langemann K. *J Am Chem Soc* 1997;119:9130.
- xxix. (a) White JD, Hrciar P, Yokoshi AFT. *J Am Chem Soc* 1998;120:7359. (b) Martin SF, Liao Y, Chen HJ, Patzel M, Ramser MN. *Tetrahedron Lett* 1994;35:6005.
- xxx. Buchwald SL, LaMaire SJ, Nielsen RB, Watson BT, King SM. *Org Synth* 1993;71:77.
- xxxi. Wipf P, Janjic J, Stephenson CRJ. *Org Biomol Chem* 2004;2:443. [PubMed: 14770219]
- xxxii. (a) Theus V, Schinz H. *Helv Chem Acta* 1956;39:1290. (b) Aitken RA, Atherton JI. *J Chem Soc, Perkin Trans 1* 1994:1281.

Figure 1.
Biologically active isoindolinones.

Scheme 1.

Proposed intramolecular addition of alkenylalanes to *N*-acyliminium ions. A: Zirconium-Catalyzed, Water-Accelerated Carboalumination or Hydrozirconation/Transmetalation; B: Elimination; C: Addition/Ring Closure.

Scheme 2.
Synthesis of lactam precursors.

Scheme 3.
Attempted cascade cyclization of **8** and **9**.

Scheme 4.

GC analysis of the water-accelerated carboalumination of heptyne in the presence and absence of **8**.

Scheme 5.

Attempted hydrozirconation and medium ring formation.

Scheme 6.

Addition to succinimide-derived iminium ion precursor.

Scheme 7.

Attempted medium ring formation via ring closing metathesis.

Scheme 8.

Application of ring closing metathesis to generate 5- and 7-membered fused ring systems.

Table 1

Reaction optimization for the addition of hexenylzirconocene to lactams 13–15.

Entry	Conditions	R (lactam acetal)	17 [%]
1	AgClO_4 , 24 h, rt	OMe (13)	Trace ^a
2	Me_2Zn , 12 h, rt	OMe (13)	-
3	Me_3Al , 12 h, rt	OMe (13)	43 ^b
4	Me_3Al , 4 h, rt	OAc (14)	30
5	Me_3Al , 4 h, rt	OPiv (15)	81
6	AgClO_4 , 12 h, rt	OPiv (15)	Complex mixture

^aProduct was formed as a mixture of alkene isomers.^bStarting material observed after 12 h.

Table 2

Variation of alkyne substrates.

Entry	R (alkyne)	Product [%]
1	C_4H_9 (16)	17 [81]
2	$n\text{-C}_6\text{H}_{11}$ (18)	19 [79]
3	$\text{CH}_2\text{CH}_2\text{OTBDPS}$ (20)	21 [71]
4	$\text{CH}_2\text{CH}_2\text{N}(\text{CO}_2\text{Me})\text{Ts}$ (22)	23 [62]

Table 3Water-accelerated carboalumination – addition to lactam **15**.

Entry	R	Product [%]
1	C_4H_9 (16)	27 [77]
2	Ph (28)	29 [77] ^a

^aProduct contains ~5% of a carbometallation regioisomer.