Phase I Trial of Daily Oral Polyphenon E in Patients With Asymptomatic Rai Stage 0 to II Chronic Lymphocytic Leukemia

Tait D. Shanafelt, Tim G. Call, Clive S. Zent, Betsy LaPlant, Deborah A. Bowen, Michelle Roos, Charla R. Secreto, Asish K. Ghosh, Brian F. Kabat, Mao-Jung Lee, Chung S. Yang, Diane F. Jelinek, Charles Erlichman, and Neil E. Kay

From the Department of Internal Medicine, Division of Hematology; Division of Biostatistics, and Departments of Immunology and Oncology, Mayo Clinic, Rochester, MN; and the Department of Chemical Biology, and the Ernest Mario School of Pharmacy Rutgers, the State University of New Jersey, Piscataway, NJ.

Submitted November 26, 2008; accepted February 9, 2009; published online ahead of print at www.jco.org on May 26, 2009.

Supported by grants from the National Cancer Institute (Grants No. CA113408 and CA6912), the CLL Global Research Foundation, CLL Topics, Commonwealth Foundation for Cancer Research, Polyphenon E International, and the Mayo Clinic Cancer Center.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Clinical Trials repository link available on JCO.org.

Corresponding author: Tait D. Shanafelt, MD, Mayo Clinic, 200 First Street SW, Rochester, MN 55905; e-mail: shanafelt.tait@mayo.edu.

© 2009 by American Society of Clinical Oncology

0732-183X/09/2723-3808/\$20.00

DOI: 10.1200/JCO.2008.21.1284

ABSTRACT

Purpose

To define the optimal dose of Polyphenon E for chronic daily administration and tolerability in patients with chronic lymphocytic leukemia (CLL).

Patients and Methods

Previously untreated patients with asymptomatic Rai stage 0 to II CLL were eligible for participation. Polyphenon E with a standardized dose of epigallocatechin-3-gallate (EGCG) was administered using the standard phase I design with three to six patients per dose level (range, 400 to 2,000 mg by mouth twice a day). Trough plasma EGCG levels were measured 1 month after initiation of therapy. Response was classified using the National Cancer Institute (NCI) Working Group (WG) Criteria.

Results

Thirty-three eligible patients were accrued to dose levels 1 to 8. The maximum-tolerated dose was not reached. The most common adverse effects included transaminitis (33%, all grade 1), abdominal pain (30% grade 1, 0% grade 2, and 3% grade 3), and nausea (39% grade 1 and 9% grade 2). One patient experienced an NCI WG partial remission. Other signs of clinical activity were also observed, with 11 patients (33%) having a sustained $\geq 20\%$ reduction in absolute lymphocyte count (ALC) and 11 (92%) of 12 patients with palpable adenopathy experiencing at least a 50% reduction in the sum of the products of all nodal areas during treatment. Trough plasma EGCG levels after 1 month of treatment ranged from 2.9 to 3,974 ng/mL (median, 40.4 ng/mL).

Conclusion

Daily oral EGCG in the Polyphenon E preparation was well tolerated by CLL patients in this phase I trial. Declines in ALC and/or lymphadenopathy were observed in the majority of patients. A phase II trial to evaluate efficacy using 2,000 mg twice a day began in November 2007.

J Clin Oncol 27:3808-3814. © 2009 by American Society of Clinical Oncology

INTRODUCTION

Chronic lymphocytic leukemia (CLL) is the most common subtype of leukemia in the United States and is presently an incurable disease. Most patients with CLL are now identified with early-stage disease when they are found to have an elevated lymphocyte count on a blood count obtained for other reasons.¹ Unfortunately, 50% of patients with early-stage disease have a biologically aggressive form of CLL that progresses²⁻⁵ and leads to premature death.² Unlike most other malignant disorders in which diagnosis at an early stage increases the probability that a patient can be cured, treatment of unselected early-stage CLL

patients with cytotoxic chemotherapy is associated with increased toxicity but no increase in survival.^{6,7} Thus, there is no effective intervention to alter the course of disease for patients with early-stage CLL at the present time.⁸

Green tea has long been touted as a health-promoting substance. Recent scientific investigations have identified the active chemical compounds in green tea designated tea polyphenols or catechins.^{9,10} Epigallocatechin-3-gallate (EGCG) is the major catechin in tea. A number of epidemiologic studies have linked consumption of green tea to decreased risk of cancer, and animal models have supported green tea's ability to prevent tumorigenesis.^{9,10} A number of mechanism have been

proposed for this effect of green tea including antiangiogenic properties,^{11,12} cell cycle arrest,^{9,10,13} effects on folate metabolism,¹⁴ effects on DNA damage,¹⁵ inhibition of telomerase,^{16,17} reductions in level of antiapoptotic proteins,¹⁸ proteasome inhibition,^{19,20} generation of reactive oxygen species,^{14,21-24} and demethylating/epigenetic effects.²⁵⁻²⁷ Minimal toxicity ($\leq$ grade 1) has been observed in clinical trials administering EGCG to healthy adults at doses of 800 mg/d.²⁸ In other trials of green tea extracts in patients with advanced malignancy, doses as high as 1 g/m² three times per day were tolerated for up to 6 months.²⁹

We recently investigated the effect of EGCG on CLL B cells *in vitro*.³⁰ EGCG induced caspase-dependent death in eight of 10 CLL samples.³⁰ Treatment of CLL B cells with EGCG at doses as low as 325 ng/mL induced significant downregulation of antiapoptotic proteins (MCL1 and XIAP) known to increase the resistance of CLL B cells to apoptosis.³⁰ EGCG also decreased vascular endothelial growth factor (VEGF) receptor (R1 and R2) phosphorylation and blocked VEGF-induced increases in MCL1 and XIAP.

After the publication of these findings, some patients with CLL and other low-grade lymphomas began using over-the-counter products containing tea polyphenols of their own initiative. We observed that several of these patients with low-grade B-cell malignancies had an objective clinical response.³¹ These observations, together with the preclinical findings³⁰ and the known favorable toxicity profile of EGCG in human patients,²⁸ provided a strong rationale for a clinical trial of EGCG in patients with asymptomatic Rai stage 0 to II CLL.

PATIENTS AND METHODS

Patient Eligibility

Previously untreated patients with asymptomatic Rai stage 0 to II CLL who did not meet National Cancer Institute Working Group (NCI WG)⁸ criteria to initiate chemotherapy were eligible for participation. The study was designed as a phase I/II trial with results of the phase I study reported here. Patients were required to have a confirmed diagnosis of CLL by standard criteria.⁸ Mantle cell lymphoma was excluded in all patients by fluorescent *in situ* hybridization (FISH) assessing for a t(11;14).

The eligibility criteria required participants to have an Eastern Cooperative Oncology Group performance status of 0 to 2, a life expectancy of $\geq$ 6 months, adequate renal function (creatinine $\leq$ 1.5 $\times$ the upper limit of normal or creatinine clearance $\geq$ 40 mL/min), and a total bilirubin less than 1.5 $\times$ the upper limit of normal (unless a result of Gilbert's disease). Concurrent chemotherapy, immunotherapy, radiotherapy, or corticosteroid treatment was not allowed. Patients were also not eligible if they had used over-the-counter green tea or green tea extracts with medicinal intent within 8 weeks of registration (half-life of EGCG is approximately 4 hours²⁸). The protocol was reviewed and approved by the Mayo Clinic Institutional Review Board and registered with the National Institutes of Health (clinicaltrials.gov). All patients provided written informed consent before study enrollment in accordance with the Declaration of Helsinki.

Protocol Treatment

Polyphenon E capsules (Polyphenon Pharma, New York, NY) containing approximately 200 mg of EGCG were supplied by the NCI. The phase I portion of the trial was designed to have up to eight dose levels using the standard phase I design of three to six patients per dose level. In December 2005, the NCI requested that dose escalation be temporarily suspended at dose level 2 to allow the US Food and Drug Administration (FDA) to review the results of animal studies evaluating EGCG toxicity. In the interim, additional patients were accrued at dose level 2 to better define the tolerability of 800 mg twice per day.

The trial was originally designed to administer Polyphenon E in the fasting state. In February 2006, the FDA temporarily suspended all human trials of EGCG to allow additional review of toxicity data. This led to interruption of treatment for regulatory reasons for eight patients on dose levels 1 and 2 after 2 to 5 months of treatment. The FDA later allowed trials of Polyphenon E to resume but mandated that all US clinical trials administer drug with food. Accordingly, the protocol was modified, and Polyphenon E was administered with food for dose levels 3 to 8. Response in the eight patients affected by this interruption was assessed on the date of interruption, at which point these individuals were censored for response analysis (six patients went off study, and two later resumed treatment after approximately a 5-month interruption).

All study participants were provided a medication diary to indicate the time and quantity of medication usage, which was reviewed at each follow-up visit. Patients remained on active treatment (dose range, 400 to 2,000 mg by mouth twice daily) for up to 6 months and were evaluated once every 4 weeks by physical examination and laboratory testing. Treatment was discontinued in the event of progressive disease, as defined by the NCI WG criteria,⁸ or excessive toxicity. At the completion of 6 months of active treatment, patients went to observation. With the approval of the treating hematologist, patients who had not experienced disease progression and who desired to remain on EGCG were provided EGCG-containing capsules at their assigned dose level for up to 12 additional months.

Toxicity was graded using NCI Common Terminology Criteria for Adverse Events version 3.0. Because there is a low threshold for toxicity in the treatment of patients not meeting standard criteria for progressive disease, cycle 1 adverse events more than grade 1 attributed to Polyphenon E that did not respond to supportive care were considered dose-limiting toxicity (DLT). Transient toxicity more than grade 1 that responded to supportive care was recorded as an adverse event but was not considered dose limiting. In general, for grade 2 adverse events attributed to Polyphenon E, therapy was held until symptoms resolved to $\leq$ grade 1 and then reinitiated along with supportive care measures at the same dose level. Because of the mandate of the FDA, the notable exception to this approach was the response to any elevation in transaminases (AST or ALT), in which case Polyphenon E was held for grade 1 adverse events regardless of attribution until these values returned to normal. Once the patient had normal levels, Polyphenon E was then reinitiated at the same dose level. If grade 1 transaminitis recurred, Polyphenon E was held and reinitiated at the next lower dose once AST or ALT values had returned to normal. Regardless of attribution, patients with $\geq$ grade 2 transaminitis were required by the FDA to permanently discontinue study treatment. For grade 3 to 4 adverse events attributed to study treatment or recurrent grade 2 events, Polyphenon E was held until symptoms resolved to $\leq$ grade 1 and then reinitiated at the next lower dose level along with supportive care measures. The maximum-tolerated dose (MTD) was defined as the dose level below the lowest dose that induced a DLT in at least one third of patients (at least two of six patients) after one cycle of therapy.

Risk Stratification Parameters

All patients underwent a comprehensive CLL prognostic evaluation including assessment of CD38 and ZAP-70 expression, FISH detectable cytogenetic defects, and *IgVH* gene mutation testing as previously described.³²⁻³⁴

Criteria for Response

Best response during the 6 months of active therapy was evaluated using the NCI WG criteria.⁸ Given the favorable toxicity profile of Polyphenon E in healthy adults¹² and the intention to evaluate the efficacy of this agent to delay or prevent disease progression in CLL, we also evaluated an additional response category termed biologic response. This category was prospectively defined in the study protocol after discussion and approval of this end point by the NCI because of recognition that anticancer botanicals such as Polyphenon E may work through unique, noncytotoxic mechanisms. These effects could be clinically beneficial by stabilizing disease rather than causing tumor regression. To be classified a biologic response, a reduction in the absolute lymphocyte count (ALC) of more than 20% from the pretreatment level for at least 2 months or a $\geq$ 30% reduction in all palpable lymphadenopathy without meeting the NCI WG criteria for partial remission (PR) was required.

Plasma Polyphenol Levels

Trough (approximately 12 hours after last dose) total plasma EGCG levels were measured at the end of the first month of therapy. Blood was collected into Vacutainer tubes (BD, Franklin Lakes, NJ) containing sodium heparin, and plasma was removed after centrifugation. Plasma was then aliquoted into cryotubes containing a small aliquot of ascorbate-EDTA solution as described by Lee et al³⁵ and stored at -80°C until analysis. Plasma levels of EGCG were then measured in the laboratory of the authors (C.S.Y.) using an established high-performance liquid chromatography procedure with a Coulchem (ESA, Chelmsford, MA) electrode array system.³⁶

Statistical Analysis

The primary outcome for this phase I trial was the number of DLTs and determination of the MTD, with clinical response the secondary outcome. Confirmed clinical responses (NCI WG complete remission or PR) on two consecutive evaluations at least 4 weeks apart and biologic responses, as defined earlier, were used as measures of clinical response and were summarized by simple descriptive statistics. Differences in response by key patient characteristics (ie, ZAP-70 status) were compared using the Fisher's exact test. Correlations between plasma EGCG level and dose/response were evaluated with the Spearman rank coefficient and the Wilcoxon rank sum test where appropriate.

RESULTS

Patient Demographics

Thirty-six patients were enrolled onto the study between August 2005 and September 2007. Median time from diagnosis to registration was 30.5 months (range, 0 to 170 months). Three patients were considered ineligible for evaluation because they took Polyphenon E incorrectly during cycle 1 (eg, took half the prescribed dose), which precluded accurate assessment of toxicity at the prescribed dose. The clinical characteristics of the 33 eligible patients are listed in Table 1. Most patients had favorable prognostic profiles on FISH, ZAP-70,

CD38, and *IgVH* gene mutation analysis at study entry consistent with the eligibility requirements that patients be asymptomatic and have earlier stage disease.

Toxicity and Tolerability

Median overall compliance with the prescribed dose of Polyphenon E as assessed using pill diaries was 99% (range, 55% to 104%). The MTD was not reached after completing accrual to dose levels 1 to 8. Only two patients experienced a DLT. A grade 2 dysphagia considered to be a DLT was observed during cycle 1 in one of six patients enrolled at dose level 4. In the second case, a patient on dose level 8 experienced grade 2 sweating, flatulence, abdominal distention, and nausea during cycle 1, which were all considered dose limiting. The number of DLTs by dose level is shown in Table 2.

Non-dose-limiting adverse effects during the 6 months of active treatment were generally mild. When examining all adverse events at least possibly attributed to therapy, five patients (15%) reported a maximum of a grade 2 event, and two patients (6%) reported a grade 3 event. The frequencies of specific adverse effects at least possibly attributed to therapy by Common Terminology Criteria for Adverse Events classification are listed in Table 3. Twenty-one patients (64%) completed six cycles of active therapy. Twelve patients discontinued therapy early; two refused further treatment, three experienced an adverse event, two experienced progression, one reported a new primary malignancy (cholangiocarcinoma), and four elected not to resume treatment when the temporary FDA suspension of all EGCG trials ended (see Patients and Methods).

Response to Therapy

One NCI WG PR was observed at dose level 8. In addition, a majority of patients had a reduction in ALC (Table 4). Reductions in ALC were transient in some patients, whereas in others, a steady, sustained stepwise reduction was observed throughout the 6 months of active therapy. Among the 15 patients (45%) who had a $\geq 20\%$ reduction in ALC, 11 (73%) had a sustained decrease $\geq 20\%$ for at least 2 months that fulfilled the criteria for biologic response. In a

Table 1. Patient Demographics and Clinical Characteristics

Patient Demographics and Clinical Characteristics	No. of Patients (N = 33)	%
Age, years		
Median	62	
Range	41-76	
Male	24	73
ALC, $\times 10^9/\text{L}$		
Median	35.1	
Range	5.4-355	
Rai stage		
0	18	55
I	11	33
II	4	12
ZAP-70 $\geq 20\%$	8	24
CD38* $\geq 30\%$	4	13
<i>IgVH</i> † unmutated	8	24
FISH		
del(13q14.2)	20	61
Normal	9	27
Trisomy 12	4	12

Abbreviations: ALC, absolute lymphocyte count; FISH, fluorescent in situ hybridization.

*Available for 32 of 33 patients.

†Although *IgVH* mutation testing was performed in all patients, two patients could not be sequenced and hence had uninterpretable *IgVH* test results.

Table 2. DLT

Dose Level	Dose (mg bid)	Total No. of Patients	No. of Patients With DLT
1	400	3	0
2	800	6*	0
3	1,000	3	0
4	1,200	6	1
5	1,400	3	0
6	1,600	3	0
7	1,800	3	0
8	2,000	6	1

NOTE. Cycle 1 adverse events $>$ grade 1 attributed to Polyphenon E that did not respond to supportive care were considered DLTs. Transient adverse events that responded to supportive care were recorded as adverse events but were not considered dose limiting (see Table 3).

Abbreviations: DLT, dose-limiting toxicity; bid, twice per day.

*Although no DLTs were observed among the first three patients accrued to dose level 2, the National Cancer Institute (NCI) temporarily suspended dose escalation to allow the US Food and Drug Administration to review the results of animal studies evaluating epigallocatechin-3-gallate toxicity. During this suspension, three additional patients were accrued at dose level 2 to better define tolerability at this dose level before receiving approval from the NCI to resume dose escalation (see Patients and Methods).

Table 3. Adverse Effects at Least Possibly Related to Treatment

Adverse Effect*	Grade 1		Grade 2		Grade 3	
	No. of Patients	%	No. of Patients	%	No. of Patients	%
Nausea	13	39	3	9	0	0
Abdominal pain	10	30	0	0	1	3
Transaminitis	11	33	0	0	0	0
Anorexia	9	27	2	6	0	0
Diarrhea	6	18	2	6	1	3
Dyspepsia	12	36	0	0	0	0
Flatulence	7	21	2	6	0	0
Fatigue	8	24	1	3	0	0
Hyperglycemia	1	3	0	0	0	0

*Classified according to the National Cancer Institute Common Terminology Criteria of Adverse Events.

per-protocol analysis of patients who completed six cycles of therapy, the ALC at the completion of therapy was less than baseline in 11 (52%) of 21 patients. Among the 12 patients with palpable adenopathy at enrollment, 11 (92%) experienced at least a 50% reduction in the sum of the products of all palpable nodal areas at some point during treatment. Overall, 18 (55%) of 33 patients fulfilled the criteria for a biologic response based on attaining either a sustained $\geq 20\%$ decline in ALC or a $\geq 30\%$ reduction in the sum of the products of all nodal areas at some point during the 6 months of active treatment. Achieving a biologic response appeared to be related to dose level; two (17%) of 12 patients treated on dose levels 1 to 3 achieved a biologic response compared with 16 (76%) of 21 patients treated on dose levels 4 to 8 (Table 5).

Correlative Prognostic Factors and Treatment Outcomes

The proportions of patients with a PR or biologic response by each prognostic parameter are listed in Table 6. Although no differences in response were observed based on *IgVH*, ZAP-70, or CD38, individuals with trisomy 12 on FISH appeared less likely to achieve a response ($P = .04$).

Plasma Polyphenon Levels

Trough total plasma EGCG levels after 1 month of therapy ranged from 2.9 to 3,974 ng/mL (median, 40.4 ng/mL). Plasma levels did not clearly correlate with dose, the likelihood of attaining a PR/biologic response, or the degree of reduction in ALC after 1 month of

treatment (Fig 1), suggesting that sensitivity to Polyphenon E may relate more to characteristics of the leukemic clone rather than the EGCG plasma level achieved.

DISCUSSION

Daily oral EGCG in the Polyphenon E preparation was well tolerated up to the maximum dose tested (2,000 mg orally twice per day) in this phase I trial for patients with asymptomatic Rai stage 0 to II CLL. The most severe toxicity was $\leq$ grade 1 for more than 80% of patients, and only two (6%) of 33 patients experienced $\geq$ grade 3 toxicity. Importantly, signs of clinical activity were observed, with one patient achieving an NCI WG PR and more than 50% of study patients attaining a sustained decline in ALC of $\geq 20\%$ and/or a $\geq 50\%$ reduction in lymphadenopathy at some point during treatment. Clinical activity seemed to be related to dose because more than 75% of patients treated on dose levels 4 to 8 (1,200 to 2,000 mg twice per day) achieved at least a biologic response compared with less than 20% of patients treated on dose levels 1 to 3.

Despite extensive data demonstrating that 50% to 70% of patients with asymptomatic, earlier stage CLL will experience progression and require treatment,^{2,6,37} such patients are presently managed with observation.³⁸ The phase I data presented here suggests that Polyphenon E may have clinical activity in these patients, and the fact that this treatment is very well tolerated suggests that Polyphenon E merits additional evaluation as a disease-stabilizing agent. With the

Table 4. Best Response in ALC and Nodes

Response	No. of Patients	%
Best reduction in ALC		
At least 10% decline	26	79
At least 20% decline	15	45
At least 30% decline	10	30
At least 40% decline	5	15
At least 50% decline	2	6
Best reduction in nodes		
At least 50% reduction in sum of products	11*	92

Abbreviation: ALC, absolute lymphocyte count.

*Twelve patients had palpable lymphadenopathy at study entry.

Table 5. Response by Dose Level

Dose Level	Dose (mg bid)	No. of Patients	No. of Patients With Partial or Biologic Response
1	400	3	1
2	800	6	0
3	1,000	3	1
4	1,200	6	4
5	1,400	3	3
6	1,600	3	2
7	1,800	3	2
8	2,000	6	5

Abbreviation: bid, twice per day.

Table 6. Biologic Response by Prognostic Parameters

Prognostic Parameter	No. of Patients	Patients With at Least Biologic Response		P
		No.	%	
ZAP-70				.70
Negative (< 20%)	25	13	52	
Positive (≥ 20%)	8	5	63	
CD38				1.0
Negative (< 30%)	28	15	54	
Positive (≥ 20%)	4	2	50	
IgVH				1.0
Mutated	23	13	57	
Unmutated	8	4	50	
FISH				.04
del(13q14.2)	20	11	55	
Normal	9	7	78	
Trisomy 12	4	0	0	

Abbreviation: FISH, fluorescent in situ hybridization.

exception of trisomy 12, the clinical activity observed with Polyphenon E in the present trial did not seem to be influenced by ZAP-70, CD38, or IgVH gene mutation status. This observation suggests that Polyphenon E may have activity in Rai stage 0 to II patients whose prognostic parameters suggest they are at high risk for disease progression.³⁹

The mechanism(s) by which EGCG affects CLL B cells in vivo is unknown. In vitro studies of CLL B cells show that EGCG decreases VEGF receptor phosphorylation and downregulates the expression of MCL-1 and XIAP.³⁰ Indeed, EGCG seems to have greater ability to antagonize MCL-1 than other BCL-2 family member inhibitory compounds in development as CLL therapies including gossypol, apogossypol, and ABT 737.⁴⁰ Other studies suggest that EGCG is a potent inhibitor of BCL-2 at nanomolar concentrations.¹⁸ Although these mechanisms are of particular interest because of the importance of BCL-2 family members,⁴¹⁻⁴³ and particularly MCL-1,⁴⁴⁻⁴⁷ in CLL

B-cell apoptotic resistance, numerous other mechanisms of action have been proposed for the antitumor effects of EGCG observed in vitro.^{9-27,40,48}

Because in vitro studies suggest EGCG may have activity against a wide range of human malignancies including prostate, colon, lung, and breast cancer,⁴⁹⁻⁵² the results of this phase I trial may have relevance for clinical studies of EGCG in other diseases. In this respect, it should be noted that the median age of patients in our study was more than 60 years old and that 70% of participants were men. Although the MTD was not reached in our trial, we chose not to escalate beyond the 2,000 mg twice per day dose level because the high number of capsules required to achieve this dose (10 capsules twice per day) made further dose escalation problematic. It may be possible to escalate beyond this dose level if a more concentrated preparation becomes available. Notably, the plasma levels attained on dose levels 3 to 8 in our study were achieved administering Polyphenon E in the fed state, whereas fasting administration would likely allow patients to achieve similar or higher plasma levels at lower Polyphenon E doses.⁵³

The reason for the wide variation in trough plasma EGCG levels observed in our study is unknown. This variation could relate to differences in absorption when administered with food (eg, total calories, proportion of fat/protein/carbohydrate), metabolism, or protein binding. It is also notable that trough plasma EGCG levels did not clearly relate to the clinical activity of Polyphenon E observed among CLL patients. This observation could indicate other pharmacokinetic parameters (eg, peak levels, area under the curve, and so on) or characteristics of the leukemic clone may relate to the clinical effects of Polyphenon E in CLL patients more strongly than trough plasma levels.

In conclusion, daily oral EGCG in the Polyphenon E preparation was well tolerated at doses up to 2,000 mg twice per day for up to 6 months in patients with asymptomatic Rai stage 0 to II CLL. Declines in ALC and lymphadenopathy during Polyphenon E therapy were observed in the majority of patients. A phase II trial evaluating the efficacy of Polyphenon E (2,000 mg twice per day) in patients with asymptomatic Rai stage 0 to II CLL was initiated in November 2007.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: None **Consultant or Advisory Role:** Neil E. Kay, Polyphenon E International **Stock Ownership:** None **Honoraria:** Neil E. Kay, Polyphenon E International **Research Funding:** Tait D. Shanafelt, Polyphenon E International; Chung S. Yang, Polyphenon E International; Neil E. Kay, Polyphenon E International **Expert Testimony:** None **Other Remuneration:** None

AUTHOR CONTRIBUTIONS

Conception and design: Tait D. Shanafelt, Neil E. Kay

Financial support: Tait D. Shanafelt, Charles Erlichman, Neil E. Kay

Fig 1. Plasma levels of epigallocatechin-3-gallate (EGCG) and response. Figure shows the relationship between absolute lymphocyte count (ALC) reduction at 1 month (x-axis) and trough plasma EGCG level (y-axis) at 1 month. EGCG plasma levels are shown on a log normal scale as a result of the wide variation in plasma levels.

Administrative support: Tait D. Shanafelt, Michelle Roos, Charles Erlichman, Neil E. Kay

Provision of study materials or patients: Tait D. Shanafelt, Tim G. Call, Clive S. Zent, Deborah A. Bowen, Neil E. Kay

Collection and assembly of data: Tait D. Shanafelt, Tim G. Call, Clive S. Zent, Betsy LaPlant, Deborah A. Bowen, Michelle Roos, Charla R.

Secreto, Asish K. Ghosh, Brian F. Kabat, Mao-Jung Lee, Chung S. Yang, Diane F. Jelinek, Neil E. Kay

Data analysis and interpretation: Tait D. Shanafelt, Tim G. Call, Clive S. Zent, Betsy LaPlant, Deborah A. Bowen, Brian F. Kabat, Chung S. Yang, Diane F. Jelinek, Charles Erlichman, Neil E. Kay

Manuscript writing: Tait D. Shanafelt

Final approval of manuscript: Tait D. Shanafelt, Tim G. Call, Clive S. Zent, Betsy LaPlant, Deborah A. Bowen, Michelle Roos, Charla R.

Secreto, Asish K. Ghosh, Brian F. Kabat, Mao-Jung Lee, Chung S. Yang, Diane F. Jelinek, Charles Erlichman, Neil E. Kay

REFERENCES

- Shanafelt T, Byrd J, Call T, et al: Initial management of newly diagnosed, early stage, chronic lymphocytic leukemia: An update for the internist. *Ann Intern Med* 145:435-447, 2006
- Call T, Phylly R, Noel P, et al: Incidence of chronic lymphocytic leukemia in Olmsted County, Minnesota, 1935 through 1989, with emphasis on changes in initial stage at diagnosis. *Mayo Clin Proc* 69:323-328, 1994
- Catovsky D, Fooks J, Richards S: Prognostic factors in chronic lymphocytic leukaemia: The importance of age, sex and response to treatment in survival. *Br J Haematol* 72:141-149, 1989
- Shanafelt TD, Geyer S, Kay N: Prognosis at diagnosis: Integrating molecular biologic insights into clinical practice for patients with CLL. *Blood* 103:1202-1210, 2004
- Knauf WU, Langenmayer I, Mohr BEB, et al: Serum levels of soluble CD23, but not soluble CD25, predict disease progression in early stage B-cell chronic lymphocytic leukemia. *Leuk Lymphoma* 27:523-532, 1997
- Dighiero G, Maloum K, Desablens B, et al: Chlorambucil in indolent chronic lymphocytic leukemia: French Cooperative Group on Chronic Lymphocytic Leukemia. *N Engl J Med* 338:1506-1514, 1998
- Shustik C, Mick R, Silver R, et al: Treatment of early chronic lymphocytic leukemia: Intermittent chlorambucil versus observation. *Hematol Oncol* 6:7-12, 1988
- Cheson BD, Bennett JM, Grever M, et al: National Cancer Institute-sponsored Working Group guidelines for chronic lymphocytic leukemia: Revised guidelines for diagnosis and treatment. *Blood* 87:4990-4997, 1996
- Yang CS: Inhibition of carcinogenesis by tea. *Nature* 389:134-135, 1997
- Yang C, Maliakal P, Meng X: Inhibition of carcinogenesis by tea. *Annu Rev Pharmacol Toxicol* 42:25-54, 2002
- Lamy S, Gingras D, Beliveau R: Green tea catechins inhibit vascular endothelial growth factor receptor phosphorylation. *Cancer Res* 62:381-385, 2002
- Jung Y, Kim M, Shin B, et al: EGCG, a major component of green tea, inhibits tumor growth by inhibiting VEGF induction in human colon carcinoma cells. *Br J Cancer* 84:844-850, 2001
- Lin J: Cancer chemoprevention by tea polyphenols through modulating signal transduction pathways. *Arch Pharm Res* 25:561-575, 2002
- Navarro-Peran E, Cabezas-Herrera J, Garcia-Canovas F, et al: The antifolate activity of tea catechins. *Cancer Res* 65:2059-2064, 2005
- Ahmad N, Feyes DK, Nieminen AL, et al: Green tea constituent epigallocatechin-3-gallate and induction of apoptosis and cell cycle arrest in human carcinoma cells. *J Natl Cancer Inst* 89:1881-1886, 1997
- Naasani I, Seimiya H, Tsuruo T: Telomerase inhibition, telomere shortening, and senescence of cancer cells by tea catechins. *Biochem Biophys Res Commun* 249:391-396, 1998
- Naasani I, Oh-Hashi F, Oh-Hara T, et al: Blocking telomerase by dietary polyphenols is a major mechanism for limiting the growth of human cancer cells in vitro and in vivo. *Cancer Res* 63:824-830, 2003
- Leone M, Zhai D, Sareth S, et al: Cancer prevention by tea polyphenols is linked to their direct inhibition of antiapoptotic Bcl-2-family proteins. *Cancer Res* 63:8118-8121, 2003
- Nam S, Smith DM, Dou QP: Ester bond-containing tea polyphenols potentially inhibit proteasome activity in vitro and in vivo. *J Biol Chem* 276:13322-13330, 2001
- Landis-Piwowar KR, Wan SB, Wiegand RA, et al: Methylation suppresses the proteasome-inhibitory function of green tea polyphenols. *J Cell Physiol* 213:252-260, 2007
- Shammas M, Kole H, Batchu R, et al: Specific killing of multiple myeloma cancer cells by epigallocatechin-3-gallate extracted from green tea. *Blood* 104:2461, 2004 (abstr)
- Furukawa A, Oikawa S, Murata M, et al: Epigallocatechin gallate causes oxidative damage to isolated and cellular DNA. *Biochem Pharmacol* 66:1769-1778, 2003
- Azam S, Hadi N, Khan N, et al: Prooxidant property of green tea polyphenols epicatechin and epigallocatechin-3-gallate: Implications for anticancer properties. *Toxicol In Vitro* 18:555-561, 2004
- Nakagawa H, Hasumi K, Woo J, et al: Generation of hydrogen peroxide primarily contributes to the induction of FE(II) dependent apoptosis in Jurkat cells by epigallocatechin gallate. *Carcinogenesis* 25:1567-1574, 2004
- Fang MZ, Wang Y, Ai N, et al: Tea polyphenol (-)-epigallocatechin-3-gallate inhibits DNA methyltransferase and reactivates methylation-silenced genes in cancer cell lines. *Cancer Res* 63:7563-7570, 2003
- Fang M, Chen D, Yang CS: Dietary polyphenols may affect DNA methylation. *J Nutr* 137:223S-228S, 2007
- Berlitch JB, Liu C, Love WK, et al: Epigenetic and genetic mechanisms contribute to telomerase inhibition by EGCG. *J Cell Biochem* 103:509-519, 2008
- Chow H-HS, Cai Y, Hakim IA, et al: Pharmacokinetics and safety of green tea polyphenols after multiple-dose administration of epigallocatechin gallate and Polyphenon E in healthy individuals. *Clin Cancer Res* 9:3312-3319, 2003
- Pisters KM, Newman RA, Coldman B, et al: Phase I trial of oral green tea extract in adult patients with solid tumors. *J Clin Oncol* 19:1830-1838, 2001
- Lee YK, Bone ND, Strega AK, et al: VEGF receptor phosphorylation status and apoptosis is modulated by a green tea component, epigallocatechin-3-gallate (EGCG), in B-cell chronic lymphocytic leukemia. *Blood* 104:788-794, 2004
- Shanafelt TD, Lee YK, Call TG, et al: Clinical effects of oral green tea extracts in four patients with low grade B-cell malignancies. *Leuk Res* 30:707-712, 2006
- Shanafelt TD, Geyer SM, Bone ND, et al: CD49d expression is an independent predictor of overall survival in patients with chronic lymphocytic leukaemia: A prognostic parameter with therapeutic potential. *Br J Haematol* 140:537-546, 2008
- Dewald G, Brockman S, Paternoster S, et al: Chromosome anomalies detected by interphase fluorescence in hybridization: Correlation with significant biological features of chronic lymphocytic leukemia. *Br J Haematol* 121:287-295, 2003
- Jelinek DF, Tschumper RC, Geyer SM, et al: Analysis of clonal B-cell CD38 and immunoglobulin variable region sequence status in relation to clinical outcome for B-cell chronic lymphocytic leukaemia. *Br J Haematol* 115:854-861, 2001
- Lee MJ, Wang ZY, Li H, et al: Analysis of plasma and urinary tea polyphenols in human subjects. *Cancer Epidemiol Biomarkers Prev* 4:393-399, 1995
- Lee MJ, Prabhu S, Meng X, et al: An improved method for the determination of green and black tea polyphenols in biomatrices by high-performance liquid chromatography with coulometric array detection. *Anal Biochem* 279:164-169, 2000
- Hamblin TJ, Davis Z, Gardiner A, et al: Unmutated Ig V(H) genes are associated with a more aggressive form of chronic lymphocytic leukemia. *Blood* 94:1848-1854, 1999
- Hallek M, Cheson BD, Catovsky D, et al: Guidelines for the diagnosis and treatment of chronic lymphocytic leukemia: A report from the International Workshop on Chronic Lymphocytic Leukemia updating the National Cancer Institute-Working Group 1996 guidelines. *Blood* 111:5446-5456, 2008
- Rassenti LZ, Jain S, Keating MJ, et al: Relative value of ZAP-70, CD38, and immunoglobulin mutation status in predicting aggressive disease in chronic lymphocytic leukemia. *Blood* 112:1923-1930, 2008
- Zhai D, Jin C, Satterthwait AC, et al: Comparison of chemical inhibitors of antiapoptotic Bcl-2-family proteins. *Cell Death Differ* 13:1419-1421, 2006
- Reed JC, Kitada S, Kim Y, et al: Modulating apoptosis pathways in low-grade B-cell malignancies using biological response modifiers. *Semin Oncol* 29:10-24, 2002
- Robertson LE, Plunkett W, McConnell K, et al: Bcl-2 expression in chronic lymphocytic leukemia and its correlation with the induction of apoptosis and clinical outcome. *Leukemia* 10:456-459, 1996
- Schimmer A, Munk-Pedersen I, Minden M, et al: Bcl-2 and apoptosis in chronic lymphocytic leukemia. *Curr Treat Options Oncol* 4:211-218, 2003
- Pepper C, Lin TT, Pratt G, et al: Mcl-1 expression has in vitro and in vivo significance in chronic lymphocytic leukemia and is associated with other

poor prognostic markers. *Blood* 112:3807-3817, 2008

45. Longo PG, Laurenti L, Gobessi S, et al: The Akt/Mcl-1 pathway plays a prominent role in mediating antiapoptotic signals downstream of the B-cell receptor in chronic lymphocytic leukemia B cells. *Blood* 111:846-855, 2008

46. Bannerji R, Kitada S, Flinn I, et al: Apoptotic-regulatory and complement -protecting protein expression in chronic lymphocytic leukemia: Relationship to in vivo rituximab resistance. *J Clin Oncol* 21:1466-1471, 2003

47. Hussain SR, Cheney CM, Johnson AJ, et al: Mcl-1 is a relevant therapeutic target in acute and

chronic lymphoid malignancies: Down-regulation enhances rituximab-mediated apoptosis and complement-dependent cytotoxicity. *Clin Cancer Res* 13:2144-2150, 2007

48. Zhang Q, Tang X, Lu Q, et al: Green tea extract and (–)-epigallocatechin-3-gallate inhibit hypoxia- and serum-induced HIF-1 α protein accumulation and VEGF expression in human cervical carcinoma and hepatoma cells. *Mol Cancer Ther* 5:1227-1238, 2006

49. Ju J, Lu G, Lambert JD, et al: Inhibition of carcinogenesis by tea constituents. *Semin Cancer Biol* 17:395-402, 2007

50. Khan N, Mukhtar H: Tea polyphenols for health promotion. *Life Sci* 81:519-533, 2007

51. Khan N, Mukhtar H: Multitargeted therapy of cancer by green tea polyphenols. *Cancer Lett* 269: 269-280, 2008

52. Khan N, Afaq F, Mukhtar H: Cancer chemoprevention through dietary antioxidants: Progress and promise. *Antioxid Redox Signal* 10:475-510, 2008

53. Chow H-HS, Hakim IA, Vining D, et al: Effects of dosing condition on the oral bioavailability of green tea catechins after single-dose administration of Polyphenon E in healthy individuals. *Clin Cancer Res* 11:4627-4633, 2005