Of patients who received post reconstructive radiotherapy, 66% had lymph node involvement. This patient group can readily be identified by sentinel lymph node biopsy prior to reconstruction and counselled/treated accordingly using the options I described in my original article. Indeed, the authors indicate the potential benefits of delayed-immediate reconstruction in such circumstances.

It is also interesting to note that almost all patients had a free TRAM flap reconstruction (96%). This high percentage would be unusual in the UK. As this is an autologous tissue reconstruction, the extent of the problems described may not be applicable to patients with implant only or LD flap and implant reconstructions. The implant-based reconstructions tend to sit prominent on the chest wall in the supine position and the field arrangements may not carry the same risks to the surrounding tissues as with an autologous reconstruction.

There is also no matching of the control group for body mass index (presumably higher than average in the treatment group as 96% had an autologous procedure), chest wall morphology or original breast size, which all may affect radiation field arrangement.

The issue of inclusion of surrounding viscera in tangential fields for radiotherapy to the chest wall following mastectomy is not limited to those having immediate reconstruction; similar problems are encountered when giving radiotherapy following breast conservation.

In respect of the psychological issues, there is limited prospective data as commented by Wallace. Prospective trials of psychological assessments pose their own particular difficulties; randomisation is not usually possible in this setting and patient preference introduces strong selection bias. It is interesting to note that the comparison between the two patient groups was based on only 10 delayed reconstruction patients.⁵ The patient groups are also dissimilar in terms of age and use of chemotherapy.

'Those opting for immediate reconstruction have a higher incidence of psychosocial impairment and functional disability' is hardly surprising in the pre-operative setting compared to those undergoing a delayed procedure in view of a recent diagnosis of a breast malignancy and recognised as such by the authors.⁴

There are many aspects to consider when counselling patients with regard to the issues surrounding immediate breast reconstruction. The patient must be fully informed as to the processes involved through discussions with surgeon, breast-care nurse and family. Reviews of photographic examples and possibly meetings with patients who have already had breast reconstruction combined with written information are all important in this process. Providing the treatment plan is oncologically safe and discussed in the MDT setting and patients appropriately counselled then high standards of treatment and satisfaction should be possible.

References

1. Motwani SB, Strom EA, Schechter NR, Butler CE, Lee GK, Langstein HN *et al.* The impact of immediate breast reconstruction on the technical delivery of postmastectomy radiotherapy. *Int J Radiat Oncol Biol Phys* 2006; **66**: 76–82.
2. The Royal College of Radiologists' Clinical Oncology information network. Guidelines on the non-surgical management of breast cancer. *Clin Oncol* 1999; **11**: S104.
3. Harcourt DM, Rumsey NJ, Ambler NR, Cawthorn SJ, Reid CD, Maddox PR *et al.* The psychological effect of mastectomy with or without breast reconstruction: a prospective, multicenter study. *Plast Reconstr Surg* 2003; **111**: 1060–8.
4. Roth RS, Lowery JC, Davis J, Wilkins EG. Quality of life and affective distress in women seeking immediate versus delayed breast reconstruction after mastectomy for breast cancer. *Plast Reconstr Surg* 2005; **116**: 993–1002.

LETTER

doi 10.1308/003588408X321846

Medical devices and medical training

NV TODD, A GHOLKAR

Regional Neurosciences Centre, Newcastle General Hospital, Newcastle, UK

CORRESPONDENCE TO

NV Todd, Consultant Neurosurgeon and Spinal Surgeon, Regional Neurosciences Centre, Newcastle General Hospital, Westgate Road, Newcastle NE4 6BE, UK
E: nick.todd@nuth.nhs.uk

There are mandatory standards for the management of medical devices.^{1,2}

Junior doctors must have specific training for the medical devices they use, which must be documented. At first glance, it seems a major undertaking formally to train every junior doctor in the use of every medical device they currently use. However, junior doctors do receive oral training in using medical devices and much of what we are being asked to do can be achieved simply, by formally acknowledging skills doctors currently have. For surgical trainees this can be based upon their log books. Using neurosurgery as an example: no consultant is going to allow a junior doctor to perform a craniotomy (*e.g.* to evacuate a haematoma) if he or she is not competent to use a perforator or craniotome (powerful air/electric drills). *Ergo*, any junior doctor who can perform a craniotomy unaided can be signed off to use those medical devices. A list of neurosurgical devices can be linked to the operations that junior doctors currently perform; examples are given in Table 1. As trainees become more proficient, more devices become authorised. This requires minimal increased bureaucracy and is applicable to all junior doctors in training.

Table 1 Examples of operations linked to particular neuro-surgical devices

Operation	Medical device used
Ventricular drain	Perforator
Evacuation haematoma	Craniotome
Microdiscectomy	Operating microscope
Tumour surgery	Cavitron
Cervical spine surgery	Cervical plates/screws

We will need formal standards for all surgeons (including consultants). Prior to using a device that is new to that surgeon or if a wholly new device is introduced into a department, formal and documented training is required. We might recommend the following prior to operating solo: (i) attendance at a manufacturer's training course; (ii) assisting a competent surgeon; or (iii) being assisted by a competent surgeon.

A record of medical device training must be maintained: surgeons must operate within their documented competencies.

References

1. Department of Health. *Standards for Better Health*. London: DH, 2006.
2. Medicines and Healthcare Products Regulatory Agency. *Managing Medical Devices*. Device Bulletin DB 2006(05). London: MHRA, 2006.

COMMENT ON

doi 10.1308/003588408X242051

A Al-Allak, S Sarasin, S Key, G Morris-Stiff. Wedding rings are not a significant source of bacterial contamination following surgical scrubbing. *Ann R Coll Surg Engl* 2008; **90**: 133–5

doi 10.1308/003588408X321819

Microbial load and decontamination in the theatre suite

SA CAIRNS¹, ET HEGARTY²

¹Welsh Centre for Burns and Plastic Surgery, Morriston Hospital, Swansea, UK

²Department of Orthopaedic Surgery, Whipps Cross Hospital, London, UK

CORRESPONDENCE TO

SA Cairns, Welsh Centre for Burns and Plastic Surgery, Morriston Hospital, Swansea, UK

E: scottcairns@doctors.org.uk

This paper represents the latest in a line of publications down-playing the role of individual pre-operative preparation;^{1–4} we

would counsel caution. While individually each aspect (*e.g.* wearing hats, masks or overshoes) may be proven not to increase bacterial counts in operating theatres, we are concerned that cumulatively they may undermine the ethos of theatre being as sterile as possible. Which of us would choose to be operated on, or assisted by, a surgeon or scrub nurse who is hatless, ring-clad and wearing outdoor shoes?

As surgeons given the privilege of operating on patients, our primary concern should be our duty of care to the patient. It is our belief all possible sterile precautions should be undertaken, with a doctrine of such precautions being mandatory.

References

1. Hubble MJ, Weale AE, Perez JV, Bowker KE, MacGowan AP, Bannister GC. Clothing in laminar-flow operating theatres. *J Hosp Infect* 1996; **32**: 1–7.
2. Humphreys H, Marshall RJ, Ricketts VE, Russell AJ, Reeves DS. Theatre over-shoes do not reduce operating theatre floor bacterial counts. *J Hosp Infect* 1991; **17**: 117–23.
3. Bischoff WE, Tucker BK, Wallis ML, Reboussin BA, Pfaller MA, Hayden FG *et al.* Preventing the airborne spread of *Staphylococcus aureus* by persons with the common cold: effect of surgical scrubs, gowns, and masks. *Infect Control Hosp Epidemiol* 2007; **28**: 1148–54.
4. Lipp A, Edwards P. Disposable surgical face masks: a systematic review. *Can Oper Room Nurs J* 2005; **23**: 20–1, 24–5, 33–8.

AUTHORS' RESPONSE

doi 10.1308/003588408X321828

A AL-ALLAK¹, S SARASIN², S KEY², G MORRIS-STIFF²

¹Department of Surgery, Princess of Wales Hospital, Bridgend, UK

²Department of Surgery, Royal Glamorgan Hospital, Ynysmaerdy, Llantrisant, Rhondda Cynon Taf, UK

CORRESPONDENCE TO

Asmaa Al-Allak, Department of Surgery, Princess of Wales Hospital, Coity Road, Bridgend CF31 1RQ, UK

E: asmaa@doctors.org.uk

Whilst we agree whole-heartedly that sterile conditions are a must in theatre, we are sure that the authors would also agree that, in this era of evidence-based medicine when there is no evidence or even negative evidence for a practice that is taken for granted, then its validity must be questioned. We would suggest that many such 'traditional' practices are re-evaluated and would greatly appreciate the performance of further studies to improve the quality of the available scientific evidence.