injuries managed non-operatively, to allow surgeons both to manage and advise these patients, based on an increasingly established evidence base.

References

1. Strange GR. *Paediatric Emergency Medicine*. Section 2: Trauma Care. P 106.
2. Tugnoli G, Casali M, Villani S, Biscardi A, Baldoni F. The treatment of splenic injuries from splenectomy to non-operative management : our experience on 429 case. *Ann Ital Chir* 2003; **74**: 37–41, discussion 41–3.
3. Crawford RS, Tabbara M, Sheridan R, Spaniolas K, Velmahos GC. Early discharge after nonoperative management for splenic injuries : increased patient risk caused by late failure? *Surgery* 2007; **142**: 337–42.
4. Chen LY, Shih HC, Wu JJ, Wen YS, Huang MS, Huang CI *et al*. The role of diagnostic algorithms in the management of blunt splenic injury. *J Chin Med Assoc* 2005; **68**: 373–8.
5. Yang J, Gao JM, Jean-Claude B. Non-operative management of adult blunt splenic injuries. *Chin J Traumatol* 2006; **9**: 246–8.
6. Lunca S, Romedea N, Morosanu C, Mihalache C, Mihalache S. Traumatic injuries to the spleen. *Rev Med Chir Soc Med Nat Iasi* 2005; **109**: 281–5.
7. Schweizer W, Bohlen L, Dennison A, Blumgart LH. Prospective study in adults of splenic preservation after traumatic rupture. *Br J Surg* 1992; **79**: 1330–3.
8. Sanders MN, Civil I. Adult splenic injuries: treatment patterns and predictive indicators. *Aust NZ J Surg* 1999; **69**: 430–2.
9. Dulchavsky SA, Lucas CE, Ledgerwood AM, Grabow D. Wound healing of the injured spleen with and without splenorrhaphy. *J Trauma* 1987; **27**: 1155–60.
10. Pachter HL, Guth AA, Hofstetter SR, Spencer FC. Changing patterns in the management of splenic trauma: the impact of non-operative management. *Ann Surg* 1998; **227**: 708–19.
11. Lyass S, Sela T, Lebensart P, Muggia-Sullam M. Follow-up imaging studies of blunt splenic injury: do they influence management ? *Isr Med Assoc J* 2001; **3**: 731–3.
12. Shafi S, Gilbert JC, Irish MS, Glick PL, Caty MG, Azizkhan RG. Follow up imaging studies in children with splenic injuries. *Clin Pediatr (Phil)* 1999; **38**: 273–5.
13. Davies JM, Barnes R, Milligan D. Update of guidelines for the prevention and treatment of infection in patients with an absent or dysfunctional spleen. *J R Coll Phys* 2002; **2**: 440–3.
14. Mourtzoukou EG, Pappas G, Peppas G, Falagas ME. Vaccination of asplenic or hyposplenic adults. *Br J Surg* 2008; **95**: 273–80.

AUTHORS' RESPONSE

doi 10.1308/003588408X321846

S SINHA, SVV RAJA, MH LEWIS

Department of Surgery, Royal Glamorgan Hospital, Ynysmaerdy, Llantrisant, UK

CORRESPONDENCE TO

Surajit Sinha, 6 Ducane Walk, Plymouth PL6 5WE, UK
E: sinhasurajit@hotmail.com

We thank Harwood and Maw for their interest in our paper and would sympathise with many of their comments and criticisms. In reply to their specific questions.

We wonder what criteria were used to allow the patients to be safely discharged from hospital and why there was such a wide range of hospital stay?

Answer: Our study was retrospective reporting typical in-patient management of splenic injuries. With regard to hospital stay, this is notoriously variable and not confined always to the patient's surgical well-being. Often, it includes home circumstances and home care. Specifically, our patients were managed by experienced general surgical consultants and the decision as to discharge was at the discretion of individual clinicians involved in the cases. Obviously, there was a variation including the grade of splenic trauma and associated injuries. All patients discharged were haemodynamically stable with stable haemoglobin levels and clinical examinations.

With regard to general advice given to the patients following discharge from hospital and specifically their resumption of driving, normal daily activities and contact sports, etc.?

Answer: Again, such advice was at the discretion of experienced consultant general surgeons. All patients appeared to be advised against heavy physical activity and, specifically, contact sports for 3 months. With regard to bed rest, this varied from consultant to consultant and certainly the senior author of this article was keen that patients were not bed-bound for weeks on end with associated risks of deep venous thrombosis and pulmonary embolism.

Were there any cases of delayed rupture of patients from the Royal Glamorgan Hospital?

Answer: Out of 11 non-operative patients in our study, one became unstable with increasing abdominal pain, tachycardia, hypotension and dropped haemoglobin on the third day of a second admission, 2 or 3 weeks following the initial trauma. She underwent urgent laparotomy and splenectomy. At laparotomy there was identified a large posterior splenic tear with haemoperitoneum. Her postoperative recovery was then uneventful.

We wonder what criteria were used for determining which imaging modality to use and the timing and number of follow-up images in their patients?

Answer: Within the Royal Glamorgan Hospital there is no set policy with regard to post splenic injury from the radiological department. The type of imaging, timing and number of follow-up scans were decided by the treating consultant and this varied. We attempted, in our paper, to highlight the large doses of radiation exposure in several patients having repeat scans and the senior author was keen to use ultrasound in preference to CT.

Sinha and colleagues did not comment on the outcome of the follow-up radiological investigations in their reported series or if there were any cases of late rupture in the non operative group. Thus the role of follow up imaging remains controversial.

Answer: This question has been partially answered above. However, we agree routine follow-up imaging in the stable patient without clinical suspicion of bleeding is debatable. Of the 10 patients managed non-operatively that did not require splenectomy, 6 were followed up with serial ultrasound scans and 4 with repeat CT scans. Four patients who had ultrasound scans and 2 patients who had CT showed resolving haematoma and localised fluid around the spleen which subsequently resolved. None of our patients developed pseudoaneurysms or progressive haematoma.

COMMENT ON

doi 10.1308/003588408X242105

JW Moor, P Murray, J Inwood, D Gouldesbrough, C Bem. Diagnostic biopsy of lymph nodes of the neck, axilla and groin: rhyme, reason or chance. *Ann R Coll Surg Engl* 2008; **90**: 221–5

doi 10.1308/003588408X321837

Reducing the number of open node biopsies carried out for benign disease

PETER A BRENNAN, DUNCAN BAYNE, ELIZABETH TILLEY
Head and Neck Unit, Queen Alexandra Hospital, Portsmouth, UK

CORRESPONDENCE TO

Peter A Brennan, Consultant Maxillofacial Surgeon and Professor of Surgery, Head and Neck Unit, Queen Alexandra Hospital, Portsmouth PO6 3LY, UK. E: Peter.brennan@porthosp.nhs.uk

We read with interest this study on diagnostic lymph node biopsy, which found that 45% of cervical nodes had histologically benign disease following open biopsy. It was suggested that the level of care may be representative within many hospitals in the UK. With the increasing use of high-frequency ultrasound, combined with fine needle aspiration cytology where appropriate, the number of diagnostic open biopsies for benign lymphadenopathy should actually be very small. Nodes with worrying features on ultrasound¹ should have an ultrasound-guided fine needle aspiration or, if appropriate (e.g. suspicious for lymphoma), an open biopsy.

A review of over 300 referrals to our head and neck fast-track clinic with cervical lymphadenopathy (run jointly between maxillofacial and ENT departments) found that the number of patients who have had node biopsy, which was subsequently found to be reactive on definitive histology, was less than 8%. Almost all patients have ultrasound, some in a joint ultrasound clinic with both surgeon and radiologist present at the time of imaging. This has significantly reduced the number of open node biopsies carried out for benign disease, confirming that the results reported by Moor and colleagues might not be representative of units in the UK after all.

Reference

1. Ahuja AT, Ying M. Sonographic evaluation of cervical lymph nodes. *AJR Am J Roentgenol* 2005; **184**: 1691–9.

AUTHORS' RESPONSE

doi 10.1308/003588408X321846

JAMES W MOOR, CHRIS BEM
Department of Otorhinolaryngology Head and Neck Surgery,
Bradford Royal Infirmary, Bradford, UK

CORRESPONDENCE TO

James Moor, Department of Otorhinolaryngology Head and Neck Surgery, Bradford Royal Infirmary, Duckworth Lane, Bradford BD9 6RJ, UK. E: jim@docmoor.freeserve.co.uk

We thank Brennan and colleagues for their comment in supporting our conclusion that dedicated neck lump clinics will achieve a higher yield of diagnostic neck node biopsies. Our paper referred to a time when there was no dedicated neck lump clinic and when there was limited use of fine needle aspiration cytology and ultrasound scanning. It was written to illustrate the problems that arise when there are no defined pathways of care and, as we concluded, may still represent a level of care in the absence of dedicated neck lump clinics.

COMMENT ON

doi 10.1308/003588408X261816

JM Wilson, R Baines, ED Babu, CJ Kelley. A role for topical insulin in the management of problematic surgical wounds *Ann R Coll Surg Engl* 2008; **90**: 160

doi 10.1308/003588408X321846

The search for 'wound solutions'

D BAYNE
Department of Plastic Surgery, The Mountbatten Plastic Surgery Unit, Royal Hospital Haslar, Gosport, Hampshire, UK

CORRESPONDENCE TO

D Bayne, Department of Plastic Surgery, The Mountbatten Plastic Surgery Unit, Royal Hospital Haslar, Gosport, Hampshire PO12 2AA, UK. E: paperscorrespondence@inbox.com

As plastic surgeons, we are frequently referred problem wounds and sympathise with the authors' difficulties. We have had experience with the use of topically applied insulin yet, for this case, we are sceptical if it had any wound healing effect. In the after therapy photograph, the