Letters and comments

Contributors to this section are asked to make their comments brief and to the point. Letters should comply with the instructions to authors at <http://www.rcseng.ac.uk/publications/annals/authorinstructions.html>. Tables and figures should only be included if absolutely essential and no more than five references should be given. The editor reserves the right to shorten letters and subedit contributions to ensure clarity.

COMMENT ON

doi 10.1308/003588408X242042

PG Sorelli, NS El-Masry, PM Dawson, NA Theodorou. The dedicated emergency surgeon: towards consultant-based acute surgical admissions. *Ann R Coll Surg Engl* 2008; **90**: 104–8

doi 10.1308/003588408X321576

Costs of additional cover

JG MOSLEY

Department of General Surgery, Leigh Infirmary, Leigh, UK

CORRESPONDENCE TO

JG Mosly, Consultant General and Vascular Surgeon, Leigh Infirmary, The Avenue, Leigh WN7 1HS, UK
E: john.mosly@wwl.nhs.uk

Sorelli and colleagues have shown that if one employs an extra general surgeon then more of the emergency admissions will be operated upon more quickly and their length of stay in hospital will be reduced, thereby improving their quality of care. However, the savings of £90,000 per annum must be set against the cost of the additional surgeon. The additional surgeon is being employed from 9 am until 5 pm, five days a week. If it is on ten programmed activities (PAs), he or she will only be available for emergency care three and a half days a week, so to cover the rest of the time, vacation and study leave, one imagines that Imperial College will actually have to employ two consultants or alternatively pay the additional consultant 12.5 PAs per week to work 9 am until 5 pm, Monday to Friday. The consultant's salary will, therefore, be considerably greater than the £90,000

saved by reduced in-patient stay. There is absolutely nothing wrong in employing extra staff to improve patients' quality of care; however, this would generally be associated with increased expenditure, as in this case.

COMMENT ON

doi 10.1308/003588408X242033

S Sinha, SVV Raja, MH Lewis. Recent changes in the management of blunt splenic trauma: effect on splenic trauma patients and hospital implications. *Ann R Coll Surg Engl* 2008; **90**: 109–12

doi 10.1308/003588408X321602

Blunt splenic trauma

TC KÖNIG, NRM TAI, MS WALSH

Trauma Service, Royal London Hospital, London, UK

CORRESPONDENCE TO

TC König, Specialist Registrar in General Surgery, Royal London Hospital, Whitechapel, London E1 1BB, UK
E: tomkonig@hotmail.com

Since January 2004, 105 patients (2.5% of total trauma admissions) were seen having suffered splenic injury; of these, 85 (81%) suffered blunt injury. Mortality was 19% patients (20 of 105). The overall median Injury Severity Score (ISS) was 35. Fifty-five patients had a laparotomy (52%). The total number of splenectomies performed was 34 (a rate of 32%) and the median ISS in this group was 40. Of the patients who survived, 24 had a splenectomy and 16 had a laparotomy and splenic preservation surgery (packing, spleen seen to have stopped bleeding and requiring no intervention). Ten of the patients who died had a splenectomy and 5 had a laparotomy and the spleen was preserved. The median ISS in the group treated conservatively was 27. Only 4 patients had an isolated splenic injury. Five patients underwent splenic artery embolisation with good effect.

The high ISS in those undergoing splenectomy re-emphasises the surgical dilemma of whether or not to attempt preservation of the damaged spleen in light of injuries to other body regions.

The paper highlights the problem with splenic injury grading from CT scanning and we feel that further efforts should