Published in final edited form as:

Pharmacol Biochem Behav. 2009 September ; 93(3): 291–299. doi:10.1016/j.pbb.2009.02.008.

Now or Later? An fMRI study of the effects of endogenous opioid blockade on a decision-making network

Charlotte A. Boettiger¹, Elizabeth A. Kelley², Jennifer M. Mitchell^{2,3}, Mark D'Esposito⁴, and Howard L. Fields^{2,3}

¹ Department of Psychology, Biomedical Research Imaging Center, and Curriculum in Neurobiology, University of North Carolina, Chapel Hill

² Ernest Gallo Clinic and Research Center, Emeryville, CA

³ Department of Neurology, University of California, San Francisco

⁴ Department of Psychology, Helen Wills Neuroscience Institute, and Henry J. Wheeler Brain Imaging Center, University of California, Berkeley

Abstract

Previously, we found that distinct brain areas predict individual selection bias in decisions between small immediate (“*Now*”) and larger delayed rewards (“*Later*”). Furthermore, such selection bias can be manipulated by endogenous opioid blockade. To test whether blocking endogenous opioids with Naltrexone (NTX) alters brain activity during decision-making in areas predicting individual bias, we compared fMRI BOLD signal correlated with *Now* versus *Later* decision-making after acute administration of NTX (50 mg) or placebo. We tested abstinent alcoholics and control subjects in a double-blind two-session design. We defined regions of interest (ROI) centered on activation peaks predicting *Now* versus *Later* selection bias. NTX administration significantly increased BOLD signal during decision-making in the right lateral orbital gyrus ROI, an area where enhanced activity during decision-making predicts *Later* bias. Exploratory analyses identified additional loci where BOLD signal during decision-making was enhanced (left orbitofrontal cortex, left inferior temporal gyrus, and cerebellum) or reduced (right superior temporal pole) by NTX. Additional analyses identified sites, including the right lateral orbital gyrus, in which NTX effects on BOLD signal predicted NTX effects on selection bias. These data agree with opioid receptor expression in human frontal and temporal cortices, and suggest possible mechanisms of NTX’s therapeutic effects.

Keywords

alcoholism; executive function; human; naltrexone; orbitofrontal

1. Introduction

The non-selective opioid antagonist naltrexone (NTX) is one of few drugs with U.S. Food and Drug Administration (FDA) approval for the treatment of alcoholism. NTX is also used to treat

CORRESPONDING AUTHOR: Charlotte A. Boettiger, Ph.D., Psychology Department, Davie Hall, CB#3270, University of North Carolina, Chapel Hill, NC 27599-3270, cab@unc.edu, ph: 919-962-2119, fax: 919-962-2537.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

other substance addictions and impulse-control disorders. NTX reduces ethanol consumption in subjects with a history of alcohol abuse (Anton et al., 1999; Davidson et al., 1999; Heidbreder, 2005; Hernandez-Avila et al., 2006; O'Brien et al., 1996), and in animals trained to self-administer ethanol (Boyle et al., 1998; Stromberg et al., 1998). Moreover, NTX reduces drug-seeking behavior triggered by drug-cue exposure in animal models (Ciccocioppo et al., 2003; Ciccocioppo et al., 2002; Liu and Weiss, 2002) and alcohol-cue induced craving in human alcoholics (Monti et al., 1999; O'Malley et al., 2002; Rohsenow et al., 2000). However, the mechanisms of these clinically important actions of NTX are not well understood. NTX has highest affinity for μ -opioid receptors, but substantial affinity for κ - and δ -opioid receptors (Kreek, 1996). Through its modulation of the endogenous opioids that control midbrain dopaminergic neurons, NTX alters forebrain release of dopamine (DA) (Herz, 1995; Margolis et al., 2006; Spanagel et al., 1992). Through blocking opioid disinhibition of ventral tegmental area (VTA) neurons, NTX reduces ethanol-induced release of DA in the nucleus accumbens (Altshuler et al., 1980; Harris and Erickson, 1979), which contributes to the motivation to consume alcohol. Based on this neurobiological action of NTX, candidate mechanisms for its therapeutic action in alcoholics include an attenuation of the rewarding effects of alcohol, for which there is experimental support (Sinclair, 2001; Swift et al., 1994; Volpicelli et al., 1995). There is also evidence that NTX can increase the aversive effects of ethanol (Davidson et al., 1999; de Wit et al., 1999; McCaul et al., 2000; Mitchell et al., 2005a). However, NTX can also significantly reduce alcohol craving in alcoholics during abstinence (Monti et al., 1999; O'Malley et al., 2002; Rohsenow et al., 2000), and attenuate thoughts and behaviors associated with drinking (Anton et al., 1999). Thus, NTX may act, in part, via alteration of specific cognitive processes. One possibility that has received experimental support is that NTX reduces the tendency to choose impulsively (Kieres et al., 2004; Mitchell et al., 2007; O'Malley et al., 2002). In fact, NTX relieves a wide variety of impulse control disorders (Grant, 2005; Kim et al., 2001; Marrazzi et al., 1995; Raymond et al., 2002; Symons et al., 2004). These data all suggest that endogenous opioids promote impulsivity, and that blocking endogenous opioid activity with NTX reduces impulsiveness.

To address whether NTX alters brain activity in regions known to predict individual bias toward immediate rewards, we used a NTX challenge in the context of functional magnetic resonance imaging (fMRI) while human subjects performed a modified delay discounting (DD) task. In a previous fMRI study of this task (Boettiger et al., 2007), we found that distinct brain areas predict individual selection bias. These areas include the dorsal prefrontal cortex, the posterior parietal cortex, the orbitofrontal cortex (OFC), the parahippocampal gyrus adjacent to the amygdala, the posterior cerebellum, and the inferior and middle temporal gyri. The task is comprised of numerous trials in which subjects are instructed to choose between two amounts of money, a smaller amount available "*Now*" (e.g. "\$80 TODAY") or a larger amount available "*Later*" (e.g. "\$100 in 1 Month"; Figure 1). We have quantified individual selection bias in this task as an "Impulsive Choice Ratio" (ICR) (Mitchell et al., 2005b). In a previous pharmacology study, we found that such selection bias can be manipulated by endogenous opioid blockade (Mitchell et al., 2007). To test the hypothesis that blocking endogenous opioid signaling with NTX would alter regional brain activity that predicts individual selection bias during decision-making, we compared the fMRI BOLD signal correlated with *Now* versus *Later* decision-making after acute administration of NTX (50 mg) or placebo. Subjects were either abstinent alcoholics or control subjects with no history of substance abuse, and they were tested in a double-blind randomized two-session crossover design.

For the purpose of analyzing brain activity between sessions, we defined regions of interest (ROI) as 10mm spheres centered on activation peaks that predict *Now* vs. *Later* selection bias (Boettiger et al., 2007), comparing the BOLD signal during decision-making within these ROI between NTX and placebo sessions. We also conducted exploratory mapwise analyses to identify other brain areas in which BOLD signal during decision-making was enhanced or

reduced by NTX administration. Finally, given the widespread evidence for variability in NTX response, we conducted additional analyses to identify sites in which NTX effects on BOLD signal predicted NTX effects on selection bias.

NTX is not effective in treating all alcoholics, thus, determining the moderators of NTX's effects is a critical goal of current research. Investigation of genetic moderators has of NTX's effects have largely focused on the A118G (ASP40) variant of the μ -opioid receptor gene (OPRM1). This OPRM1 variant produces a μ -opioid receptor with greater affinity for β -endorphin (Bond et al., 1998), suggesting a gain-of-function effect. However, more recent data indicate that the ASP40 variant impairs transcription and thus expression of the receptor protein, resulting in a loss-of-function effect (Zhang et al., 2005). Laboratory studies demonstrate that this OPRM1 variant determines NTX's effect on physiological responses to alcohol cue exposure (McGeary et al., 2006) and to alcohol-induced high (Ray and Hutchison, 2007). Moreover, clinical trials have yielded some data demonstrating that the ASP40 variant predicts therapeutic response to NTX (Anton et al., 2008; Oslin et al., 2003); although, other studies have failed to replicate this finding (Gelernter et al., 2007). Thus, we also collected genetic material from these subjects to test whether the A118G variant of the OPRM1 gene also modulates NTX effects on immediate reward bias or underlying neural circuit activity. In addition to the A118G variant of the OPRM1 gene, we also tested several other single nucleotide polymorphisms (SNPs) that have been associated in other studies with addiction and/or impulsivity.

2. Methods

2.1. Participants

Participants were either abstinent alcoholics (AA; $n = 9$) or control subjects with no history of substance abuse (CS; $n = 10$). All participants were healthy, right-handed volunteers (8 females; mean age: 28.3 ± 5.8), and were paid for their participation. Participants were screened for psychoactive drug use (Biotechnostix, Inc.), including alcohol (Lifeloc Technologies, Inc.) at the start of each session. Five additional participants were tested, but were excluded from fMRI analyses due to excess head motion or equipment failure. All subjects had at least a high school education, and the two groups did not differ in terms of socio-economic status, age, IQ, and years of education. Alcohol addiction severity (prior to sobriety for AA group) was assessed via the Alcohol Use Disorders Identification Test (AUDIT; (Saunders et al., 1993)), with mean AUDIT scores of 19 ± 7 and 5 ± 2 , for the AA and CS groups, respectively (scores ≥ 8 indicate problem drinking). AA subjects self-reported a minimum of 2 weeks sobriety at the time of recruitment (mean sobriety = 2.5 yrs). All participants gave written, informed consent, in accordance with the guidelines of the University of California, Berkeley Committee for the Protection of Human Subjects. Subject payment was not dependent upon choice behavior in the task.

2.2. Experimental paradigm

Each subject completed two sessions separated by a minimum of 48 h. At the start of each session, subjects gave written, informed consent and after completing screening for contraindications for NTX, subjects were administered either a 50mg NTX capsule or an identical placebo capsule. Experimenter and subject were blind to capsule content; capsule order was counter-balanced across subjects. Three hours after pill ingestion, subjects performed a delay discounting task in the context of an fMRI scanning session. For the task, subjects made a series of choices between *Now* and *Later* hypothetical monetary rewards. Later amounts were \$2, \$5, \$10, \$20, or \$100, at 1 of 5 future delays: 1 WEEK, 2 WEEKS, 1 MONTH, 3 MONTHS, or 6 MONTHS. The *Now* option was always a lesser amount available TODAY. *Now* and *Later* options randomly appeared on the right or left side. Subjects were given task instructions

and a brief practice session to familiarize them with the delay discounting (DD) task. Subjects performed the DD task within an fMRI scanner, while we collected continuous whole-brain bold oxygen level dependent (BOLD) signal. Scanning comprised 8 functional scans of 37 trials each (32 choice trials, and 5 “null” trials); trial type order was pseudorandom and unique for each subject (total scanning duration ~1 hour). For each trial, subjects were shown a cue that indicated how to select between the two subsequently displayed hypothetical reward options, each option was a dollar amount available at a point in time (Figure 1); these two alternatives are considered as “*Now*” and “*Later*”. The *Now* option was available “TODAY” and was the lesser of the two amounts by a percentage randomly selected from the set: {30%, 15%, 10%, 5%}. Right-left position of each option was randomized across trials. Subjects indicated their choice by pressing a button on an MRI-compatible keypad. Instruction cues included: WANT (W), DON’T WANT (DW), SOONER, and LARGER, the latter two cues are considered together as “CONTROL” (CON). In the W condition, subjects chose their preferred option. In the DW condition, subjects were asked to make the same evaluation and press the button corresponding to the opposite choice. On CON trials, subjects selected the option reflecting either the sooner time point or larger amount of money, depending on the instruction cue. CON trial performance verified subject comprehension and compliance with task instructions. The DW condition provides a gross assessment of motor impulsiveness, by comparing inferred choices in the DW trials as a function of time with choices in the W condition as a function of time. This comparison rules out spurious impulsive choice as a result of unintended motor responses. Trial types frequencies were weighted with ratios of 1/2 for the W condition and 1/6 each for each of the other conditions. The order of inter-trial intervals was fixed across subjects. As the choice condition followed the cue condition at a fixed delay for each trial, “null” trials were included in each run (5 trials per run); in these trials, the instruction cue appeared and no options followed. Subjects saw each of the 120 possible choices two or three times, although not necessarily in the same trial type context. All choices made in the task were based on hypothetical rewards, although subjects were instructed to choose as if they would actually receive their choices.

2.3. Analysis of behavioral data

Single factor between-group comparisons were made via unpaired t-tests. Within group comparisons of condition were made via paired t-tests. All t-tests were 2-tailed and were conducted using Excel or SPSS. Multi-factorial comparisons were made using mixed repeated-measures ANOVA in SPSS, with group as a between subjects factor. Where sphericity assumptions were violated, a Greenhouse-Geisser non-sphericity correction was applied. For each subject, we calculated an impulsive choice ratio (ICR), which is the proportion of *Now* choices relative to all W condition choices made. Reward preference in DW trials was inferred to be the unselected option. From these DW responses, we calculated an inferred ICR (*i*ICR) for each delay time and took the sum of $|ICR - iICR|$ across all delays as a gross index of motor error. To ensure the validity of parametric statistical tests, we used an arcsine-root transformation of ICR data prior to statistical comparisons.

2.4. Imaging

T2*-weighted images (EPI) were acquired on a 4T Varian Inova whole body magnetic resonance scanner (Palo Alto, CA) equipped with a TEM send-receive radio frequency (RF) head coil, using a 1-shot gradient echo EPI sequence (TR = 2200 ms, TE = 28 ms, flip angle = 20°) to detect blood oxygen level dependent (BOLD) contrast. The sequence employed phase map correction to reduce Nyquist ghosts. To facilitate coverage of the ventral/orbital PFC, sampling of k-space was in the order positive to negative (De Panfilis and Schwarzbauer, 2005). 40 coronal slices (3.5mm thick w/0.5mm gap) per volume were obtained (FOV = 22.4×22.4, in-plane resolution = 3.5mm). Each fMRI acquisition was preceded by 22s of dummy gradient RF pulses to achieve steady-state tissue magnetization and minimize startle-

induced motion. Low-resolution T1-weighted co-planar images were acquired for each participant. In addition, a magnetization-prepared fast low-angle shot high-resolution (MPFLASH) T1-weighted image was acquired for the purposes of normalizing data to standardized space. E-Prime software (PST, Inc., Pittsburgh, PA) synchronized the stimulus display with the fMRI acquisition and collected subject responses via an MRI-compatible keypad. An LCD projector (Epson, Long Beach, CA) projected stimuli onto a rear projection screen (Stewart, Torrance, CA), which the subjects viewed via a mirror mounted within the head coil.

2.5. Image Analysis

fMRI data were processed offline with a combination of in-house software, SPM2 (Wellcome Dept. of Imaging Neuroscience, London, UK), and the Artifact Detection Tools package (<http://web.mit.edu/swg/software.htm>). Images were first reconstructed into Cartesian space, and sinc interpolated in time to correct for timing differences in slice acquisition. Data were next motion corrected using a six-parameter, rigid-body, least-squares algorithm (Friston et al., 1995), realigning all volumes to the first EPI volume, then smoothed with a 7 mm full-width half-maximum (FWHM) Gaussian smoothing kernel. Each subject's coplanar and MPFLASH anatomical images were co-registered to the first EPI volume, and then normalized to Montreal Neurological Institute (MNI) space and resampled to 2 mm³ isotropic voxels. The functional data were analyzed in an event-related manner based on the modified general linear model as implemented in the statistical package SPM2. Each subject's model included a design matrix of regressors of interest, which were δ functions positioned at the onset time of each experimental event type (ready; instruction cue; choice pair) convolved with a canonical hemodynamic response function (HRF). Noise spikes were rarely detected in the data (ArtRepair2 Toolbox); these were modeled as covariates of no interest with duration of 2.2 s and were not convolved with an HRF. Data were high-pass filtered using a cut-off of 128 s, and serial autocorrelations were corrected with a restricted maximum likelihood algorithm using a second-order autoregressive model (AR2 + white noise).

For each subject, linear contrast images were calculated from the regressor parameter estimates; contrast images were then normalized into MNI space using the parameters derived from the co-registered MPFLASH. Contrast maps were smoothed resulting in a total smoothing kernel of 10mm FWHM. To test whether Naltrexone (NTX) would alter brain activity during subjective decision-making, we compared the mean contrast values of (WANT – CON) from the NTX and placebo (PBO) sessions within predefined regions of interest (ROI) from both NTX and PBO sessions. ROI were 10mm³ spheres centered on coordinates identified in a previous study (Boettiger et al., 2007). We also subtracted NTX and PBO session contrasts to assess mapwise drug effects; the subtracted contrasts were then subject to both simple regression and one-sample t-test analyses, as implemented in the SnPM3 analysis package (<http://www.sph.umich.edu/ni-stat/SnPM/>). To control for multiple comparisons, statistical non-parametric maps are corrected for Type I error ($\alpha < 0.001$; whole brain volume), determined by random permutation of the NTX-PBO subjective decision-making contrast parameter estimates (with either ICR change or subject) in 5000 new analyses (Nichols and Hayasaka, 2003; Nichols and Holmes, 2002). Activation peaks were localized within defined anatomical regions, identified from the T1-weighted anatomical images obtained from each subject, with reference to standard brain atlases (Duvernoy et al., 1991; Schmahmann et al., 1999; Tzourio-Mazoyer et al., 2002).

2.6. Genetic Analysis

DNA extraction and analysis was conducted according to standard methods on samples obtained from subjects providing informed consent. The Ernest Gallo Clinic and Research Center Genomics Core carried out genotyping of the following polymorphisms with

polymerase chain reaction using TaqMan® technology (Applied Biosystems): adrenergic alpha-2A-receptor (ADRA2A; rs1800544), ankyrin repeat and kinase domain containing 1 (ANKK1/DRD2 Taq I; rs1800497), brain-derived neurotrophic factor (BDNF; rs6265), catechol-*O*-methyltransferase (COMT; rs4680), dopamine D2 receptor (DRD2; rs6277), dopamine D4 receptor (DRD4; rs1800955), kappa-opioid receptor (OPRK1; rs1051660), mu-opioid receptor (OPRM1; rs1799971), and tryptophan hydroxylase 1 (TPH1; rs1800532).

To estimate which genotypes had the greatest predictive value for the NTX effects on ICR and ROI activity, linear multiple regression analyses were carried out using SPSS. For each multiple regression analysis, we entered genotypes stepwise, divided into five blocks. The blocks were as follows: block 1 – OPRM1; block 2 – OPRK1; block 3 – COMT, DRD2 TaqI, DRD2_7a, DRD4; block 4 – ADRA2, TPH1; block 5 – BDNF.

3. Results

3.1. Sample Characteristics

The AA and CS groups were matched in terms of demographic variables. Specifically, the groups did not differ in terms of education level, age, SES, IQ, gender composition, or ethnic composition (Table 1). However, as expected, the AA subjects reported significantly higher levels of alcohol abuse and alcohol-related problems (Table 1). A family history of alcohol abuse was also more common among the AA group (Table 1).

3.2. NTX effects on behavior

Replicating our previous finding (Mitchell et al., 2007), a mixed design repeated measures ANOVA found no significant main effect of drug ($F_{1,17}=1.07$, $p=0.341$) or group ($F_{1,17}=0.01$, $p=0.941$) on RTs, although, as expected, we did observe a significant effect of trial type ($F_{1,24}=49.1$, $p<0.001$) (Table 2). This latter effect reflected increasing RTs from the CON to WANT to DW trial types. We also found no significant drug $\times$ group interaction effect on RT ($F_{1,17}=2.14$, $p=0.161$).

A mixed design repeated measures ANOVA considering Impulsive Choice Ratio (ICR) as the dependent variable found a significant main effect of group ($F_{1,17}=10.11$, $p<0.001$), but not of NTX ($F_{1,17}=1.22$, $p=0.286$) on ICR. Replicating our previous results (Boettiger et al., 2007; Mitchell et al., 2005b; Mitchell et al., 2007), the group effect was due to significantly higher ICRs among the AA group relative to the CS group. We also found no significant drug $\times$ group interaction effect on ICR ($F_{1,17}=0.48$, $p=0.496$).

3.3. NTX effects on brain activity during subjective preference decisions

In a previous fMRI study, we found that during decisions between small, immediate (“Now”) versus larger, delayed (“Later”) rewards the magnitude of the fMRI BOLD signal scales with individual immediate reward bias in several brain areas (Boettiger et al., 2007). Here we first tested whether acute NTX administration alters brain activity during subjective decision-making in these areas. We then conducted an exploratory analysis to identify other areas in the brain in which activity during decision-making is significantly altered. Finally, across the whole brain, we tested for correlations between NTX’s effect on brain activity during *Now* versus *Later* decision-making and its effect on choice behavior.

3.3.1. Effects within choice-predicting ROI—For the Region of interest (ROI) analyses, we defined the ROIs as 10 mm³ spheres centered on the coordinates identified in our previous study (Boettiger et al., 2007). These areas each predict individual immediate reward selection bias during *Now* versus *Later* decision-making. These areas can be divided according to those in which activity during such decisions is directly correlated with *Now* bias (“*Now* areas”) and

one in which activity during such decisions is inversely correlated with *Now* bias (“*Later*” area). For each ROI, the mean parameter estimates (β values) for a subjective decision-making contrast (WANT-CON) were extracted from both the NTX and the placebo sessions and then subject to further statistical analyses.

3.3.1.1. “Now” Areas: To test for significant effects of NTX on BOLD activity during decision-making in *Now* ROIs, we conducted paired *t*-tests with ROI β values as dependent variables. For the three ROIs that were the focus of our previous study (posterior parietal cortex, superior frontal gyrus, and parahippocampal gyrus), we did not observe a significant effect of NTX on brain activity subjective decision-making (t_{18} , minimum p value = 0.548). We observed similar effects in additional areas in the temporal lobe and cerebellum (see Table 3), in which activity during *Now* versus *Later* decisions is also known to correlate with *Now* bias (t_{18} , minimum p value = 0.489). These results indicate that any effect that NTX may have on temporal discounting behavior are not mediated by actions in any of these brain areas.

3.3.1.2. “Later” Area: We next tested for a significant effect of NTX on BOLD activity during decision-making in the *Later* ROI in the OFC, again using a paired *t*-test with ROI β values as dependent variables. Here we observed that during NTX sessions, brain activity during subjective decision-making was significantly elevated (Figure 2; t_{18} , $p = 0.045$). Together with our previous finding that activity in the OFC during *Now* versus *Later* decisions inversely correlates with *Now* bias, this result suggests that NTX may reduce impulsive decision-making via effects on the OFC.

3.3.1.3. Group interaction with NTX effects on ROI: Two-way factorial ANOVA with ROI β values as dependent variables failed to identify a significant group β drug interaction for any of the *Now* or *Later* ROIs (maximum $F_{1,17} = 1.44$, $p = 0.256$). These findings suggest that NTX does not exert differential brain effects based on alcohol use history in areas that predict *Now* versus *Later* bias.

3.3.1.4. Genotype interaction with NTX effects: **3.3.1.4.1. Effects on choice behavior:** Multiple linear regression analysis was used to determine whether any of the SNPs we assayed had significant predictive value in terms of the NTX effect on ICR. We found that a single SNP had significant predictive power: the Val66Met locus of BDNF ($r=0.52$, $t_{15}=-2.27$, $p=0.04$). That is, electing *Now* more often during the NTX session was associated with fewer Val alleles at position 66 in the BDNF protein. No other genotype factor added into the model added significantly more predictive power than BDNF genotype alone for the NTX effect on ICR. A two-way factorial ANOVA with ICR as dependent variable confirmed a significant interaction between drug condition and BDNF genotype ($F_{1,13} = 3.99$, $p = 0.045$; Figure 3). There was no significant main effect of BDNF genotype on choice behavior ($p > 0.8$).

3.3.1.4.2. Effects on brain activity during subjective choice: We performed an identical multiple linear regression analysis using the OFC ROI β values as the dependent variable to test whether any of the tested genotypes held significant predictive value for NTX’s effect. This procedure found that none of the SNPs tested significantly predicted the NTX effect on OFC activity during *Now* versus *Later* choices.

3.3.2. Mapwise analysis of NTX effects

3.3.2.1. Sites in which NTX raised or lowered activity: In addition to our ROI-based analyses, we conducted an exploratory mapwise analysis of NTX’s effect on BOLD signal during subjective decision-making. We found that NTX elevated activity during *Now* versus *Later* decisions in the left OFC, the left inferotemporal cortex, and bilateral sites in the cerebellum (Figure 4A). In addition, we found that NTX depressed activity during *Now* versus

Later decision-making in the right superior temporal gyrus (Figure 4B). See Table 4 for peak coordinates.

3.3.2.2. Correlation between NTX effects on brain and on behavior: Our final analysis employed simple regression analysis to identify brain areas in which NTX's effects on decision-making activity predicted NTX's effect on behavior. To accomplish this, we entered the choice condition drug contrasts (NTX – PBO) from each subject into a map-wise simple regression analysis. This correlation analysis exploited the intersubject variability in NTX behavioral effects to identify differences in NTX effects on the BOLD signal during *Now* versus *Later* choices. Using the ICR change of each subject as a between subjects statistical regressor, we found a significant positive correlation between ICR change and session difference in activity between during preference-based decision-making in a constellation of areas (Table 5). Notably, one such site was in the right lateral OFC (Figure 5A, Table 5), falling within the OFC ROI investigated above. As can be seen in the regression plot of Figure 5B, the NTX effect on OFC activity during *Now* versus *Later* decisions was inversely correlated with the NTX effect on ICR. In other words, elevated OFC activity on NTX was associated with an increased the likelihood of selecting the larger, delayed reward (Figure 5B).

6. Discussion

6.1. NTX modulation of OFC function

Determining the functional effects of NTX on the human brain is an important step in understanding the bases for variability in its efficacy. This information may not only help predict whether an individual will respond to NTX, it may also help identify which of NTX's actions are critical to its therapeutic effect. This latter information may prove especially valuable in driving the development of new treatments for alcoholism and related disorders, as it could suggest which brain regions are critical therapeutic targets and thus help guide or validate animal experiments. To date, this study is only the second published report about the effects of NTX in the brains of people with a history of alcohol dependence. Here we found that among areas that predict individual bias towards immediate rewards, a single area was modulated by NTX during decision-making: the OFC. Interestingly, the other recent study found that NTX modulates OFC responses to alcohol cues in alcohol dependent subjects (Myrick et al., 2008). Together these findings highlight the OFC as a key brain area affected by NTX.

The OFC is thought to play a critical role in moderating impulsive choice (Mobini et al., 2002; Rudebeck et al., 2006), and in representing subjective value during choice tasks (Izquierdo et al., 2004; Padoa-Schioppa and Assad, 2006; Roesch and Olson, 2004; Schoenbaum and Roesch, 2005). OFC damage impairs the ability of animals to change their responses to stimuli that are no longer reinforced (Dias et al., 1996; McAlonan and Brown, 2003; Mishkin, 1964; Ostlund and Balleine, 2007; Schoenbaum and Roesch, 2005; Schoenbaum et al., 2007; Tait and Brown, 2007). Likewise, patients with damage to the OFC, due either to focal lesions (Rolls et al., 1994) or to degenerative frontotemporal dementia (Rahman et al., 1999), also show impairment in changing their responses when the "rules" for correct responding change. Human neuroimaging studies also support a role for the OFC in overcoming learned response contingencies (Boettiger CA, 2004; Cools et al., 2002; Elliott et al., 2000; O'Doherty et al., 2004; Remijnse et al., 2006).

Such inability to refrain from previously preferred actions in response to formerly rewarding cues that are no longer reinforced is highly reminiscent of the transition from flexible to habitual responding that is characteristic of drug addiction. It should come as no surprise then, that neuroimaging studies have repeatedly found abnormal OFC functioning in substance abusers (Boettiger et al., 2007; Dom et al., 2005; Ersche et al., 2005; London et al., 2000; Volkow and

Fowler, 2000). The present data suggest that NTX may support the long-term decision-making critical to recovery from alcoholism by increasing activity in the OFC. Determining whether this biomarker is an addiction vulnerability or relapse risk factor is an important future question.

6.2. Immediate reward bias and endogenous opioids

These data are consistent with our previous finding that blockade of endogenous opioids with NTX can alter immediate reward bias and that this effect is highly variable across individuals (Mitchell et al., 2007). The present results provide new data regarding possible neurobiological sources for this variability. First, we found that a common polymorphism in the BDNF gene predicted individual NTX effect on immediate reward bias. Second, we identified brain regions in which the difference in activity during *Now* versus *Later* decisions between NTX and placebo sessions predicted changes in choice behavior. These areas included the right lateral OFC, an area in which high activity during *Now* versus *Later* decisions predicts *Later* reward selection bias. Consistent with this finding, here we show an association of NTX elevated OFC activity with reduced selection of immediate rewards. The two primary targets of NTX, μ -opioid receptors and kappa-opioid receptors, are both expressed at high levels in the OFC (Gorelick et al., 2005; Staley et al., 1997; Zubieta et al., 1999; Zubieta et al., 1996), thus NTX may be directly mediating its effects on OFC activity via local receptors. However, many of the OFC's inputs, including the dorsal prefrontal cortex, temporal cortex, the amygdala, and the VTA, also show rich expression of opioid receptors (Gorelick et al., 2005; Maurer et al., 1983; Staley et al., 1997; Zubieta et al., 1999; Zubieta et al., 1996), thus NTX's effect on OFC activity may alternatively be mediated indirectly through one or more of these areas. For example, opioid agonists directly inhibit cortically projecting VTA dopamine neurons in rats (Margolis et al., 2006), thus NTX's actions at opioid receptors in the VTA could increase dopaminergic input to the OFC, resulting in the heightened OFC activity during decision-making that we observed. However, we do not currently know how endogenous opioids that affect the release of DA alter the BOLD signal in the human OFC. Rat studies indicate that whereas μ -opioids in the VTA increase cortical and striatal DA, κ -opioids inhibit striatal dopamine, since NTX blocks both receptors, its effect on cortical DA levels is uncertain (Herz, 1995; Margolis et al., 2006; Spanagel et al., 1992) and could depend on the relative level of μ -versus κ -mediated actions. Relative μ -versus κ -effects would be expected to differ in subjects with low circulating levels of endogenous μ -receptor ligands, such as alcoholics and their offspring (Dai et al., 2005; del Arbol et al., 1995; Govoni et al., 1983; Vescovi et al., 1992), or in those with low levels of μ -receptor expression, as is found in subjects with low frontal DA levels due to their COMT genotype (Berthele et al., 2005), and in subjects with the A118G polymorphism of the μ -opioid receptor (Zhang et al., 2005). Combining PET imaging of NTX binding with behavioral studies of immediate reward bias shifts may shed light on this issue.

Clinical and pre-clinical data support the hypothesis that opioids regulate immediate reward bias. In rats, NTX reverses morphine induced immediate reward bias (Kieres et al., 2004). In humans, NTX shifts preference from immediate alcohol reward to delayed monetary reward in a lab bar setting (O'Malley et al., 2002). Moreover, NTX attenuates a variety of impulsive behaviors, including pathological gambling (Kim et al., 2001), binge eating in bulimics (Marrazzi et al., 1995), compulsive sexual behavior (Raymond et al., 2002), self-injurious behavior (Symons et al., 2004), and kleptomania (Grant, 2005). Together these results suggest that opioids promote impulsivity, and that opioid receptor blockade via NTX may improve self-control. The fact that immediate reward bias is reduced by acute elevation of DA levels (de Wit et al., 2002; Wade et al., 2000), further suggests that NTX may alter immediate reward bias via indirect effects on DA release, an idea consistent with studies demonstrating opioid regulation of dopamine neurons (Berthele et al., 2005; Margolis et al., 2006; Ostrowski et al., 1982; Sesack and Pickel, 1992).

6.3. BDNF and the endogenous opioid system

The Met allele of the BDNF Val66Met polymorphism is associated with reduced activity-dependent release of BDNF, impaired episodic memory, and reduced Hippocampal function (Egan et al., 2003). It has also been associated with increased anxiety (Chen et al., 2006), increased risk of eating disorders and schizophrenia, and reduced risk of substance abuse disorders (Gratacos et al., 2008). Although endogenous opioids are known to regulate BDNF expression (Zhang et al., 2007; Zhang et al., 2006), it remains unclear how variation in the BDNF system could alter response to endogenous opioid blockade. One recent study however indicates that genetic variability in the BDNF gene alters response to methadone treatment in opioid addicts (de Cid et al., 2008), providing some precedence for variability in opioid system responses based on BDNF genotype.

6.4. Limitations

6.4.1. Sample size—One limitation of the results reported here is the small sample size. This limitation is particularly critical for interpreting the genetic association data. Thus, the finding that BDNF genotype predicts NTX effects on impulsive decision-making should be considered as pilot data, which await confirmation in a larger scale study. It is also critical to acknowledge that our small sample size may have resulted in false negatives with regard to the moderating effects of the other SNPs tested.

6.4.2. Acute dose of NTX—A second limitation of the present study is the use of a single acute dose of NTX. Typical therapeutic use of NTX entails daily dosing, which may result in rather different responses to the drug. However, clinical data demonstrate equivalent or greater efficacy of acute NTX dosing, relative to daily maintenance, in reducing excessive alcohol intake (Hernandez-Avila et al., 2006).

6.4.3. No clinical outcome measures—A final limitation of the present study is the lack of any clinical outcome measures. Thus, we are unable to say whether the NTX effects on decision-making behavior or on brain activity during decision-making are directly correlated with NTX's therapeutic effects. Future studies that incorporate this decision-making task into a clinical trial would shed light on this issue.

6.5. Summary

The present findings identify actions of NTX in the human brain that may contribute to its therapeutic effect in the treatment of alcoholism. Moreover, OFC activity during subjective decisions may be a key therapeutic target for alcoholism medication development. Individual differences in NTX effects on the OFC may contribute to the variable clinical efficacy of NTX. Individual differences in BDNF genotype may also contribute to the variability in NTX's effects on behavior.

Acknowledgments

This work was sponsored by the Department of the Army under Award No. W81XWH-07-1-0430, the Wheeler Center for the Neurobiology of Addiction, and Award Number KL2RR025746 from the National Center for Research Resources (CAB). The authors acknowledge the expert technical assistance of V.C. Tavares, M. Robertson, and G. Josslyn.

References

Altshuler HL, Phillips PE, Feinhandler DA. Alteration of ethanol self-administration by naltrexone. *Life sciences* 1980;26:679–688. [PubMed: 6767889]

- Anton RF, Moak DH, Waid LR, Latham PK, Malcolm RJ, Dias JK. Naltrexone and cognitive behavioral therapy for the treatment of outpatient alcoholics: results of a placebo-controlled trial. *The American journal of psychiatry* 1999;156:1758–1764. [PubMed: 10553740]
- Anton RF, Oroszi G, O'Malley S, Couper D, Swift R, Pettinati H, Goldman D. An evaluation of mu-opioid receptor (OPRM1) as a predictor of naltrexone response in the treatment of alcohol dependence: results from the Combined Pharmacotherapies and Behavioral Interventions for Alcohol Dependence (COMBINE) study. *Archives of general psychiatry* 2008;65:135–144. [PubMed: 18250251]
- Berthele A, Platzer S, Jochim B, Boecker H, Buettner A, Conrad B, Riemenschneider M, Toelle TR. COMT Val108/158Met genotype affects the mu-opioid receptor system in the human brain: evidence from ligand-binding, G-protein activation and preproenkephalin mRNA expression. *NeuroImage* 2005;28:185–193. [PubMed: 16040257]
- Boettiger CA, Mitchell JM, Tavares VC, Robertson M, Joslyn G, D'Esposito M, Fields HL. Immediate reward bias in humans: fronto-parietal networks and a role for the catechol-O-methyltransferase 158 (Val/Val) genotype. *J Neurosci* 2007;27:14383–14391. [PubMed: 18160646]
- Boettiger, CASA.; Sun, FT.; D'Esposito, M. Society for Neuroscience Abstracts. San Diego: 2004. Identifying Frontal Networks For Replacing Well-Established Response Associations; p. 664.611
- Bond C, LaForge KS, Tian M, Melia D, Zhang S, Borg L, Gong J, Schluger J, Strong JA, Leal SM, Tischfield JA, Kreek MJ, Yu L. Single-nucleotide polymorphism in the human mu opioid receptor gene alters beta-endorphin binding and activity: possible implications for opiate addiction. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:9608–9613. [PubMed: 9689128]
- Boyle AE, Stewart RB, Macenski MJ, Spiga R, Johnson BA, Meisch RA. Effects of acute and chronic doses of naltrexone on ethanol self-administration in rhesus monkeys. *Alcoholism, clinical and experimental research* 1998;22:359–366.
- Chen Z-Y, Jing D, Bath KG, Ieraci A, Khan T, Siao C-J, Herrera DG, Toth M, Yang C, McEwen BS, Hempstead BL, Lee FS. Genetic Variant BDNF (Val66Met) Polymorphism Alters Anxiety-Related Behavior. *Science* 2006;314:140–143. [PubMed: 17023662]
- Ciccocioppo R, Lin D, Martin-Fardon R, Weiss F. Reinstatement of ethanol-seeking behavior by drug cues following single versus multiple ethanol intoxication in the rat: effects of naltrexone. *Psychopharmacology* 2003;168:208–215. [PubMed: 12664190]
- Ciccocioppo R, Martin-Fardon R, Weiss F. Effect of selective blockade of mu(1) or delta opioid receptors on reinstatement of alcohol-seeking behavior by drug-associated stimuli in rats. *Neuropsychopharmacology* 2002;27:391–399. [PubMed: 12225696]
- Cools R, Clark L, Owen AM, Robbins TW. Defining the neural mechanisms of probabilistic reversal learning using event-related functional magnetic resonance imaging. *J Neurosci* 2002;22:4563–4567. [PubMed: 12040063]
- Dai X, Thavundayil J, Gianoulakis C. Differences in the peripheral levels of beta-endorphin in response to alcohol and stress as a function of alcohol dependence and family history of alcoholism. *Alcoholism, clinical and experimental research* 2005;29:1965–1975.
- Davidson D, Palfai T, Bird C, Swift R. Effects of naltrexone on alcohol self-administration in heavy drinkers. *Alcoholism, clinical and experimental research* 1999;23:195–203.
- de Cid R, Fonseca F, Gratacos M, Gutierrez F, Martin-Santos R, Estivill X, Torrens M. BDNF variability in opioid addicts and response to methadone treatment: preliminary findings. *Genes, brain, and behavior*. 2008
- De Panfilis C, Schwarzbauer C. Positive or negative blips? The effect of phase encoding scheme on susceptibility-induced signal losses in EPI. *NeuroImage* 2005;25:112–121. [PubMed: 15734348]
- de Wit H, Enggasser JL, Richards JB. Acute administration of d-amphetamine decreases impulsivity in healthy volunteers. *Neuropsychopharmacology* 2002;27:813–825. [PubMed: 12431855]
- de Wit H, Svenson J, York A. Non-specific effect of naltrexone on ethanol consumption in social drinkers. *Psychopharmacology* 1999;146:33–41. [PubMed: 10485962]
- del Arbol, JL.; Aguirre, JC.; Raya, J.; Rico, J.; Ruiz-Requena, ME.; Miranda, MT. Alcohol. Vol. 12. Fayetteville; N.Y: 1995. Plasma concentrations of beta-endorphin, adrenocorticotrophic hormone, and cortisol in drinking and abstinent chronic alcoholics; p. 525-529.

- Dias R, Robbins TW, Roberts AC. Dissociation in prefrontal cortex of affective and attentional shifts. *Nature* 1996;380:69–72. [PubMed: 8598908]
- Dom G, Sabbe B, Hulstijn W, van den Brink W. Substance use disorders and the orbitofrontal cortex: systematic review of behavioural decision-making and neuroimaging studies. *Br J Psychiatry* 2005;187:209–220. [PubMed: 16135857]
- Duvernoy, HM.; Cabanis, EA.; Vannson, JL. Wien. New York: Springer-Verlag; 1991. The human brain : surface, three-dimensional sectional anatomy and MRI.
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, Zaitsev E, Gold B, Goldman D, Dean M, Lu B, Weinberger DR. The BDNF val66met Polymorphism Affects Activity-Dependent Secretion of BDNF and Human Memory and Hippocampal Function. *Cell* 2003;112:257–269. [PubMed: 12553913]
- Elliott R, Dolan RJ, Frith CD. Dissociable functions in the medial and lateral orbitofrontal cortex: evidence from human neuroimaging studies. *Cereb Cortex* 2000;10:308–317. [PubMed: 10731225]
- Ersche KD, Fletcher PC, Lewis SJ, Clark L, Stocks-Gee G, London M, Deakin JB, Robbins TW, Sahakian BJ. Abnormal frontal activations related to decision-making in current and former amphetamine and opiate dependent individuals. *Psychopharmacology* 2005;180:612–623. [PubMed: 16163533]
- Friston KJ, Ashburner J, Frith CD, Poline JB, Heather JD, Frackowiak RSJ. Spatial registration and normalization of images. *Human Brain Mapping* 1995;3:165–189.
- Gelernter J, Gueorguieva R, Kranzler HR, Zhang H, Cramer J, Rosenheck R, Krystal JH. Opioid receptor gene (OPRM1, OPRK1, and OPRD1) variants and response to naltrexone treatment for alcohol dependence: results from the VA Cooperative Study. *Alcoholism, clinical and experimental research* 2007;31:555–563.
- Gorelick DA, Kim YK, Bencherif B, Boyd SJ, Nelson R, Copersino M, Endres CJ, Dannals RF, Frost JJ. Imaging brain mu-opioid receptors in abstinent cocaine users: time course and relation to cocaine craving. *Biological psychiatry* 2005;57:1573–1582. [PubMed: 15953495]
- Govoni S, Bosio A, Di Monda E, Fazzari G, Spano PF, Trabucchi M. Immunoreactive met-enkephalin plasma concentrations in chronic alcoholics and in children born from alcoholic mothers. *Life sciences* 1983;33:1581–1586. [PubMed: 6633159]
- Grant JE. Outcome study of kleptomania patients treated with naltrexone: a chart review. *Clinical neuropharmacology* 2005;28:11–14. [PubMed: 15711433]
- Gratacos M, Soria V, Urretavizcaya M, Gonzalez JR, Crespo JM, Bayes M, de Cid R, Menchon JM, Vallejo J, Estivill X. A brain-derived neurotrophic factor (BDNF) haplotype is associated with antidepressant treatment outcome in mood disorders. *The pharmacogenomics journal* 2008;8:101–112. [PubMed: 17505499]
- Harris RA, Erickson CK. Alteration of ethanol effects by opiate antagonists. *Currents in alcoholism* 1979;5:17–28. [PubMed: 755620]
- Heidbreder C. Recent Advances in the Pharmacotherapeutic Management of Drug Dependence and Addiction. *Current Psychiatry Reviews* 2005;1:45–67.
- Hernandez-Avila CA, Song C, Kuo L, Tennen H, Armeli S, Kranzler HR. Targeted versus daily naltrexone: secondary analysis of effects on average daily drinking. *Alcoholism, clinical and experimental research* 2006;30:860–865.
- Herz, A. The Pharmacology of Opioids. Germany: Harwood; 1995. Endogenous opioids and addiction; p. 445-462.
- Izquierdo A, Suda RK, Murray EA. Bilateral orbital prefrontal cortex lesions in rhesus monkeys disrupt choices guided by both reward value and reward contingency. *J Neurosci* 2004;24:7540–7548. [PubMed: 15329401]
- Kieres AK, Hausknecht KA, Farrar AM, Acheson A, de Wit H, Richards JB. Effects of morphine and naltrexone on impulsive decision making in rats. *Psychopharmacology* 2004;173:167–174. [PubMed: 14752586]
- Kim SW, Grant JE, Adson DE, Shin YC. Double-blind naltrexone and placebo comparison study in the treatment of pathological gambling. *Biological psychiatry* 2001;49:914–921. [PubMed: 11377409]
- Kreek MJ. Opiates, opioids and addiction. *Mol Psychiatry* 1996;1:232–254. [PubMed: 9118348]

- Liu X, Weiss F. Additive effect of stress and drug cues on reinstatement of ethanol seeking: exacerbation by history of dependence and role of concurrent activation of corticotropin-releasing factor and opioid mechanisms. *J Neurosci* 2002;22:7856–7861. [PubMed: 12223538]
- London ED, Ernst M, Grant S, Bonson K, Weinstein A. Orbitofrontal cortex and human drug abuse: functional imaging. *Cereb Cortex* 2000;10:334–342. [PubMed: 10731228]
- Margolis EB, Lock H, Chefer VI, Shippenberg TS, Hjelmstad GO, Fields HL. Kappa opioids selectively control dopaminergic neurons projecting to the prefrontal cortex. *Proceedings of the National Academy of Sciences of the United States of America* 2006;103:2938–2942. [PubMed: 16477003]
- Marrazzi MA, Markham KM, Kinzie J, Luby ED. Binge eating disorder: response to naltrexone. *Int J Obes Relat Metab Disord* 1995;19:143–145. [PubMed: 7735342]
- Maurer R, Cortes R, Probst A, Palacios JM. Multiple opiate receptor in human brain: an autoradiographic investigation. *Life sciences* 1983;33 (Suppl 1):231–234. [PubMed: 6319868]
- McAlonan K, Brown VJ. Orbital prefrontal cortex mediates reversal learning and not attentional set shifting in the rat. *Behavioural brain research* 2003;146:97–103. [PubMed: 14643463]
- McCaul ME, Wand GS, Eissenberg T, Rohde CA, Cheskin LJ. Naltrexone alters subjective and psychomotor responses to alcohol in heavy drinking subjects. *Neuropsychopharmacology* 2000;22:480–492. [PubMed: 10731623]
- McGeary JE, Monti PM, Rohsenow DJ, Tidey J, Swift R, Miranda R Jr. Genetic moderators of naltrexone's effects on alcohol cue reactivity. *Alcoholism, clinical and experimental research* 2006;30:1288–1296.
- Mishkin, M. Perseveration of central sets after frontal lesions in monkeys. In: Mishkin, M.; JW; Akert, K., editors. *The frontal granular cortex and behavior*. New York: McGraw-Hill; 1964. p. 219–241.
- Mitchell, JM.; Bergren, LJ.; Chen, KS.; Fields, HL. *Society for Neuroscience*. Washington, DC: 2005a. Prior ethanol consumption predicts aversion elicited by the opioid antagonist naltrexone in rats.
- Mitchell JM, Fields HL, D'Esposito M, Boettiger CA. Impulsive responding in alcoholics. *Alcoholism, clinical and experimental research* 2005b;29:2158–2169.
- Mitchell JM, Tavares VC, Fields HL, D'Esposito M, Boettiger CA. Endogenous opioid blockade and impulsive responding in alcoholics and healthy controls. *Neuropsychopharmacology* 2007;32:439–449. [PubMed: 17047667]
- Mobini S, Body S, Ho MY, Bradshaw CM, Szabadi E, Deakin JF, Anderson IM. Effects of lesions of the orbitofrontal cortex on sensitivity to delayed and probabilistic reinforcement. *Psychopharmacology* 2002;160:290–298. [PubMed: 11889498]
- Monti PM, Rohsenow DJ, Hutchison KE, Swift RM, Mueller TI, Colby SM, Brown RA, Gulliver SB, Gordon A, Abrams DB. Naltrexone's effect on cue-elicited craving among alcoholics in treatment. *Alcoholism, clinical and experimental research* 1999;23:1386–1394.
- Myrick H, Anton RF, Li X, Henderson S, Randall PK, Voronin K. Effect of Naltrexone and Ondansetron on Alcohol Cue-Induced Activation of the Ventral Striatum in Alcohol-Dependent People. *Archives of general psychiatry* 2008;65:466–475. [PubMed: 18391135]
- Nichols T, Hayasaka S. Controlling the familywise error rate in functional neuroimaging: a comparative review. *Statistical Methods in Medical Research* 2003;12:419–446. [PubMed: 14599004]
- Nichols TE, Holmes AP. Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Hum Brain Mapp* 2002;15:1–25. [PubMed: 11747097]
- O'Brien, CP.; Volpicelli, LA.; Volpicelli, JR. *Alcohol*. Vol. 13. Fayetteville; N.Y: 1996. Naltrexone in the treatment of alcoholism: a clinical review; p. 35–39.
- O'Doherty J, Dayan P, Schultz J, Deichmann R, Friston K, Dolan RJ. Dissociable roles of ventral and dorsal striatum in instrumental conditioning. *Science* 2004;304:452–454. [PubMed: 15087550]
- O'Malley SS, Krishnan-Sarin S, Farren C, Sinha R, Kreek MJ. Naltrexone decreases craving and alcohol self-administration in alcohol-dependent subjects and activates the hypothalamo-pituitary-adrenocortical axis. *Psychopharmacology* 2002;160:19–29. [PubMed: 11862370]
- Oslin DW, Berrettini W, Kranzler HR, Pettinati H, Gelernter J, Volpicelli JR, O'Brien CP. A functional polymorphism of the mu-opioid receptor gene is associated with naltrexone response in alcohol-dependent patients. *Neuropsychopharmacology* 2003;28:1546–1552. [PubMed: 12813472]
- Ostlund SB, Balleine BW. Orbitofrontal cortex mediates outcome encoding in Pavlovian but not instrumental conditioning. *J Neurosci* 2007;27:4819–4825. [PubMed: 17475789]

- Ostrowski NL, Hatfield CB, Caggiula AR. The effects of low doses of morphine on the activity of dopamine-containing cells and on behavior. *Life sciences* 1982;31:2347–2350. [PubMed: 7162350]
- Padoa-Schioppa C, Assad JA. Neurons in the orbitofrontal cortex encode economic value. *Nature* 2006;441:223–226. [PubMed: 16633341]
- Rahman S, Sahakian BJ, Hodges JR, Rogers RD, Robbins TW. Specific cognitive deficits in mild frontal variant frontotemporal dementia. *Brain* 1999;122 (Pt 8):1469–1493. [PubMed: 10430832]
- Ray LA, Hutchison KE. Effects of naltrexone on alcohol sensitivity and genetic moderators of medication response: a double-blind placebo-controlled study. *Archives of general psychiatry* 2007;64:1069–1077. [PubMed: 17768272]
- Raymond NC, Grant JE, Kim SW, Coleman E. Treatment of compulsive sexual behaviour with naltrexone and serotonin reuptake inhibitors: two case studies. *International clinical psychopharmacology* 2002;17:201–205. [PubMed: 12131605]
- Remijne PL, Nielen MM, van Balkom AJ, Cath DC, van Oppen P, Uylings HB, Veltman DJ. Reduced orbitofrontal-striatal activity on a reversal learning task in obsessive-compulsive disorder. *Archives of general psychiatry* 2006;63:1225–1236. [PubMed: 17088503]
- Roesch MR, Olson CR. Neuronal activity related to reward value and motivation in primate frontal cortex. *Science* 2004;304:307–310. [PubMed: 15073380]
- Rohsenow DJ, Monti PM, Hutchison KE, Swift RM, Colby SM, Kaplan GB. Naltrexone's effects on reactivity to alcohol cues among alcoholic men. *Journal of abnormal psychology* 2000;109:738–742. [PubMed: 11195999]
- Rolls ET, Hornak J, Wade D, McGrath J. Emotion-related learning in patients with social and emotional changes associated with frontal lobe damage. *Journal of neurology, neurosurgery, and psychiatry* 1994;57:1518–1524.
- Rudebeck PH, Walton ME, Smyth AN, Bannerman DM, Rushworth MF. Separate neural pathways process different decision costs. *Nature neuroscience* 2006;9:1161–1168.
- Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction (Abingdon, England)* 1993;88:791–804.
- Schmahmann JD, Doyon J, McDonald D, Holmes C, Lavoie K, Hurwitz AS, Kabani N, Toga A, Evans A, Petrides M. Three-Dimensional MRI Atlas of the Human Cerebellum in Proportional Stereotaxic Space. *NeuroImage* 1999;10:233–260. [PubMed: 10458940]
- Schoenbaum G, Roesch M. Orbitofrontal cortex, associative learning, and expectancies. *Neuron* 2005;47:633–636. [PubMed: 16129393]
- Schoenbaum G, Saddoris MP, Stalnaker TA. Reconciling the roles of orbitofrontal cortex in reversal learning and the encoding of outcome expectancies. *Annals of the New York Academy of Sciences* 2007;1121:320–335. [PubMed: 17698988]
- Sesack SR, Pickel VM. Dual ultrastructural localization of enkephalin and tyrosine hydroxylase immunoreactivity in the rat ventral tegmental area: multiple substrates for opiate-dopamine interactions. *J Neurosci* 1992;12:1335–1350. [PubMed: 1348271]
- Sinclair JD. Evidence about the use of naltrexone and for different ways of using it in the treatment of alcoholism. *Alcohol and alcoholism (Oxford, Oxfordshire)* 2001;36:2–10.
- Spanagel R, Herz A, Shippenberg TS. Opposing tonically active endogenous opioid systems modulate the mesolimbic dopaminergic pathway. *Proceedings of the National Academy of Sciences of the United States of America* 1992;89:2046–2050. [PubMed: 1347943]
- Staley JK, Rothman RB, Rice KC, Partilla J, Mash DC. Kappa2 opioid receptors in limbic areas of the human brain are upregulated by cocaine in fatal overdose victims. *J Neurosci* 1997;17:8225–8233. [PubMed: 9334398]
- Stromberg, MF.; Casale, M.; Volpicelli, L.; Volpicelli, JR.; O'Brien, CP. Alcohol. Vol. 15. Fayetteville; N.Y.: 1998. A comparison of the effects of the opioid antagonists naltrexone, naltrindole, and beta-funaltrexamine on ethanol consumption in the rat; p. 281–289.
- Swift RM, Whelihan W, Kuznetsov O, Buongiorno G, Hsuing H. Naltrexone-induced alterations in human ethanol intoxication. *The American journal of psychiatry* 1994;151:1463–1467. [PubMed: 8092339]

- Symons FJ, Thompson A, Rodriguez MC. Self-injurious behavior and the efficacy of naltrexone treatment: a quantitative synthesis. *Mental retardation and developmental disabilities research reviews* 2004;10:193–200. [PubMed: 15611982]
- Tait DS, Brown VJ. Difficulty overcoming learned non-reward during reversal learning in rats with ibotenic acid lesions of orbital prefrontal cortex. *Annals of the New York Academy of Sciences* 2007;1121:407–420. [PubMed: 17698991]
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B, Joliot M. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage* 2002;15:273–289. [PubMed: 11771995]
- Vescovi PP, Coiro V, Volpi R, Giannini A, Passeri M. Plasma beta-endorphin, but not met-enkephalin levels are abnormal in chronic alcoholics. *Alcohol and alcoholism (Oxford, Oxfordshire)* 1992;27:471–475.
- Volkow ND, Fowler JS. Addiction, a disease of compulsion and drive: involvement of the orbitofrontal cortex. *Cereb Cortex* 2000;10:318–325. [PubMed: 10731226]
- Volpicelli JR, Watson NT, King AC, Sherman CE, O'Brien CP. Effect of naltrexone on alcohol "high" in alcoholics. *The American journal of psychiatry* 1995;152:613–615. [PubMed: 7694913]
- Wade TR, de Wit H, Richards JB. Effects of dopaminergic drugs on delayed reward as a measure of impulsive behavior in rats. *Psychopharmacology* 2000;150:90–101. [PubMed: 10867981]
- Zhang H, Shi YG, Woods JH, Watson SJ, Ko MC. Central kappa-opioid receptor-mediated antidepressant-like effects of nor-Binaltorphimine: behavioral and BDNF mRNA expression studies. *European journal of pharmacology* 2007;570:89–96. [PubMed: 17601558]
- Zhang H, Torregrossa MM, Jutkiewicz EM, Shi YG, Rice KC, Woods JH, Watson SJ, Ko MC. Endogenous opioids upregulate brain-derived neurotrophic factor mRNA through delta- and micro-opioid receptors independent of antidepressant-like effects. *The European journal of neuroscience* 2006;23:984–994. [PubMed: 16519663]
- Zhang Y, Wang D, Johnson AD, Papp AC, Sadee W. Allelic expression imbalance of human mu opioid receptor (OPRM1) caused by variant A118G. *The Journal of biological chemistry* 2005;280:32618–32624. [PubMed: 16046395]
- Zubieta JK, Dannals RF, Frost JJ. Gender and age influences on human brain mu-opioid receptor binding measured by PET. *The American journal of psychiatry* 1999;156:842–848. [PubMed: 10360121]
- Zubieta JK, Gorelick DA, Stauffer R, Ravert HT, Dannals RF, Frost JJ. Increased mu opioid receptor binding detected by PET in cocaine-dependent men is associated with cocaine craving. *Nature medicine* 1996;2:1225–1229.

Figure 1. Illustration of behavioral paradigm

The temporal sequence of events is depicted for an example WANT trial. Illumination of a fixation cross (“Ready”) indicated the start of each trial. The instruction cue (in this case “WANT”) was then displayed for 4.4 s, informing the subject as to the upcoming trial type. The two options (*Now* and *Later*) then appeared while the instruction cue remained on the screen. The specific dollar amounts (from \$1 to \$100) and the time of availability (from “Today” to “6 months” later) varied across trials, as did the differences between the two options. The options remained on the screen for 4.4 s, although subjects had 6.6 s to indicate their choice.

Figure 2. NTX significantly elevates activity during decision-making in the OFC
Plot of the mean *Now* vs. *Later* subjective decision-making parameter estimates for the OFC ROI as a function of drug condition. Plots reflect mean $\pm$ S.D. NTX, naltrexone; PBO, placebo.

Figure 3. Genotype at the BDNF 66 locus predicts NTX effect on ICR

Ratio of impulsive choices as a function of genotype and drug condition. There was a significant interaction effect between drug condition and genotype at the BDNF 66 locus on impulsive choice probability. The 66^{Met/Met} genotype was associated with greater *Now* bias on NTX. Met/Met, 66^{Met/Met}; Val/Met, 66^{Val/Met}; Val/Val, 66^{Val/Val}; BDNF, brain derived neurotrophic factor.

Figure 4. Mapwise analyses of NTX effects

Statistical non-parametric *t*-maps contrasting [WANT-CON] activity in NTX and placebo sessions, overlaid on a standard T1-weighted anatomical image. Criterion threshold was based on permutation-based voxelwise correction of FWE ($p < 0.001$), with a minimum cluster size of 5 contiguous voxels. A) Areas more active during subjective decision-making on NTX. B) The indicated area in the superior temporal gyrus was less active during subjective decision-making on NTX. Cr1, cerebellum crus 1; Cr2, cerebellum crus 2; 7b, cerebellum lobule 7b; IT, inferotemporal cortex; OFC, orbitofrontal cortex; STG, superior temporal gyrus.

Figure 5. NTX effect on the OFC predicts NTX effect on choice behavior

Statistical non-parametric t -maps depicting results of a mapwise simple regression analysis. A) A site in the OFC is shown in which NTX's effect on activity during decision-making significantly correlated with NTX's effect on choice behavior. B) Plot of each subject's NTX session change in *Now* vs. *Later* subjective decision-making parameter estimate as a function of change in ICR for the peak of the region displayed in B; peak regression statistic displayed in plot. Units are in standard deviations, mean = 0.

Table 1**Subject demographics**

Values are reported as mean $\pm$ standard deviation. Reported *p*-values reflect the results of unpaired two-tailed comparison between groups.

Participants	AA (<i>n</i> = 9)	CS (<i>n</i> =10)	<i>t</i> ₁₇	<i>p</i> value
<i>General</i>				
Age (yrs)	30 $\pm$ 8	27 $\pm$ 3	1.49	<i>ns</i>
Education (Hollingshead subscore)	6 $\pm$ 1	6 $\pm$ 1	1.77	<i>ns</i>
Hollingshead SES	51 $\pm$ 9	50 $\pm$ 2	0.34	<i>ns</i>
Shipley Institute of Living Scale (raw)	65 $\pm$ 8	72 $\pm$ 8	1.70	<i>ns</i>
Gender (# female)	5	3		<i>ns</i> [†]
Ethnicity (# non-white)	2	1		<i>ns</i> [†]
<i>Alcohol-related</i>				
AUDIT	19 $\pm$ 7	5 $\pm$ 2	6.30	< 0.001
DUSI-I (%)	75 $\pm$ 15	13 $\pm$ 18	8.06	< 0.001
FTQ (#)	3 $\pm$ 2	1 $\pm$ 1	2.99	< 0.01

AA, Abstinent alcoholic; CS, Control subject; SES, socioeconomic status; AUDIT, Alcohol Use Disorders Identification Test; DUSI-I, Drug Use Screening Inventory, Domain I; FTQ, Family Tree Questionnaire;

[†]
p-value represents results of χ^2 test.

ns: *p* > 0.05

Table 2**NTX effects on behavior**

Changes in reaction times and ICR in the delay discounting task following 50mg naltrexone (NTX), relative to the placebo session.

Participants	AA (n = 9)	CS (n=10)	All (n=19)
<i>Reaction Times Changes (%)</i>			
CON	-6 ± 9	8 ± 10	1 ± 12
W	-5 ± 15	3 ± 14	-1 ± 15
DW	-4 ± 16	3 ± 17	0 ± 16
<i>ICR Changes (%)</i>	5 ± 14	3 ± 14	4 ± 14

AA, Abstinent alcoholic; CS, Control subject; W, WANT trials; DW, DON'T WANT trials; CON, Control trials (SOONER & LARGER); ICR, Impulsive choice ratio. All data given as mean ± standard deviation.

Table 3**Peak coordinates for ROI analyses**

ROI anatomical region, hemisphere, and coordinates of ROI center based on the Montreal Neurological Institute (MNI) coordinate system value are given.

Brain region	Hemisphere	MNI coordinates [x, y, z] (mm)
<i>Now ROI centers</i>		
Supramarginal gyrus (parietal lobe)	Right	66, -42, 44
Cerebellum, posterior lobe (lobule VIII)	Right	36, -52, -62
Middle temporal gyrus	Right	56, -52, -2
Inferior temporal gyrus	Right	68, -50, -12
Parahippocampal gyrus (Gyrus ambiens, Entorhinal area, Semilunar gyrus)	Right	18, -4, -28
Superior frontal gyrus	Left	-22, 50, 44
<i>Later ROI center</i>		
Lateral orbital gyrus	Right	58, 36, -10

Table 4**Peak coordinates for sites of activity modulation by NTX**

Anatomical region, hemisphere, and peak coordinates based on the Montreal Neurological Institute (MNI) coordinate system value are given.

Brain region	Hemisphere	MNI coordinates [x, y, z] (mm)
NTX <i>increased</i> activity during decision-making		
Lateral orbital gyrus	Left	-40, 46, -11
Inferior temporal gyrus	Left	-37, 0, -41
Cerebellum, crus I	Left	-37, -42, -35
Cerebellum, lobule VIIb	Left	-44, -42, -49
Cerebellum, crus I	Right	45, -46, -35
Cerebellum, crus II	Left	-20, -76, -45
NTX <i>decreased</i> activity during decision-making		
Superior temporal gyrus	Right	63, 10, -8

Table 5**Peak coordinates for sites in which NTX co-modulated BOLD and ICR**

Anatomical region, hemisphere, and peak coordinates based on the Montreal Neurological Institute (MNI) coordinate system value are given.

Brain region	Hemisphere	MNI coordinates [x, y, z] (mm)
NTX BOLD effect inversely related to ICR effect		
Middle frontal gyrus	Right	40, 46, 6
Lateral orbital gyrus	Right	51, 38, -16
Superior temporal pole	Left	-47, 12, -14
Middle temporal pole	Left	-27, 9, -37
Inferior frontal junction	Right	45, 8, 28
Inferior temporal gyrus	Right	50, 4, -38
Middle temporal gyrus	Left	-54, 2, -22
Supplementary motor area	Right	15, 1, 72
Cerebellum, lobule VI	Right	36, -43, -32
	Right	32, -64, -21
NTX BOLD effect directly related to ICR effect		
Middle orbital gyrus	Left	-20, 57, -14
Inferior temporal gyrus	Left	-59, -9, -29
Middle temporal gyrus	Left	-68, -38, 5
Intraparietal sulcus	Left	-44, -45, 61