Published in final edited form as:

Org Lett. 2009 May 21; 11(10): 2117–2119. doi:10.1021/ol9005757.

A Simple One-pot Conversion of Aldehydes and Ketones to Enals

Petr Valenta, Natalie A. Drucker, Jeffrey W. Bode, and Patrick J. Walsh*

P. Roy and Diana T. Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104-6323

Abstract

A simple and efficient method to convert aldehydes into α,β -unsaturated aldehydes with a two-carbon homologation is presented. Hydroboration of ethoxy acetylene with $\text{BH}_3\cdot\text{SMe}_2$ generates tris(ethoxyvinyl) borane. Transmetalation with diethylzinc, addition to aldehydes or ketones, and acidic workup affords enals. When the addition is quenched with anilinium hydrochloride, 1,2-dithioglycol, or acetic anhydride the unsaturated imine, dithiolane, or 1,1-diacetate is isolated in high yield. These transformations can be performed in a one-pot procedure.

The field of organocatalysis has witnessed explosive growth in the last decade and will continue to be a dynamic area of chemical research.¹ A wide variety of organocatalysts and reaction classes have been recently advanced, and many of these are based on α,β -unsaturated aldehydes as substrates. As a result, enals have arguably become the single most important class of substrates for development and applications of organocatalytic transformations² while remaining a mainstay of metal-based catalytic asymmetric processes.^{1a,3} To keep pace with the ever-increasing demand for enals, current synthetic methods must be reevaluated and optimized or supplanted with new, direct and more efficient approaches. Herein we disclose a simple and practical yet broadly applicable one-pot method for the conversion of aldehydes and ketones into α,β -unsaturated aldehydes.

A practical synthesis of enals would meet the following criteria: it must be general, employ commercially available and inexpensive reagents, use common laboratory techniques, be functional group compatible, and be tolerant of α -stereogenic centers in enantioenriched aldehydes and ketones. Furthermore, the transformation should be amenable to a one-pot procedure to enable rapid generation of an array of enals.

Existing methods for conversion of aldehydes and ketones to enals do not satisfy these criteria (Scheme 1). For example, cross aldol reactions with acetaldehyde enolates are not general, yielding self-condensation products with aliphatic aldehyde partners (Scheme 1, **A**).⁴ This difficulty can be circumvented with *C*-silylated imines (**B** and **C**),⁵ but these imines must be prepared, silylated, and distilled, adding additional steps. Wittig,⁶ Horner-Emmons,⁷ and Peterson-type reagents⁸ are generally not compatible with base-sensitive functional groups and/or can require additional synthetic steps to generate enals (**D**).

Methods employing commercially available ethoxy acetylene are most attractive (Scheme 1, **E** and **F**). Hydrostannylation of ethoxy acetylene provides a vinylstannane. Transmetallation with *n*-BuLi leads to a highly reactive vinylolithium that readily adds to aldehydes and ketones to afford enals on acidic workup.⁹ The drawback of such methods is the incompatibility of the vinylolithium with most functional groups. Suzuki introduced a method that combines the two-steps above into one and circumvents the highly reactive vinylolithium intermediate in **E**.¹⁰ Thus, hydrozirconation of ethoxy acetylene, in situ transmetallation of the vinylzirconocene intermediate with catalytic AgClO₄, addition to aldehydes and elimination provides enals (**F**). Although this method has advantages over those outlined above, it is not economically viable because Schwartz's reagent is prohibitively expensive (>\$2,000/mol from Aldrich).

Our approach to α,β -unsaturated aldehydes involves hydroboration of ethoxy acetylene with BH₃•SMe₂ to generate the tris(vinyl) borane (Scheme 2).¹¹ This intermediate can be prepared on scale and stored for months under a nitrogen atmosphere or used directly in situ. Transmetallation from boron to zinc at -78 °C and addition to aldehydes or ketones affords zinc alkoxy enol ethers,¹² which readily undergo elimination on workup with 2M HCl.

The broad substrate scope of this tandem reaction is illustrated in Table 1. Aliphatic (entries 1–3), aromatic (entries 4 and 8), and heteroaromatic aldehydes (entries 5–7) and conjugated enals (entries 9 and 10) and ynals (entry 11) undergo homologation to afford enals with 72–96% yield. Particularly noteworthy is the successful conversion of phenyl acetaldehyde to the expected enal in 89% yield, despite the acidic nature of its α -hydrogens (entry 3).

Ketone substrates were more challenging and exhibited variable results. Cyclohexanone (entry 12) was an excellent substrate and furnished the enal in 86% yield. Given the low reactivity of benzophenone and the mild nature of organozinc reagents,¹³ we were surprised to isolate 54% yield of the enal (entry 13). Unsymmetrical ketones, such as acetophenone (entry 14), gave little *E:Z* selectivity (3:1) unless the two groups flanking the carbonyl were significantly different in size, such as *tert*-butyl vs. methyl (*E:Z*=15:1, 53% yield, entry 15). To examine the selectivity of the vinylzinc intermediate toward aldehydes vs. ketones, a 1:1 mixture of cyclohexane carboxaldehyde and cyclohexanone were subjected to one equivalent ethoxy vinylzinc reagent. The more reactive aldehyde was converted to the enal product and the ketone was left untouched.

In the case of *p*-phthalaldehyde the di(enal) was isolated in 70% yield (entry 16). The *o*-derivative (entry 17), however, furnished the monoenal in 54% yield, most likely because the second aldehyde forms an internal hemiacetal derivative after the first addition, preventing further reaction.

Enantioenriched aldehydes are important starting materials and intermediates in synthesis. We therefore examined two optically active aldehyde substrates each with a stereocenter flanking the carbonyl. Benzyl protected 2-hydroxy propanal and Garner's aldehyde were converted to enals in 88 and 94% yield, respectively (entries 18 and 19). Both enals had >98% *ee* after isolation and purification, indicating the organozinc intermediate in this process is indeed mild.

In the development of practical and useful methods, scalability must be demonstrated. The homologation of cyclohexenecarboxaldehyde was therefore performed on a 10 mmol scale, affording the dialenal in 95% yield (entry 9).

Although enals are very important synthetic intermediates, derivatized enals are often desired. Under the assumption that the mechanism of conversion of hydroxy enol ethers to enals proceeds via oxocarbenium ions, we rationalized that these intermediates could be trapped by other nucleophiles. Thus, after addition of cinnamaldehyde to the vinylzinc to generate the alkoxide, 3 equivalents of anilinium hydrochloride were added leading to the imine in 90%

isolated yield (Scheme 3). Likewise, trapping with 2 equivalents 1,2-dithioglycol afforded the unsaturated 1,3-dithiolane in 98% yield. Unsaturated 1,1-diacetates are useful intermediates in palladium allylation chemistry.¹⁴ The diacetate was isolated in 92% yield after treatment of the alkoxide intermediate with aqueous HCl, removal of the volatile materials under reduced pressure and addition of 7 equivalents acetic anhydride and catalytic FeCl₃ (5 mol %).

In summary, we have developed a simple and efficient one-pot procedure that enables rapid access to α - β -unsaturated aldehydes from aldehydes and ketones. The advantages of this method include its extensive substrate scope, functional group compatibility, tolerance of stereocenters *alpha* to carbonyl groups, and low cost. The method can also be employed in the one-pot synthesis of synthetically valuable unsaturated aldimines, dithiolanes, and 1,1-diacetates. We anticipate this method will facilitate advances in organocatalysis with enal substrates.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by the NIH (National Institute of General Medical Sciences, GM58101) and NSF (CHE-065210). P.V. thanks the Isabel and Alfred Bader Foundation for a graduate fellowship.

References

1. Walsh, P.J.; Kozlowski, M.C. *Fundamentals of Asymmetric Catalysis*. Sausalito: University Science Books; 2008. (b) Enders D, Grondal C, Hüttel MR. *Angew. Chem., Int. Ed* 2007;46:1570–1581. (c) Dalko PI, Moisan L. *Angew. Chem., Int. Ed* 2004;43:5138–5175. (d) Lelais G, MacMillan DWC. *Aldrichimica Acta* 2006;39:79–87. (f) Seayad J, List B. *Org. Biomol. Chem* 2005;3:719–724. [PubMed: 15731852]
2. (a) Sohn SS, Rosen EL, Bode JW. *J. Am. Chem. Soc* 2004;126:14370–14371. [PubMed: 15521753] (b) Burstein C, Glorius F. *Angew. Chem., Int. Ed* 2004;43:6205–6208. (c) He M, Bode JW. *Org. Lett* 2005;7:3131–3134. [PubMed: 15987223] (d) Cabrera S, Reyes E, Aleman J, Milelli A, Kobbelgaard S, Jørgensen KA. *J. Am. Chem. Soc* 2008;130:12031–12037. [PubMed: 18698775] (e) Huang Y, Walji AM, Larsen CH, MacMillan DWC. *J. Am. Chem. Soc* 2005;127:15051–15053. [PubMed: 16248643] (f) Chen YK, Yoshida M, MacMillan DWC. *J. Am. Chem. Soc* 2006;128:9328–9329. [PubMed: 16848457] (g) Wilson RM, Jen WS, MacMillan DWC. *J. Am. Chem. Soc* 2005;127:11616–11617. [PubMed: 16104734] (h) Yang JW, Hechavarria Fonseca MT, List B. *J. Am. Chem. Soc* 2005;127:15036–15037. [PubMed: 16248637]
3. (a) Kim HY, Salvi L, Carroll PJ, Walsh PJ. *J. Am. Chem. Soc* 2009;131:954–962. [PubMed: 19154172] (b) Rowley Kelly A, Lurain AE, Walsh PJ. *J. Am. Chem. Soc* 2005;127:14668–14674. [PubMed: 16231920]
4. Mahata PK, Barun O, Ila H, Junjappa H. *Synlett* 2000;9:1345–1347.
5. (a) Gaudemar M, Bellassoued M. *Tetrahedron Lett* 1990;31:349–352. (b) Bellassoued M, Majidi A. *J. Org. Chem* 1993;58:2517–2522.
6. (a) Zhou G, Hu Q-Y, Corey EJ. *Org. Lett* 2003;5:3979–3982. [PubMed: 14535758] (b) Hudson RDA, Manning AR, Gallagher JF, Garcia MH, Lopes N, Asselberghs I, Boxel RV, Persoons A, Lough AJ. *J. Organomet. Chem* 2002;655:70–88. (c) Mladenova M, Ventelon L, Blanchard-Desce M. *Tetrahedron Lett* 1999;40:6923–6926. (d) Daubresse N, Francesch C, Rolando C. *Tetrahedron* 1998;54:10761–10770. (e) Cristau H-J, Gasc M-B. *Tetrahedron Lett* 1990;31:341–344. (f) Fraser-Reid B, Molino BF, Magdzinski L, Mootoo DR. *J. Org. Chem* 1987;52:4505–4511. (g) Cresp TM, Sargent MV, Vogel P. *J. Chem. Soc., Perkin Trans. 1* 1974:37–41.
7. (a) Michel P, Rassat A, Daly JW, Spande TF. *J. Org. Chem* 2000;65:8908–8918. [PubMed: 11149832] (b) Paterson I, Cumming JG, Ward RA, Lambole S. *Tetrahedron* 1995;51:9393–9412. (c) Nicolaou

- KC, Harter MW, Gunzner JL, Nadin A. *Liebigs Ann.-Recl* 1997;7:1283–1301. (d) Still WC, Gennari C. *Tet. Lett* 1983;24:4405–4408. (e) Ando K. *Tet. Lett* 1995;36:4105–4108.
8. (a) Corey EJ, Enders D, Bock MG. *Tetrahedron Lett* 1976;1:7–10. (b) Martin SF, Dodge JA, Burgess LE, Limberakis C, Hartmann M. *Tetrahedron* 1996;52:3229–3246.
9. Wollenberg RH, Albizati KF, Peries R. *J. Am. Chem. Soc* 1977;99:7365–7367.
10. (a) Maeta H, Hashimoto T, Hasegawa T, Suzuki K. *Tetrahedron Lett* 1992;33:5965–5968. (b) Maeta H, Suzuki K. *Tetrahedron Lett* 1993;34:341–344.
11. Miyaoura N, Maeda K, Sugimoto H. *J. Org. Chem* 1982;47:2117–2120.
12. Jeon S-J, Chen YK, Walsh PJ. *Org. Lett* 2005;7:1729–1732. [PubMed: 15844892]
13. García C, Libra ER, Carroll PJ, Walsh PJ. *J. Am. Chem. Soc* 2003;125:3210–3211. [PubMed: 12630865]
14. Trost BM, Lee CB, Weiss JM. *J. Am. Chem. Soc* 1995;117:7247–7248.

Scheme 1.
Synthesis of Enals

Scheme 2.
Our Synthesis of Enals

Org Lett. Author manuscript; available in PMC 2010 May 21.

entry	product	yield (%) ^a
11		94
87		87

yield (%)^ayield (%)^a entry product

86

12

94

iii) aq. HCl

yield (%)^a

yield (%)^a entry product

89

13

54

yield (%)^a

product

entry

yield (%)^a83^b

14

96

53^c

15

92

yield (%)^a

product

entry

yield (%)^a

70

16

91

yield (%)^ayield (%)^a entry product

72

17

54

Org Lett. Author manuscript; available in PMC 2010 May 21.

yield (%)^a

yield (%)^a

entry

product

88 (98% ee)^d

18

93

94 (98% ee)^d

19

95

95

^a Isolated yield.

^b *E:Z* ratio 1:3 (¹H NMR).

^c *E:Z* ratio >15:1 (¹H NMR).

^d *ee* determined by HPLC.