Published in final edited form as:

Biochemistry. 2008 September 9; 47(36): 9486–9496. doi:10.1021/bi800847a.

Structural Investigation of a Phosphorylation-Catalyzed, Isoaspartate-Free, Protein Succinimide: Crystallographic Structure of Post-Succinimide His15Asp Histidine-Containing Protein^{†,‡}

Scott Napper^{*,§}, Lata Prasad^{||}, and Louis T. J. Delbaere^{||}

[§]Vaccine and Infectious Disease Organization, University of Saskatchewan, Saskatoon, Saskatchewan S7N 5E3, Canada

^{||}Department of Biochemistry, University of Saskatchewan, Saskatoon, Saskatchewan S7N 5E5, Canada

Abstract

Aspartates and asparagines can spontaneously cyclize with neighboring main-chain amides to form succinimides. These succinimides hydrolyze to a mixture of isoaspartate and aspartate products. Phosphorylation of aspartates is a common mechanism of protein regulation and increases the propensity for succinimide formation. Although typically regarded as a form of protein damage, we hypothesize succinimides could represent an effective mechanism of phosphoaspartate autophosphatase activity, provided hydrolysis is limited to aspartate products. We previously reported the serendipitous creation of a protein, His15Asp histidine-containing protein (HPr), which undergoes phosphorylation-catalyzed formation of a succinimide whose hydrolysis is seemingly exclusive for aspartate formation. Here, through the high-resolution structure of postsuccinimide His15Asp HPr, we confirm the absence of isoaspartate residues and propose mechanisms for phosphorylation-catalyzed succinimide formation and its directed hydrolysis to aspartate. His15Asp HPr represents the first characterized protein example of an isoaspartate-free succinimide and lends credence to the hypothesis that intramolecular cyclization could represent a physiological mechanism of autophosphatase activity. Furthermore, this indicates that current strategies for succinimide evaluation, based on isoaspartate detection, underestimate the frequencies of these reactions. This is considerably significant for evaluation of protein stability and integrity.

Post-translation alterations of protein structure, whether through deliberate enzyme-mediated modification or spontaneous covalent damage, have the ability to influence structure and activity. Analogous to how different strategies of enzymatic modification are associated with specific types of amino acids, particular residues also have unique propensities for spontaneous alteration based on their chemical characteristics. For example, aspartates and asparagines can undergo intramolecular cyclization to form a five-member succinimide ring through the nucleophilic attack of the adjacent peptide bond amide nitrogen on the side-chain carbonyl of either residue (1–3). Succinimides are unstable and undergo rapid hydrolysis to a mixture of

[†]This work was supported by funding from the Natural Sciences and Engineering Research Council to S.N., and L.T.J.D. is a Tier 1 Research Chair in Structural Biochemistry.

[‡]Published with permission of the Director of Vaccine and Infectious Disease Organization as journal series 494.

© 2008 American Chemical Society

*To whom correspondence should be addressed: Vaccine and Infectious Disease Organization, 120 Veterinary Rd., University of Saskatchewan, Saskatoon, Saskatchewan S7N 5E3, Canada. Telephone: (306) 966-1546. Fax: (306) 966-7478. scott.napper@usask.ca.

aspartate and isoaspartate products (Figure 1A). Whether the succinimide was derived from aspartate or asparagine, the hydrolysis products are the same such that the reaction represents deamidation of asparagines and isomerization of aspartates.

Formation of isoaspartates has significant potential for disruption of protein structure and function through both the introduction of an additional methylene unit (CH₂) into the main chain and the reduction of the length of the side chain, which can influence its ability to participate in stabilizing hydrogen bonds and salt bridges. Furthermore, deamidation of asparagine changes the net charge at the reacting position which can further compromise protein stability. The physiological significance of isoaspartate formation is supported by the presence of the highly conserved enzyme protein isoaspartate methyltransferase (PIMT),¹ which methylates isoaspartates in both peptide and protein contexts (4,5). Seemingly consistent with a general repair function, PIMT is present in a broad range of prokaryotic and eukaryotic organisms (6,7) and recognizes isoaspartyl residues in most amino acid sequence contexts (8). PIMT catalyzes transfer of a methyl group from *S*-adenosylmethionine to the α -carboxyl group of an isoaspartate to stimulate re-formation of the succinimide. Subsequent hydrolysis of this ring results in a fraction of the original isoaspartates being converted to aspartates.

However, as isoaspartates remain the predominant product of succinimide hydrolysis, multiple rounds of repair are required for complete conversion of an isoaspartate to aspartate. Furthermore, this represents true repair only when the initiating residue was an aspartate as asparagines are not regenerated. The consequences of PIMT deficiency have been reported for a number of species and include fatal seizures in mice (9), poor dauer stage survival for *Caenorhabditis elegans* (10), and increased sensitivity to environmental stressors in *Escherichia coli* (11). All of these phenotypes are an apparent consequence of accumulation of isoaspartyl-containing proteins.

Succinimide formation is the rate-limiting step in the transformation of self-reactive asparagines and aspartates with half-lives of days to weeks (12). These rates are influenced by the structural and sequence context of the reacting residue and in particular the identity of the residue in the immediate C-terminal position (12). The greatest rates of cyclization are observed when the reacting residue is followed by a glycine, which likely reflects the structural flexibility imparted by this residue on the main chain. Main-chain flexibility is believed to be a key structural determinant for succinimide propensity with cyclization most frequently observed in regions of undefined secondary structure (1). Specifically, succinimide formation is favored when the main chain of the reacting residues is able to adopt a conformation of approximately -120° (1). This is a rare conformation for native structures and likely limits the self-reactivity of most asparagines and aspartates.

As a dual consequence of their slow rates of formation and chemical instability, succinimides are difficult to directly observe. As such, most strategies for quantification of these reactions focus on characterization of the isoaspartate product rather than the succinimide itself. For example, a number of techniques employing electron capture dissociation mass spectrometry have been described for detection and quantification of isoaspartates (13,14). These approaches target either diagnostic fragment ions (14) or the incorporation of H₂¹⁸O (15–17). Alternatively, and more commonly, enzymatic assays, based on the PIMT enzyme, for quantification of isoaspartate content in peptides and proteins have been established (18). In this capacity, PIMT has found application as a bioreagent for quantification of the transformation of aspartates and

¹Abbreviations: HPr, histidine-containing protein; PIMT, protein isoaspartyl methyltransferase; PTS, phosphoenolpyruvate:sugar phosphotransferase system; PS-His15Asp HPr, post-succinimide His15Asp HPr; pI, isoelectric point; rmsd, root-mean-square deviation; DSIP, Iso-Delta Sleep-Inducing Peptide; NMR, nuclear magnetic resonance; TEMED, N,N,N',N'-tetramethylethylenediamine; NCS, noncrystallographic symmetry.

asparagines through succinimide-based mechanisms, most notably as a commercially available kit (ISOQUANT Promega).

On the basis of their stability and relative ease of detection, isoaspartates are a logical biomarker for succinimide events. However, this approach is likely to underestimate the frequency of ring formation as isoaspartyl residues do not represent an exclusive or consistent product of succinimide breakdown. Isoaspartates typically account for 65–85% of the residues emerging from succinimide hydrolysis, and this ratio can vary in different protein contexts (2). The variability of the ratio of products which emerge from succinimide hydrolysis likely reflects structural limitations imposed by the protein that influence patterns of succinimide hydrolysis.

Patterns of succinimide hydrolysis are sensitive to the structural context in which they are contained as parameters that limit access of an incoming water molecule to either of the succinimide carbonyl groups influence the ratio of products formed. Hydrolysis at the side-chain succinimide carbonyl produces an aspartate, while hydrolysis at the main-chain carbonyl results in an isoaspartate (Figure 1A). It is therefore theoretically possible for protein structure to restrict isoaspartate formation by blocking access of incoming water molecules to the main-chain succinimide carbonyl group. As such, while an isoaspartate offers strong evidence of succinimide occurrence, it is presumptuous to interpret the absence of isoaspartates as an absolute indicator of the absence of succinimide formation. The demonstration of a protein which undergoes isoaspartate-free succinimide hydrolysis would offer reason to conceptually separate succinimides from isoaspartates. Our fortuitous creation of a protein in which succinimide formation is catalyzed by phosphorylation, and appears to undergo exclusive hydrolysis to an aspartate, seems to provide such an example and offers an ideal system for investigating such particulars of succinimide behavior.

The phosphoenolpyruvate:sugar phosphotransferase system (PTS) functions through a series of phosphotransfer events from enzyme I to histidine-containing protein (HPr) to a sugar-species IIA protein (19). HPr is a single-domain, nonglycosylated protein with no disulfide bonds. Depending upon the bacterial strain of origin, HPr varies in size from 6.7 to 15 kDa and adopts a highly conserved open-faced β -sandwich composed of four antiparallel β -strands and three α -helices (20). HPr undergoes phosphorylation at the N^{δ1} position of His15 (21), but this active center histidine can be mutated to aspartate, His15Asp HPr, while the ability to phosphotransfer from enzyme I and to enzyme IIA^{glucose} via a phosphoaspartyl intermediate is retained (22). Other PTS phosphotransfer proteins have shown similar potential for substitution of active site histidines with the general trend of histidines which are phosphorylated at the N^{δ2} position being functionally replaced with aspartate and histidines which are phosphorylated at the N^{ε2} position being functionally replaced with glutamate (23). This likely reflects the importance of conserved positioning of the phosphoacceptor atom.

Phosphorylation of the active site aspartate of His15Asp HPr leads to a spontaneous internal rearrangement with the formation of an unphosphorylated protein species with a pI higher than that of the original unmodified protein. This species can be isolated by ion exchange chromatography for detailed investigation. Our group has demonstrated that the high-pI species exhibits (a) changes in mass consistent with loss of a water molecule, (b) a +1 charge relative to the unmodified protein, (c) sensitivity to basic conditions, and (d) the presence of a nonstandard peptide bond at the phosphoacceptor position (22). All of these observations are consistent with succinimide formation. In the absence of phosphorylation, formation of the high-pI species is not observed, indicating that the rearrangement is phosphorylation-catalyzed. The ability of phosphorylation of an aspartate to catalyze succinimide formation was not unanticipated as covalent modification of aspartyl side chains is known to increase reactivity. For example, the β -methyl ester derivatives, as formed by the PIMT enzyme, have been shown to form succinimides 24000 times faster than the corresponding unmodified

aspartate residues (2). Modification of the carboxyl group of aspartic acid through phosphorylation would also be expected to increase the rate of ring formation by providing a better leaving group.

While succinimide theory would predict the rapid accumulation of isoaspartate residues as a consequence of dephosphorylation via a succinimide intermediate, we instead observed within PS-His15Asp HPr (a) retention of activity of the protein after multiple rounds of phosphorylation, (b) retention of a standard peptide bond at the reacting position, and (c) the fact that the post-succinimide His15Asp HPr does not function as a PIMT substrate (22). These observations are consistent with retention of aspartate at position 15 and are therefore inconsistent with typical succinimide hydrolysis.

In the original work, we proposed two alternate explanations for the apparent absence of isoaspartates following resolution of the high-pI form of His15Asp HPr. First, the absence of isoaspartates could result from the formation of a novel five-member isoimide ring in which aspartates emerge as the exclusive product of ring hydrolysis (Figure 1B). Alternatively, steric restraints imposed by the protein of succinimide hydrolysis could exclude the isoaspartyl product.

From crystallographic determinations of wild-type and His15Asp *E. coli* HPr, Athmer et al. (24) observed that the C-terminal residue, Glu85, is positioned to limit access of an incoming water molecule to the main-chain carbonyl of position 15. As the required point of hydrolysis for formation of isoaspartate, this offers a structural rationalization for the breakdown of the high-pI species of His15Asp HPr exclusively to aspartate. This was confirmed through mutational analysis in which deletion of the C-terminal residue (His15Asp Δ 85 HPr) did not influence the ability of the protein to undergo phosphorylation or formation of the high-pI species but did permit rapid accumulation of isoaspartates.

An attempt to determine the crystallographic structure of the succinimide-containing intermediate of His15Asp HPr proved unsuccessful as the succinimide was hydrolyzed during crystallization. The structure revealed an aspartate at position 15; however, due to the nonphysiological pH (pH 4.5) and high salt concentrations of the crystallization conditions, it was not possible to conclusively say whether this would also reflect the products which would form under physiological conditions (22).

The example of phosphorylation-catalyzed succinimide formation in His15Asp HPr, with subsequent hydrolysis exclusively to aspartate, is sufficiently extraordinary to warrant further structural investigation to verify the absence of isoaspartates as well as to offer structural explanation for catalyzed ring formation and directed hydrolysis to aspartate. With these objectives in mind, we present the high-resolution structure of post-succinimide His15Asp HPr (PS-His15Asp HPr). This protein was generated from hydrolysis of the isolated high-pI species at physiological pH and salt concentrations. The unambiguous electron density and absence of alternate positions of the active site main-chain atoms verify the exclusive production of aspartate residues following succinimide hydrolysis. These findings are further confirmed through N-terminal sequencing and PIMT assays. This definitively confirms the ability of the protein structure to influence, and even eliminate, isoaspartate formation during succinimide hydrolysis. From this high-resolution structure, we offer structural rationalizations for both catalyzed succinimide formation and exclusive hydrolysis to aspartate through limitation of attack at the main-chain succinimide carbonyl group.

These findings are biologically significant, in general, as they offer strong evidence to support the hypothesis that succinimide formation may serve as an effective mechanism for removal of modifying groups from aspartates. Furthermore, that succinimide formation can occur in a manner independent of isoaspartate formation suggests the estimated frequency of succinimide

events is likely underestimated both *in vivo* and *in vitro*. This is significant both in evaluation of protein and peptide integrity and in assigning potential biological roles to these rearrangements.

MATERIALS AND METHODS

Protein Expression and Purification

His15Asp HPr was overexpressed in *E. coli* strain ESK108, F' trp thi rpsL ptsH465 recA56, using the puc19(ptsH) plasmid under the control of the ptsH promoter (25). Homogeneous protein was produced as previously described (25). Enzyme I was purified by methods previously described (26).

Phosphorylation of His15Asp HPr

His15Asp HPr (3 mg) was phosphorylated at 37 °C for 60 min in 10 mM potassium phosphate buffer (pH 7.0) with 5 mM phosphoenolpyruvate, 5 mM MgCl₂, and 0.3 mg of enzyme I.

Isolation of the Succinimide-Containing His15Asp HPr Intermediate

The phosphorylation reaction mixture was separated by anion exchange chromatography using a Mini-Q-Sepharose column and an Amersham Pharmacia Biotech Gradifrac system at 4 °C. The reaction mixture was loaded with 10 mM citrate-phosphate buffer (pH 4.6) and the column eluted at 2 mL/min with a 20 mL gradient to 0.07 M NaCl in the same buffer. Fractions (0.5 mL) were collected, and protein elution was monitored at 214 nm. Isoelectric focusing confirmed the ability of the ion exchange protocol to separate the three forms of the protein, phosphorylated (lower pI), unphosphorylated, and cyclized (higher pI). The isolated high-pI species was dialyzed overnight against phosphate-buffered saline (pH 7.4). Following dialysis, this form of the protein migrated the same as unmodified His15Asp HPr, indicating hydrolysis of the succinimide. This species was used for structural investigations.

Isoelectric Focusing

Isoelectric focusing gels were constructed using a Bio-Rad capillary tray. Gels were created by dissolving 1.1 g of sucrose in 10.3 mL of 5% acrylamide with subsequent addition of ampholytes (600 µL of 20% pH 3.0–10.0, 40% pH 3.5–5.0, and 40% pH 5.0–7.0). Gels were set by the addition of 50 µL of 5% ammonium persulfate, 25 µL of riboflavin 5-phosphate (2.5 mg/mL), and 5.0 µL of TEMED. The mixture was poured under the glass plate and allowed to set under a light source for 30 min. Polymerized gels were removed from the tray, and electrode wicks soaked in 1 M NaOH (cathode) and 1 M phosphoric acid (anode) were positioned at either side of the gel. The gel was placed, glass side down, on a precooled (4 °C) flat bed Pharmacia IEF apparatus, and samples (20 µL) were loaded and run for 2 h at 8 W with a 1000 V upper limit.

Crystallization and Data Collection

PS-His15Asp HPr was crystallized by the hanging drop vapor diffusion method at 14 °C. Crystals formed in 0.1 M citrate-phosphate buffer (pH 4.4) and 20–25% saturated ammonium sulfate. X-ray diffraction data were collected using a flash-cooled crystal on the 14-BMC beamline at BioCARS, Advanced Photon Source (Argonne National Laboratory, Chicago, IL). The total oscillation range was 180°, using radiation with a wavelength of 0.9984 Å. The crystal–detector distance was 75 mm. The crystal diffracted to 1.0 Å resolution with an overall completeness of 71%. The data were indexed and scaled with DENZO and SCALEPACK (27). The crystal space group was identified as *P*2₁ on the basis of symmetry and systematically absent reflections.

Solution and Refinement of the Crystallographic Structure

The structure of the His15Asp mutant in space group $P2_1$ was determined by molecular replacement using PHASER (28) with the Ser46Asp mutant of HPr (PDB entry 1OPD) as the starting model. The structure was refined using Refmac5 in the CCP4 package. Between refinements, the model was visually adjusted and water molecules were added using the molecular graphics program COOT (29). Non-crystallographic symmetry (NCS) restraints were invoked between molecules A and B in the asymmetric unit during the initial stages of refinement, and the NCS restraints were relaxed in the later stages of refinement. All molecular figures were generated with Pymol (30). This structure has been submitted to the RCSB Protein Data Bank as RCSB entry rcsb046605 and PDB entry 3CCD.

Molecular Modeling

The phosphorylated Asp15 and the succinimide ring at position 15 were initially modeled visually, and the modeled protein(s) was subjected to energy minimization using MacroModel (31). The Merck molecular force field was used with water as the solvent medium, and the minimization was carried out by the molecular dynamics simulated annealing method.

Methyltransferase Assays

Methyltransferase assays were performed using the ISOQUANT isoaspartate detection kit (Promega) according to the manufacturer's specifications for the radioactive protocol. For each experiment, 50 pmol of the Iso-Delta Sleep-Inducing Peptide (DSIP) reference standard and either 200 pmol of His15Asp HPr and PS-His15Asp HPr or 100 pmol of wild-type HPr were analyzed. Percentage isoaspartate contents were calculated as (sample counts per minute – blank counts per minute) $\times$ (pico-moles of reference standard)/(reference standard counts per minute – blank counts per minute) $\times$ (picomoles of protein). PIMT assays were performed on both the intact proteins and the proteolytic fragments generated by treating the samples with a 1:50 (w/w) ratio of trypsin to substrate for 4 h at 30 °C. Digestions were halted with the addition of phenylmethanesulfonyl fluoride to a final concentration of 100 mM.

N-Terminal Sequencing

Protein sequencing was performed using an Applied Biosystems Inc. model 471A sequencer equipped with a model MG5 microgradient pump and a blot cartridge for polyvinylidene difluoride-type membranes, and data were analyzed using an Applied Biosystems Inc. 601A data system. Sequencing was carried out by Dr. S. MacKenzie (Plant Biotechnology Institute, National Research Council of Canada, Saskatoon, SK).

RESULTS

Phosphorylation of His15Asp Generates Three Protein Species

Incubation of His15Asp HPr under phosphorylation conditions generates three species of the protein, one with a pI higher than that of the unmodified protein and the other with a pI lower than that of the unmodified protein (Figure 2). On the basis of previous experiments, we have demonstrated that the lower-pI species corresponds to the phosphorylated form of the protein (22).

Crystallographic Structure

PS-His15Asp HPr crystallizes in space group $P2_1$ with two molecules in the asymmetric unit and the following unit cell dimensions: $a = 33.65 \text{ \AA}$, $b = 42.30 \text{ \AA}$, $c = 48.30 \text{ \AA}$, and $\beta = 89.95^\circ$. The two molecules (A and B) in the asymmetric unit are identical, the root-mean-square deviation (rmsd) between the two molecules being $0.04 \text{ \AA}$ (Figure 3A). All further discussions refer to molecule A only. Notably, this is distinct from the unit cell adopted by the

unmodified His15Asp HPr which, under identical crystallization conditions, crystallized in space group $P2_1$ with one molecule in the asymmetric unit and the following unit cell dimensions: $a = 25.37 \text{ \AA}$, $b = 45.34 \text{ \AA}$, $c = 27.62 \text{ \AA}$, and $\beta = 104.0^\circ$.

The tertiary structure of PS-His15Asp HPr was determined as described in Materials and Methods. Data collection and refinement parameters are listed in Table 1. Due to its small size and relative ease of crystallization, HPr has been extensively characterized structurally and employed as a model for various structure–activity investigations. The structure we report here is the highest-resolution determination performed to date. The overall structure of PS-His15Asp HPr is consistent with previous crystallographic investigations of *E. coli* HPr, including the wild type (32), His15Asp HPr (22), and Ser46Asp HPr (33). Overall, the structure forms an open-faced β -sandwich composed of four antiparallel β -strands and three helices. The four strands form a single, pleated β -sheet on one face of the protein, and the three α -helices lie on one side of the sheet. The active center is situated in a relatively unstructured turn between β -strand 1 and α -helix 1.

Overall Conformation

The overall structure of PS-His15Asp HPr is in close agreement with our previous structural determination of the unmodified form of His15Asp HPr. As a relative measure of structural deviations between the structures, it should be noted that the rmsd between the two molecules of PS-His15Asp HPr in the asymmetric unit was only $0.04 \text{ \AA}$ (Figure 3A) whereas comparison of the main-chain atoms in the pre- and post-succinimide structures reveals a rmsd of $0.7 \text{ \AA}$ (Figure 3B). In particular, there are three noted regions of structural deviation between the pre and post-succinimide forms of His15Asp HPr: the minor helix B, the C-terminal residue Glu85, and the active center.

Helix B is a small, irregular helix composed of residues 46–53, where residues 46 and 47 conform to an α -helical conformation, residue 48 is a transition point, and residues 49–53 are in a 3_{10} -helical conformation. Apart from the active site, the α -carbon atoms of these residues have the greatest variability in HPr structural determinations and the flexibility of this region is believed to be important to the physiological functions of HPr. A highly conserved Ser46 N-caps this helix and is phosphorylated in Gram-positive bacteria to regulate HPr activity (34). While HPrs from Gram-negative bacteria are not modified at this site, the functional consequences can be mimicked with a Ser46Asp mutation that results in activation of the protein (33). One of the most significant structural alterations of phosphorylation of Ser46 or its mutation to aspartate is the promotion of formation of a more idealized α -helix as a result of the stabilizing interaction of these negatively charged groups with the helix dipole (33).

In the post-succinimide His15Asp structure, a more idealized α -helix is observed in terms of the hydrogen bonds that are formed and their geometry throughout the helix. Furthermore, in the post-succinimide structure, the main-chain carbonyl group of Phe48 is able to form a hydrogen bond with the main-chain amide of Gln51, whereas in the pre-succinimide, the carbonyl group of Phe48 hydrogen bonds with water. The stabilization of this helix in the post-succinimide structure is believed to be a consequence of a sulfate anion from the crystallization solution that is in the proximity of Ser46 which causes the differential positioning within helix B of the pre- and post-succinimide forms of His15Asp HPr and illustrates the sensitivity of this region of the protein to electrostatic influence (Figure 4A).

The second region of structural deviation is at the C-terminus. The positioning and involvement of the C-terminus in the active site of HPr have been a point of disagreement between structural and functional investigations. While the involvement of the C-terminus in the active center is supported by NMR investigations of HPr (35) as well as most, but not all (36), crystallographic investigations, mutation or deletion of the C-terminal residues does not have a significant

impact on HPr phosphotransfer function (25). It does, however, have a significant impact on the production of isoaspartyl residues during phosphorylation-catalyzed succinimide formation and hydrolysis in His15Asp HPr. We have previously demonstrated that this residue functions as a steric restraint to limit the attack of a water molecule on the main-chain carbonyl at position 15 and thereby prevent isoaspartate formation (24). In spite of the difference in the positions of the α -carbons of the C-terminal region, the α -carboxyl is similarly positioned in each of the structures to limit access to the main-chain carbonyl of residue 15 (Figure 4B). The similar positioning of this region in two different crystallographic investigations of His15Asp HPr, which crystallized in the same space group, but with different unit cell dimensions, to minimize the potential influence of packing interactions, further supports the proposed role of this region of the protein in preventing isoaspartyl formation.

The final region of structural deviation between the pre- and post-succinimide structures of His15Asp HPr is in the stretch of amino acids linking β -strand 1 to α -helix 1, which corresponds to the active site region. While the positioning of this region is generally well conserved in the various crystallographic determinations of *E. coli* HPr, even for different space groups, NMR investigations indicate the potential for alternate positioning of this region (35,37). The shift corresponds to a general repositioning of the main chain to bring it closer to the core of the protein as well as a tightening of the cleft that contains the active site region. The cusps of this cleft are defined by Pro11 and Pro18; in the pre-succinimide structure, the distance between the α -carbons of these residues is 11.85 Å, whereas in the post-succinimide structure, this closes to 9.97 Å (Figure 4C).

Structure of the Active Site

Changes in conformation at the active site region are obviously of interest as they may indicate the presence of isoaspartyl residues. In spite of the moderate repositioning within the active center region, the electron density of the main-chain atoms is very well defined with no indication of alternate placements, in particular those that would be required for the inclusion of an additional methylene group as necessitated by an isoaspartate (Figure 5). Both the side-chain and main-chain atoms of residues 15 and 16 were maximally refined at 100% occupancy with mean *B*-factors of 8.01 and 8.46 Å², respectively, for each molecule in the asymmetric unit. The average mean *B*-factor value of 8.23 Å² for this region is lower than the average mean *B*-factor for the protein of 9.71 Å². Collectively, this high-resolution structure shows no evidence for the presence of an isoaspartyl group in post-succinimide His15Asp HPr.

N-Terminal Sequencing

N-Terminal sequencing depends on the presence of standard peptide bonds and is unable to proceed through succinimide or isoaspartyl residues. The modification of His15Asp HPr occurs sufficiently close to the N-terminus to permit N-terminal sequence analysis to provide a qualitative measure of nonstandard residues at the phosphoacceptor site. For His15Asp HPr, the anticipated sequence was obtained through the active center; however, for high-pI His15Asp HPr, the sequencing reaction failed after the fourteenth round, indicating the presence of a nonstandard peptide bond. This is consistent with the presence of a succinimide. Hydrolysis of this species to generate PS-His15Asp HPr, under the same conditions used the protein for crystallographic investigations, re-established the ability to sequence past the reacting residue. This is supportive of the presence of an aspartate rather than an isoaspartate (Figure 6A).

Protein Isoaspartate Methyltransferase Assays

PIMT assays were performed to provide quantitative analysis of isoaspartate content in the wild type, His15Asp HPr, and PS-His15Asp HPr. For these three proteins, an identical baseline content of isoaspartyl residues is observed, with fewer than five isoaspartates per 100 protein

molecules (Figure 6B). A low frequency of isoaspartate generation is anticipated for HPr as the protein has been shown to deamidate at Asn12 and Asn38 (38). That wild-type HPr, which has a histidine at the phosphoacceptor position which is unable to undergo succinimide or isoaspartate formation, has isoaspartate content identical to that of His15Asp and PS-His15Asp HPr indicates these isoaspartates are independent of the phosphoacceptor site.

On the basis of the hypothesis that structural restraints of the active site may impede isoaspartate formation, similar steric interference may limit the ability for PIMT to recognize isoaspartates in this region. However, proteolytic digestion of these proteins did not increase the number of isoaspartates detected, indicating that protein structure is not interfering with its ability to function as a PIMT substrate (Figure 6B).

Conserved Positioning of the Phosphoacceptor Site

The ability of the aspartate to act as a functional substitution for the His15 N δ 1 phosphorylation event is rationalized on the basis of the conserved geometric positioning of phosphoacceptor atoms of aspartate and histidine. Consistent with this hypothesis, one of the side-chain carboxyl group oxygens of His15Asp occupies essentially the same position as the N δ 1 atom of the imidazole ring of the wild-type structure for both the pre- and post-succinimide forms of His15Asp (Figure 7). The electron density of position 15 clearly corresponds to that of an aspartate group with all of the non-hydrogen atoms of the side chain clearly defined. There is no additional electron density for which we cannot account near this position to suggest the presence of an isoaspartyl side chain.

The Conformation of the Active Site Is Consistent with Succinimide Formation

The formation of a succinimide within a protein is generally limited by the unique conformations required for side-chain reactivity with the main chain. Specifically, succinimide formation is favored when the main-chain ψ angles and side-chain χ_1 dihedral angles are approximately -120° and 60° , respectively (1). For the PS-His15Asp HPr, these angles are -169° and 60° , respectively. The amide nitrogen of residue 16 is positioned only 3.5 Å from the carboxyl oxygen of the phosphoacceptor Asp15. This positioning is predictive of succinimide formation and offers a partial structural rationalization of the ability of the protein to undergo rapid succinimide formation. However, that the high-pI intermediate is not observed in the absence of phosphorylation indicates that the favorable conformation, in the absence phosphorylation, is insufficient to initiate succinimide formation.

Phosphorylation-Catalyzed Succinimide Formation

The ability of phosphorylation to catalyze succinimide formation in His15Asp HPr likely occurs as a dual function of the phosphoryl group providing a better leaving group as well as the ability of phosphorylation to induce conformations which favor succinimide formation. The NMR structure of phosphorylated wild-type HPr demonstrates that the phosphoryl group forms hydrogen bonds with the main-chain amides of residues 16 and 17 (37). On the basis of the conserved positioning of the carboxyl oxygens of Asp15 with the phosphoryl-accepting site, it is reasonable to assume similar interactions occur with the phosphoacyl group of phosphorylated His15Asp. On the basis of conservation of conformation in phosphorylated His15Asp, the ψ and χ_1 angles are -169° and 60° , respectively, which are highly conducive to succinimide formation. Furthermore, the side-chain carbonyl group is brought within 2.4 Å of the amide nitrogen of residue 16 which is again highly favorable for succinimide formation (Figure 8A).

In addition to offering a more favorable conformation of the reactive side chain for succinimide formation, the anticipated interaction between the phosphoacyl group and the main chain is also expected to initiate a concerted chemical reaction in which extraction of protons from the

main-chain amides by the phosphoryl group renders the phosphoryl group an even better leaving group, further increasing rates of succinimide formation (Figure 8B).

DISCUSSION

Traditionally, protein succinimides are regarded as a form of nonspecific damage affecting asparagines and aspartates which are vulnerable on the basis of their sequence and structural contexts. This vantage point is supported by numerous examples of proteins which have been shown to undergo spontaneous and physiological formation of isoaspartates, including calbindin (39), calmodulin (40), somatotropin (41), and collagen (42), as well as the detrimental effect these nonstandard residues have on protein function. The presence of an apparent repair enzyme, PIMT, which is widely expressed in most organisms supports the physiological occurrence and importance of isoaspartyl formation through succinimide intermediates.

It has alternatively been suggested that succinimide formation may serve as a molecular, biological clock (43–45). This could serve to indicate protein age for regulating longevity, not unlike PEST sequences (46) or the N-end rule (47), or could serve to trigger biological events by providing alternative protein isoforms of distinct functions. From this perspective, the deliberate placement of aspartates or asparagines within susceptible regions of the polypeptide would cause predictable rates of isoaspartate formation to predetermine protein lifespan or initiate changes in protein function. Importantly, however, the considerations of succinimide formation as a form of protein damage, or as a molecular clock, are both influenced by the dogma that succinimide formation and isoaspartate production are inseparably linked. Our demonstration of isoaspartate-free succinimide cycling expands and complicates the potential physiological roles for succinimide formation as well as their detection for protein quality assurance purposes.

First, the example of His15Asp HPr demonstrates that succinimide formation represents an effective strategy for removal of phosphoryl groups from modified aspartic acid residues and could represent a mechanism of autophosphatase activity for the numerous proteins which are regulated through phosphorylation of aspartyl residues. Specifically, signal transduction in microorganisms and plants frequently occurs through two-component systems which involve His-Asp phosphorelay networks of sensory histidine protein kinases and response regulator proteins (48,49). The phosphorylated response regulators have intrinsic autophosphatase activities to limit the duration of the phosphorylated state. These autophosphatase activities vary by 4 orders of magnitude for different response regulators depending on their physiological function (50,51).

The mechanism of response regulator autophosphatase activity has generally been assumed to proceed as a reversal of the phosphohistidine reaction with water replacing the imidazole side chain. Alternately, through succinimide formation, the phosphoryl group could be displaced by the intramolecular nucleophilic attack at the β -carboxyl group. There are two notable examples to support this model. First, a mass spectrometric investigation of the phosphorylation of OmpR, the response regulator that mediates osmotic responses in *E. coli*, relieved a third species, in addition to the unphosphorylated and phosphorylated OmpR proteins, with a mass approximately 18 Da lower than that of the unmodified protein (52). This dehydrated form of the protein was dependent upon phosphorylation, so it presumably arose as a consequence of dephosphorylation. An identical dehydrated species was observed as a consequence of phosphorylation of the chemotaxis response regulator CheY (53), although succinimide formation does not appear to be a principle source of autophosphatase activity for this particular response regulator (54).

Phosphorylated aspartate residues are also observed outside of the response regulators, specifically, within the haloacid dehydrogenase superfamily, which includes the P-type ATPases and the dehalogenases. For most dehalogenase enzymes, the reaction mechanism involves a covalent intermediate formed with an active site Asp residue. This intermediate is hydrolyzed via attack on the Asp as demonstrated by ^{18}O incorporation experiments (55,56) and is consistent with succinimide formation during hydrolysis of the covalent intermediate.

In addition to a potential role in serving as a source of autophosphatase activity of phosphoaspartyl residues, the demonstration of succinimide hydrolysis exclusively to an aspartate also is significant for the consideration of these modifications as a measure of protein damage. Experimentally, it is easier to quantify isoaspartates than succinimides in proteins. As a result, and as a consequence of the assumption that succinimide formation is inseparable from isoaspartate formation, strategies for quantification of these reactions are based on the occurrence of isoaspartate. Our demonstration of isoaspartate-independent succinimide hydrolysis indicates these approaches likely underestimate the frequency of this modification. This could be significant in the evaluation of the structural integrity of peptides and proteins. This is particularly important as it has been shown that deamidation can significantly influence the quality of vaccines (57,58) and other peptide and protein pharmaceuticals (59,60).

ACKNOWLEDGMENT

We thank Robert Henning for assisting with the data collection on the BioCARS beamline at the Advanced Photon Source. Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Basic Energy Sciences, Office of Science, under Contract W-31-109-Eng-38. Use of the BioCARS Sector 14 was supported by the National Institutes of Health, National Center for Research Resources, under Grant RR07707.

REFERENCES

1. Clarke S. Propensity for spontaneous succinimide formation from aspartyl and asparaginyl residues in cellular proteins. *Int. J. Pept. Protein Res* 1987;30:808–821. [PubMed: 3440704]
2. Clarke, S.; Stephenson, RC.; Lowenson, JD. Stability of Protein Pharmaceuticals: Chemical and Physical Pathways of Protein Degradation. Ahern, TJ.; Manning, MC., editors. New York: Plenum Press; 1992. p. 1-29.
3. Wright HT. *CRC Crit. Rev. Biochem. Mol. Biol* 1991;27:1–52. [PubMed: 1678690]
4. McFadden PN, Clarke S. Methylation at d-aspartyl residues in erythrocytes: Possible step in the repair of aged membrane proteins. *Proc. Natl. Acad. Sci. U.S.A* 1982;79:2460–2664. [PubMed: 6123997]
5. McFadden PN, Clarke S. Conversion of isoaspartyl peptides to normal peptides: Implications for the cellular repair of damaged proteins. *Proc. Natl. Acad. Sci. U.S.A* 1987;84:2595–2599. [PubMed: 3472227]
6. O'Connor CM. Regulation and Sub-cellular Distribution of a Protein Methyltransferase and its Damaged Aspartyl Substrate Sites in Developing *Xenopus* Oocytes. *J. Biol. Chem* 1987;261:10398–10403.
7. Li C, Clarke S. A Protein Methyltransferase Specific for Altered Aspartyl Residues is Important in *Escherichia coli* Stationary-Phase Survival and Heat-shock Resistance. *Proc. Natl. Acad. Sci. U.S.A* 1992;89:9885–9889. [PubMed: 1409717]
8. Aswad DW, Johnson BA. The Unusual Substrate Specificity of Eukaryotic Protein Carboxyl Methyltransferase. *Trends Biochem. Sci* 1987;12:155–158.
9. Kim E, Lowenson JD, MacLaren DC, Clarke S, Young SG. Deficiency of a Protein-repair Enzyme Results in Accumulation of Altered Proteins, Retardation of Growth and Fatal Seizures in Mice. *Proc. Natl. Acad. Sci. U.S.A* 1997;94:6132–6137. [PubMed: 9177182]
10. Niewmierzycka A, Clarke S. Do damaged proteins accumulate in *Caenorhabditis elegans* isoaspartate methyltransferase (pcm-1) deletion mutants? *Arch. Biochem. Biophys* 1999;364:209–218. [PubMed: 10190976]

11. Visick JE, Cai H, Clarke S. The l-isoaspartyl Protein Repair Methyltransferase Enhances Survival of Aging *Escherichia coli* Subjected to Secondary Environmental Stresses. *J. Bacteriol* 1998;180:2623–2629. [PubMed: 9573145]
12. Brennan TV, Clarke S. Effect of Adjacent Histidine and Cysteine Residues on the Spontaneous Degradation of Asparaginyl- and Aspartyl-containing Peptides. *Int. J. Pept. Protein Res* 1995;45:547–553. [PubMed: 7558585]
13. Cournoyer JJ, Lin C, Bowman MJ, O'Connor PB. Quantifying the relative abundance of isoaspartyl residues in deamidated proteins by electron capture dissociation. *J. Am. Soc. Mass Spectrom* 2007;18:48–56. [PubMed: 16997569]
14. Cournoyer JJ, Lin C, O'Connor PB. Detecting deamidation products in proteins by electron capture dissociation. *Anal. Chem* 2006;78:1264–1271. [PubMed: 16478121]
15. Xiao G, Bondarenko PV, Jacob J, Chu GC, Chelius D. ¹⁸O Labeling Method for Identification of Succinimides in Proteins. *Anal. Chem* 2007;79:2714–2721. [PubMed: 17313184]
16. Cournoyer JJ, Pittman JL, Ivleva VB, Fallows E, Waskell L, Costello C, O'Connor PB. Deamidation: Differentiation of aspartate from isoaspartate products in peptides by electron capture dissociation. *Protein Sci* 2007;14:452–463. [PubMed: 15659375]
17. Xiaojuan L, Cournoyer J, Lin C, O'Connor PB. Use of ¹⁸O Labels to Monitor Deamidation during Protein and Peptide Sample Processing. *J. Am. Soc. Mass Spectrom* 2008;19:855–864. [PubMed: 18394920]
18. Johnson BA, Aswad DW. Optimal conditions for the use of protein l-isoaspartyl methyltransferase in assessing the isoaspartate content of peptides and proteins. *Anal. Biochem* 1991;192:384–391. [PubMed: 1827964]
19. Meadow ND, Fox DK, Roseman S. The bacterial phosphoenolpyruvate: Glycose phosphotransferase system. *Annu. Rev. Biochem* 1990;59:497–542. [PubMed: 2197982]
20. Jia Z, Quail JW, Delbaere LTJ, Waygood EBW. Structural Comparison of the Histidine-Containing Phosphocarrier Protein HPr. *Biochem. Cell Biol* 1994;72:202–217. [PubMed: 7840940]
21. Waygood EB, Erickson EE, El-Kabbani OAL, Delbaere LTJ. Characterization of phosphorylated histidine-containing protein (HPr) of the bacterial phosphoenolpyruvate:sugar phosphotransferase system. *Biochemistry* 1985;24:6938–6945. [PubMed: 3907706]
22. Napper S, Delbaere LTJ, Waygood EB. The Aspartate Replacement of the Active Site Histidine in Histidine-containing, HPr, of the *Escherichia coli* Phosphoenolpyruvate: Sugar Phosphotransferase System Can Accept and Donate a Phosphoryl Group. *J. Biol. Chem* 1999;274:21776–21782. [PubMed: 10419492]
23. Napper S, Brokx SJ, Pally E, Kindrachuk J, Delbaere LTJ, Waygood EB. Substitution of Aspartate and Glutamate for Active Center Histidines in the *Escherichia coli* Phosphoenolpyruvate: Sugar Phosphotransferase System Maintain Phosphotransfer Potential. *J. Biol. Chem* 2001;276:41588–41593. [PubMed: 11551914]
24. Athmer L, Kindrachuk K, Georges F, Napper S. The Influence of Protein Structure on the Products Emerging from Succinimide Hydrolysis. *J. Biol. Chem* 2002;277:30502–30507. [PubMed: 12068021]
25. Anderson JW, Bhanot P, Georges F, Klevit RE, Waygood EB. Involvement of the carboxy-terminal in the active site of the histidine-containing protein, HPr, of the phosphoenolpyruvate:sugar phosphotransferase system of *Escherichia coli*. *Biochemistry* 1991;30:9601–9607. [PubMed: 1680393]
26. Anderson JW, Pullen K, Georges F, Klevit RE, Waygood EB. The involvement of the arginine 17 residue in the active site of the histidine-containing protein, HPr, of the phosphoenolpyruvate:sugar phosphotransferase system of *Escherichia coli*. *J. Biol. Chem* 1993;268:12325–12333. [PubMed: 8509371]
27. Otwinowski, Z.; Minor, W. Processing of X-Ray Diffraction Data Collected in Oscillation Mode. In: Carter, CW., Jr; Sweet, RM., editors. *Methods in Enzymology; Macromolecular Crystallography Part A*. San Diego: Academic Press; 1997. p. 307–325.
28. Read RJ. Pushing the boundaries of molecular replacement with maximum likelihood. *Acta Crystallogr* 2001;D57:1373–1382.

29. Emsley P, Cowtan K. COOT: Model-building tools for molecular graphics. *Acta Crystallogr* 2004;D60:2126–2132.
30. DeLano, WL. The PyMOL Molecular Graphics System. 2002. <http://www.pymol.org>
31. Mohamadi F, Richards N, Guida W, Liskamp R, Lipton M, Caufield C, Chang G, Hendrickson T, Still W. MacroModel: An Integrated Software System for Modeling Organic and Bioorganic Molecules Using Molecular Mechanics. *J. Comput. Chem* 1990;11:440.
32. Jia Z, Quail JW, Waygood EB, Delbaere LTJ. The 2.0 Å Resolution Structure of *Escherichia coli* Histidine-containing Phosphocarrier Protein HPr. A Redetermination. *J. Biol. Chem* 1993;268:22490–22501. [PubMed: 8226757]
33. Napper S, Anderson W, Georges F, Quail JW, Delbaere LTJ, Waygood EB. Mutation of Serine 46 to Aspartate in the Histidine-Containing Protein of *Escherichia coli* Mimics the Inactivation by Phosphorylation of Serine-46 in HPrs from Gram-Positive Bacteria. *Biochemistry* 1996;35:11260–11267. [PubMed: 8784179]
34. Reizer J, Romano AH, Deutscher J. The role of phosphorylation of HPr, a phosphocarrier protein of the phosphotransferase system, in the regulation of carbon metabolism in Gram-positive bacteria. *J. Cell. Biochem* 1993;51:19–24. [PubMed: 8432739]
35. Van Nuland NA, Hangyi IW, van Schaik RC, Berendsen HJ, van Gunsteren WF, Scheek RM, Robillard GT. The high-resolution structure of the histidine-containing phosphocarrier protein HPr from *Escherichia coli* determined by restrained molecular dynamics from nuclear magnetic resonance nuclear Overhauser effect data. *J. Mol. Biol* 1994;237:544–559. [PubMed: 8158637]
36. Prasad L, Waygood EB, Lee JS, Delbaere LT. The 2.5 Å resolution structure of the jel42 Fab fragment/HPr complex. *J. Mol. Biol* 1998;280:829–845. [PubMed: 9671553]
37. van Nuland NA, Boelens R, Scheek RM, Robillard GT. High-resolution structure of the phosphorylated form of the histidine-containing protein HPr from *Escherichia coli* determined by restrained molecular dynamics from NMR-NOE data. *J. Mol. Biol* 1995;246:180–193. [PubMed: 7853396]
38. Sharma S, Hammen PK, Anderson JW, Leung A, Georges F, Hengstenberg W, Klevit RE, Waygood EB. Deamidation of HPr, a phosphocarrier protein of the phosphoenolpyruvate: sugar phosphotransferase system, involves asparagine 38 (HPr-1) and asparagine 12 (HPr-2) in isoaspartyl acid formation. *J. Biol. Chem* 1993;268:17695–17704. [PubMed: 8349654]
39. Chazin WJ, Kordel J, Thulin E, Hofmann T, Drakenberg T, Forsen S. Identification of an Isoaspartyl Linkage Formed Upon Deamidation of Bovine Calbindin D9K and Structural Characterization by 2D ¹H NMR. *Biochemistry* 1989;28:8646–8653. [PubMed: 2605213]
40. Ota IM, Clarke S. Enzymatic Methylation of l-Isoaspartyl Residue Derived from Aspartyl Residue in Affinity-purified Calmodulin. *J. Biol. Chem* 1989;264:54–60. [PubMed: 2642479]
41. Violand BN, Schlittler MR, Kolodziej EW, Toren PC, Cabonce MA, Siegel NR, Duffin KL, Zobel JF, Smith CE, Tou JS. Isolation and Characterization of Procine Somatropin Containing a Succinimide in Place of Aspartate. *Protein Sci* 1992;1:1634–1641. [PubMed: 1363933]
42. Lanthier J, Desrosiers RR. Protein l-isoaspartyl methyltransferase repairs abnormal aspartyl residues accumulated in vivo in type-I collagen and restores cell migration. *Exp. Cell Res* 2004;293:96–105. [PubMed: 14729060]
43. Reissner KJ, Aswad DW. Deamidation and isoaspartate formation in proteins: Unwanted alterations or surreptitious signals? *Cell. Mol. Life Sci* 2003;60:1281–1295. [PubMed: 12943218]
44. McKerrow JH, Robinson AB. Primary Sequence Dependence of the Deamidation of Rabbit Muscle Aldolase. *Science* 1973;183:85. [PubMed: 4808790]
45. Robinson NE, Robinson AB. Molecular Clocks. *Proc. Natl. Acad. Sci. U.S.A* 2001;98:944–949. [PubMed: 11158575]
46. Rogers S, Wells R, Rechsteiner M. Amino acid sequences common to rapidly degraded proteins: The PEST hypothesis. *Science* 1986;234:364–368. [PubMed: 2876518]
47. Bachmair A, Finley D, Varshavsky A. *In vivo* half-life of a protein is a function of its amino-terminal residue. *Science* 1986;234:179–186. [PubMed: 3018930]
48. Urao T, Yamaguchi-Shinozaki K, Shinozaki K. Two-component systems in plant signal transduction. *Trends Plant Sci* 2000;5:67–74. [PubMed: 10664616]

49. Stock JB, Stock AM, Mottonen JM. Signal transduction in bacteria. *Nature* 1990;344:395–400. [PubMed: 2157156]
50. Hess JF, Oosawa K, Kaplan N, Simon MI. Phosphorylation of three proteins in the signaling pathway of bacterial chemotaxis. *Cell* 1988;53:79–87. [PubMed: 3280143]
51. Zapf J, Madhusudan M, Grimshaw CE, Hoch JA, Varughese KI, Whiteley JM. A source of response regulator autophosphatase activity: The critical role of a residue adjacent to the SpoOF autophosphorylation active site. *Biochemistry* 1998;37:7725–7732. [PubMed: 9601032]
52. Kenney LJ, Bauer MD, Silhavy TJ. Phosphorylation-dependent conformational changes in OmpR, an osmo-regulatory DNA-binding protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A* 1995;92:8866–8870. [PubMed: 7568033]
53. Napper S, Wolanin PM, Webre DJ, Kindrachuk J, Waygood B, Stock JB. Intramolecular rearrangements as a consequence of the dephosphorylation of phosphoaspartate residues in proteins. *FEBS Lett* 2003;538:77–80. [PubMed: 12633856]
54. Wolanin PM, Webre DJ, Stock JB. Mechanism of phosphatase activity in the chemotaxis response regulator CheY. *Biochemistry* 2003;42:14075–14082. [PubMed: 14636076]
55. Liu JQ, Kurihara T, Ichiyama S, Miyagi M, Tsunasawa S, Kawasaki H, Soda K, Esaki N. Reaction mechanism of fluoroacetate dehalogenase from *Moraxella* sp. *B. J. Biol. Chem* 1998;273:30897–30902. [PubMed: 9812982]
56. Liu JQ, Kurihara T, Miyagi M, Esaki N, Soda K. Reaction mechanism of l-2-haloacid dehalogenase of *Pseudomonas* sp. YL. Identification of Asp10 as the active site nucleophile by ¹⁸O incorporation experiments. *J. Biol. Chem* 1995;270:18309–18312. [PubMed: 7629151]
57. Ribot WJ, Powell BS, Ivins BE, Little SF, Johnson WM, Hoover TA, Norris SL, Adamovicz JJ, Friedlander AM, Andrews GP. Comparative vaccine efficacy of different isoforms of recombinant protective antigen against *Bacillus anthracis* spore challenge in rabbits. *Vaccine* 2006;24:3469–3476. [PubMed: 16519970]
58. Chen W, Ede NJ, Jackson DC, McCluskey J, Purcell AW. CTL recognition of an altered peptide associated with asparagine bond rearrangement. Implications for immunity and vaccine design. *J. Immunol* 1996;157:1000–1005. [PubMed: 8757603]
59. Jars MU, Hvass A, Waaben D. Insulin aspart (AspB28 human insulin) derivatives formed in pharmaceutical solutions. *Pharm. Res* 2002;19:621–628. [PubMed: 12069164]
60. Meyer JD, Ho B, Manning MC. Effects of conformation on the chemical stability of pharmaceutically relevant polypeptides. *Pharm. Biotechnol* 2002;13:85–107. [PubMed: 11987755]

FIGURE 1.

Mechanism of succinimide and isoimide formation and hydrolysis. (A) Succinimide formation and hydrolysis. Mechanism of succinimide formation and hydrolysis from an asparagine and a phosphoaspartate. A similar mechanism occurs from aspartate, with the distinction being release of a water molecule rather than a phosphoryl group upon ring formation. Also shown are the products which emerge from succinimide hydrolysis with aspartates being produced from hydrolysis at the side-chain carbonyl and isoaspartates from hydrolysis at the main-chain carbonyl. (B) Isoimide formation and hydrolysis. Hydrolysis of an isoimide ring results in aspartyl products independent of the point of attack of the incoming water molecule.

FIGURE 2.

Phosphorylation of His15Asp generates three species of the protein. His15Asp HPr (3 mg) was phosphorylated at 37 °C for 60 min in 10 mM potassium phosphate buffer (pH 7.0) with 5 mM phosphoenolpyruvate, 5 mM MgCl₂, and 0.3 mg of enzyme I. Protein species were separated by isoelectric focusing: lane 1, unphosphorylated His15Asp HPr; lane 2, His15Asp HPr under phosphorylating conditions with the generation of both the phosphorylated and high-pI species; lane 3, isolated high-pI His15Asp HPr; and lane 4, isolated high-pI His15Asp HPr dialyzed overnight against PBS.

FIGURE 3.

Regions of structural deviation between His15Asp HPr and PS-His15Asp HPr. Root-mean-square deviations were calculated for the α -carbons using the LSQKAB program from CCP4: (A) rmsds of the α -carbons of the two molecules within the asymmetric unit of PS-His15Asp HPr and (B) rmsds of the α -carbons of pre- and post-succinimide PS-His15Asp HPr.

FIGURE 4.

Structural deviation between His15Asp and PS-His15Asp HPr. These structures were created with Pymol (30), and in each case, molecules with green carbon-carbon bonds represent the PS-His15Asp HPr whereas molecules with blue carbon-carbon bonds represent His15Asp HPr. Structural comparisons are presented for the following. (A) Minor helix B. Formation of a more idealized helix in His15Asp HPr is believed to be a consequence of a sulfate anion from the crystallization solution. (B) C-Terminus. The similar positioning of this region in two different crystallographic investigations of His15Asp HPr supports the proposed role in preventing isoaspartyl formation. (C) Active center. This involves a general repositioning of

the main chain to bring it closer to the core of the protein as well as a tightening of the cleft that contains the active site region.

FIGURE 5.

Stereographic representation of electron density of the active site region. Electron density of the active site region of post-succinimide His15Asp HPr at 1 Å resolution. The clear electron density for Asp15 as well as the absence of additional electron density that is unaccounted for in the active center region argues against the possibility of an isoaspartyl residue at position 15.

A)

	Residue						
	10	11	12	13	14	15	16
His15Asp	A	P	N	G	L	D	T
High pI His15Asp	A	P	N	G	L	NS	NS
PS-His15Asp	A	P	N	G	L	D	T

No detectable signal (NS).

FIGURE 6.

Amino acid analysis and methyltransferase assays for detection of isoaspartyls in PS-His15Asp HPr. (A) Amino acid sequencing. The inability for N-terminal sequencing to proceed past nonstandard peptide bonds, such as succinimides or isoaspartyls, provides a qualitative assessment for the presence of these residues. The inability to sequence past the 14th residue of high-pI His15Asp HPr indicates the presence of a succinimide, while detection of correct sequences in positions 15 and 16 of PS-His15Asp HPr indicates breakdown to an aspartyl product. Not shown are the first 10 residues of each protein. (B) Methyltransferase assays were performed using the ISOQUANT isoaspartate kit (Promega) according to the manufacturer's instructions. Isoaspartate content is expressed as a percentage of modified residues per 100

molecules of protein. Isoaspartate contents were determined for the wild type, His15Asp HPr, and PS-His15Asp HPr for both intact and proteolytic fragments, to eliminate any influence of protein structure on the ability to act as a PIMT substrate. Each data point represents separate PIMT assay.

FIGURE 7.

Stereographic comparison of the phosphoacceptor site of the wild type, His15Asp HPr, and PS-His15Asp HPr. This figure was created with Pymol (30). Molecules with green carbon–carbon bonds represent the post-succinimide structure, molecules with blue carbon–carbon bonds the pre-succinimide structure, and molecules with magenta carbon–carbon bonds wild-type HPr. For both the preand post-succinimide His15Asp structures, the carboxyl group oxygen occupies a position similar to that of the N^{δ1} atom of the phosphoacceptor His15 residue of wild-type HPr.

FIGURE 8.

Structural mechanism for succinimide-based autophosphatase activity. (A) Structural isomers of phosphorylation and succinimide-based dephosphorylation of His15Asp are modeled on the basis of the crystallographic structure. (B) General mechanism for catalyzed succinimide formation from a phosphoaspartate with subsequent hydrolysis to aspartate.

Table 1
Data Collection and Refinement Statistics of Post-Succinimide His15Asp HPr

Data Collection Statistics			
unit cell dimensions			
<i>a</i> (Å)	33.65	resolution range (Å)	48.3-1.0 (1.06-1.0)
<i>b</i> (Å)	42.30	no. of reflections measured	331691
<i>c</i> (Å)	48.30	no. of unique reflections	73053 (944)
<i>β</i> (deg)	89.95	<i>R</i> _{sym} ^a	0.021 (0.15)
no. of molecules in asymmetric unit	2	completeness (%)	71 (23)
space group	<i>P</i> 2 ₁	redundancy	2.6
		mean <i>I</i> / <i>σ</i> (<i>I</i>)	16.0 (4.0)
Refinement Statistics ^b			
resolution limits (Å)			
<i>R</i> _{free}	26.3-1.0 (1.03-1.0)	rmsd from ideal geometry	
<i>R</i> _{work}	0.20 (0.26)	bond distances (Å)	0.01
<i>R</i> _{cryst}	0.19 (0.26)	bond angles (deg)	1.2
total no. of reflections	0.19	dihedral angles (deg)	5.9
no. of non-H atoms	57397	improper angles (deg)	0.08
no. of water molecules	1318	Ramachandran statistics	
mean <i>B</i> factor for protein atoms (Å ²)	198	most favored (%)	90.7
	9.71	favored (%)	9.3
		additional allowed (%)	0
		disallowed (%)	0

^a $R_{\text{sym}} = \sum (I_{\text{hkl}} - \langle I_{\text{hkl}} \rangle) / \langle I_{\text{hkl}} \rangle$, where $\langle I_{\text{hkl}} \rangle$ is the average intensity over symmetry-related reflections and I_{hkl} is the observed intensity.
^b $R_{\text{value}} = |F_o| - |F_c| / |F_o|$, where F_o and F_c are the observed and calculated structure factors, respectively. For R_{free} , the sum is done on the test set reflections (5% of the total reflections). For R_{work} , the sum is done on the remaining reflections. For R_{cryst} , the sum is done on all reflections included in the resolution range.