Published in final edited form as:

Tetrahedron Lett. 2009 August 12; 50(32): 4571–4574. doi:10.1016/j.tetlet.2009.05.067.

Euryjanicin A: a new cycloheptapeptide from the Caribbean marine sponge *Prosuberites laughlini*

Jan Vicente, Brunilda Vera, Abimael D. Rodríguez*, Idaliz Rodríguez-Escudero, and Raphael G. Raptis

Department of Chemistry, University of Puerto Rico, P.O. Box 23346, UPR Station, San Juan, Puerto Rico 00931-3346

Abstract

A new proline-containing cycloheptapeptide, euryjanicin A (1), has been isolated from the marine sponge *Prosuberites laughlini* indigenous to Puerto Rico, and its structure established by an X-ray crystal structure determination. The absolute configuration of each amino acid residue was determined by Marfey's method.

Keywords

Prosuberites laughlini; Cycloheptapeptide; Caribbean marine sponge; *Eurypon laughlini*; X-ray crystallography; Peptides; Macrocycles

Marine sponges, especially those found in tropical ocean areas, continue to represent the single most prolific source of structurally novel natural products of marine origin.¹ Sponge-derived secondary metabolites have also attracted considerable attention because of their relevant and diverse pharmacological actions.² During the last 25 years a large body of evidence has been accumulated to indicate that certain marine sponges (especially sponges belonging to the class Demospongiae) also contain amino acid derivatives, usually as trace components, with significant cancer cell growth inhibitory properties.³ Examples of such biologically active peptides are the axinellins,⁴ kapakahines,⁵ microsclerodermins,⁶ polytheonamides,⁷ papuamides,⁸ stylisins,⁹ hymenamides,¹⁰ wainunuamide,¹¹ and dominicin.¹² Additional noteworthy examples include the laxaphycins¹³ and discodermins¹⁴ (they inhibit tumor promotion), calyculin¹⁵ (tumor promoting) and the anti-thrombin cyclotheonamides.¹⁶

As part of our ongoing campaign to find new natural products with potential anticancer activity from marine organisms found near Puerto Rico, a small library of sponge extracts from this location was recently screened for anti-cancer activity. Considerable activity was detected in the crude extract of *Prosuberites laughlini* (Díaz, Alvarez & Van Soest, 1987) (Order Hadromerida, Family Suberitidae).¹⁷ The sponge was collected in April 2006 at a depth of about 15 m by scuba from *El Natural* reef, off the municipality of Aguadilla, in the northwestern coast of Puerto Rico (18°27'46.90"N, 67°10'06.11"W). A freeze dried sample (623 g) was repeatedly extracted with MeOH (16 L), and the extracts were combined, concentrated in vacuo (83 g), and partitioned between H₂O (1.5 L) and EtOAc (3 × 1 L), followed by *n*-butanol (3 ×

*Corresponding author. Tel.: +1 787 764 0000 Ext. 4799; fax: +1 787 756 8242; e-mail: abrodriguez@uprrp.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 L). The EtOAc and *n*-butanol extracts were concentrated under vacuum to give 6.1 g and 14.8 g of residue, respectively. Significant anti-cancer activity was detected in the EtOAc fraction; upon treatment with 1.0 mg/mL extract, reduction of cell viability was 15, 10, and 19% for DU145 prostate cancer, A2058 melanoma, and MDA-MB-435s breast cancer types, respectively. The EtOAc extract was subjected to normal-phase silica gel (200 g) column chromatography (CC) using a mixture of CHCl₃ and MeOH which had been previously saturated with dry NH₃ (80/20). Similar fractions were pooled together on the basis of TLC analysis. Subsequent evaluation of the fraction eluting first (2.8 g) by ¹H and ¹³C NMR indicated that this pooled fraction contained a mixture of small peptides. Further fractionation was performed by normal-phase silica gel (100 g) CC with a mixture of hexane and EtOAc (95/5). Close inspection by NMR indicated that sub-fractions 10 (183 g) and 11 (156 mg) appeared to contain most of the peptides observed in the initial NMR analysis. Purification of sub-fraction 10 was achieved by C18 silica gel (5 g) reversed-phase CC using a mixture of MeOH and H₂O (85/15) as eluent; this yielded the new compound euryjanicin A (**1**, 44 mg, 0.05%) as well as the known cyclic octapeptide dominicin (**2**, 33 mg, 0.04%).¹² Also isolated was a new cycloheptapeptide, which is currently still under investigation and will be reported separately.

Euryjanicin A (**1**),¹⁸ was obtained as optically active, $[\alpha]_D^{20} -18.0^\circ$ (*c* 1.0, CHCl₃), colorless crystals that showed a pseudomolecular ion peak at *m/z* 827 in the ESIMS spectrum. The molecular formula of **1** was determined to be C₄₄H₅₈N₈O₈ by the HRESIMS [*m/z* 827.4439 (M+H)⁺, C₄₄H₅₉N₈O₈, Δ -1.7 mDa], requiring 20 sites of unsaturation. The intense IR absorptions at 3316 and 1644 cm⁻¹ were attributed to amino and amide carbonyl groups, respectively, and the UV absorptions at λ_{max} 281 (ε 1800) and 290 (ε 1600) nm indicated the presence of aromatic ring(s). The ¹H NMR spectrum of **1** (Figure 1A) was indicative of a peptide. Standard amino acid analysis of the acid hydrolysate of **1** revealed the presence of one mole each of valine (Val), phenylalanine (Phe), tryptophan (Trp), serine (Ser), and isoleucine (Ile), and two moles of proline (Pro) residues, which subsequently, were all confirmed to be of L-configuration by Marfey's derivatization and subsequent HPLC analysis with standards.^{19,20} Euryjanicin A (**1**) was negative to a ninhydrin test, which together with its solubility in organic solvents and the 20 degrees of unsaturation, indicated a cyclic structure.

The crystals of **1** gave a sharp, well-resolved peak when analyzed by reversed-phase HPLC using a variety of solvent systems, in addition to a sharp melting point range (mp 174–176 °C) and a single, clean molecular ion in the HRESI. Despite these indications of purity, the

initial structural approaches to euryjanicin A (**1**) employing high field 2D-NMR techniques were complicated by the existence of a mixture of slowly interconverting conformers in varying proportions depending on the solvent used (e.g., CDCl₃, DMSO-*d*₆, acetone-*d*₆, MeOH-*d*₄, pyridine-*d*₅, MeCN-*d*₃, benzene-*d*₆). Attempts to simplify the ¹H NMR spectra by cooling (to -15 °C) and heating (to 60 °C), or adjusting the pH by addition of TFA or Et₃N, did not alleviate the problem, and typically resulted in further loss of resolution which precluded structural studies. Likewise, many of the resonance lines in the ¹³C NMR spectra were doubled, tripled, or broadened when the spectra were recorded in the same solvents (Figure 1B). Because the 1D and 2D NMR spectra of euryjanicin A in these organic solvents were complex and difficult to interpret, only a handful of NMR spectral assignments could be made with complete confidence.²¹ Therefore, efforts to determine the amino acid sequence by 2D NMR experiments were abandoned.

Crystals suitable for X-ray crystallographic analysis were obtained by slow recrystallization from MeOH(NH₃). The results obtained from this analysis (Figure 2) defined the relative stereochemistry at all centers, and the absolute configuration shown was selected to agree with the known amino acid centers.²² The geometry of the peptidic linkages about the proline residues was also assigned on the basis of the X-ray diffraction method, which showed that the two proline peptide bonds were *cis*. The *cis* peptide bonds and bifurcated transannular hydrogen bonds between the C44 carbonyl oxygen (O8) and N4-H (2.50 Å) and N5-H (2.43 Å) set the sharp turn conformation of the macrocyclic ring. An additional hydrogen bond (2.18 Å) was pinpointed between primary hydroxyl oxygen O4 of the Ser⁴ residue and N6-H. This turn results in the side chains of the only two aromatic amino acids, L-Trp¹ and L-Phe⁵, being projected in the same direction. The absolute structure **1** of euryjanicin A was thereby completed and thus confirmed as *cyclo*(Trp¹-Pro²-Ile³-Ser⁴-Phe⁵-Val⁶-Pro⁷). The fact that other investigators have isolated similar cyclic peptides from extracts of cultured marine bacteria and fungi, suggests a possible microbial or symbiotic origin for metabolites euryjanicin A (**1**) and dominicin (**2**).²³

Unlike the EtOAc extract, strong anti-tubercular activity was detected in the *n*-butanol partition fraction; thus a small portion of the crude extract (2 g) was subjected to C18 silica gel (50 g) reversed-phase CC using a mixture of MeOH and H₂O (85/15). Similar fractions were combined on the basis of TLC and ¹H NMR analyses. The first (most polar) fraction to elute out (0.90 g) was further chromatographed over normal-phase silica gel (100 g, dry packing), initially with a mixture of CHCl₃ and MeOH(NH₃) (90/10), and later using gradients of increasing polarity. This procedure yielded the known compounds hymenidin (**3**, 66 mg, 0.08%) and monobromoisophakellin (**4**, 57 mg, 0.07%).^{24,25}

Pure euryjanicin A (**1**) and dominicin (**2**) were evaluated for their inhibitory activity against cell viability in melanoma (A2058), breast cancer (MDA-MB-435s), and prostate cancer (DU145). Surprisingly, none of them displayed significant inhibitory activity.⁹ On the other hand, in the National Cancer Institute's one-dose (10^{-5} M) 60-cell-line assay, euryjanicin A (**1**) inhibited a non-small cell lung and renal cancer cell line: NCI-H522 displayed growth at 72% of the mean and UO-31 at 73% of the mean. Dominicin (**2**), in turn, inhibited a leukemia (SR) and a renal (UO-31) cancer cell line, displaying growths at 80% and 32% of the mean, respectively. Preliminary tests for anti-tuberculosis activity of alkaloids **3** and **4** demonstrated that these compounds significantly inhibited the growth of *Mycobacterium tuberculosis* H₃₇Rv (MIC values = 6.1 and 64.0 μ g/mL, respectively). On the other hand, alkaloids **3** and **4** lacked significant anti-plasmodial activity as their IC₅₀s against the malaria parasite *Plasmodium falciparum* (chloroquine-resistant W2 clone) were ≥ 10 μ g/mL.

Acknowledgments

We thank Dr. Ileana I. Rodríguez for helping out during the collection of *P. laughlini*, Dr. Hong Zhao for assistance with the X-ray data, and the UPR-MARC and UPR-RISE Fellowship Programs for financial support to JV and BV, respectively. Sponge extracts were screened for anti-cancer activity by Dr. Sangkil Nam at the City of Hope Beckman Research Institute and for anti-TB activity by Dr. Yuehong Wang at the Institute for Tuberculosis Research of the University of Illinois. Anti-cancer, anti-mycobacterial, and anti-malarial bioassays of the pure natural products were conducted at the National Cancer Institute (NCI), the Institute for Tuberculosis Research of the UIC, and the Instituto de Investigaciones Avanzadas y Servicios de Alta Tecnología (Panama), respectively. Major financial support was provided by the NIH-MBRS SCORE program (Grant S06GM08102) of the University of Puerto Rico.

References

1. Blunt JW, Copp BR, Hu WP, Munro MHG, Northcote PT, Prinsep MR. Nat Prod Rep 2009;26:170–244. and previous articles in this series. [PubMed: 19177222]
2. Keyzers RA, Davies-Coleman MT. Chem Soc Rev 2005;34:355–365. [PubMed: 15778769]
3. Fusetani N, Matsunaga S. Chem Rev 1993;93:1793–1806.
4. Randazzo A, Dal Piaz F, Orru S, Debitus C, Roussakis C, Pucci P, Gomez-Paloma L. Eur J Org Chem 1998;2659–2665.
5. (a) Nakao Y, Yeung BKS, Yoshida WY, Scheuer PJ. J Am Chem Soc 1995;117:8271–8272. (b) Yeung BKS, Nakao Y, Kinnel RB, Carney JR, Yoshida WY, Scheuer PJ, Kelly-Borges M. J Org Chem 1996;61:7168–7173. [PubMed: 11667621]
6. Qureshi A, Colin PL, Faulkner DJ. Tetrahedron 2000;56:3679–3685.
7. Hamada T, Matsunaga S, Yano G, Fusetani N. J Am Chem Soc 2005;127:110–118. [PubMed: 15631460]
8. Ford PW, Gustafson KR, McKee TC, Shigematsu N, Maurizi LK, Pannell LK, Williams DE, de Silva ED, Lassota P, Allen TM, Van Soest R, Andersen RJ, Boyd MR. J Am Chem Soc 1999;121:5899–5909.
9. Mohammed R, Peng J, Kelly M, Hamann MT. J Nat Prod 2006;69:1739–1744. [PubMed: 17190452]
10. (a) Kobayashi J, Tsuda M, Nakamura T, Mikami Y, Shigemori H. Tetrahedron 1993;49:2391–2402. (b) Tsuda M, Shigemori H, Mikami Y, Kobayashi J. Tetrahedron 1993;49:6785–6796. (c) Tsuda M, Sasaki T, Kobayashi J. Tetrahedron 1994;50:4667–4680.
11. Tabudravu J, Morris LA, van den Bosch JJK, Jaspars M. Tetrahedron Lett 2001;42:9273–9276.
12. Williams DE, Patrick BO, Behrisch HW, Van Soest R, Roberge M, Andersen RJ. J Nat Prod 2005;68:327–330. [PubMed: 15787430]
13. Bonnard I, Rolland M, Salmon JM, Debiton E, Barthomeuf C, Banaigs B. J Med Chem 2007;50:1266–1279. [PubMed: 17323939]
14. (a) Matsunaga S, Fusetani N, Konosu S. J Nat Prod 1985;48:236–241. [PubMed: 3839260] (b) Matsunaga S, Fusetani N, Konosu S. Tetrahedron Lett 1984;25:5165–5168. (c) Matsunaga S, Fusetani N, Konosu S. Tetrahedron Lett 1985;26:855–856.
15. Kato Y, Fusetani N, Matsunaga S, Hashimoto K, Koseki K. J Org Chem 1988;53:3930–3932.

16. Fusetani N, Matsunaga S, Matsumoto H, Takebayashi Y. *J Am Chem Soc* 1990;112:7053–7054.
17. This sponge was originally described as *Eurypon laughlini*.
18. Euryjanicin A (**1**): colorless crystals, mp 174–176 °C; $[\alpha]_D^{20}$ –18.0° (*c* 1.0, CHCl₃); $\nu_{\max}$ (thin film) 3316, 2961, 2928, 1644, 1531, 1454, 1348, 750, 702 cm^{–1}; UV (MeOH) $\lambda_{\max}$ 211 (Σ 12500), $\lambda_{\max}$ 274 (Σ 1700), $\lambda_{\max}$ 281 (Σ 1800), $\lambda_{\max}$ 290 (Σ 1600) nm; ¹H (CDCl₃, 500 MHz) and ¹³C NMR (CDCl₃, 125 MHz) spectra, see Figure 1; HRESIMS *m/z* 827.4439 [M+H]⁺ (calcd for C₄₄H₅₉N₈O₈, 827.4456).
19. Pure euryjanicin A (0.5 mg) was hydrolyzed in 6N HCl (100 μ L) at 110 °C for 12 h. The cooled reaction mixture was evaporated to dryness, and traces of HCl were removed from the residual hydrolysate by repeated evaporation from H₂O (3 $\times$ 0.5 mL) using N₂ gas.
20. Marfey P. *Carlsberg Res Commun* 1984;49:591–596.
21. The NMR spectra of dominicin (**2**) recorded in a variety of solvents also resulted in poor resolution. Acceptable NMR spectra, however, could be obtained using pyridine-*d*₅ as the solvent (see reference 12). Slowly interconverting conformations have been observed in other cyclic peptides such as schizotrin A and pahayokolides; see: (a) Pergament I, Carmeli S. *Tetrahedron Lett* 1994;35:8473–8476. (b) An T, Krishnaswamy T, Kumar S, Wang M, Liu L, Lay JO, Liyanage R, Berry J, Gantar M, Marks V, Gawley RE, Rein KS. *J Nat Prod* 2007;70:730–735. [PubMed: 17432902]
22. Single-crystal X-ray diffraction measurements were made on a Bruker SMART 1 K CCD diffractometer with graphite-monochromated Mo K α radiation (λ = 0.71073 Å). Crystals were mounted on a glass fiber. Crystal data for **1** at 293(2) K: C₄₄H₅₈N₈O₈, *M_r* = 826.98, orthorhombic, space group P2₁2₁2₁, *a* = 13.7230(13), *b* = 17.275(2), *c* = 19.026(3) Å, β = 90°, *V* = 4510.3(9) Å³, *Z* = 4, ρ_{calc} = 1.218 Mg/m^{–3}, *F*₀₀₀ = 1768, μ = 0.085 mm^{–1}. Data collection and reduction: crystal size, 0.39 $\times$ 0.38 $\times$ 0.16 mm, θ range = 1.59–28.28°, 32,630 reflections collected, 10,396 independent reflections (*R*_{int} = 0.0304), final *R* indices [*I* > 2 σ (*I*): *R*₁ = 0.0580, *wR*₂ = 0.1418 for 541 variable parameters, GOF = 1.034. The crystallographic data for euryjanicin A (**1**) have been deposited at CCDC under the registry number 721666. They can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK; fax: + 44 1223 336033; e-mail: deposit@ccdc.cam.ac.uk].
23. (a) Tan LT, Cheng XC, Jensen PR, Fenical W. *J Org Chem* 2003;68:8767–8773. [PubMed: 14604342]
(b) Renner MK, Shen YC, Cheng XC, Jensen PR, Frankmoelle W, Kauffman CA, Fenical W, Labbovsky E, Clardy J. *J Am Chem Soc* 1999;121:11273–11276.
24. Kobayashi J, Ohizumi Y, Nakamura H, Hirata Y. *Experientia* 1986;42:1176–1177. [PubMed: 3770140]
25. Assmann M, Köck M. *Z Naturforsch* 2002;57:153–156.

Figure 1. ¹H (500 MHz) (A) and ¹³C (125 MHz) (B) NMR spectra of euryjanicin A (1) in CDCl₃ at 25 °C. The doubling and tripling of resonances observed in the NMR spectra is attributed to the existence of slow conformational equilibria.

Figure 2.

Perspective drawing of the X-ray crystal structure of euryjanicin A (**1**). Hydrogen atoms have been omitted for clarity and selected atoms are labeled. Carbon atoms are indicated in black, nitrogens in blue, and oxygens in red.