Published in final edited form as:

Inflamm Bowel Dis. 2008 December ; 14(12): 1715–1735. doi:10.1002/ibd.20501.

Leukocyte Adhesion Molecules in Animal Models of Inflammatory Bowel Disease

Jesus Rivera-Nieves^{*,2}, Gezahegn Gorfu[‡], and Klaus Ley[‡]

^{*} Mucosal Inflammation Program, Division of Gastroenterology, Department of Internal Medicine, University of Colorado Health Sciences Center, Denver Colorado 80206

[‡] La Jolla Institute for Allergy and Immunology, Division of Inflammation Biology, La Jolla, CA 92037, USA

Introduction

Recent advances in our understanding of the fundamental mechanisms of seemingly unrelated diseases, i.e., rheumatoid arthritis (RA), Type 1 diabetes mellitus, psoriasis, multiple sclerosis, and inflammatory bowel disease (IBD), have demonstrated that they share similar pathogenetic mechanisms. These diseases are immune-mediated chronic inflammatory conditions, characterized by inappropriate and sustained recruitment of inflammatory cells into affected tissues, resulting in chronic tissue damage and loss of function^{1–3}.

The fate of the inflammatory cells differs according to cell type. Neutrophils migrate to sites of inflammation and undergo apoptosis. Some monocytes remain in inflamed tissues for a few days and others become permanent residents. By contrast, lymphocytes not only migrate but recirculate from the blood into tissues and through the lymphatics back to lymphatic tissues and blood, constantly patrolling for foreign antigens. Once they encounter an antigen in the context of an antigen presenting cell, naïve T cells proliferate and acquire effector functions, along with a repertoire of surface molecules (i.e., adhesion molecules, cytokine and chemokine receptors) that allow them to recognize counter-receptors/ligands expressed in specific vascular beds. Using these surface molecules, lymphocytes are able to recirculate thousands of times back to areas with a similar microenvironment to where they first encountered their cognate antigen^{4, 5}. These unique capabilities (i.e., memory acquisition and recirculation) are essential for the perpetuation of chronic inflammatory processes, including IBD. The molecules involved in the recirculation of lymphocytes have therefore attracted a great deal of interest regarding their potential as therapeutic targets. Some of these have crossed from the bench into the clinical arena, being currently in clinical use^{6–8}.

In IBD, the inflammatory process is characterized by heavy leukocytic infiltration of the intestinal lamina propria (LP), leading to fibrosis and loss of function^{9–11}. Lymphocytes that produce cytokines such as IL-12, IFN- γ , tumor necrosis factor- α (TNF- α), IL-23 and IL-17¹² all play an important role in chronic intestinal inflammation^{9–13}.

The success of the anti-TNF- α strategy in IBD¹⁴ has led to the systematic study of anti-inflammatory cytokines and the development of antibody-based strategies to modulate the overall cytokine balance^{15, 16}. Unfortunately, the therapeutic efficacy of some of these newer

2Address correspondence and reprint requests to: Dr. Jesus Rivera-Nieves, Mucosal Inflammation Program, Division of Gastroenterology, Department of Internal Medicine, University of Colorado Health Sciences Center, Biochemistry Research Building Room 742A, 4200 E. 9th Ave SE, B146, Denver, CO 80206, e-mail address: jesus.rivera-nieves@uchsc.edu.

cytokine-targeted therapies (e.g. IL-10, IL-11 blockade) has been limited^{15, 16}. A neutralizing antibody against the IL-12 p40 subunit, shared by IL-12 and IL-23, has shown promise¹⁷. Alternative therapies that target other pathways of the chronic inflammatory process may be directed at interfering with lymphocyte recirculation to the intestine by targeting specific adhesion molecules, their ligands, chemokines or their receptors^{18–20}. Using this approach, two monoclonal antibodies against integrin α_4 and $\alpha_L\beta_2$ have been approved by the FDA for the treatment of MS, CD and psoriasis (i.e. natalizumab, efalizumab)^{6, 18}. However, many of the basic mechanisms that account for their clinical efficacy remain to be elucidated. This limited knowledge has likely contributed to the occurrence of serious adverse events in clinical practice^{21, 22}.

The Leukocyte Adhesion Cascade

Leukocytes primarily migrate from the blood into the tissues across the walls of post-capillary venules. Surface molecules on specialized venular endothelial cells play a crucial role. These adhesion molecules not only serve as mechanical anchors, but also confer tissue specificity to the recruitment process through their selective patterns of expression by vascular beds²³. Myeloid cells and lymphocytes share some of the steps in the adhesion cascade, but there are also significant differences²⁴. This review focuses primarily on lymphocyte recruitment to the intestine and how this process has been targeted for therapeutic purposes in animal models of colitis and ileitis that mimic aspects of either UC or CD respectively.

Leukocyte recruitment

Several major classes of leukocyte adhesion molecules are involved in leukocyte recruitment, including the selectins and their glycoprotein ligands, integrins and immunoglobulin-superfamily molecules. They are all type I transmembrane glycoproteins that span the cell membrane only once. The structural and functional aspects of these adhesion molecules have been extensively discussed elsewhere^{25,26–28}.

The process of leukocyte recruitment to a site of inflammation encompasses the engagement and efficient arrest of leukocytes onto the vascular endothelium and their subsequent transmigration^{4, 23, 29}. This sequence is composed of several major steps, capture, rolling, activation and firm adhesion (Figure 1).

Capture

Capture is defined as the formation of the first molecular bond or tether between the circulating leukocyte and the vascular endothelium. Close proximity between the two is required. Capture is distinguishable from stable rolling and is mediated by L- and P-selectins. In inflamed venules *in vivo*, leukocyte attachment also involves other mechanisms, such as secondary capture through leukocyte-leukocyte interaction^{30, 31}. Most neutrophils start rolling as they exit capillaries³², whereas nai-ve lymphocytes capture in high endothelial venules (HEV) of peripheral lymph nodes (PLN) and other lymphoid tissues⁵. Little is known about capture of effector/memory lymphocytes, although selectins and CD44 may be involved^{33, 34}.

Rolling

If leukocyte capture is followed by the formation of new molecular bonds before the initial molecular bonds dissociate, a stable rolling movement is established^{35, 36}. This is a flow-driven downstream movement of the cell, during which it is in continuous contact with the vessel wall. Nai-ve lymphocyte rolling in HEV is mainly mediated by lymphocyte L-selectin binding to sulfated carbohydrate-containing ligands expressed on endothelial cells. Rolling has at least two distinct consequences for the cell: (i) it facilitates stable leukocyte arrest (firm adhesion); and (ii) it drastically reduces leukocyte velocity (to between 1 and 100 $\mu\text{m/s}$),

increasing the duration of exposure to the endothelial surface and to chemokines and other activating signals present there. Neutrophils arrest after gradually slowing down³⁷, whereas naïve lymphocytes arrest immediately upon encountering chemokines³⁸. Effector T cells can arrest through the chemokine receptor CXCR3 *in vitro*, but the molecules involved in arrest *in vivo* remain unknown³⁹.

Firm adhesion

Activation of the rolling lymphocyte can be triggered by the binding of a chemokine to a heptahelical transmembrane receptor on the leukocyte surface and can result in firm adhesion^{23, 40}. In flow chamber systems, this process is exceptionally rapid^{41, 42}. Prior studies, focused on lymphocyte function-associated antigen (LFA)-1, have shown that firm adhesion requires binding of integrin receptors in their active conformation to their endothelial ligands⁴³. Both $\alpha_4\beta_1$ (VLA-4) and $\alpha_4\beta_7$ integrins have been shown to mediate firm adhesion to their respective ligands Vascular Cell Adhesion Molecule (VCAM)-1 and Mucosal Addressin Cell Adhesion Molecule (MAdCAM)-1^{44, 45}.

Families of adhesion molecules involved in leukocyte recruitment

The selectins

The selectins are a family of transmembrane mammalian lectins expressed on the surface of leukocytes (L-selectin), endothelial cells (P-, E-selectins), and platelets (P-selectin)²⁷. The selectins contain an N-terminal extracellular domain with structural homology to calcium-dependent lectins, followed by a domain homologous to epidermal growth factor; and two to nine consensus repeats (CR), similar to sequences found in complement regulatory proteins (Figure 2). Each selectin inserts into the cell membrane via a hydrophobic transmembrane domain and possesses a short cytoplasmic tail. Each of the three selectins can independently mediate leukocyte rolling, given the appropriate conditions. The three selectins share approximately 50% sequence homology among the extracellular portions of the molecule, but there is no significant homology in the transmembrane and cytoplasmic domains^{27, 46}.

All three selectins are involved in the trafficking of granulocytes and lymphocytes to sites of inflammation. However, the role of each selectin differs in the physiological sequence of events, and the importance of each selectin varies in different models of inflammation. Importantly, L-selectin is essential for the homing of lymphocytes to peripheral lymph nodes⁴⁷, and plays an accessory role in lymphocyte homing to Peyer's patches (PP), a component of the gut-associated lymphatic tissue⁴⁷. High endothelial venules (HEV) in these tissues constitutively express functional ligands for L-selectin on their surface. The importance of L-selectin in lymphocyte homing is emphasized by the decreased size of lymph nodes in L-selectin-deficient mice⁴⁷ and a delay in lymphocyte homing to Peyer's patches⁴⁷. In PP, L-selectin synergizes with β_7 integrins, since absence of both L-selectin and integrins completely abrogates lymphocyte homing to these structures⁴⁸.

T cell activation results in proteolytic shedding of L-selectin⁴⁹. This pattern of surface molecule expression (CD44^{high}, L-selectin^{low}) is characteristic of effector memory T cells which recirculate to effector sites, while limiting their traffic through secondary lymphoid organs⁵. However, subpopulations of memory cells (i.e. central memory, T_{CM})⁵⁰ retain or re-express L-selectin to recirculate through inductive sites and potentially to tertiary lymphoid organs present at sites of chronic inflammation, where functional ligands for L-selectin are expressed.

P-selectin

P-selectin (CD62P) contains nine consensus repeats (CR) and extends approximately 40 nm from the endothelial surface (Figure 2). P-selectin is constitutively synthesized by megakaryocytes and endothelial cells, but is not expressed on the surface of resting platelets or endothelial cells. Rather, it is stored in secretory vesicles called Weibel-Palade bodies in endothelial cells, and α -granules in platelets^{51, 52}.

These granules are triggered to fuse with the plasma membrane by activators such as histamine, thrombin, complement C5a, calcium ionophores, and adenosine diphosphate (ADP), resulting in expression of P-selectin on the cell surface⁵³. Maximal levels of P-selectin are detected within minutes at the cell surface, followed by rapid internalization and degradation or recycling⁵¹. In the mouse, activation of endothelial cells by cytokines prolongs the surface expression of P-selectin^{54–56}. Yet in humans, expression of P-selectin is not induced by cytokine stimulation, possibly a result of structural differences in the P-selectin promoter⁵⁷.

The transient interactions between P-selectin and P-selectin Glycoprotein Ligand (PSGL)-1 allow leukocytes to roll along the venular endothelium. P- and E-selectins tend to have overlapping functions in TNF- α -stimulated venules. Indeed, in mice deficient for P-selectin, it is necessary to block E-selectin function to significantly reduce rolling, and in E-selectin knockouts, an antibody against P-selectin must be introduced to reduce rolling. Correspondingly, no leukocyte rolling is observed in E-selectin/P-selectin double deficient mice treated with TNF- α ⁵⁸. Nonetheless, although P- and E-selectins seem to have redundant functions, observations of rolling flux fractions and rolling velocities indicate that under physiological conditions, P-selectin is responsible for early rolling, while E-selectin allows slow rolling and adhesion^{37, 59}. In mice deficient for P-selectin, rolling is absent initially but returns after 1–2 hours⁶⁰. Endothelial cell expression of P-selectin is elicited within minutes by inflammatory mediators such as histamine, thrombin, or phorbol esters. The expression is short-lived, reaching its peak after only ten minutes. In mice, additional synthesis of P-selectin is brought about within two hours by cytokines such as IL-1 or TNF- α ^{54–56}.

E-selectin

E-selectin is expressed by endothelial cells after activation by IL-1 or TNF- α ^{61, 62}, but unlike P-selectin it is not expressed by platelets. On the surface of cultured endothelial cells, it is only expressed transiently for a period of 2 to 6 hours. Subsequently, E-selectin can either be re-internalized or shed from the endothelial surface⁶³. *In vivo*, the expression of E-selectin appears to be more prolonged in some organs⁶⁴ and is constitutive in skin microvessels^{65,66}. As discussed earlier, the role of E-selectin during rolling is partially redundant with that of P-selectin. E-selectin-deficient mice have a reduced number of firmly adherent leukocytes in response to local chemoattractants⁶⁷ or cytokine stimulation⁶⁸, thus E-selectin participates in the conversion of rolling to firm adhesion. E-selectin ligation of PSGL-1 triggers neutrophil activation through spleen tyrosine kinase (Syk)⁶⁹. E-selectin is of particular importance in skin inflammation, as it supports the recruitment of skin-specific T lymphocytes⁷⁰. The ligand (s) responsible for E-selectin-mediated rolling interactions include PSGL-1, CD44 and E-selectin ligand-1 (ESL-1) (Figure 2)^{71–73}. In addition to glycoproteins, glycolipids can also support E-selectin dependent rolling *in vitro*⁷⁴. E-selectin mediates much slower rolling than P-selectin, because it triggers partial LFA-1 activation⁶⁹. These rolling velocities range between less than 5 $\mu\text{m/s}$ ^{59, 75} and 15 $\mu\text{m/s}$ ⁶⁷. The molecular dissociation rate or off-rate of E-selectin is very similar to that of P-selectin⁷⁶. However, the velocity of E-selectin-mediated rolling is remarkably invariant with wall shear rate, presumably because E-selectin engagement triggers LFA-1 activation⁶⁹. A role of E-selectin and its ligands has been demonstrated for lymphocyte homing, particularly to skin. Cutaneous-homing effector/memory T cells express

a carbohydrate epitope (i.e. cutaneous associated lymphocyte antigen (CLA)) which decorates glycoproteins such as PSGL-1 and CD43 and is associated with E-selectin ligand activity ⁷⁷.

L-selectin

L-selectin, a 74–100 kDa molecule also known as CD62L, is constitutively expressed at the tips of microfolds or microvilli on granulocytes, monocytes, and most circulating lymphocytes. Lymphocytes preferentially express it when in the naïve state. In addition, it is expressed by central memory T cells ⁵⁰, and most natural killer cells ²⁷, as well as by many immature hematopoietic cells in the bone marrow. L-selectin expression on neutrophils correlates negatively with the age of the individual cell, suggesting that neutrophils may lose L-selectin as they mature in the bone marrow ⁷⁸ and as they age in the bloodstream ⁷⁹. Although most lymphocytes in the cortex of secondary lymphoid organs express L-selectin, the activated cells in the germinal centers do not ²⁷. This indicates that expression of L-selectin is at least transiently reduced from activated lymphocytes. L-selectin is proteolytically cleaved from the surface of leukocytes after activation by a variety of stimuli ^{80–83}. For neutrophils, activating agents that induce L-selectin shedding include inflammatory chemokines such as IL-8, complement factors such as C5a, bacterial peptides such as formyl-methionyl-leucyl-phenylalanine (fMLP), and lipid mediators such as platelet activating factor (PAF) or leukotriene B4 (LTB4) ⁸⁴. *In vitro*, lymphocytes shed L-selectin when they are stimulated with phorbol esters or by cross-linking of CD3 ⁸⁵. *In vivo*, soluble L-selectin is present in the plasma at high concentrations ⁸⁶, most of which is derived from lymphocytes ⁸⁷. L-selectin is responsible for naïve lymphocyte homing to PLN and MLN, but is also expressed by central memory T cells, which may play a role for the maintenance of IBD and other chronic inflammatory conditions ⁸⁸.

Selectin Ligands

PSGL-1—PSGL-1 is the best-characterized selectin ligand (Fig. 2) ^{89, 90}. It is a type 1 surface dimeric glycoprotein that is expressed on the microprocesses of virtually all leukocytes. PSGL-1 is also expressed in certain endothelial cells within the walls of the small intestine ⁹¹, and inflamed colon ⁹² as well as in cultured umbilical vein and microvascular endothelial cells ^{93, 94}. Certain post-translational modifications are necessary for PSGL-1 to function as a selectin ligand, including tyrosine sulfation ^{95, 96}, sialylation ⁹⁷, decoration with core 2 oligosaccharides ⁹⁸ and, perhaps most importantly, fucosylation ⁹⁶. Fucosylation is regulated by the specific, inducible fucosyl transferase (FTVII), which plays a key role in the generation of functional selectin ligands ^{98–101}. T-helper 1 (Th1)-polarized T cells preferentially express functional P- and E-selectin ligands ^{102–105}. Indeed the expression of FTVII is modulated by differentiation into Th1 and Th2 phenotypes ³³. FTVII is constitutively expressed by neutrophils, monocytes and eosinophils ^{99, 100}. The trafficking of neutrophils is severely restricted in mice that lack FTVII ¹⁰⁶, although unlike mice that lack both E- and P-selectin, these mice do not develop spontaneous disease ^{58, 107}. The importance of selectin ligands is dramatically illustrated by a rare human disease, Leukocyte Adhesion Deficiency-II (LAD-II). A fucosylation defect in these patients renders selectin ligands inactive, causing an inability of leukocytes to roll ¹⁰⁸. These patients have significant inflammatory pathology ¹⁰⁹. The binding of PSGL-1 to P-selectin and the functional implications thereof have been demonstrated *in vivo* ^{110, 111}. PSGL-1 also binds L-selectin ^{112, 113}, and appears to be responsible for most ¹¹⁴ though not all ¹¹⁵ of the interactions between flowing and already adherent leukocytes. Although all lymphocytes express PSGL-1, this molecule does not only function as a selectin ligand, but also binds the chemokines CCL19 and CCL21 ¹¹⁶.

Other selectin ligands—Other glycoproteins serve as ligands for L-selectin, including glycosylation-dependent cell adhesion molecule-1 (GlyCAM-1) ¹¹⁷, Nepmucin, Endomucin ²⁸ CD34 ¹¹⁸, and podocalyxin ¹¹⁹. Most L-selectin ligands expressed by high endothelial

venules contain sulfated determinants that are identified using the carbohydrate-binding monoclonal antibody MECA-79^{120, 121}. MECA-79 binds to a sulfation-dependent epitope and can block L-selectin binding¹²⁰. This epitope overlaps with sialyl 6-sulfo Lewis X, an L-selectin recognition determinant¹²². Both PNAd and sialyl 6-sulfo Lewis X are absent in mice deficient for two N-acetylglucosamine-6-O-sulfotransferases (GlcNAc6ST-1 and GlcNAc6ST-2) resulting in reduced lymphocyte homing and adhesion in HEV^{122, 123}. MECA-79 binding requires sulfation. Recent studies have identified the molecular structure of its epitope as an extended core 1 structure containing GlcNAc-6-SO₄ on O-linked glycans¹²⁴. The enzymes HECGlcNAc-6-sulfotransferases (HEC-6ST also known as GST, L-selectin ligand sulfotransferase, (LSST)) are responsible for post-translational modifications which are essential for L-selectin binding. HEC-6ST expression is highly restricted to HEV, and in humans and mice a sulfotransferase with restricted expression to the intestine has been identified¹²⁵. Little is known about its role in inflammation, but two recent studies have demonstrated its induction by TNF in human bronchial mucosa and lymphotoxin in an animal model of lymphoid neogenesis in the pancreas^{126, 127}.

Another ligand for L-selectin is mucosal addressin adhesion molecule-1 (MAdCAM-1), expressed on the surface of high endothelial venules of Peyer's patches and mucosal lymph nodes^{128, 129}. MAdCAM-1 isolated from mesenteric lymph nodes of young mice supports L-selectin dependent lymphocyte rolling in a flow chamber assay¹³⁰. A role for L-selectin in lymphocyte rolling in Peyer's patch high endothelial venules has also been demonstrated by intravital microscopy, but it is not clear whether all L-selectin ligand activity in this system is attributable to MAdCAM-1^{44, 48}.

For E-selectin, a 250 kDa potential ligand was precipitated from bovine $\gamma\delta$ T cells¹³¹, whereas on myeloid cells, PSGL-1, CD44 and E-selectin Ligand-1 (ESL-1), a molecule with homology to fibroblast growth factor receptor, are E-selectin ligands^{71,72}. In addition, CD43 might be a relevant ligand for E-selectin on skin-homing activated T cells^{132,133}.

Strategies to target selectins therapeutically—Several selectin inhibitors have been developed and tested in preclinical models and clinical trials. Carbohydrate-based selectin inhibitors of the sialyl Lewis^x type, which inhibit all three selectins at high concentrations, have unfavorable pharmacokinetics, low affinity and short half lives¹³⁴. Drug candidates within this family (e.g. OC229648, Efomycine and CY1503) have been tested in preclinical models of peritonitis, psoriasis and ischemia-reperfusion injury. Antibodies to selectins including antibodies that block more than one selectin have been developed and humanized. Protein Design Laboratories has tested a humanized version of an anti-L-selectin antibody (HuDREG-55), in patients with trauma (www.pdl.com) and in psoriasis. An antibody against both P- and E-selectin (Hu EP5C7) has been tested in a baboon stroke model, while specific antibodies against E- (i.e. CDP850) and P-selectin (CY1747) have been tested in psoriasis and preclinically in a model of ischemia-reperfusion injury. A recombinant truncated form of a PSGL-1-immunoglobulin fusion protein has shown promise as a selectin inhibitor¹³⁵ (mainly aimed at P- and L-selectin) (www.wyeth.com). This molecule is currently in clinical trials for liver and renal transplantation (<http://www.ysthera.com/news/070502.html>). Small-molecule inhibitors of selectins known as glycomimetics have also been developed¹³⁶. A small-molecule inhibitor of selectin function (i.e. TBC-1269, Bimosiamose), developed by Texas Biotechnology (www.tbc.com) has been tested as an inhaled formulation for pediatric asthma and in psoriasis¹³⁷. Positive results were reported from a phase II clinical trial in patients with chronic obstructive pulmonary disease (COPD) (http://www.revotar.de/news_and_press.php).

Integrins

Integrins are a large superfamily of heterodimeric transmembrane glycoproteins that allow attachment of cells to extracellular matrix proteins or to ligands on other cells. Integrins contain large (α) and small (β) subunits of 120–170 kDa and 90–100 kDa, respectively. Integrins contain binding sites for the divalent cations Mg^{2+} and Ca^{2+} , which are necessary for their adhesive function. Distinct families of mammalian integrins form through the association of specific α subunits with different β subunits²⁵ (Figure 3). The families of greatest importance for leukocyte adhesion in inflammation and immunity are the β_2 , α_4 and β_7 families of adhesion molecules. Unlike the selectins, they establish stable protein-protein interactions and mediate arrest of cells on the vessel wall. However, some integrins have also been shown to participate in rolling¹³⁸. Integrins are also important signaling molecules involved in many cellular processes¹³⁹. In addition, several β_1 integrins are involved in lymphocyte interaction with ECM proteins²⁵.

β_2 Integrins (CD18)—Integrins containing the β_2 subunit (CD18) are exclusively expressed on bone marrow-derived cells. β_2 integrins undergo a conformational change upon cellular activation (inside-out signaling), which is necessary for ligand binding^{139, 140}. Activation is initiated by the binding of a chemokine or other chemoattractant to its heptahelical G-protein coupled receptor (GPCR) on the leukocyte surface^{41, 141}. Ligation of the GPCR results in integrin activation within milliseconds¹³⁹.

β_2 integrins include four different heterodimers: CD11a/CD18 (LFA-1) is the predominant β_2 integrin on lymphocytes and neutrophils; CD11b/CD18 (Mac-1) is expressed by monocytes, granulocytes and some NK cells; CD11c/CD18 (p150, p95) is present on monocytes and dendritic cells. CD11d/CD18 is expressed primarily by myelomonocytic cell lines and is predominantly found on macrophages and granulocytes in the red pulp of the spleen¹⁴². In humans¹⁴³ and mice¹⁴⁴, null or hypomorphic mutations in the *Itgb2* gene encoding the β_2 (CD18) integrin subunit result in the genetic disorder leukocyte adhesion deficiency-1 (LAD-1)¹⁴³. Patients with LAD-1 have recurrent bacterial infections due to ineffective recruitment of granulocytes in response to infections¹⁴³. This spontaneous human disease dramatically illustrates the important role played by CD18 integrins in innate immunity.

LFA-1 binds to ICAM-1 and ICAM-2 on the endothelium^{145, 146}. Its crucial role has been demonstrated in many cellular processes, such as migration, antigen presentation, and cell proliferation. Mice homozygous for a null mutation in the gene encoding the LFA-1 α chain have, amongst other defects, a mild defect in lymphocyte homing to secondary lymphoid organs¹⁴⁷. Targeted blockade of the integrin α_L subunit (CD11a) of LFA-1 (i.e. efalizumab) has been approved by the FDA for the treatment of chronic plaque psoriasis⁷. LFA-1 shares several characteristics and functions with Mac-1. LFA-1 has a predominant role for adhesion, Mac-1 is necessary for respiratory burst and certain forms of phagocytosis^{148, 149}. Mac-1 has no known function on lymphocytes. The other β_2 integrins (i.e. $\alpha_x\beta_2$ (CD11c/CD18) and $\alpha_d\beta_2$ (CD11d/CD18) are not expressed on lymphocytes.

α_4 Integrins (CD49d)—An important member of the α_4 family of integrins for leukocyte adhesion and trafficking is very late antigen-4 (VLA-4, CD49d/CD29, $\alpha_4\beta_1$)^{150, 151}. VLA-4 binds to its ligand VCAM-1, and is chiefly responsible for lymphocyte and monocyte adhesion to vascular endothelium. Studies in autoimmune encephalitis (EAE) have demonstrated a critical role for VLA-4 for lymphocyte homing into the CNS. VCAM-1, which is expressed at very low levels in CNS microvessels, is highly induced by proinflammatory stimuli in EAE and in patients with MS¹⁵².

$\alpha_4\beta_7$ is a pivotal integrin involved in gut-homing. This integrin binds to mucosal addressin cell adhesion molecule-1 (MAdCAM-1)^{128, 129} and plays a critical role in homing of lymphocytes

to Peyer's patches by supporting their binding to high endothelial venules. Both $\alpha_4\beta_1$ and $\alpha_4\beta_7$ can be activated to a high-avidity state¹⁵³. The β_7 integrin null mouse shows reduced homing of nai-ve lymphocytes to Peyer's patches⁴⁸, which is even further reduced in double mutant mice lacking both β_7 integrin and L-selectin^{31,154}. Homozygous null mutations for integrins α_4 and β_1 lead to embryonic lethality¹⁵⁵. α_4 -null chimeric mice have normal thymus cellularity and circulating T cells at birth, yet after a few weeks the thymus becomes atrophic^{156, 157}. This data supports an important role for α_4 integrins for the maintenance of thymocyte populations after birth. As integrin β_7 deficiency does not have a comparable effect it is then likely that the deficiency for $\alpha_4\beta_1$ integrin is responsible for this finding.

Integrin $\alpha_E\beta_7$ (CD103)

Integrin $\alpha_E\beta_7$ (CD103) was first identified by a monoclonal antibody that recognizes lymphocytes in intestinal tissue (i.e. human mucosal lymphocyte antigen-1, HML-1). The 175kDa alpha E chain pairs exclusively with the β_7 chain also present in $\alpha_4\beta_7$. $\alpha_E\beta_7$ is expressed by over 90% of intraepithelial lymphocytes and subsets of dendritic cells (DC). Its role in lymphocyte traffic remains unclear, however recently it has emerged as an important marker of CD4⁺ and CD8⁺ regulatory T cells as well as for DC subsets responsible for imprinting gut tropism to T cells^{158–160}.

Immunoglobulins

Members of the immunoglobulin superfamily share structural and genetic features with immunoglobulin molecules, and each contains at least one immunoglobulin domain (Figure 4). An immunoglobulin domain consists of two β -pleated sheets held together by a disulfide bond. Some molecules of the immunoglobulin superfamily are expressed on the vascular endothelium, where they act as counter-receptors for leukocyte integrins. Three of these immunoglobulins are of particular importance for leukocyte adhesion in inflammatory bowel disease. Intercellular adhesion molecule-1 (ICAM-1) or CD54, Vascular Cell Adhesion Molecule-1 (VCAM-1) or CD106, and Mucosal Addressin Cell Adhesion Molecule-1 (MAdCAM-1) all play important roles in the dysregulated trafficking of leukocytes that occurs during IBD.

ICAM-1

ICAM-1 is a homodimeric molecule constitutively expressed on a variety of cell types including resting endothelial cells. Its expression on endothelial cells is up-regulated by stimulation with cytokines¹⁶¹. ICAM-1 is one of the principal ligands for the β_2 -integrins LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18)¹⁶², although in the context of transmigration it seems that CD11a predominantly binds to ICAM-1, whereas CD11b binds to multiple ligands¹⁶³. Under resting conditions, the affinities of LFA-1 and Mac-1 for ICAM-1 are low¹³⁹, but activation of leukocytes causes conformational changes in LFA-1 and Mac-1 that greatly increases their affinities for ICAM-1. Although ICAM-1 cannot support rolling on its own, isolated LFA-1 I domain supports rolling on ICAM-1¹⁶⁴. On neutrophils, E-selectin engagement can induce partially activated LFA-1 to mediate LFA-1-dependent rolling on ICAM-1⁶⁹. An antisense oligonucleotide that blocks the translation of ICAM-1 mRNA (i.e. Alicaforsen) has been evaluated in IBD, yet its efficacy has been limited¹⁶⁵.

ICAM-2

ICAM-2, like ICAM-1 is a homodimeric molecule¹⁴⁶ Different from ICAM-1 which has five Ig-like domains, ICAM-2 has only two (figure 4). It is expressed constitutively on endothelial cells, including HEV, on certain leukocytes and on megakaryocytes and platelets¹⁶⁶. The expression of ICAM-2, different from ICAM-1 does not appear to be upregulated by inflammatory stimuli, thus it is thought to be involved mostly during constitutive trafficking.

ICAM-2-deficient mice have normal leukocyte homing and maturation and do not suffer from spontaneous inflammatory conditions¹⁶⁷.

VCAM-1—VCAM-1 (CD106) is a type-1 transmembrane protein that contains either six or seven immunoglobulin domains^{168, 169} and is expressed on both large and small vessels after stimulation by cytokines^{170–172}. Expression on resting endothelial cells is very low or absent. On cultured endothelial cells, sustained expression of VCAM-1 lasts at least 24 hours. VCAM-1 may also be found on epithelial cells, dendritic cells, Kupffer cells and on smooth muscle cells within atherosclerotic lesions^{172–174}. VCAM-1 is primarily an endothelial ligand for very late antigen-4 (VLA-4 or $\alpha_4\beta_1$) and promotes the adhesion of lymphocytes, monocytes, eosinophils, and basophils. VCAM-1-deficiency results in embryonic lethality. A conditional VCAM-1-mutant mouse that lacks VCAM-1 on endothelial cells showed a lymphocyte homing defect to bone marrow and impaired humoral responses^{175, 176}. Mice hypomorphic for VCAM-1 expression were protected from atherosclerosis

MAdCAM-1—MAdCAM-1 contains two immunoglobulin domains and a mucin domain¹²⁸, and is the main ligand for $\alpha_4\beta_7$ integrin¹²⁹. It is expressed at high levels by HEV of Peyer's patches and in lamina propria venules¹²⁸. The frequency of MAdCAM-1 positive vessels is increased in human IBD¹⁷⁷ and in animal models of colitis¹⁷⁸. It is also present in the inflamed pancreas¹⁷⁹ and may be inducible in other locations by proinflammatory cytokines such as TNF- α and IL-1¹⁸¹.

In addition to binding to $\alpha_4\beta_7$ integrin, MAdCAM-1 isolated from young mice also supports rolling through L-selectin¹³⁰. This double role of MAdCAM-1 may underlie the synergistic effects of L-selectin and $\alpha_4\beta_7$ integrin in lymphocyte homing to gut-associated lymphoid tissue^{31, 44}. MAdCAM-1 is up-regulated in the IL-10 knockout mouse, which develops spontaneous colitis¹⁷⁸, and in intestinal lesions of patients with IBD¹⁷⁷ supporting an important role in chronic inflammatory conditions.

Animal Models of Colitis

As there are excellent reviews in the topic¹⁸² we will only briefly introduce models in which the anti-adhesion strategy has been evaluated with the goal of attenuating inflammation. Three broad categories of colitic models have been used to explore this strategy. A spontaneous model (i.e., the cotton top tamarin model), four chemically induced models (i.e. acetic acid-, 2,4,6-trinitrobenzenesulfonic acid (TNBS), dextran sulfate sodium (DSS) and formalin-induced colitic models) and two others that develop colitis due to deficient regulatory responses (i.e. CD45Rb^{high} CD4 lymphocyte transfer, IL10-deficient mice).

Cotton-top tamarins

A wasting syndrome was reported in cotton-top tamarins, an endangered South American marmoset, in 1976. An association between colitis and colonic adenocarcinoma in CTT was reported in 1981¹⁸³ and their chronic colitis formally characterized by Madara in 1985¹⁸⁴. Similar to patients with UC, CTT develop recurrent spontaneous flares of colitis with distortion of the mucosal architecture, suggesting chronic dysregulated immune responses. No identifiable pathogens to which the histologic findings may be attributed were identified. Like human IBD, colitis in CTT responds to treatment with the 5-ASA compound sulfasalazine¹⁸⁴.

Chemically Induced models

The administration of diverse chemical agents such as: sulfated polysaccharides (e.g. carrageenan, amylopectin sulfate, dextran sulfate (DSS)), rectal instillation of diluted acetic

acid, TNBS, and intravenous injection of immune complexes followed by chemical irritation of the colon, all result in the development of colitis. A common shared mechanism for all of these models is the disruption of the epithelial barrier with increased exposure of intestinal microflora to the intestinal immune system. The observation that administration of such distinct chemical irritants result in similar histologic findings suggests that the colitis is rather an unspecific, stereotypic response to injury (125). The original disturbance does not determine the specific nature of the resulting lesions but likely serves as a trigger of common immunologic responses. The inflammatory response in all of these models is self-limited and resolves spontaneously upon removal of the injurious agent. Therefore these models do not accurately replicate the specific chronic pathogenetic mechanisms that mediate disease maintenance in human ulcerative colitis or colonic CD.

Acetic acid

Intrarectal administration of acetic acid (AA) results in barrier dysfunction and induces diffuse colitis in a dose-dependent manner. At the tissue level, AA induces diffuse ulcers of the distal colon with alterations in crypt architecture with occasional transmural inflammatory infiltrates that vary according to the severity of the process ¹⁸⁵. As an acute colitis, it is better suited to address the early pathogenetic mechanisms of IBD lesions.

2,4,6-trinitrobenzenesulfonic acid (TNBS)

Intracolonic co-administration of 50% ethanol, which breaches the mucosal barrier along with the hapten TNBS induces a sustained colitis that persists up to 8 weeks. The inflammatory infiltrate is mixed, composed of neutrophils, macrophages and lymphocytes. Granulomas are present in over 50% of animals. Disruption of the barrier by ethanol and TNBS are both required for the induction of colitis and administration of one or the other fails to induce disease ¹⁸⁶. As the inflammatory process last for up to 2 months additional information may be obtained beyond the very early developmental stages of colitis.

Dextran sulfate sodium (DSS)

DSS administered orally is a reproducible and technically simple way to induce colitis in rodents. Differential susceptibility to DSS has been reported in several mouse strains, yet the agent induces some degree of colitis in all mice. DSS disrupts the mucosal barrier and induces a colitic process that is more dependent on innate immune responses, as it develops even in the absence of T, B and NK cells ¹⁸⁷. Yet in mice with an intact immune system there is eventual activation of T cells responses, which is superimposed on the primary innate responses induced by the chemical agent.

Formalin-induced

This model is part of an early family of IBD animal models induced by deposition of immune-complexes in the colon, triggered by the mucosal injury that results after administration of dilute formalin. The lesion is mostly localized to the mucosa with damage to the epithelium, loss of the goblet cells and a predominant acute inflammatory infiltrate ¹⁸⁸. Although some of these features are shared with human IBD, the resultant inflammatory process is transient and resolves spontaneously.

Colitic models due to deficient regulatory responses

CD45RB^{high} Transfer Model—Powrie et al described in 1994 that transfer of CD45RB^{high} cells into immunodeficient recombinate activating gene (RAG)-deficient or severe combined immunodeficient (SCID) mice results in the development of colitis ¹⁸⁹. Most naïve T cells are CD45RB^{high} whereas regulatory T cells are not within this subset. Co-transfer of the CD45RB^{low} cells, which contain regulatory T cell subsets, prevents and cures

colitis. The bacterial flora remains an important factor in this model, as transfer of CD45RB^{high} cells into gnotobiotic recipients results in attenuated disease¹⁹⁰. Different from previously discussed models, adaptive immune responses are fundamental in CD45RB^{high} colitis, enabling the assessment of the role of adhesion molecules expressed by pathogenic T cells and the contribution of specific T cells subsets to the disease process.

IL10-deficient mouse

IL-10-deficient mice develop chronic transmural colitis¹⁹¹. The disease is characterized by T-helper (Th)1 responses during early stages and by Th2 responses during late stages. Lymphocyte development and antibody responses are normal. Histological findings in the colon include mucosal hyperplasia, mixed inflammatory infiltrates and aberrant expression of major histocompatibility complex class II molecules on epithelia. The bacterial flora remains an important player, as mice kept under specific pathogen-free conditions develop only local inflammation within the proximal colon. Thus IBD in this model results from dysregulated immune responses likely triggered by bacterial antigens. IL-10 is probably an essential regulatory cytokine for the maintenance of intestinal homeostasis. The inflammatory process is chronic and therefore suited to address mechanisms related to the chronic stages of IBD.

Adhesion Molecules as Therapeutic Targets in Animal Models of Colitis

Interference with lymphocyte recirculation to the intestine by targeting specific adhesion molecules and chemokine receptors is an attractive strategy for drug development in IBD¹⁹. This approach has already resulted in the development of several agents that have reached clinical use, e.g., natalizumab (Biogen/Elan Pharmaceuticals, www.tysabri.com) which targets the α_4 integrin, shared by the $\alpha_4\beta_1$ and $\alpha_4\beta_7$ integrins. MLN02 (Millenium Pharmaceuticals, www.mlnm.com/rd/inflammation/candidates/mln02.asp) targets specifically the $\alpha_4\beta_7$ integrin. Most recently Traficet EN (Chemocentryx Corporation), which targets the small-intestinal-specific chemokine receptor CCR9 has been tested in IBD. The therapeutic success of natalizumab has been overshadowed by the appearance of serious complications²¹. Numerous studies have targeted adhesion molecules in animal models of IBD (Table 1), yet much more remains to be learned. A greater knowledge of the basic mechanisms that underlie dysregulated chronic inflammatory trafficking, as well as the differences between trafficking to the small or large intestine will be required for the further development of the anti-adhesion strategy in IBD.

Targeting selectins in murine models of Colitis

Given the critical role of the selectins in mediating the initial steps of leukocyte recruitment to inflammatory sites, they would appear to be obvious potential therapeutic targets in IBD. Yet few reports attempted to target these molecules in animal models of IBD or reported disease attenuation by targeting a selectin. In the earliest report by Podolsky *et al.*, colitic Cotton-top tamarins, which suffer from a spontaneous relapsing colitis reminiscent of UC, were administered two different antibodies against E-selectin. Despite close attention to dosing, half-life and tissue levels, these antibodies had no significant attenuating effect on the severity of the colitis¹⁹². In a subsequent report, colitis was induced in P-selectin-deficient, and P- and E-selectin double-deficient mice using acetic acid or TNBS. Damage scores were not different in the P-selectin-deficient mice when compared to wild-type mice, whereas the double-deficient mice showed enhanced leukocyte recruitment and more severe disease¹⁹³. More recently, Sans and colleagues reported a series of experiments that explored the role of the selectins in the TNBS-induced rat model of colitis¹⁹⁴. Antibodies against E-, L- and P-selectin decreased rolling fluxes at certain time points, while the anti-P-selectin monoclonal antibody also decreased the number of adherent leukocytes. Yet no attenuating effect on the severity of the colitis was observed, even with the anti-P-selectin antibody. By contrast, the following year

the same group assessed the effect of an anti-P-selectin antibody and P-selectin-deficiency in dextran sulfate sodium (DSS)-induced colitis in mice, and observed an attenuating effect of the severity of the colitis for both ¹⁹⁵. These authors propose that this discrepancy may result from inter-species variability, different experimental approaches or distinct pathophysiological mechanisms related to the different ways of inducing colitis in the different models. In agreement with this study, anti-P-selectin antibodies interfered with leukocyte rolling and adhesion in DSS-treated mice ¹⁹⁶ and attenuated TNBS-colitis in rats ¹⁹⁷.

Selectin Ligands in colitis

We are aware of only one study that targeted selectin ligands in a mouse model of colitis. Rijcken and colleagues treated mice with an antibody against PSGL-1 (2PH1), and studied its effects on rolling and adhesion through intravital microscopy. In addition they assessed its effect on colonic inflammation by means of myeloperoxidase assays, a surrogate marker for neutrophil infiltration and histological assessment of the severity of colitis ¹⁹⁸. Blockade of PSGL-1 with 2PH1 reduced the number of rolling leukocytes, increased rolling velocities and attenuated colitis, supporting an important role for PSGL-1 for leukocyte recruitment in this model. Recently PSGL-1 deficient CD4⁺ T cells were shown to induce colitis in RAG^{-/-} mice indistinguishable to that of WT cells ¹⁹⁹, whereas the expression of PSGL-1 in colonic endothelium was induced by DSS administration ⁹². The functional role of PSGL-1 in colitis remains unclear.

Targeting integrins in murine models of Colitis

β_2 Integrins in colitis

The first report of an attenuating effect by targeting an integrin in an animal model of IBD appeared in 1992 when Wallace and colleagues administered an anti-CD18 mAb to rabbits before and after induction of colitis by TNBS. Both neutrophilic infiltration and epithelial injury improved with this strategy, regardless of whether the antibody was administered prior to or 12h after TNBS administration ²⁰⁰. This result was confirmed in TNBS colitis in rats using 2 antibodies against CD11b/CD18, administered 2 hrs prior to and 3 days after induction of colitis ²⁰¹. Immune complex colitis induced by formalin administration in rabbits also responded to an antagonist of CD11b/CD18 ²⁰². Recently, Grisham and coworkers have reported attenuated colitis induced by CD4⁺ T cells isolated from LFA-1 (CD11a/CD18)-deficient (*Itgal*^{-/-}) mice, and attenuated colitis when DSS was administered to CD18 (*Itgb2*^{-/-}) and CD11a null mice ^{203,204}. An antibody against complement receptor 3, which serves as a ligand for $\alpha_M\beta_2$ (Mac-1) and $\alpha_X\beta_2$ integrins ameliorated TNBS- and CD45RB^{high}-induced colitides ²⁰⁵.

α_4 integrins in colitis

Targeted blockade of the α_4 integrins in an animal model of IBD was first described by Podolsky and colleagues in cotton-top tamarins. An anti-human α_4 integrin (HP1/2) known to interfere with the interaction of VLA-4 with VCAM-1 or a saline placebo control were administered daily for 7 days to colitic animals. Colonic biopsies obtained before and after treatment demonstrated that HP1/2 attenuated acute activity but had no effect on the chronic inflammatory infiltrates ¹⁹². An antibody that targeted a combinatorial epitope on the $\alpha_4\beta_7$ integrin was then successfully used to attenuate colitis in cotton-top tamarins. Following a similar protocol to that of Podolsky et al, cotton-top tamarins were randomized to receive daily injections for 7 days of anti- $\alpha_4\beta_7$ or an isotype control, with acquisition of colonic biopsies at three time points and close monitoring of a clinical effect on stool consistency. Significant histological improvement was observed. Clinically cotton-top tamarins showed improvement in stool consistency during the first 24h after administration of the antibody. This therapeutic

response persisted for over 2 weeks after completion of the therapeutic regimen²⁰⁶. Monoclonal antibodies against $\alpha_4\beta_7$ integrin and MAdCAM-1 blocked lymphocyte recruitment and reduced the severity of colonic inflammation in the CD45Rb^{high} transfer model of colitis, one of the best studied IBD murine models²⁰⁷. Similarly, anti- MAdCAM-1 antibodies significantly reduced colonic injury and decreased the number of β_7 integrin-positive cells in the colonic mucosa in mice with DSS-induced colitis^{208, 209}.

Targeting immunoglobulins in murine models of colitis

ICAM-1 and VCAM-1 mRNA levels are increased in inflamed intestine of both TNBS colitic mice^{194, 210} and IL-10-deficient mice²¹¹; in the latter mRNA levels for MAdCAM-1 were also increased. Similarly, in the CD45Rb^{high} colitic model, mRNA levels for the three immunoglobulins increased²¹². MAdCAM-1 is up-regulated in chronically inflamed small and large intestine of patients with UC and CD^{177, 179}. Antibodies against MAdCAM-1 ameliorate DSS-induced colitis²¹³.

In TNBS-induced colitis, ICAM-1 was slightly upregulated and VCAM-1 was upregulated at least 8-fold, as measured by an intravenous radiolabeled antibody strategy useful to assess adhesion molecule expression on the endothelial surface²¹⁴. Intravital microscopy of colonic venules revealed increased leukocyte rolling and adhesion. An antibody to VCAM-1 was able to relieve some of the symptoms of disease and also reduced leukocyte adhesion. This is probably the strongest mechanistic evidence suggesting an involvement of VCAM-1 in intestinal inflammation, although TNBS-induced colitis differs significantly from human IBD. An anti-ICAM-1 antibody also attenuated TNBS colitis in rats¹⁹⁷. Colonization by *Lactobacillus casei* attenuated leukocyte recruitment to the colon, potentially by preventing the upregulation of ICAM-1 induced by TNBS²¹⁵. Evidence for increased adhesion molecule expression in humans comes from a study based on cultured human intestinal endothelial cells isolated from IBD patients, which support increased adhesion of monocytic cells (U937) and neutrophils compared with controls^{216, 217}, although a molecular mechanism was not identified. In therapeutic studies, antibodies to ICAM-1 protected rats from acute colitis induced by acetic acid and DSS^{218, 219, 220}. In addition, an ICAM-1 antisense oligonucleotide both prevented and reversed DSS colitis in mice²²¹ and ICAM-1-deficient mice showed decreased mortality and attenuated colitis induced by DSS²²². The endothelial molecule appears to be more important than leukocyte ICAM-1, as CD4⁺ T cells isolated from ICAM-1-deficient mice induced colitis in *Rag1*^{-/-} or *Rag2*^{-/-} mice indistinctly from that induced by WT mice¹⁹⁹.

Animal Models of Chronic Ileitis

Although there is a body of literature describing the physiology of lymphocyte trafficking, our knowledge of leukocyte recruitment during conditions of chronic inflammation as seen in patients with IBD is limited. This is especially true in inflammatory cell trafficking to the small intestine, in part due to the lack of animal models that develop disease in the terminal ileum. However, two murine models that develop chronic ileitis similar to that characteristic of CD have now been described, i.e., SAMP1/Yit and TNF Δ ARE mice^{223, 224}. This has facilitated investigation of the adhesion pathways that are involved in lymphocyte trafficking specifically to the small intestine under conditions of chronic inflammation. Both the SAMP1/YitFc and TNF Δ ARE models of chronic ileitis show many of the features that are characteristic of CD^{182, 225}, including the chronic and unrelenting course, the critical role of T lymphocytes and the specific localization of the inflammatory process to the small intestine²²⁶.

The SAMP1/Yit model of chronic ileitis

In 1998, Matsumoto *et al.* described a mouse strain that develops chronic inflammation predominantly in their terminal ilea and truly recapitulated many of the histopathological findings of CD in humans ²²⁴. Unlike all the previously described models of IBD, intestinal inflammation develops spontaneously in the small intestine, in the absence of chemical, immunologic or genetic manipulation. These mice were derived from the SAMP1 strain of senescent-accelerated mice (SAM), which originated from AKR inbred mice ²²⁷. SAMP1/Yit mice represent a sub-line of the SAMP1 strain, uniquely identified by the development of spontaneous skin lesions that correlate with intestinal inflammation ²²⁴. After selective inbreeding for the presence of skin lesions, followed by transfer from conventional to specific pathogen-free conditions, the early senescence phenotype was lost and the presence of discontinuous chronic ileitis was first described ²²⁴.

After over 30 generations of brother-sister matings at the University of Virginia, and the appearance of several phenotypic differences not present in the donor colony, a sub-strain was designated SAMP1/YitFc ^{228, 229}. Similarly to the original strain, these mice have 100% disease penetrance, and exhibit patchy, transmural inflammation that is primarily localized to the terminal ileum, with formation of granulomata, as well as dense infiltration of the terminal ileal lamina propria by lymphocytes, neutrophils and macrophages. In addition these mice develop extraintestinal manifestations that target many of the organs also affected by human IBD. These include pyoderma-like skin lesions, ocular involvement and perianal disease, which is similar to the perianal manifestations observed in humans with CD, including fissures, rectal prolapse, fistulae and abscess formation ²²⁹.

Immunologically, the ileitis is characterized by increased production of IFN- γ and TNF α from lymphocytes by 4 weeks, predating any histological evidence of inflammation ^{228, 229}. The role of Th17 cells in the pathogenesis of the disease remains to be tested. The disease is mediated by lymphocytes that infiltrate the lamina propria and mesenteric lymph nodes, display an activated phenotype, and adoptively transfer disease to SCID mice ²²⁸. The adoptive transfer of SAMP1/YitFc CD4⁺T cells predominantly induces ileitis, different from that transferred by CD45Rb^{high} T cells that cause colitis, which is thought to reflect the inherent ability of SAMP1/YitFc T cells to recirculate preferentially to the small intestine. The disease in these mice also responds to therapeutics currently used in the treatment of CD, including corticosteroids, and monoclonal antibodies against TNF α ²³⁰ and α 4 integrins, all of which ameliorate disease severity ²³¹.

The TNF Δ ARE model

A second model of chronic ileitis was described by Kontoyiannis *et al.* in 1998 ²²³. The TNF Δ ARE model develops both a predominantly Crohn's-like chronic ileitis similar to that of SAMP1/Yit mice, and deforming rheumatoid arthritis-like joint disease in both heterozygous and homozygous mice. Like SAMP1/Yit, the disease in TNF Δ ARE mice responds to immunoblockade of TNF- α , supporting the relevance of this model to the human diseases. TNF Δ ARE mice were developed by targeted deletion of 69 base pairs of AU-rich elements in the 3' UTR region of the gene that encodes TNF- α . This deletion results in increased stability of the mRNA for TNF- α , increased synthesis of TNF- α protein and elevated levels of systemic TNF- α ²²³. Interestingly, the increased levels of TNF- α do not result in uniform pan-enteritis, but predominantly in ileitis. Homozygous TNF Δ ARE^{+/+} mice develop severe ileitis and arthritis, and die at an early age. Heterozygous mice (^{+/−}) develop disease by 8 weeks-of-age that worsens progressively, yet they reach sexual maturity and procreate. A related strain (i.e. B6.129S-Tnf^{tm2Gkl}/Jarn, MGI: 3720980) on the BL6 background has been generated by over 20 generations of continuous backcrosses between TNF Δ ARE mice on mixed genetic background (i.e. C57BL6 and 129S6), generated as previously described ²²³ to C57BL6/J mice.

The C57BL/6/J genetic background did not alter the localization, time course or severity of the intestinal inflammation. The defined C57BL/6 genetic background greatly facilitates immunologic studies that would be difficult or impossible on the original strain with a mixed genetic background.

Trafficking pathways in models of ileitis

Dissecting the lymphocyte-trafficking pathways in chronic murine ileitis may help optimize the current therapeutics in clinical use in IBD, as well as to identify new potential therapeutic targets. To that effect we have targeted many of the adhesion molecules and chemokines using these two models of ileitis.

L-selectin in ileitis—The role played by L-selectin in nai-ve T-cell trafficking has been relatively well characterized²³², whereas the role it plays during induction or maintenance of chronic inflammation is incompletely understood.

Expression of L-selectin is often considered to correlate with antigenic inexperience and to be indicative of the nai-ve state. Although nai-ve T cells indeed express L-selectin, it is also expressed in central memory T cells⁵⁰ at other times. L-selectin is re-expressed after lymphocyte activation in *in vitro* systems, and subpopulations of memory and effector lymphocytes express L-selectin^{33, 50, 233}. Therefore, L-selectin may continue to play a role in the recruitment of antigen-experienced subpopulations of lymphocytes important for the pathogenesis of chronic inflammatory diseases, where L-selectin ligands (expressed in PNAd⁺ and MAdCAM-1⁺ vessels) have been demonstrated within effector sites. PNAd⁺ vessels are consistently found within inducible lymphoid aggregates or follicles (ILF), also known as tertiary lymphoid organs (TLO)^{124, 127}. Thus, recruitment of L-selectin-expressing T cells likely occurs at these sites, and interference with these pathways may be of therapeutic value^{122, 123}.

Targeting selectins in ileitis

The models of chronic ileitis have allowed us to assess for the first time whether the trafficking pathways to the small and large intestine were shared or distinct from those to other intestinal segments under conditions of dysregulated inflammation. Systematic targeting of the different families of adhesion molecules with neutralizing antibodies against single molecules or in combination was performed in an adoptive transfer model in which CD4⁺ T cells from SAMP1/YitFc mice are adoptively transferred into MHC-matched SCID mice. The predominant disease in the recipients is ileitis, not colitis. Single (data not shown) or combined blockade of P-, E- and/or L-selectin failed to attenuate the ileitis in SCID mice or in SAMP1/YitFc mice with established ileitis⁹¹ (Figure 5). However, administration of the anti-L-selectin antibody MEL-14 prior to the development of inflammation had a protective effect, suggesting a requirement for L-selectin during the initiation stages of the disease (Rivera-Nieves, J, unpublished results).

Targeting selectin ligands in ileitis

PNAd

The selectins possess a C-type lectin domain at their amino terminus that interacts with carbohydrate determinants expressed by ligands expressed on high endothelial venules (HEV) and PSGL-1 expressed by adherent leukocytes²⁶. Indeed, all of the L-selectin ligands identified to date (i.e., CD34, GlyCAM-1, sgp200, PSGL-1, endomucin, MAdCAM-1) contain carbohydrate moieties¹³⁴. The carbohydrate post-translational modifications found on HEVs that are necessary for L-selectin recognition include sialylation, fucosylation and carbohydrate sulfation¹²⁴. Chronic inflammatory processes induce the appearance of HEV-like vessels in

numerous human diseases and animal models of chronic inflammation²³⁴. A likely role of these inflammation-induced structures is to support recruitment of L-selectin-expressing lymphocytes directly into effector sites, where during chronic inflammation numerous tertiary lymphoid organs appear¹²⁷.

The MECA-79 monoclonal antibody stains HEVs and blocks L-selectin-dependent lymphocyte adhesion²³⁵. Glycoproteins carrying the PNAd epitope are recognized as relevant L-selectin ligands, as no additive effects on lymphocyte homing was observed after simultaneous blockade of PNAd and L-selectin. The MECA-79 epitope is found in PP, MLN, PLN but not within the intestinal LP, except in chronic inflammation²³⁶. Yet this pattern is not restricted to the intestine and PNAd reactivity has been reported in over 20 immune-mediated animal models and human conditions, including asthma, diabetes, psoriasis, RA, Hashimoto's thyroiditis, Grave's disease, cutaneous lymphoma and allograft rejection¹²⁴.

Immunohistochemical studies in SCID mice before and after CD4⁺ T cell adoptive transfer from donor SAMP1/YitFc mice demonstrated that the PNAd epitope is expressed after the development of inflammation in these mice. However, single or combined blockade of MAdCAM-1 (MECA-367) and PNAd (MECA-79) provided no therapeutic benefit compared to mice treated with isotype-matched control antibodies⁹¹.

PSGL-1

Inoue and colleagues described for the first time the therapeutic effect of neutralization of PSGL-1 with the monoclonal antibody 2PH1 in SAMP1/Yit mice²³⁷. Administration of 2PH1 to mice with established disease significantly reduced infiltration of the LP by CD4, CD8 and CD68⁺ subsets, and the villus height was partially restored, although no effect on the hypertrophy of the muscularis (a chronic structural change) was noted. Their emphasis was on the role of PSGL-1 on monocyte recruitment to the intestine; however when bone marrow chimeric SAMP1/Yit mice that lacked PSGL-1 in their hematopoietic compartment were generated, their disease was equal in severity to that of PSGL-1-sufficient SAMP1/Yit mice suggesting that PSGL-1 on monocytes was not the critical molecule. The small intestinal microvasculature expressed PSGL-1 and a therapeutic effect of blockade of PSGL-1 was demonstrated in SAMP mice using a different antibody (i.e. 4RA10)⁹¹. It is likely that the therapeutic effect of this strategy is due to its interference with multiple recruitment pathways (Figure 6). Interfering with PSGL-1 would block rolling on P-selectin expressing endothelial cells, rolling on platelets, platelet-leukocyte interactions and endothelial-platelet interactions.

Integrins in chronic murine ileitis

The importance of the α_4 integrins in lymphocyte trafficking is well established in the literature⁴. $\alpha_4\beta_7$ and $\alpha_4\beta_1$ integrins are crucial for the dichotomy in migration to gut or to non-intestinal sites in physiological recruitment. In this setting, $\alpha_4\beta_7$ -expressing cells migrate preferentially to the gut, whereas lymphocytes expressing the $\alpha_4\beta_1$ integrin (VLA-4) preferentially traffic to non-intestinal sites^{235, 238}. Expression of these integrins under physiological conditions is reciprocal, and gut-homing $\alpha_4\beta_7^+$ cells tend to be $\alpha_4\beta_1^{\text{neg}}$ or low and vice versa^{235, 238}. Recent clinical trials with a humanized mAb to α_4 integrins in patients with Crohn's disease have suggested that α_4 integrins may be relevant therapeutic targets^{18, 20}. Yet while targeting both α_4 integrins ($\alpha_4\beta_7$, $\alpha_4\beta_1$) through its shared α_4 moiety is effective in CD, a more specific mAb that targeted $\alpha_4\beta_7$ specifically was effective in UC but not in CD²³⁹. This data supports divergent trafficking pathways between small and large bowels. Further understanding of potential mechanisms of induction of small intestinal specific homing have recently emerged with the discovery that retinoic acid is essential for the imprinting of T cells with a small intestinal homing phenotype^{240,241, 242}.

Expansion of the β_7 integrin-expressing CD4⁺T cell population and of MAdCAM-1 after T cell transfer suggests a role for these molecules in the pathogenesis of chronic ileitis. Yet, neutralizing monoclonal antibodies against the β_7 integrin subunit (FIB-504) or $\alpha_4\beta_7$ integrin (DATK-32) were ineffective in reducing the severity of acute or chronic infiltrates, compared to isotype antibody-treated SCID mice with chronic ileitis⁸⁸. In agreement a β_7 -deficient SAMP1/Yit substrain recently generated develops ileitis indistinguishable from that of β_7 -sufficient SAMP1/Yit mice (Gorfu, G. Rivera-Nieves, J, McDuffie, M., Pizarro TT, Cominelli, F and Ley K, unpublished observations). To attenuate ileitis, blockade of at least two integrins might be required, for example using an anti- α_4 integrin antibody, which interferes with both $\alpha_4\beta_7$ and $\alpha_4\beta_1$ integrins⁸⁸.

The immunoglobulins in chronic murine ileitis

The expression of ICAM-1 and VCAM-1 was increased in intestinal microvessels of SAMP1/YitFc mice and the related SCID adoptive transfer model mice in the presence of inflammation, consistent with their induction under conditions of chronic inflammation. In addition, combined blockade of ICAM-1 and VCAM-1 was able to attenuate ileitis in adoptively transferred SCID mice, whereas blockade of either alone was not sufficient²³¹. This data supports the hypothesis that the trafficking pathways under conditions of chronic inflammation are highly redundant and resistant to therapeutic intervention. This is further supported by the complete lack of efficacy by the same combined intervention in SAMP1/Yit mice, in which the inflammatory process develops over many months with the likely acquisition of highly redundant trafficking pathways²³¹.

In a separate set of experiments, neutralization of MAdCAM-1 was found to be ineffective at attenuating ileitis in SCID mice, despite clear induction of this molecule with the development of chronic inflammation. It is likely that T cells continue to home to the intestine using VCAM-1/ $\alpha_4\beta_1$ and ICAM-1/LFA-1 after the MAdCAM-1/ $\alpha_4\beta_7$ pathway has been neutralized. This is supported by the efficacy of the simultaneous blockade of α_4 integrins (clone PS2) and ICAM-1²³¹, which targets multiple mechanisms of arrest⁸⁸.

Future Directions

Combination anti-adhesion molecule therapies

During the past decade, the success of anti-TNF- α strategies has revolutionized the treatment of IBD²⁴³. Yet, only about 70% of patients respond to this therapy, driving the continued search for other anti-inflammatory strategies for the treatment of the remaining 30% of patients who do not respond and for those that respond initially but eventually fail. Therefore, alternative biological therapies that target other pathways of the chronic inflammatory cascade must be evaluated in CD^{14, 243}. Interference with leukocyte recirculation to the intestine by targeting specific adhesion molecules has resulted in the development of agents that have advanced into clinical use, e.g., Natalizumab (Biogen/Elan Pharmaceuticals, www.tysabri.com) and MLN02 (Millenium Pharmaceuticals, www.mlnm.com/rd/inflammation/candidates/mln02.asp), which has been evaluated in CD and UC²³⁹.

A pivotal role for $\alpha_4\beta_7$ integrin/MAdCAM-1 in lymphocyte trafficking to the gut has become dogma under physiological conditions^{4, 235}, fueling the evolution of these anti- α_4 strategies in IBD. The evidence for a similar non-redundant role in chronic gut inflammation is less clear. Antibody-blocking studies in animals and humans suggest efficacy^{192, 207–209}, but specificity is unclear. Despite the rapid progression from the bench to bedside of the anti- α_4 therapeutic strategy^{18, 192}, much remains to be learned regarding its fundamental mechanism of action. Our data in the ileitis models support the hypothesis that compensatory mechanisms allow continued trafficking after neutralization of the $\alpha_4\beta_7$ integrin/MAdCAM-1 pathway that is

pivotal for physiologic and pathologic traffic to the colon. Further dissection of the adhesive pathways in chronic ileitis may help to optimize current therapies already in clinical use for CD¹⁸, minimize risks and potentially to further understand the mechanisms of inflammatory lymphocyte recruitment to other organs as well.

Chemokines as therapeutic targets

Recent studies have shown that early blockade of CCL25 using a small molecule antagonist (CCX282, Traficet-EN, Chemocentryx Corp.) effectively prevented ileitis in TNFΔARE mice, confirming a critical role for this molecule during the induction phase of chronic ileitis (Zheng Wei, Chemocentryx Corp., personal communication). However when we targeted this molecule in a different model of ileitis (SAMP1/Yit), the strategy was only efficacious during early disease²⁴⁴. Targeting this and other chemokines alone or in combination with anti-adhesion strategies may be promising in subsets of patients with IBD. Small molecule approaches may be particularly useful as adjunctive therapies.

Conclusions and Implications for future anti-adhesion strategies

Physiological leukocyte recruitment is a highly regulated process with a limited number of decision points along every step of the adhesion cascade^{4, 23, 24}. Expression of specific combinations of adhesion molecules on lymphocyte subpopulations determines their capacity to reach specific tissues, where cognate endothelial ligands are often restrictively expressed (e.g. MAdCAM-1: gut vs. VCAM-1: elsewhere). This results in orderly recruitment that begins and terminates physiological inflammatory responses. Under conditions of chronic inflammation, continuous dysregulated production of pro-inflammatory cytokines results in inappropriately increased adhesion molecule expression (e.g. MAdCAM-1), as well as aberrant expression of molecules not normally expressed by specific lymphocyte subpopulations and tissues, thus increasing the chances of lymphocytes contacting appropriate endothelial ligands. Therefore, when one adhesion molecule is blocked, others compensate for its deficiency. These redundant adhesive pathways favor adhesiveness over tissue specificity, resulting in failure of termination of the inflammatory response and perpetuation of dysregulated leukocyte recruitment. Further dissection of redundant adhesive pathways in chronic ileitis may help to optimize therapies already in clinical use and potentially develop new strategies for the treatment of CD that in two thirds of patients involves the small intestine.

References

1. Davidson A, Diamond B. Autoimmune diseases. *N Engl J Med* 2001;345:340–50. [PubMed: 11484692]
2. Marrack P, Kappler J, Kotzin BL. Autoimmune disease: why and where it occurs. *Nat Med* 2001;7:899–905. [PubMed: 11479621]
3. O'Shea JJ, Ma A, Lipsky P. Cytokines and autoimmunity. *Nat Rev Immunol* 2002;2:37–45. [PubMed: 11905836]
4. Springer TA. Traffic signals on endothelium for lymphocyte recirculation and leukocyte emigration. *Annu Rev Physiol* 1995;57:827–72. [PubMed: 7778885]
5. von Andrian UH, Mackay CR. T-cell function and migration. Two sides of the same coin. *N Engl J Med* 2000;343:1020–34. [PubMed: 11018170]
6. Lebowitz M, Tyring SK, Hamilton TK, Toth D, Glazer S, Tawfik NH, Walicke P, Dummer W, Wang X, Garovoy MR, Pariser D. A novel targeted T-cell modulator, efalizumab, for plaque psoriasis. *N Engl J Med* 2003;349:2004–13. [PubMed: 14627785]
7. Simmons DL. Anti-adhesion therapies. *Curr Opin Pharmacol* 2005;5:398–404. [PubMed: 15955735]
8. Davenport RJ, Munday JR. Alpha4-integrin antagonism - an effective approach for the treatment of inflammatory diseases? *Drug Discov Today* 2007;12:569–76. [PubMed: 17631252]

9. Xavier RJ, Podolsky DK. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448:427–34. [PubMed: 17653185]
10. Fiocchi C. Inflammatory bowel disease: etiology and pathogenesis. *Gastroenterology* 1998;115:182–205. [PubMed: 9649475]
11. Podolsky DK. Inflammatory bowel disease (1). *N Engl J Med* 1991;325:928–37. [PubMed: 1881418]
12. Weaver CT, Hatton RD, Mangan PR, Harrington LE. IL-17 family cytokines and the expanding diversity of effector T cell lineages. *Annu Rev Immunol* 2007;25:821–52. [PubMed: 17201677]
13. Kullberg MC, Jankovic D, Feng CG, Hue S, Gorelick PL, McKenzie BS, Cua DJ, Powrie F, Cheever AW, Maloy KJ, Sher A. IL-23 plays a key role in *Helicobacter hepaticus*-induced T cell-dependent colitis. *J Exp Med* 2006;203:2485–94. [PubMed: 17030948]
14. Targan SR, Hanauer SB, van Deventer SJ, Mayer L, Present DH, Braakman T, DeWoody KL, Schaible TF, Rutgeerts PJ. A short-term study of chimeric monoclonal antibody cA2 to tumor necrosis factor alpha for Crohn's disease. Crohn's Disease cA2 Study Group. *N Engl J Med* 1997;337:1029–35. [PubMed: 9321530]
15. Sands BE, Bank S, Sninsky CA, Robinson M, Katz S, Singleton JW, Miner PB, Safdi MA, Galandiuk S, Hanauer SB, Varilek GW, Buchman AL, Rodgers VD, Salzberg B, Cai B, Loewy J, DeBruin MF, Rogge H, Shapiro M, Schwertschlag US. Preliminary evaluation of safety and activity of recombinant human interleukin 11 in patients with active Crohn's disease. *Gastroenterology* 1999;117:58–64. [PubMed: 10381910]
16. Schreiber S, Nikolaus S, Malchow H, Kruis W, Lochs H, Raedler A, Hahn EG, Krummenerl T, Steinmann G. Absence of efficacy of subcutaneous antisense ICAM-1 treatment of chronic active Crohn's disease. *Gastroenterology* 2001;120:1339–46. [PubMed: 11313303]
17. Mannon PJ, Fuss JJ, Mayer L, Elson CO, Sandborn WJ, Present D, Dolin B, Goodman N, Groden C, Hornung RL, Quezado M, Yang Z, Neurath MF, Salfeld J, Veldman GM, Schwertschlag U, Strober W. Anti-interleukin-12 antibody for active Crohn's disease. *N Engl J Med* 2004;351:2069–79. [PubMed: 15537905]
18. Ghosh S, Goldin E, Gordon FH, Malchow HA, Rask-Madsen J, Rutgeerts P, Vyhnaek P, Zadorova Z, Palmer T, Donoghue S. Natalizumab for active Crohn's disease. *N Engl J Med* 2003;348:24–32. [PubMed: 12510039]
19. van Assche G, Rutgeerts P. Antiadhesion molecule therapy in inflammatory bowel disease. *Inflamm Bowel Dis* 2002;8:291–300. [PubMed: 12131614]
20. von Andrian UH, Engelhardt B. Alpha4 integrins as therapeutic targets in autoimmune disease. *N Engl J Med* 2003;348:68–72. [PubMed: 12510047]
21. Alvarez-Cermeno JC, Masjuan J, Villar LM. Progressive multifocal leukoencephalopathy, natalizumab, and multiple sclerosis. *N Engl J Med* 2005;353:1744–6. [PubMed: 16240475]author reply 1744–6
22. Bommakanti S, Patil A, Eshoa C, Chitambar CR. Case reports: efalizumab-associated lymphoproliferative disease. *J Drugs Dermatol* 2007;6:646–8. [PubMed: 17668531]
23. Butcher EC. Leukocyte-endothelial cell recognition: three (or more) steps to specificity and diversity. *Cell* 1991;67:1033–6. [PubMed: 1760836]
24. Ley K, Laudanna C, Cybulsky MI, Nourshargh S. Getting to the site of inflammation: the leukocyte adhesion cascade updated. *Nat Rev Immunol* 2007;7:678–89. [PubMed: 17717539]
25. Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell* 2002;110:673–87. [PubMed: 12297042]
26. Sperandio M. Selectins and glycosyltransferases in leukocyte rolling in vivo. *Febs J* 2006;273:4377–89. [PubMed: 16956372]
27. Kansas GS. Selectins and their ligands: current concepts and controversies. *Blood* 1996;88:3259–87. [PubMed: 8896391]
28. Rosen SD. Ligands for L-selectin: homing, inflammation, and beyond. *Annu Rev Immunol* 2004;22:129–56. [PubMed: 15032576]
29. Arfors KE, Lundberg C, Lindbom L, Lundberg K, Beatty PG, Harlan JM. A monoclonal antibody to the membrane glycoprotein complex CD18 inhibits polymorphonuclear leukocyte accumulation and plasma leakage in vivo. *Blood* 1987;69:338–40. [PubMed: 3539230]

30. Eriksson EE, Xie X, Werr J, Thoren P, Lindbom L. Importance of primary capture and L-selectin-dependent secondary capture in leukocyte accumulation in inflammation and atherosclerosis in vivo. *J Exp Med* 2001;194:205–18. [PubMed: 11457895]
31. Kunkel EJ, Ramos CL, Steeber DA, Muller W, Wagner N, Tedder TF, Ley K. The roles of L-selectin, beta 7 integrins, and P-selectin in leukocyte rolling and adhesion in high endothelial venules of Peyer's patches. *J Immunol* 1998;161:2449–56. [PubMed: 9725243]
32. Ley K. Adhesion molecules and the recruitment of leukocytes in postcapillary venules. *Microvascular Research: Biology and Pathology* 2005;321–325.
33. Ley K, Kansas GS. Selectins in T-cell recruitment to non-lymphoid tissues and sites of inflammation. *Nat Rev Immunol* 2004;4:325–35. [PubMed: 15122198]
34. Siegelman MH, Stanescu D, Estess P. The CD44-initiated pathway of T-cell extravasation uses VLA-4 but not LFA-1 for firm adhesion. *J Clin Invest* 2000;105:683–91. [PubMed: 10712440]
35. Tozeren A, Ley K. How do selectins mediate leukocyte rolling in venules? *Biophys J* 1992;63:700–9. [PubMed: 1420908]
36. Hammer DA, Apte SM. Simulation of cell rolling and adhesion on surfaces in shear flow: general results and analysis of selectin-mediated neutrophil adhesion. *Biophys J* 1992;63:35–57. [PubMed: 1384734]
37. Kunkel EJ, Dunne JL, Ley K. Leukocyte arrest during cytokine-dependent inflammation in vivo. *J Immunol* 2000;164:3301–8. [PubMed: 10706723]
38. Shamri R, Grabovsky V, Gauguier JM, Feigelson S, Manevich E, Kolanus W, Robinson MK, Staunton DE, von Andrian UH, Alon R. Lymphocyte arrest requires instantaneous induction of an extended LFA-1 conformation mediated by endothelium-bound chemokines. *Nat Immunol* 2005;6:497–506. [PubMed: 15834409]
39. Piali L, Weber C, LaRosa G, Mackay CR, Springer TA, Clark-Lewis I, Moser B. The chemokine receptor CXCR3 mediates rapid and shear-resistant adhesion-induction of effector T lymphocytes by the chemokines IP10 and Mig. *Eur J Immunol* 1998;28:961–72. [PubMed: 9541591]
40. Lorant DE, Patel KD, McIntyre TM, McEver RP, Prescott SM, Zimmerman GA. Coexpression of GMP-140 and PAF by endothelium stimulated by histamine or thrombin: a juxtacrine system for adhesion and activation of neutrophils. *J Cell Biol* 1991;115:223–34. [PubMed: 1717478]
41. Alon R, Grabovsky V, Feigelson S. Chemokine induction of integrin adhesiveness on rolling and arrested leukocytes local signaling events or global stepwise activation? *Microcirculation* 2003;10:297–311. [PubMed: 12851647]
42. Campbell JJ, Hedrick J, Zlotnik A, Siani MA, Thompson DA, Butcher EC. Chemokines and the arrest of lymphocytes rolling under flow conditions. *Science* 1998;279:381–4. [PubMed: 9430588]
43. Kim M, Carman CV, Springer TA. Bidirectional transmembrane signaling by cytoplasmic domain separation in integrins. *Science* 2003;301:1720–5. [PubMed: 14500982]
44. Bargatze RF, Jutila MA, Butcher EC. Distinct roles of L-selectin and integrins alpha 4 beta 7 and LFA-1 in lymphocyte homing to Peyer's patch-HEV in situ: the multistep model confirmed and refined. *Immunity* 1995;3:99–108. [PubMed: 7542550]
45. Rose DM, Han J, Ginsberg MH. Alpha4 integrins and the immune response. *Immunol Rev* 2002;186:118–24. [PubMed: 12234367]
46. Bevilacqua MP, Nelson RM. Selectins. *J Clin Invest* 1993;91:379–87. [PubMed: 7679406]
47. Arbones ML, Ord DC, Ley K, Ratech H, Maynard-Curry C, Otten G, Capon DJ, Tedder TF. Lymphocyte homing and leukocyte rolling and migration are impaired in L-selectin-deficient mice. *Immunity* 1994;1:247–60. [PubMed: 7534203]
48. Wagner N, Lohler J, Kunkel EJ, Ley K, Leung E, Krissansen G, Rajewsky K, Muller W. Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. *Nature* 1996;382:366–70. [PubMed: 8684468]
49. Butterfield K, Fathman CG, Budd RC. A subset of memory CD4+ helper T lymphocytes identified by expression of Pgp-1. *J Exp Med* 1989;169:1461–6. [PubMed: 2564418]
50. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 1999;401:708–12. [PubMed: 10537110]

51. McEver RP, Beckstead JH, Moore KL, Marshall-Carlson L, Bainton DF. GMP-140, a platelet alpha-granule membrane protein, is also synthesized by vascular endothelial cells and is localized in Weibel-Palade bodies. *J Clin Invest* 1989;84:92–9. [PubMed: 2472431]
52. Berman CL, Yeo EL, Wencel-Drake JD, Furie BC, Ginsberg MH, Furie B. A platelet alpha granule membrane protein that is associated with the plasma membrane after activation. Characterization and subcellular localization of platelet activation-dependent granule-external membrane protein. *J Clin Invest* 1986;78:130–7. [PubMed: 2941452]
53. Foreman KE, Vaporciyan AA, Bonish BK, Jones ML, Johnson KJ, Glovsky MM, Eddy SM, Ward PA. C5a-induced expression of P-selectin in endothelial cells. *J Clin Invest* 1994;94:1147–55. [PubMed: 7521884]
54. Gotsch U, Jager U, Dominis M, Vestweber D. Expression of P-selectin on endothelial cells is upregulated by LPS and TNF-alpha in vivo. *Cell Adhes Commun* 1994;2:7–14. [PubMed: 7526954]
55. Jung U, Ley K. Regulation of E-selectin, P-selectin, and intercellular adhesion molecule 1 expression in mouse cremaster muscle vasculature. *Microcirculation* 1997;4:311–9. [PubMed: 9219223]
56. Sanders WE, Wilson RW, Ballantyne CM, Beaudet AL. Molecular cloning and analysis of in vivo expression of murine P-selectin. *Blood* 1992;80:795–800. [PubMed: 1379089]
57. Pan J, Xia L, McEver RP. Comparison of promoters for the murine and human P-selectin genes suggests species-specific and conserved mechanisms for transcriptional regulation in endothelial cells. *J Biol Chem* 1998;273:10058–67. [PubMed: 9545353]
58. Bullard DC, Kunkel EJ, Kubo H, Hicks MJ, Lorenzo I, Doyle NA, Doerschuk CM, Ley K, Beaudet AL. Infectious susceptibility and severe deficiency of leukocyte rolling and recruitment in E-selectin and P-selectin double mutant mice. *J Exp Med* 1996;183:2329–36. [PubMed: 8642341]
59. Kunkel EJ, Ley K. Distinct phenotype of E-selectin-deficient mice. E-selectin is required for slow leukocyte rolling in vivo. *Circ Res* 1996;79:1196–204. [PubMed: 8943958]
60. Ley K, Bullard DC, Arbones ML, Bosse R, Vestweber D, Tedder TF, Beaudet AL. Sequential contribution of L- and P-selectin to leukocyte rolling in vivo. *J Exp Med* 1995;181:669–75. [PubMed: 7530761]
61. Bevilacqua MP, Pober JS, Mendrick DL, Cotran RS, Gimbrone MA Jr. Identification of an inducible endothelial-leukocyte adhesion molecule. *Proc Natl Acad Sci U S A* 1987;84:9238–42. [PubMed: 2827173]
62. Bevilacqua MP, Stengelin S, Gimbrone MA Jr, Seed B. Endothelial leukocyte adhesion molecule 1: an inducible receptor for neutrophils related to complement regulatory proteins and lectins. *Science* 1989;243:1160–5. [PubMed: 2466335]
63. Subramaniam M, Koedam JA, Wagner DD. Divergent fates of P- and E-selectins after their expression on the plasma membrane. *Mol Biol Cell* 1993;4:791–801. [PubMed: 7694691]
64. Henseleit U, Steinbrink K, Goebeler M, Roth J, Vestweber D, Sorg C, Sunderkotter C. E-selectin expression in experimental models of inflammation in mice. *J Pathol* 1996;180:317–25. [PubMed: 8958812]
65. Keelan ET, Licence ST, Peters AM, Binns RM, Haskard DO. Characterization of E-selectin expression in vivo with use of a radiolabeled monoclonal antibody. *Am J Physiol* 1994;266:H278–90. [PubMed: 7508207]
66. Picker LJ, Kishimoto TK, Smith CW, Warnock RA, Butcher EC. ELAM-1 is an adhesion molecule for skin-homing T cells. *Nature* 1991;349:796–9. [PubMed: 1705666]
67. Ley K, Allietta M, Bullard DC, Morgan S. Importance of E-selectin for firm leukocyte adhesion in vivo. *Circ Res* 1998;83:287–94. [PubMed: 9710121]
68. Milstone DS, Fukumura D, Padgett RC, O'Donnell PE, Davis VM, Benavidez OJ, Monsky WL, Melder RJ, Jain RK, Gimbrone MA Jr. Mice lacking E-selectin show normal numbers of rolling leukocytes but reduced leukocyte stable arrest on cytokine-activated microvascular endothelium. *Microcirculation* 1998;5:153–71. [PubMed: 9789256]
69. Zarbock A, Lowell CA, Ley K. Spleen tyrosine kinase Syk is necessary for E-selectin-induced alpha (L)beta(2) integrin-mediated rolling on intercellular adhesion molecule-1. *Immunity* 2007;26:773–83. [PubMed: 17543554]

70. Picker LJ, Warnock RA, Burns AR, Doerschuk CM, Berg EL, Butcher EC. The neutrophil selectin LECAM-1 presents carbohydrate ligands to the vascular selectins ELAM-1 and GMP-140. *Cell* 1991;66:921–33. [PubMed: 1716182]
71. Hidalgo A, Peired AJ, Wild MK, Vestweber D, Frenette PS. Complete identification of E-selectin ligands on neutrophils reveals distinct functions of PSGL-1, ESL-1, and CD44. *Immunity* 2007;26:477–89. [PubMed: 17442598]
72. Katayama Y, Hidalgo A, Chang J, Peired A, Frenette PS. CD44 is a physiological E-selectin ligand on neutrophils. *J Exp Med* 2005;201:1183–9. [PubMed: 15824084]
73. Steegmaier M, Levinovitz A, Isenmann S, Borges E, Lenter M, Kocher HP, Kleuser B, Vestweber D. The E-selectin-ligand ESL-1 is a variant of a receptor for fibroblast growth factor. *Nature* 1995;373:615–20. [PubMed: 7531823]
74. Alon R, Feizi T, Yuen CT, Fuhlbrigge RC, Springer TA. Glycolipid ligands for selectins support leukocyte tethering and rolling under physiologic flow conditions. *J Immunol* 1995;154:5356–66. [PubMed: 7537307]
75. Jung U, Norman KE, Scharffetter-Kochanek K, Beaudet AL, Ley K. Transit time of leukocytes rolling through venules controls cytokine-induced inflammatory cell recruitment in vivo. *J Clin Invest* 1998;102:1526–33. [PubMed: 9788965]
76. Smith MJ, Berg EL, Lawrence MB. A direct comparison of selectin-mediated transient, adhesive events using high temporal resolution. *Biophys J* 1999;77:3371–83. [PubMed: 10585960]
77. Robert C, Kupper TS. Inflammatory skin diseases, T cells, and immune surveillance. *N Engl J Med* 1999;341:1817–28. [PubMed: 10588968]
78. Van Eeden S, Miyagashima R, Haley L, Hogg JC. L-selectin expression increases on peripheral blood polymorphonuclear leukocytes during active marrow release. *Am J Respir Crit Care Med* 1995;151:500–7. [PubMed: 7531098]
79. Van Eeden SF, Bicknell S, Walker BA, Hogg JC. Polymorphonuclear leukocytes L-selectin expression decreases as they age in circulation. *Am J Physiol* 1997;272:H401–8. [PubMed: 9038962]
80. Jutila MA, Rott L, Berg EL, Butcher EC. Function and regulation of the neutrophil MEL-14 antigen in vivo: comparison with LFA-1 and MAC-1. *J Immunol* 1989;143:3318–24. [PubMed: 2553811]
81. Kishimoto TK, Jutila MA, Berg EL, Butcher EC. Neutrophil Mac-1 and MEL-14 adhesion proteins inversely regulated by chemotactic factors. *Science* 1989;245:1238–41. [PubMed: 2551036]
82. Walzog B, Seifert R, Zakrzewicz A, Gaehtgens P, Ley K. Cross-linking of CD18 in human neutrophils induces an increase of intracellular free Ca²⁺, exocytosis of azurophilic granules, quantitative up-regulation of CD18, shedding of L-selectin, and actin polymerization. *J Leukoc Biol* 1994;56:625–35. [PubMed: 7525820]
83. Smalley DM, Ley K. L-selectin: mechanisms and physiological significance of ectodomain cleavage. *J Cell Mol Med* 2005;9:255–66. [PubMed: 15963248]
84. Jutila MA, Kishimoto TK, Butcher EC. Regulation and lectin activity of the human neutrophil peripheral lymph node homing receptor. *Blood* 1990;76:178–83. [PubMed: 2194589]
85. Bührer C, Berlin C, Thiele HG, Hamann A. Lymphocyte activation and expression of the human leucocyte-endothelial cell adhesion molecule 1 (Leu-8/TQ1 antigen). *Immunology* 1990;71:442–8. [PubMed: 1702752]
86. Schleiffenbaum B, Spertini O, Tedder TF. Soluble L-selectin is present in human plasma at high levels and retains functional activity. *J Cell Biol* 1992;119:229–38. [PubMed: 1382078]
87. Galkina E, Tanousis K, Preece G, Tolaini M, Kioussis D, Florey O, Haskard DO, Tedder TF, Ager A. L-selectin shedding does not regulate constitutive T cell trafficking but controls the migration pathways of antigen-activated T lymphocytes. *J Exp Med* 2003;198:1323–35. [PubMed: 14597735]
88. Rivera-Nieves J, Olson T, Bamias G, Bruce A, Solga M, Knight RF, Hoang S, Cominelli F, Ley K. L-selectin, alpha 4 beta 1, and alpha 4 beta 7 integrins participate in CD4⁺ T cell recruitment to chronically inflamed small intestine. *J Immunol* 2005;174:2343–52. [PubMed: 15699171]
89. Moore KL, Stults NL, Diaz S, Smith DF, Cummings RD, Varki A, McEver RP. Identification of a specific glycoprotein ligand for P-selectin (CD62) on myeloid cells. *J Cell Biol* 1992;118:445–56. [PubMed: 1378449]

90. Sako D, Chang XJ, Barone KM, Vachino G, White HM, Shaw G, Veldman GM, Bean KM, Ahern TJ, Furie B, et al. Expression cloning of a functional glycoprotein ligand for P-selectin. *Cell* 1993;75:1179–86. [PubMed: 7505206]
91. Rivera-Nieves J, Burcin TL, Olson TS, Morris MA, McDuffie M, Cominelli F, Ley K. Critical role of endothelial P-selectin glycoprotein ligand 1 in chronic murine ileitis. *J Exp Med* 2006;203:907–17. [PubMed: 16567389]
92. Vowinkel T, Wood KC, Stokes KY, Russell J, Tailor A, Anthoni C, Senninger N, Kriegelstein CF, Granger DN. Mechanisms of platelet and leukocyte recruitment in experimental colitis. *Am J Physiol Gastrointest Liver Physiol* 2007;293:G1054–60. [PubMed: 17884975]
93. da Costa Martins P, Garcia-Vallejo JJ, van Thienen JV, Fernandez-Borja M, van Gils JM, Beckers C, Horrevoets AJ, Hordijk PL, Zwaginga JJ. P-selectin glycoprotein ligand-1 is expressed on endothelial cells and mediates monocyte adhesion to activated endothelium. *Arterioscler Thromb Vasc Biol* 2007;27:1023–9. [PubMed: 17322099]
94. Moore KL, Patel KD, Bruehl RE, Li F, Johnson DA, Lichenstein HS, Cummings RD, Bainton DF, McEver RP. P-selectin glycoprotein ligand-1 mediates rolling of human neutrophils on P-selectin. *J Cell Biol* 1995;128:661–71. [PubMed: 7532174]
95. Pouyani T, Seed B. PSGL-1 recognition of P-selectin is controlled by a tyrosine sulfation consensus at the PSGL-1 amino terminus. *Cell* 1995;83:333–43. [PubMed: 7585950]
96. Sako D, Comess KM, Barone KM, Camphausen RT, Cumming DA, Shaw GD. A sulfated peptide segment at the amino terminus of PSGL-1 is critical for P-selectin binding. *Cell* 1995;83:323–31. [PubMed: 7585949]
97. Norgard KE, Moore KL, Diaz S, Stults NL, Ushiyama S, McEver RP, Cummings RD, Varki A. Characterization of a specific ligand for P-selectin on myeloid cells. A minor glycoprotein with sialylated O-linked oligosaccharides. *J Biol Chem* 1993;268:12764–74. [PubMed: 7685350]
98. Li F, Wilkins PP, Crawley S, Weinstein J, Cummings RD, McEver RP. Post-translational modifications of recombinant P-selectin glycoprotein ligand-1 required for binding to P- and E-selectin. *J Biol Chem* 1996;271:3255–64. [PubMed: 8621728]
99. Natsuka S, Gersten KM, Zenita K, Kannagi R, Lowe JB. Molecular cloning of a cDNA encoding a novel human leukocyte alpha-1,3-fucosyltransferase capable of synthesizing the sialyl Lewis x determinant. *J Biol Chem* 1994;269:16789–94. [PubMed: 8207002]
100. Sasaki K, Kurata K, Funayama K, Nagata M, Watanabe E, Ohta S, Hanai N, Nishi T. Expression cloning of a novel alpha 1,3-fucosyltransferase that is involved in biosynthesis of the sialyl Lewis x carbohydrate determinants in leukocytes. *J Biol Chem* 1994;269:14730–7. [PubMed: 8182079]
101. Snapp KR, Wagers AJ, Craig R, Stoolman LM, Kansas GS. P-selectin glycoprotein ligand-1 is essential for adhesion to P-selectin but not E-selectin in stably transfected hematopoietic cell lines. *Blood* 1997;89:896–901. [PubMed: 9028320]
102. Austrup F, Vestweber D, Borges E, Lohning M, Brauer R, Herz U, Renz H, Hallmann R, Scheffold A, Radbruch A, Hamann A. P- and E-selectin mediate recruitment of T-helper-1 but not T-helper-2 cells into inflamed tissues. *Nature* 1997;385:81–3. [PubMed: 8985251]
103. Knibbs RN, Craig RA, Natsuka S, Chang A, Cameron M, Lowe JB, Stoolman LM. The fucosyltransferase FucT-VII regulates E-selectin ligand synthesis in human T cells. *J Cell Biol* 1996;133:911–20. [PubMed: 8666674]
104. Lim YC, Henault L, Wagers AJ, Kansas GS, Luscinskas FW, Lichtman AH. Expression of functional selectin ligands on Th cells is differentially regulated by IL-12 and IL-4. *J Immunol* 1999;162:3193–201. [PubMed: 10092770]
105. Wagers AJ, Lowe JB, Kansas GS. An important role for the alpha 1,3 fucosyltransferase, FucT-VII, in leukocyte adhesion to E-selectin. *Blood* 1996;88:2125–32. [PubMed: 8822932]
106. Maly P, Thall A, Petryniak B, Rogers CE, Smith PL, Marks RM, Kelly RJ, Gersten KM, Cheng G, Saunders TL, Camper SA, Camphausen RT, Sullivan FX, Isogai Y, Hindsaul O, von Andrian UH, Lowe JB. The alpha(1,3)fucosyltransferase Fuc-TVII controls leukocyte trafficking through an essential role in L-, E-, and P-selectin ligand biosynthesis. *Cell* 1996;86:643–53. [PubMed: 8752218]

107. Frenette PS, Mayadas TN, Rayburn H, Hynes RO, Wagner DD. Susceptibility to infection and altered hematopoiesis in mice deficient in both P- and E-selectins. *Cell* 1996;84:563–74. [PubMed: 8598043]
108. Karsan A, Cornejo CJ, Winn RK, Schwartz BR, Way W, Lannir N, Gershoni-Baruch R, Etzioni A, Ochs HD, Harlan JM. Leukocyte Adhesion Deficiency Type II is a generalized defect of de novo GDP-fucose biosynthesis. Endothelial cell fucosylation is not required for neutrophil rolling on human nonlymphoid endothelium. *J Clin Invest* 1998;101:2438–45. [PubMed: 9616215]
109. Etzioni A, Frydman M, Pollack S, Avidor I, Phillips ML, Paulson JC, Gershoni-Baruch R. Brief report: recurrent severe infections caused by a novel leukocyte adhesion deficiency. *N Engl J Med* 1992;327:1789–92. [PubMed: 1279426]
110. Borges E, Eytner R, Moll T, Steegmaier M, Campbell MA, Ley K, Mossmann H, Vestweber D. The P-selectin glycoprotein ligand-1 is important for recruitment of neutrophils into inflamed mouse peritoneum. *Blood* 1997;90:1934–42. [PubMed: 9292527]
111. Norman KE, Moore KL, McEver RP, Ley K. Leukocyte rolling in vivo is mediated by P-selectin glycoprotein ligand-1. *Blood* 1995;86:4417–21. [PubMed: 8541529]
112. Spertini O, Cordey AS, Monai N, Giuffre L, Schapira M. P-selectin glycoprotein ligand 1 is a ligand for L-selectin on neutrophils, monocytes, and CD34+ hematopoietic progenitor cells. *J Cell Biol* 1996;135:523–31. [PubMed: 8896607]
113. Walcheck B, Moore KL, McEver RP, Kishimoto TK. Neutrophil-neutrophil interactions under hydrodynamic shear stress involve L-selectin and PSGL-1. A mechanism that amplifies initial leukocyte accumulation of P-selectin in vitro. *J Clin Invest* 1996;98:1081–7. [PubMed: 8787668]
114. Alon R, Fuhlbrigge RC, Finger EB, Springer TA. Interactions through L-selectin between leukocytes and adherent leukocytes nucleate rolling adhesions on selectins and VCAM-1 in shear flow. *J Cell Biol* 1996;135:849–65. [PubMed: 8909556]
115. Ramos CL, Smith MJ, Snapp KR, Kansas GS, Stickney GW, Ley K, Lawrence MB. Functional characterization of L-selectin ligands on human neutrophils and leukemia cell lines: evidence for mucinlike ligand activity distinct from P-selectin glycoprotein ligand-1. *Blood* 1998;91:1067–75. [PubMed: 9446670]
116. Veerman KM, Williams MJ, Uchimura K, Singer MS, Merzaban JS, Naus S, Carlow DA, Owen P, Rivera-Nieves J, Rosen SD, Ziltener HJ. Interaction of the selectin ligand PSGL-1 with chemokines CCL21 and CCL19 facilitates efficient homing of T cells to secondary lymphoid organs. *Nat Immunol* 2007;8:532–9. [PubMed: 17401367]
117. Lasky LA, Singer MS, Dowbenko D, Imai Y, Henzel WJ, Grimley C, Fennie C, Gillett N, Watson SR, Rosen SD. An endothelial ligand for L-selectin is a novel mucin-like molecule. *Cell* 1992;69:927–38. [PubMed: 1376638]
118. Baumhueter S, Singer MS, Henzel W, Hemmerich S, Renz M, Rosen SD, Lasky LA. Binding of L-selectin to the vascular sialomucin CD34. *Science* 1993;262:436–8. [PubMed: 7692600]
119. Sassetti C, Tangemann K, Singer MS, Kershaw DB, Rosen SD. Identification of podocalyxin-like protein as a high endothelial venule ligand for L-selectin: parallels to CD34. *J Exp Med* 1998;187:1965–75. [PubMed: 9625756]
120. Hemmerich S, Butcher EC, Rosen SD. Sulfation-dependent recognition of high endothelial venules (HEV)-ligands by L-selectin and MECA 79, and adhesion-blocking monoclonal antibody. *J Exp Med* 1994;180:2219–26. [PubMed: 7525849]
121. Streeter PR, Rouse BT, Butcher EC. Immunohistologic and functional characterization of a vascular addressin involved in lymphocyte homing into peripheral lymph nodes. *J Cell Biol* 1988;107:1853–62. [PubMed: 2460470]
122. Uchimura K, Rosen SD. Sulfated L-selectin ligands as a therapeutic target in chronic inflammation. *Trends Immunol* 2006;27:559–65. [PubMed: 17049924]
123. Uchimura K, Gauguier JM, Singer MS, Tsay D, Kannagi R, Muramatsu T, von Andrian UH, Rosen SD. A major class of L-selectin ligands is eliminated in mice deficient in two sulfotransferases expressed in high endothelial venules. *Nat Immunol* 2005;6:1105–13. [PubMed: 16227986]
124. Rosen SD. Endothelial ligands for L-selectin: from lymphocyte recirculation to allograft rejection. *Am J Pathol* 1999;155:1013–20. [PubMed: 10514381]

125. Lee JK, Bhakta S, Rosen SD, Hemmerich S. Cloning and characterization of a mammalian N-acetylglucosamine-6-sulfotransferase that is highly restricted to intestinal tissue. *Biochem Biophys Res Commun* 1999;263:543–9. [PubMed: 10491328]
126. Delmotte P, Degroote S, Lafitte JJ, Lamblin G, Perini JM, Roussel P. Tumor necrosis factor alpha increases the expression of glycosyltransferases and sulfotransferases responsible for the biosynthesis of sialylated and/or sulfated Lewis x epitopes in the human bronchial mucosa. *J Biol Chem* 2002;277:424–31. [PubMed: 11679593]
127. Drayton DL, Ying X, Lee J, Lesslauer W, Ruddle NH. Ectopic LT alpha beta directs lymphoid organ neogenesis with concomitant expression of peripheral node addressin and a HEV-restricted sulfotransferase. *J Exp Med* 2003;197:1153–63. [PubMed: 12732657]
128. Briskin MJ, McEvoy LM, Butcher EC. MAdCAM-1 has homology to immunoglobulin and mucin-like adhesion receptors and to IgA1. *Nature* 1993;363:461–4. [PubMed: 8502297]
129. Berlin C, Berg EL, Briskin MJ, Andrew DP, Kilshaw PJ, Holzmann B, Weissman IL, Hamann A, Butcher EC. Alpha 4 beta 7 integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1. *Cell* 1993;74:185–95. [PubMed: 7687523]
130. Berg EL, McEvoy LM, Berlin C, Bargatze RF, Butcher EC. L-selectin-mediated lymphocyte rolling on MAdCAM-1. *Nature* 1993;366:695–8. [PubMed: 7505053]
131. Walcheck B, Watts G, Jutila MA. Bovine gamma/delta T cells bind E-selectin via a novel glycoprotein receptor: first characterization of a lymphocyte/E-selectin interaction in an animal model. *J Exp Med* 1993;178:853–63. [PubMed: 7688790]
132. Matsumoto M, Shigeta A, Furukawa Y, Tanaka T, Miyasaka M, Hirata T. CD43 collaborates with P-selectin glycoprotein ligand-1 to mediate E-selectin-dependent T cell migration into inflamed skin. *J Immunol* 2007;178:2499–506. [PubMed: 17277158]
133. Fuhlbrigge RC, King SL, Sackstein R, Kupper TS. CD43 is a ligand for E-selectin on CLA+ human T cells. *Blood* 2006;107:1421–6. [PubMed: 16269612]
134. Ley K. The role of selectins in inflammation and disease. *Trends Mol Med* 2003;9:263–8. [PubMed: 12829015]
135. Wang K, Zhou X, Zhou Z, Tarakji K, Qin JX, Sitges M, Shiota T, Forudi F, Schaub RG, Kumar A, Penn MS, Topol EJ, Lincoff AM. Recombinant soluble P-selectin glycoprotein ligand-Ig (rPSGL-Ig) attenuates infarct size and myeloperoxidase activity in a canine model of ischemia-reperfusion. *Thromb Haemost* 2002;88:149–54. [PubMed: 12152656]
136. Norman KE, Anderson GP, Kolb HC, Ley K, Ernst B. Sialyl Lewis(x) (sLe(x)) and an sLe(x) mimetic, CGP69669A, disrupt E-selectin-dependent leukocyte rolling in vivo. *Blood* 1998;91:475–83. [PubMed: 9427700]
137. Kaneider NC, Leger AJ, Kuliopulos A. Therapeutic targeting of molecules involved in leukocyte-endothelial cell interactions. *Febs J* 2006;273:4416–24. [PubMed: 16956369]
138. Berlin C, Bargatze RF, Campbell JJ, von Andrian UH, Szabo MC, Hasslen SR, Nelson RD, Berg EL, Erlandsen SL, Butcher EC. alpha 4 integrins mediate lymphocyte attachment and rolling under physiologic flow. *Cell* 1995;80:413–22. [PubMed: 7532110]
139. Luo BH, Carman CV, Springer TA. Structural basis of integrin regulation and signaling. *Annu Rev Immunol* 2007;25:619–47. [PubMed: 17201681]
140. Hughes PE, Pfaff M. Integrin affinity modulation. *Trends Cell Biol* 1998;8:359–64. [PubMed: 9728397]
141. Ley K. Arrest chemokines. *Microcirculation* 2003;10:289–95. [PubMed: 12851646]
142. Van der Vieren M, Le Trong H, Wood CL, Moore PF, St John T, Staunton DE, Gallatin WM. A novel leukointegrin, alpha d beta 2, binds preferentially to ICAM-3. *Immunity* 1995;3:683–90. [PubMed: 8777714]
143. Anderson DC, Springer TA. Leukocyte adhesion deficiency: an inherited defect in the Mac-1, LFA-1, and p150,95 glycoproteins. *Annu Rev Med* 1987;38:175–94. [PubMed: 3555290]
144. Scharffetter-Kochanek K, Lu H, Norman K, van Nood N, Munoz F, Grabbe S, McArthur M, Lorenzo I, Kaplan S, Ley K, Smith CW, Montgomery CA, Rich S, Beaudet AL. Spontaneous skin ulceration and defective T cell function in CD18 null mice. *J Exp Med* 1998;188:119–31. [PubMed: 9653089]
145. Marlin SD, Springer TA. Purified intercellular adhesion molecule-1 (ICAM-1) is a ligand for lymphocyte function-associated antigen 1 (LFA-1). *Cell* 1987;51:813–9. [PubMed: 3315233]

146. Staunton DE, Dustin ML, Springer TA. Functional cloning of ICAM-2, a cell adhesion ligand for LFA-1 homologous to ICAM-1. *Nature* 1989;339:61–4. [PubMed: 2497351]
147. Schmits R, Kundig TM, Baker DM, Shumaker G, Simard JJ, Duncan G, Wakeham A, Shahinian A, van der Heiden A, Bachmann MF, Ohashi PS, Mak TW, Hickstein DD. LFA-1-deficient mice show normal CTL responses to virus but fail to reject immunogenic tumor. *J Exp Med* 1996;183:1415–26. [PubMed: 8666900]
148. Zen K, Cui LB, Zhang CY, Liu Y. Critical role of Mac-1 sialyl lewis x moieties in regulating neutrophil degranulation and transmigration. *J Mol Biol* 2007;374:54–63. [PubMed: 17919658]
149. Ross GD. Regulation of the adhesion versus cytotoxic functions of the Mac-1/CR3/alphaMbeta2-integrin glycoprotein. *Crit Rev Immunol* 2000;20:197–222. [PubMed: 10968371]
150. Hynes RO. Integrins: a family of cell surface receptors. *Cell* 1987;48:549–54. [PubMed: 3028640]
151. Hemler ME, Huang C, Schwarz L. The VLA protein family. Characterization of five distinct cell surface heterodimers each with a common 130,000 molecular weight beta subunit. *J Biol Chem* 1987;262:3300–9. [PubMed: 3546305]
152. Lee SJ, Benveniste EN. Adhesion molecule expression and regulation on cells of the central nervous system. *J Neuroimmunol* 1999;98:77–88. [PubMed: 10430040]
153. Luque A, Gomez M, Puzon W, Takada Y, Sanchez-Madrid F, Cabanas C. Activated conformations of very late activation integrins detected by a group of antibodies (HUTS) specific for a novel regulatory region (355–425) of the common beta 1 chain. *J Biol Chem* 1996;271:11067–75. [PubMed: 8626649]
154. Steeber DA, Tang ML, Zhang XQ, Muller W, Wagner N, Tedder TF. Efficient lymphocyte migration across high endothelial venules of mouse Peyer's patches requires overlapping expression of L-selectin and beta7 integrin. *J Immunol* 1998;161:6638–47. [PubMed: 9862692]
155. Hynes RO. Targeted mutations in cell adhesion genes: what have we learned from them? *Dev Biol* 1996;180:402–12. [PubMed: 8954713]
156. Arroyo AG, Taverna D, Whittaker CA, Strauch UG, Bader BL, Rayburn H, Crowley D, Parker CM, Hynes RO. In vivo roles of integrins during leukocyte development and traffic: insights from the analysis of mice chimeric for alpha 5, alpha v, and alpha 4 integrins. *J Immunol* 2000;165:4667–75. [PubMed: 11035110]
157. Arroyo AG, Yang JT, Rayburn H, Hynes RO. Differential requirements for alpha4 integrins during fetal and adult hematopoiesis. *Cell* 1996;85:997–1008. [PubMed: 8674127]
158. Coombes JL, Siddiqui KR, Arancibia-Carcamo CV, Hall J, Sun CM, Belkaid Y, Powrie F. A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. *J Exp Med* 2007;204:1757–64. [PubMed: 17620361]
159. Izcue A, Coombes JL, Powrie F. Regulatory T cells suppress systemic and mucosal immune activation to control intestinal inflammation. *Immunol Rev* 2006;212:256–71. [PubMed: 16903919]
160. Johansson-Lindbom B, Agace WW. Generation of gut-homing T cells and their localization to the small intestinal mucosa. *Immunol Rev* 2007;215:226–42. [PubMed: 17291292]
161. Dustin ML, Rothlein R, Bhan AK, Dinarello CA, Springer TA. Induction by IL 1 and interferon-gamma: tissue distribution, biochemistry, and function of a natural adherence molecule (ICAM-1). *J Immunol* 1986;137:245–54. [PubMed: 3086451]
162. Diamond MS, Staunton DE, Marlin SD, Springer TA. Binding of the integrin Mac-1 (CD11b/CD18) to the third immunoglobulin-like domain of ICAM-1 (CD54) and its regulation by glycosylation. *Cell* 1991;65:961–71. [PubMed: 1675157]
163. Shang XZ, Issekutz AC. Contribution of CD11a/CD18, CD11b/CD18, ICAM-1 (CD54) and -2 (CD102) to human monocyte migration through endothelium and connective tissue fibroblast barriers. *Eur J Immunol* 1998;28:1970–9. [PubMed: 9645379]
164. Knorr R, Dustin ML. The lymphocyte function-associated antigen 1 I domain is a transient binding module for intercellular adhesion molecule (ICAM)-1 and ICAM-3 in hydrodynamic flow. *J Exp Med* 1997;186:719–30. [PubMed: 9271587]
165. Yacyshyn B, Chey WY, Wedel MK, Yu RZ, Paul D, Chuang E. A randomized, double-masked, placebo-controlled study of alicaforsen, an antisense inhibitor of intercellular adhesion molecule 1,

- for the treatment of subjects with active Crohn's disease. *Clin Gastroenterol Hepatol* 2007;5:215–20. [PubMed: 17296530]
166. Diacovo TG, deFougerolles AR, Bainton DF, Springer TA. A functional integrin ligand on the surface of platelets: intercellular adhesion molecule-2. *J Clin Invest* 1994;94:1243–51. [PubMed: 8083366]
 167. Gerwin N, Gonzalo JA, Lloyd C, Coyle AJ, Reiss Y, Banu N, Wang B, Xu H, Avraham H, Engelhardt B, Springer TA, Gutierrez-Ramos JC. Prolonged eosinophil accumulation in allergic lung interstitium of ICAM-2 deficient mice results in extended hyperresponsiveness. *Immunity* 1999;10:9–19. [PubMed: 10023766]
 168. Osborn L, Hession C, Tizard R, Vassallo C, Lühowskyj S, Chi-Rosso G, Lobb R. Direct expression cloning of vascular cell adhesion molecule 1, a cytokine-induced endothelial protein that binds to lymphocytes. *Cell* 1989;59:1203–11. [PubMed: 2688898]
 169. Rice GE, Bevilacqua MP. An inducible endothelial cell surface glycoprotein mediates melanoma adhesion. *Science* 1989;246:1303–6. [PubMed: 2588007]
 170. Carlos TM, Schwartz BR, Kovach NL, Yee E, Rosa M, Osborn L, Chi-Rosso G, Newman B, Lobb R, et al. Vascular cell adhesion molecule-1 mediates lymphocyte adherence to cytokine-activated cultured human endothelial cells. *Blood* 1990;76:965–70. [PubMed: 1697486]
 171. Elices MJ, Osborn L, Takada Y, Crouse C, Lühowskyj S, Hemler ME, Lobb RR. VCAM-1 on activated endothelium interacts with the leukocyte integrin VLA-4 at a site distinct from the VLA-4/fibronectin binding site. *Cell* 1990;60:577–84. [PubMed: 1689216]
 172. Rice GE, Munro JM, Bevilacqua MP. Inducible cell adhesion molecule 110 (ICAM-110) is an endothelial receptor for lymphocytes. A CD11/CD18-independent adhesion mechanism. *J Exp Med* 1990;171:1369–74. [PubMed: 1691264]
 173. Manka DR, Wiegman P, Din S, Sanders JM, Green SA, Gimble LW, Ragosta M, Powers ER, Ley K, Sarembock IJ. Arterial injury increases expression of inflammatory adhesion molecules in the carotid arteries of apolipoprotein-E-deficient mice. *J Vasc Res* 1999;36:372–8. [PubMed: 10559677]
 174. Cybulsky MI, Gimbrone MA Jr. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science* 1991;251:788–91. [PubMed: 1990440]
 175. Koni PA, Joshi SK, Temann UA, Olson D, Burkly L, Flavell RA. Conditional vascular cell adhesion molecule 1 deletion in mice: impaired lymphocyte migration to bone marrow. *J Exp Med* 2001;193:741–54. [PubMed: 11257140]
 176. Leuker CE, Labow M, Muller W, Wagner N. Neonatally induced inactivation of the vascular cell adhesion molecule 1 gene impairs B cell localization and T cell-dependent humoral immune response. *J Exp Med* 2001;193:755–68. [PubMed: 11257141]
 177. Briskin M, Winsor-Hines D, Shyjan A, Cochran N, Bloom S, Wilson J, McEvoy LM, Butcher EC, Kassam N, Mackay CR, Newman W, Ringler DJ. Human mucosal addressin cell adhesion molecule-1 is preferentially expressed in intestinal tract and associated lymphoid tissue. *Am J Pathol* 1997;151:97–110. [PubMed: 9212736]
 178. Connor EM, Eppihimer MJ, Morise Z, Granger DN, Grisham MB. Expression of mucosal addressin cell adhesion molecule-1 (MAdCAM-1) in acute and chronic inflammation. *J Leukoc Biol* 1999;65:349–55. [PubMed: 10080539]
 179. Hanninen A, Taylor C, Streeter PR, Stark LS, Sarte JM, Shizuru JA, Simell O, Michie SA. Vascular addressins are induced on islet vessels during insulinitis in nonobese diabetic mice and are involved in lymphoid cell binding to islet endothelium. *J Clin Invest* 1993;92:2509–15. [PubMed: 7693764]
 180. Shelley CS, Farokhzad OC, Arnaout MA. Identification of cell-specific and developmentally regulated nuclear factors that direct myeloid and lymphoid expression of the CD11a gene. *Proc Natl Acad Sci U S A* 1993;90:5364–8. [PubMed: 8099450]
 181. Sikorski EE, Hallmann R, Berg EL, Butcher EC. The Peyer's patch high endothelial receptor for lymphocytes, the mucosal vascular addressin, is induced on a murine endothelial cell line by tumor necrosis factor-alpha and IL-1. *J Immunol* 1993;151:5239–50. [PubMed: 7693807]
 182. Strober W, Fuss IJ, Blumberg RS. The immunology of mucosal models of inflammation. *Annu Rev Immunol* 2002;20:495–549. [PubMed: 11861611]

183. Chalifoux LV, Bronson RT. Colonic adenocarcinoma associated with chronic colitis in cotton top marmosets, *Saguinus oedipus*. *Gastroenterology* 1981;80:942–6. [PubMed: 6781977]
184. Madara JL, Podolsky DK, King NW, Sehgal PK, Moore R, Winter HS. Characterization of spontaneous colitis in cotton-top tamarins (*Saguinus oedipus*) and its response to sulfasalazine. *Gastroenterology* 1985;88:13–9. [PubMed: 2856876]
185. MacPherson BR, Pfeiffer CJ. Experimental production of diffuse colitis in rats. *Digestion* 1978;17:135–50. [PubMed: 627326]
186. Morris GP, Beck PL, Herridge MS, Depew WT, Szewczuk MR, Wallace JL. Hapten-induced model of chronic inflammation and ulceration in the rat colon. *Gastroenterology* 1989;96:795–803. [PubMed: 2914642]
187. Axelsson LG, Landstrom E, Goldschmidt TJ, Gronberg A, Bylund-Fellenius AC. Dextran sulfate sodium (DSS) induced experimental colitis in immunodeficient mice: effects in CD4(+) -cell depleted, athymic and NK-cell depleted SCID mice. *Inflamm Res* 1996;45:181–91. [PubMed: 8741008]
188. Hodgson HJ, Potter BJ, Skinner J, Jewell DP. Immune-complex mediated colitis in rabbits. An experimental model *Gut* 1978;19:225–32.
189. Powrie F, Correa-Oliveira R, Mauze S, Coffman RL. Regulatory interactions between CD45RBhigh and CD45RBlow CD4+ T cells are important for the balance between protective and pathogenic cell-mediated immunity. *J Exp Med* 1994;179:589–600. [PubMed: 7905019]
190. Aranda R, Sydora BC, McAllister PL, Binder SW, Yang HY, Targan SR, Kronenberg M. Analysis of intestinal lymphocytes in mouse colitis mediated by transfer of CD4+, CD45RBhigh T cells to SCID recipients. *J Immunol* 1997;158:3464–73. [PubMed: 9120308]
191. Kuhn R, Lohler J, Rennick D, Rajewsky K, Muller W. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* 1993;75:263–74. [PubMed: 8402911]
192. Podolsky DK, Lobb R, King N, Benjamin CD, Pepinsky B, Sehgal P, deBeaumont M. Attenuation of colitis in the cotton-top tamarin by anti-alpha 4 integrin monoclonal antibody. *J Clin Invest* 1993;92:372–80. [PubMed: 7686922]
193. McCafferty DM, Smith CW, Granger DN, Kubes P. Intestinal inflammation in adhesion molecule-deficient mice: an assessment of P-selectin alone and in combination with ICAM-1 or E-selectin. *J Leukoc Biol* 1999;66:67–74. [PubMed: 10410991]
194. Sans M, Salas A, Soriano A, Prats N, Gironella M, Pizcueta P, Elena M, Anderson DC, Pique JM, Panes J. Differential role of selectins in experimental colitis. *Gastroenterology* 2001;120:1162–72. [PubMed: 11266380]
195. Gironella M, Molla M, Salas A, Soriano A, Sans M, Closa D, Engel P, Salas A, Pique JM, Panes J. The role of P-selectin in experimental colitis as determined by antibody immunoblockade and genetically deficient mice. *J Leukoc Biol* 2002;72:56–64. [PubMed: 12101263]
196. Wan MX, Riaz AA, Schramm R, Wang Y, Vestweber D, Menger MD, Thorlacius H. Leukocyte rolling is exclusively mediated by P-selectin in colonic venules. *Br J Pharmacol* 2002;135:1749–56. [PubMed: 11934816]
197. Yoshida N, Yamaguchi T, Nakagawa S, Nakamura Y, Naito Y, Yoshikawa T. Role of P-selectin and intercellular adhesion molecule-1 in TNB-induced colitis in rats. *Digestion* 2001;63 (Suppl 1): 81–6. [PubMed: 11173915]
198. Rijcken EM, Laukoetter MG, Anthoni C, Meier S, Mennigen R, Spiegel HU, Bruewer M, Senninger N, Vestweber D, Krieglstein CF. Immunoblockade of PSGL-1 attenuates established experimental murine colitis by reduction of leukocyte rolling. *Am J Physiol Gastrointest Liver Physiol* 2004;287:G115–24. [PubMed: 15001428]
199. Ostanin DV, Furr KL, Pavlick KP, Gray L, Kevil CG, Shukla D, D'Souza D, Hoffman JM, Grisham MB. T cell-associated CD18 but not CD62L, ICAM-1, or PSGL-1 is required for the induction of chronic colitis. *Am J Physiol Gastrointest Liver Physiol* 2007;292:G1706–14. [PubMed: 17332469]
200. Wallace JL, Higa A, McKnight GW, MacIntyre DE. Prevention and reversal of experimental colitis by a monoclonal antibody which inhibits leukocyte adherence. *Inflammation* 1992;16:343–54. [PubMed: 1326482]

201. Palmen MJ, Dijkstra CD, van der Ende MB, Pena AS, van Rees EP. Anti-CD11b/CD18 antibodies reduce inflammation in acute colitis in rats. *Clin Exp Immunol* 1995;101:351–6. [PubMed: 7648720]
202. Meenan J, Hommes DW, Mevissen M, Dijkhuizen S, Soule H, Moyle M, Buller HR, ten Kate FW, Tytgat GN, van Deventer SJ. Attenuation of the inflammatory response in an animal colitis model by neutrophil inhibitory factor, a novel beta 2-integrin antagonist. *Scand J Gastroenterol* 1996;31:786–91. [PubMed: 8858748]
203. Pavlick KP, Ostanin DV, Furr KL, Laroux FS, Brown CM, Gray L, Kevil CG, Grisham MB. Role of T-cell-associated lymphocyte function-associated antigen-1 in the pathogenesis of experimental colitis. *Int Immunol* 2006;18:389–98. [PubMed: 16415103]
204. Abdelbaqi M, Chidlow JH, Matthews KM, Pavlick KP, Barlow SC, Linscott AJ, Grisham MB, Fowler MR, Kevil CG. Regulation of dextran sodium sulfate induced colitis by leukocyte beta 2 integrins. *Lab Invest* 2006;86:380–90. [PubMed: 16482101]
205. Leon F, Contractor N, Fuss I, Marth T, Lahey E, Iwaki S, la Sala A, Hoffmann V, Strober W, Kelsall BL. Antibodies to complement receptor 3 treat established inflammation in murine models of colitis and a novel model of psoriasiform dermatitis. *J Immunol* 2006;177:6974–82. [PubMed: 17082612]
206. Hesterberg PE, Winsor-Hines D, Briskin MJ, Soler-Ferran D, Merrill C, Mackay CR, Newman W, Ringler DJ. Rapid resolution of chronic colitis in the cotton-top tamarin with an antibody to a gut-homing integrin alpha 4 beta 7. *Gastroenterology* 1996;111:1373–80. [PubMed: 8898653]
207. Picarella D, Hurlbut P, Rottman J, Shi X, Butcher E, Ringler DJ. Monoclonal antibodies specific for beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MAdCAM-1) reduce inflammation in the colon of scid mice reconstituted with CD45RB^{high} CD4⁺ T cells. *J Immunol* 1997;158:2099–106. [PubMed: 9036954]
208. Shigematsu T, Specian RD, Wolf RE, Grisham MB, Granger DN. MAdCAM mediates lymphocyte-endothelial cell adhesion in a murine model of chronic colitis. *Am J Physiol Gastrointest Liver Physiol* 2001;281:G1309–15. [PubMed: 11668040]
209. Farkas S, Hornung M, Sattler C, Edtinger K, Steinbauer M, Anthuber M, Schlitt HJ, Herfarth H, Geissler EK. Blocking MAdCAM-1 in vivo reduces leukocyte extravasation and reverses chronic inflammation in experimental colitis. *Int J Colorectal Dis* 2006;21:71–8. [PubMed: 15856265]
210. Sun FF, Lai PS, Yue G, Yin K, Nagele RG, Tong DM, Krzesicki RF, Chin JE, Wong PY. Pattern of cytokine and adhesion molecule mRNA in hapten-induced relapsing colon inflammation in the rat. *Inflammation* 2001;25:33–45. [PubMed: 11293664]
211. Kawachi S, Jennings S, Panes J, Cockrell A, Laroux FS, Gray L, Perry M, van der Heyde H, Balish E, Granger DN, Specian RA, Grisham MB. Cytokine and endothelial cell adhesion molecule expression in interleukin-10-deficient mice. *Am J Physiol Gastrointest Liver Physiol* 2000;278:G734–43. [PubMed: 10801266]
212. Kawachi S, Morise Z, Jennings SR, Conner E, Cockrell A, Laroux FS, Chervenak RP, Wolcott M, van der Heyde H, Gray L, Feng L, Granger DN, Specian RA, Grisham MB. Cytokine and adhesion molecule expression in SCID mice reconstituted with CD4⁺ T cells. *Inflamm Bowel Dis* 2000;6:171–80. [PubMed: 10961589]
213. Kato S, Hokari R, Matsuzaki K, Iwai A, Kawaguchi A, Nagao S, Miyahara T, Itoh K, Ishii H, Miura S. Amelioration of murine experimental colitis by inhibition of mucosal addressin cell adhesion molecule-1. *J Pharmacol Exp Ther* 2000;295:183–9. [PubMed: 10991977]
214. Sans M, Panes J, Ardite E, Elizalde JJ, Arce Y, Elena M, Palacin A, Fernandez-Checa JC, Anderson DC, Lobb R, Pique JM. VCAM-1 and ICAM-1 mediate leukocyte-endothelial cell adhesion in rat experimental colitis. *Gastroenterology* 1999;116:874–83. [PubMed: 10092309]
215. Angulo S, Llopis M, Antolin M, Gironella M, Sans M, Malagelada JR, Pique JM, Guarner F, Panes J. *Lactobacillus casei* prevents the upregulation of ICAM-1 expression and leukocyte recruitment in experimental colitis. *Am J Physiol Gastrointest Liver Physiol* 2006;291:G1155–62. [PubMed: 16901988]
216. Binion DG, West GA, Ina K, Ziats NP, Emancipator SN, Fiocchi C. Enhanced leukocyte binding by intestinal microvascular endothelial cells in inflammatory bowel disease. *Gastroenterology* 1997;112:1895–907. [PubMed: 9178682]

217. Binion DG, West GA, Volk EE, Drazba JA, Ziats NP, Petras RE, Fiocchi C. Acquired increase in leucocyte binding by intestinal microvascular endothelium in inflammatory bowel disease. *Lancet* 1998;352:1742–6. [PubMed: 9848350]
218. Wong PY, Yue G, Yin K, Miyasaka M, Lane CL, Manning AM, Anderson DC, Sun FF. Antibodies to intercellular adhesion molecule-1 ameliorate the inflammatory response in acetic acid-induced inflammatory bowel disease. *J Pharmacol Exp Ther* 1995;274:475–80. [PubMed: 7616434]
219. Taniguchi T, Tsukada H, Nakamura H, Kodama M, Fukuda K, Saito T, Miyasaka M, Seino Y. Effects of the anti-ICAM-1 monoclonal antibody on dextran sodium sulphate-induced colitis in rats. *J Gastroenterol Hepatol* 1998;13:945–9. [PubMed: 9794195]
220. Hamamoto N, Maemura K, Hirata I, Murano M, Sasaki S, Katsu K. Inhibition of dextran sulphate sodium (DSS)-induced colitis in mice by intracolonic administered antibodies against adhesion molecules (endothelial leucocyte adhesion molecule-1 (ELAM-1) or intercellular adhesion molecule-1 (ICAM-1)). *Clin Exp Immunol* 1999;117:462–8. [PubMed: 10469048]
221. Bennett CF, Kornbrust D, Henry S, Stecker K, Howard R, Cooper S, Dutson S, Hall W, Jacoby HI. An ICAM-1 antisense oligonucleotide prevents and reverses dextran sulfate sodium-induced colitis in mice. *J Pharmacol Exp Ther* 1997;280:988–1000. [PubMed: 9023316]
222. Bendjelloul F, Maly P, Mandys V, Jirkovska M, Prokesova L, Tuckova L, Tlaskalova-Hogenova H. Intercellular adhesion molecule-1 (ICAM-1) deficiency protects mice against severe forms of experimentally induced colitis. *Clin Exp Immunol* 2000;119:57–63. [PubMed: 10606964]
223. Kontoyiannis D, Pasparakis M, Pizarro TT, Cominelli F, Kollias G. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity* 1999;10:387–98. [PubMed: 10204494]
224. Matsumoto S, Okabe Y, Setoyama H, Takayama K, Ohtsuka J, Funahashi H, Imaoka A, Okada Y, Umesaki Y. Inflammatory bowel disease-like enteritis and caecitis in a senescence accelerated mouse P1/Yit strain. *Gut* 1998;43:71–8. [PubMed: 9771408]
225. Strober W, Nakamura K, Kitani A. The SAMP1/Yit mouse: another step closer to modeling human inflammatory bowel disease. *J Clin Invest* 2001;107:667–70. [PubMed: 11254665]
226. Pizarro TT, Arseneau KO, Bamias G, Cominelli F. Mouse models for the study of Crohn's disease. *Trends Mol Med* 2003;9:218–22. [PubMed: 12763527]
227. Kozaiwa K, Sugawara K, Smith MF Jr, Carl V, Yamschikov V, Belyea B, McEwen SB, Moskaluk CA, Pizarro TT, Cominelli F, McDuffie M. Identification of a quantitative trait locus for ileitis in a spontaneous mouse model of Crohn's disease: SAMP1/YitFc. *Gastroenterology* 2003;125:477–90. [PubMed: 12891551]
228. Kosiewicz MM, Nast CC, Krishnan A, Rivera-Nieves J, Moskaluk CA, Matsumoto S, Kozaiwa K, Cominelli F. Th1-type responses mediate spontaneous ileitis in a novel murine model of Crohn's disease. *J Clin Invest* 2001;107:695–702. [PubMed: 11254669]
229. Rivera-Nieves J, Bamias G, Vidrich A, Marini M, Pizarro TT, McDuffie MJ, Moskaluk CA, Cohn SM, Cominelli F. Emergence of perianal fistulizing disease in the SAMP1/YitFc mouse, a spontaneous model of chronic ileitis. *Gastroenterology* 2003;124:972–82. [PubMed: 12671894]
230. Marini M, Bamias G, Rivera-Nieves J, Moskaluk CA, Hoang SB, Ross WG, Pizarro TT, Cominelli F. TNF- α neutralization ameliorates the severity of murine Crohn's-like ileitis by abrogation of intestinal epithelial cell apoptosis. *Proc Natl Acad Sci U S A* 2003;100:8366–71. [PubMed: 12832622]
231. Burns RC, Rivera-Nieves J, Moskaluk CA, Matsumoto S, Cominelli F, Ley K. Antibody blockade of ICAM-1 and VCAM-1 ameliorates inflammation in the SAMP-1/Yit adoptive transfer model of Crohn's disease in mice. *Gastroenterology* 2001;121:1428–36. [PubMed: 11729122]
232. Bradley LM, Malo ME, Tonkonogy SL, Watson SR. L-selectin is not essential for nai-ve CD4 cell trafficking or development of primary responses in Peyer's patches. *Eur J Immunol* 1997;27:1140–6. [PubMed: 9174603]
233. Buhner C, Berlin C, Jablonski-Westrich D, Holzmann B, Thiele HG, Hamann A. Lymphocyte activation and regulation of three adhesion molecules with supposed function in homing: LECAM-1 (MEL-14 antigen), LPAM-1/2 (α 4-integrin) and CD44 (Pgp-1). *Scand J Immunol* 1992;35:107–20. [PubMed: 1370869]

234. van Zante A, Rosen SD. Sulphated endothelial ligands for L-selectin in lymphocyte homing and inflammation. *Biochem Soc Trans* 2003;31:313–7. [PubMed: 12653627]
235. Butcher EC, Williams M, Youngman K, Rott L, Briskin M. Lymphocyte trafficking and regional immunity. *Adv Immunol* 1999;72:209–53. [PubMed: 10361577]
236. Salmi M, Granfors K, MacDermott R, Jalkanen S. Aberrant binding of lamina propria lymphocytes to vascular endothelium in inflammatory bowel diseases. *Gastroenterology* 1994;106:596–605. [PubMed: 8119529]
237. Inoue T, Tsuzuki Y, Matsuzaki K, Matsunaga H, Miyazaki J, Hokari R, Okada Y, Kawaguchi A, Nagao S, Itoh K, Matsumoto S, Miura S. Blockade of PSGL-1 attenuates CD14+ monocytic cell recruitment in intestinal mucosa and ameliorates ileitis in SAMPl/Yit mice. *J Leukoc Biol* 2005;77:287–95. [PubMed: 15569697]
238. Kunkel EJ, Butcher EC. Chemokines and the tissue-specific migration of lymphocytes. *Immunity* 2002;16:1–4. [PubMed: 11825560]
239. Feagan BG, Greenberg GR, Wild G, Fedorak RN, Pare P, McDonald JW, Dube R, Cohen A, Steinhart AH, Landau S, Aguzzi RA, Fox IH, Vandervoort MK. Treatment of ulcerative colitis with a humanized antibody to the alpha4beta7 integrin. *N Engl J Med* 2005;352:2499–507. [PubMed: 15958805]
240. Iwata M, Hirakiyama A, Eshima Y, Kagechika H, Kato C, Song SY. Retinoic acid imprints gut-homing specificity on T cells. *Immunity* 2004;21:527–38. [PubMed: 15485630]
241. Mora JR, Bono MR, Manjunath N, Weninger W, Cavanagh LL, Roseblatt M, Von Andrian UH. Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells. *Nature* 2003;424:88–93. [PubMed: 12840763]
242. Mora JR, Cheng G, Picarella D, Briskin M, Buchanan N, von Andrian UH. Reciprocal and dynamic control of CD8 T cell homing by dendritic cells from skin- and gut-associated lymphoid tissues. *J Exp Med* 2005;201:303–16. [PubMed: 15642741]
243. Sandborn WJ, Hanauer SB. Infliximab in the treatment of Crohn's disease: a user's guide for clinicians. *Am J Gastroenterol* 2002;97:2962–72. [PubMed: 12492177]
244. Rivera-Nieves J, Ho J, Bamias G, Ivashkina N, Ley K, Oppermann M, Cominelli F. Antibody blockade of CCL25/CCR9 ameliorates early but not late chronic murine ileitis. *Gastroenterology* 2006;131:1518–29. [PubMed: 17101325]

Figure 1.

The original steps of the leukocyte adhesion cascade are shown in red: rolling, mediated by selectins, activation, mediated by chemokines, and arrest, mediated by integrins. This has been expanded to include additional steps: capture (or tethering), slow rolling, adhesion strengthening and spreading, intravascular crawling, and paracellular and transcellular transmigration. Crucial molecules involved are indicated in boxes. ESAM, endothelial cell-selective adhesion molecule; ICAM-1, intercellular adhesion molecule 1; JAM, junctional adhesion molecule; LFA-1, lymphocyte function-associated antigen 1 (also known as $\alpha_L\beta_2$ -integrin); Mac-1, macrophage antigen 1; MAdCAM-1, mucosal vascular addressin cell-adhesion molecule 1; PSGL-1, P-selectin glycoprotein ligand 1; PECAM-1, platelet/endothelial-cell adhesion molecule 1; PI3Kases, phosphoinositide 3-kinases; VCAM-1, vascular cell-adhesion molecule 1; VLA-4, very late antigen 4 (also known as $\alpha_4\beta_1$ -integrin)

Figure 2.

Selectins and their ligands. During the inflammatory response, leukocyte adhesion to endothelial cells is controlled by the binding of selectins to complementary carbohydrate ligands. All known selectin ligands relevant for lymphocyte trafficking are transmembrane glycoproteins, which present oligosaccharide structures to the selectins. Transient bond formations between the selectins and their ligands mediate the early steps of the adhesion cascade. All three selectins can recognize glycoproteins presenting the tetrasaccharide sialyl-LewisX (sialyl-CD15). This tetrasaccharide is found on all circulating myeloid cells and some activated T cells. It is composed of sialic acid, galactose, fucose, and N-acetyl-galactosamine. It is unclear how selectins achieve specific interactions with ligands, given this common carbohydrate recognition.

Figure 3.

The integrins are $\alpha\beta$ heterodimers; each subunit crosses the membrane once. Mammalian integrins form subfamilies that share common subunits that bind distinct ligands. Illustrated in color are the integrins that have been targeted for the treatment of human diseases. RGD: arginine-glycine-aspartic acid sequence found in some integrin ligands. C3bi: Complement 3b inactivated

Figure 4. Immunoglobulin superfamily. Members of this family contain two to seven immunoglobulin domains and serve as ligands for leukocyte integrins. ICAM-1 and ICAM-2 exist as homodimers on the cell surface

Figure 5.

Combined selectin blockade did not significantly attenuate established ileitis in SAMP1/YitFc mice. SAMP1/YitFc mice at 10 weeks of age received antibodies against P-, E- and L-selectins (mAb RB40.34, 9A9, MEL-14), their respective isotype antibodies combined or dexamethasone (Dex 100 μ g I.P) and the histological severity of the ileitis was assessed as described previously²³¹ (villus-villus architectural distortion, active- neutrophilic infiltrates, chronic-predominantly lymphocyte infiltrates and total- sum of villus active and chronic indices, mean $\pm$ S.E.M., $n=7$ mice/treatment group).

Figure 6. Role of PSGL-1 in leukocyte recruitment to small intestine

A. Effector T cells and other leukocytes (PMN, monocytes) express PSGL-1 which enables their interaction with P-selectin on activated endothelium or platelets. **B.** In addition L-selectin-expressing cells may alternatively bind to PSGL-1 on endothelium or other leukocytes. **C.** Platelets may also bind to endothelial PSGL-1 perpetuating dysregulated recruitment to the small intestine. **Modified from** Ley&Kansas 2004

Table 1Effect of anti-adhesion molecules blockade in IBD^I

IBD Model	Adhesion molecules	Effect	Reference
CTT	E-selectin	no attenuation in colitis	Podolovsky, 1993
TNBS	E-, L- and P-selectin	decrease in rolling flux; no attenuation in colitis	Sans, 2001
DSS	P-selectin	attenuation in the severity of colitis	Gironella, 2002
	PSGL-1	reduction in number of rolling leukocyte; increased rolling velocity; attenuated colitis	Rijcken, 2004
SAMP/YitFc	PSGL-1	attenuation of transfer and spontaneous ileitis	Rivera-Nieves, 2006
TNBS	β_2 integrin	decreased neutrophil infiltration and Epithelial injury; attenuation in the severity of colitis	Wallace, 1992
	$\alpha_m\beta_2$ integrin	attenuation of colitis	Palmen, 1995
TNBS-/CD45RB ^{high}	$\alpha_m\beta_2$ and $\alpha_x\beta_2$ integrin	amelioration of colitis	Leon, 2006
CTT	α_4 integrin	attenuated acute but not chronic inflammatory infiltrates	Podolovsky, 1993
	$\alpha_4\beta_7$ integrin	attenuation of colitis	Hesterberg, 1996
Gai2	α_4 integrin	exacerbation of colitis in long term treatment	Bjursten, 2005
CD45RB ^{high}	$\alpha_4\beta_7$ integrin/MAdCAM-1	blockade of lymphocyte recruitment; reduction in severity of colitis	Picarella, 1997
DSS	MAdCAM-1	Decreased number of β_7 integrin positive cells; reduction in colonic injury	Shigematsu, 2001
SAMP/Yit	MAdCAM-1	amelioration of ileitis; attenuation of established ileitis	Matsuzaki, 2005
SAMP/YitFc	$\beta_7/\alpha_4\beta_7$ integrin MAdCAM-1	no effect on the severity of acute and chronic ileitis	Rivera- Nieves, 2005

^I CTT—define

TNBS—define

DSS—dextran sodium sulfate

Gai2—G protein alpha2