Published in final edited form as:

Biol Psychiatry. 2009 July 1; 66(1): 41–46. doi:10.1016/j.biopsych.2009.01.005.

Autonomic Activation During Sleep in PTSD and Panic: A Mattress Actigraphic Study

Steven H. Woodward¹, Ned J. Arsenault¹, Karin Voelker², Tram Nguyen³, Janel Lynch³, Karyn Skultety⁴, Erika Mozer^{3,5}, Gregory A. Leskin⁶, and Javid I. Sheikh³

¹ National Center for PTSD, Dissemination and Training Division, VA Palo Alto HCS, Palo Alto, CA, USA

² San Francisco State University, Department of Psychology, San Francisco, CA, USA

³ Stanford University, Department of Psychiatry and Behavioral Sciences, Stanford, CA, USA

⁴ Institute on Aging, San Francisco, CA, USA

⁵ Memory and Aging Center, Department of Neurology, University of California, San Francisco, CA, USA

⁶ Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, CA, USA

Abstract

Background—While it has been reported that persons with posttraumatic stress disorder manifest tonic autonomic activation, the literature contains numerous counter-examples. In revisiting the question, this study employed a novel method of mattress actigraphy to unobtrusively estimate heart rate and respiratory sinus arrhythmia over multiple nights of sleep in the home.

Method—Sleep cardiac autonomic status was estimated in four diagnostic groups, posttraumatic stress disorder, panic disorder, persons comorbid for both conditions, and controls. All 59 participants were community-residing non-veterans screened for sleep apnea and period leg movement disorder with polysomnography. Heart rate and respiratory sinus arrhythmia were calculated from the kinetocardiogram signal measured via accelerometers embedded in a mattress topper. Times in bed and asleep were also estimated. Per participant data were obtained from a median of 12 nights.

Results—Both posttraumatic stress disorder and posttraumatic stress disorder/panic disorder comorbid groups exhibited significantly higher heart rates and lower respiratory sinus arrhythmia magnitudes than panic disorder participants and controls. Panic disorder participants were indistinguishable from controls. The PTSD-only group exhibited longer times in bed and longer times presumably asleep than the other three groups.

Conclusion—In this study, posttraumatic stress disorder but not panic disorder was associated with altered cardiac autonomic status during sleep. Among participants meeting criteria for PTSD alone, autonomic activation co-occurred with prolongation of actigraphic sleep.

Corresponding Author: Steven H. Woodward, PhD, National Center for PTSD, Dissemination and Training Division, Mail Code 334 PTSD, VA Palo Alto HCS, 3801 Miranda Ave, Palo Alto, CA 94304 USA, V: 650-493-5000 x22111, F: 650-617-2701, steve.woodward@va.gov.

All other authors reported no biomedical financial interests or potential conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

The literature on basal autonomic status in posttraumatic stress disorder (PTSD) contains numerous inconsistencies. Lately, this issue has acquired additional significance with the emergence of an association between basal heart rate (HR) and metabolic syndrome (1–3). In the laboratory, phasically elevated HR is reliably observed in persons with PTSD as they anticipate re-exposure to trauma reminders (4–6). HR is also elevated in the acute aftermath of trauma in persons who go on to develop PTSD (7–9). Whether this heralds a persistent elevation in those same persons is unknown as longitudinal studies have yet to be performed. Three laboratory studies in combat veterans failing to observe HR elevations all informed participants they would *not* be exposed to trauma reminders but instead to generic stressors such as orthostatic challenge, cold pressure, and a mock job interview (10–12).

Two approaches to addressing this question deserve special attention. The first, measurement during sleep, appears to provide an opportunity to assess tonic autonomic activation largely free of influences of affect and cognition. As well, sleep has been shown to be independently relevant to appetite regulation, obesity, and diabetes, all associates of metabolic syndrome (13). Woodward et al (14) recorded continuous electrocardiogram (ECG) in 56 unmedicated male combat veterans and 14 mixed-trauma, age- and gender-matched controls during laboratory sleep studies employing conventional polysomnography (PSG). HR estimates were obtained from PSG-verified sleep, and per night HRs averaged over two to four post-adaptational nights. They observed no effect of PTSD diagnosis on sleep HR, and no association between HR and hyperarousal (D-Criterion) scores obtained using the Clinician-Administered PTSD Scale (CAPS, 15). D-criterion scores were, instead, associated with electroencephalogram (EEG) power in higher frequencies. An important qualification of this result is the low rate of emergence, in the laboratory, of a key symptom of PTSD-related sleep disturbance, trauma-related nightmares (16). A second approach, ambulatory measurement, would appear to offer increased reliability and validity. Beckham et al (17) tested 117 male combat veterans with and without PTSD using ambulatory blood pressure (BP) monitoring. An analysis accounting for self-reported activity level found PTSD to be associated with a five beat-per-minute (BPM) increase in basal HR. A smaller study by Buckley et al (18) also adjusting for body mass index, smoking status, and ethnicity and found that basal HR was 6.63 BPM higher in PTSD-positive participants than in controls. While both the Beckham and Buckley studies included the sleep period, neither could verify sleep. In addition, in ambulatory BP monitoring, HR is measured solely during the cuff inflation-deflation cycle; hence, HR estimates are unavoidably confounded with responses to this mildly aversive stimulus. While these generic stimuli might not be expected to induce exaggerated responses in PTSD, it is not clear that this argument extends to the sleep periods which constituted a substantial fraction of measurement periods in both studies. In normals, BP cuff inflation during sleep often leads to awakenings (19) which are typically associated with elevations of HR and BP (20).

The current study sought to combine virtues of both ambulatory and sleep-based measurement by measuring HR and respiratory sinus arrhythmia during the sleep period for an extended period in the home. It employed a novel sleep assessment method, mattress actigraphy, which transduces large and small sleeper movements using accelerometers embedded in a mattress topper. These small movements include the kinetocardiogram (KCG), which, for movement-free periods, closely tracks the ECG (21). Because mattress actigraphy employs no body surface sensors, in-home recordings can be continued indefinitely. In the current study, a median of 12 nights of recording were obtained from participants. Conventional actigraphic sleep variables were also calculated. In addition, this study employed a sample including participants meeting criteria for PTSD, Panic Disorder (PD), both PTSD and PD, who were otherwise free of Axis I psychopathology (other than dysthymia). As such, this study represents

the first direct comparison of sleep in PTSD with PD, two anxiety-spectrum disorders sharing dysphoric episodic sympathetic activations and subjectively poor sleep (22).

Method

Participants

This research was performed under the auspices of the Stanford/VA Palo Alto Human Research Protection Program. Written informed consent was obtained from all participants passing telephone screening. If present, a sleep partner had also to agree to actigraphic recording during the in-home portion of the study. Participants were recruited from the community using radio and newspaper advertisements emphasizing disturbances of sleep due to distressing experiences, fear and anxiety, and/or nightmares and nocturnal panic attacks. The recruitment of participants denying subjective sleep complaints relied principally on personal contacts between study participants and study staff and their respective social networks. Telephone screening excluded callers if they reported current medical illness, histories of central nervous system disease or injury, a Multivariate Apnea Prediction Index (MAPI, 23) greater than 0.5, or a previously-diagnosed medical sleep disorder (e.g. obstructive sleep apnea or periodic limb movement disorder). Persons were also excluded if currently abusing alcohol or other psychotropic substances, or were unwilling to discontinue hypnotic medications. Provisional diagnoses were made during telephone screening requiring from 15 to 60 minutes. Participants passing screening traveled to the research setting and underwent structured interviews. Diagnoses of PTSD and PD were based upon the CAPS and the PD section of the Structured Clinical Interview for the DSM-IV (SCID) (24), respectively. Trauma histories were determined prior to administration of the CAPS using the Life Events Checklist (25) and Life Events Follow-Up Interview (Kaloupek and Kutter, unpublished). Typical of community samples, index traumas were mostly single incidents such as motor vehicle accidents, criminal victimizations, and exposures related to the 1989 Loma Prieta earthquake. Screening of other Axis I psychopathology was confirmed using the Mini-International Neuropsychiatric Interview (MINI) (26). Eight participants diagnosed with dysthymic disorder were retained. Participants were administered the Beck Depression Inventory (BDI-II) (27), the Beck Anxiety Inventory (BAI) (28), and the Pittsburgh Sleep Quality Index (29). The latter included the PTSD Addendum comprising ten questions focusing on nocturnal arousals, nightmares, and related anxiety.

Seventeen participants meeting screening and diagnostic criteria declined to return for a screening PSG. Ten participants who underwent PSG had apnea/hypopnea indices (AHIs) greater than 10 events per hour. Three participants were excluded due to limb movement arousal indices (PLMAIs) greater than 20 events per hour. (AHIs and PLMAIs were manually scored using the Lifeshirt records and PSG records, respectively.) Ten participants who passed the screening PSG declined to undergo in-home recording. Among those recorded at home, seven were excluded due to equipment failure. While many potential participants were lost due to sleep medical conditions, voluntary discontinuation, or equipment failure, non-completers were evenly distributed over diagnoses (14 PD, 14 PTSD, 13 PTSD+PD, 8 controls). Completing participants were paid \$300. Non-completers received pro-rated payments. Participants were asked to maintain their non-hypnotic medications. Rates of psychotropic medication use differed over diagnostic groups (controls, 0%, PD, 18%, PTSD, 27%, and PTSD +PD, 45%; $\chi^2(3) = 7.87$, $p < 0.05$). There were no associations between group and specific classes of medication: SSRI anti-depressants ($\chi^2(3) = 2.68$, n.s., 14% of total sample); non-SSRI anti-depressants ($\chi^2(3) = 4.50$, n.s., 8% of sample); anxiolytic agents ($\chi^2(3) = 5.97$, n.s., 5% of sample). No participants were taking stimulants or beta-blockers. Three tested participants reported having prescriptions for hypnotic medications which they agreed to discontinue.

Procedures

Participants came to the study site in the afternoon to complete diagnostic interviews and self-report psychometrics. After dinner, they came to the sleep laboratory to prepare for the screening PSG. The PSG montage included EEG (Cz and Pz referenced to linked ears), electro-oculogram (EOG, vertical and horizontal), submental electromyogram (EMG), bilateral anterior tibialis EMG, and oral/nasal airflow (thermistor). ECG and respiratory effort (RESP, thoracic and abdominal bands, inductive plethysmography) were recorded via a Vivometrics LifeShirt system. After completion of a pre-sleep startle protocol, participants went to sleep *ad libitum*.

Home sleep recording involved installation of the mattress actigraphy system and setting up the participant for a single night of concurrent ambulatory PSG. Ambulatory PSG employed EEG (Cz and Pz referenced to linked ears), EOG (horizontal only), submental EMG, and ECG. Participants were also fitted with an ankle actigraph (Octagonal Sleep Watch, Ambulatory Monitoring, Inc.) to be worn while in bed for the duration of the home study. Participants denying any nocturnal paroxysmal symptoms were recorded for approximately one week. Participants reporting such symptoms were recorded for up to a month. The following results are based upon mattress actigraphy data from all home nights excluding the night of concurrent ambulatory PSG.

Mattress actigraphy was recorded by four sensitive, low-noise, DC accelerometers (Silicon Designs 2010) embedded in a mattress topper that was placed on top of the participant's regular mattress. The topper was composed of a one-inch foam pad encased in bacteriostatic ticking. The accelerometers were suspended in spandex "trampolines" over three-inch diameter holes in the underlying foam to promote compliance under load. The net effect was not unlike a "pillow-top" mattress of which most participants spoke favorably, none discontinuing on the basis of discomfort. The accelerometers were connected to a 12-volt medical grade power supply (Condor MLL 250) and to a data acquisition card (Kiethley KMPCI 3807) which sampled the movement data at 600 Hz with 16 bits of amplitude resolution. Extraction of HR and RSA from the actigraphic record began with visual inspection and determination of the onset and offset of the "intended sleep period" (ISP) determined using all available data, including ambient light and sound. Automated computer-based processing began with the identification of epochs of stable body position. Within these epochs, the sensor supplying the KCG signal with the highest signal-to-noise ratio was identified. Extraction of KCG beats began with determination of the canonical (median) KCG ensemble for each epoch. These were then used as templates to identify beats as locations of high-cross correlation with the raw data. At each stage, strict artifact exclusion criteria were applied with the result that per-night estimates were based on approximately three hours of recording, on average. The aggregate time producing low-artifact KCG beat series, here labeled "KCGtime", was co-determined by time in bed and by typical KCG signal quality. The latter, in turn, was determined by the presence/absence of body movements contaminating the KCG, and by body position in relation to the sensors (prone vs supine vs lateral) which was not recorded. As with all actigraphy, epochs of low movement were used to estimate sleep vs wake, but could not provide direct information regarding sleep stage. Additional details of the estimation of HR and RSA from the KCG have been published elsewhere (21).

Twenty participants had sleeping partners for some or all nights. Participant/partner pairs were issued a double topper which included a single sensor under the thorax of the partner. While participant KCG dominated the signals at sensors under his or her own thorax, an opportunity for signal "bleed" arose when sleeping was asynchronous. All participant/partner HR time series were reviewed for evidence of cross-sleeper artifact. This effort was facilitated by the concentration of such artifacts at the beginning and ends of nights, the availability of nightly records of the presence/absence of a partner, and the fact that sleeper pairs typically had distinct

baseline HRs. Substantial cross-sleeper artifact was found in 21 nights. These were excluded and per-participant across-night median HR and RSA values recalculated. Post-exclusion, the mean deviation was -0.079 BPM, and no per-participant median diverged from the original value by more than 0.73 BPM.

Differences between diagnostic groups were tested with univariate ANOVAs followed by Fisher's LSD. Relations among continuous variables were assessed using simple and partial correlation and multiple regression. (30)

Results

Demographic, sleep-medical, and psychometric data characterizing the final sample of 59 participants is summarized in Table 1. The BDI-II exhibited a moderate effect of group ($F(3,50) = 10.3$, $p < 0.001$, partial $\eta^2 = 0.381$) and a modest effect of gender ($F(1,50) = 4.7$, $p < 0.05$, partial $\eta^2 = 0.087$). (BDI-II, BAI, and PSQI scores were missing for one PTSD participant.) Controls endorsed very little depressive mentation. Among the patient groups, both PD participants (Fisher's LSD: $p < 0.001$) and PTSD participants (Fisher's LSD: $p < 0.01$) endorsed significantly less depressive mentation than PTSD+PD participants. Female participants endorsed more than males. The BAI exhibited a moderate effect of group ($F(3,50) = 10.8$, $p < 0.001$, partial $\eta^2 = 0.393$) and a group by gender interaction ($F(3,50) = 3.6$, $p < 0.05$, partial $\eta^2 = 0.177$). Again, controls endorsed little anxious mentation relative to the anxiety-disorder groups. Among the latter, PTSD+PD participants endorsed significantly more than PTSD participants (Fisher's LSD: $p < 0.001$). The group by gender interaction originated in the observation that female PD participants endorsed significantly more anxious mentation than male PD participants ($t(11) = 2.74$, $p < 0.05$), whereas at other levels of the group factor there were no gender differences. The Pittsburgh Sleep Quality Index (PSQI) exhibited a large effect of group ($F(3,50) = 6.3$, $p < 0.001$, partial $\eta^2 = 0.273$). Again, controls reported less sleep complaint than the anxiety-disordered groups, though controls' PSQI scores were in the low clinical range. Among anxiety-disordered groups, PTSD participants reported significantly more sleep complaint than PD participants (Fisher's LSD, $p < 0.05$). The PTSD Addendum of the PSQI also exhibited a large effect of group ($F(3,50) = 13.9$, $p < 0.001$, partial $\eta^2 = 0.454$), with both PTSD participants (Fisher's LSD: $p < 0.01$) and PTSD+PD participants (Fisher's LSD: $p < 0.001$) endorsing significantly more PTSD-related sleep symptomology than PD participants. Eleven of 13 controls and 5 of 13 PD participants denied any history of trauma. Among anxiety-disorder participants endorsing a trauma history, the numbers of different trauma types reported was associated with diagnosis ($F(2,34) = 5.65$, $p < 0.01$), being significantly elevated in those meeting criteria for PTSD (Fisher's LSD, $p < 0.05$) and PTSD+PD (Fisher's LSD, $p < 0.01$) relative to PD alone. The number of different trauma types reported was not associated with gender ($F < 1$). The anxiety-disordered groups did not differ in terms of age at first trauma ($F < 1$).

Diagnostic groups were closely matched on physical and medical characteristics associated with effects on sleep other than those under consideration, including age, body mass index (BMI), apnea-hypopnea indices (AHI), and periodic limb movement arousal indices (PLMAI). Controls had slightly lower BMIs than the PTSD participants (Fisher's LSD, $p < 0.05$) and PTSD+PD (Fisher's LSD, $p < 0.05$). Females had lower MAPI scores than males ($F(1,51) = 5.5$, $p < 0.05$, partial $\eta^2 = 0.098$). Neither diagnosis nor gender was associated with differences in years of education or hours worked per week.

Additional characteristics of the mattress actigraphy sleep sample can be found in Table 2. Because controls all denied experiencing any nocturnal paroxysmal events, they were recorded for fewer nights, on average, than the patient groups ($F(3,51) = 8.3$, $p < 0.001$, partial $\eta^2 = 0.327$). PTSD participants were recorded on significantly more nights than PD participants

(Fisher's LSD: $p < 0.05$). The average ISP (calculated over all nights providing interpretable ISPs and without regard to the quality of KCG data) exhibited a main effect of diagnosis ($F(3,51) = 3.43$, $p < 0.05$, partial $\eta^2 = 0.194$). Surprisingly, PTSD participants exhibited significantly *longer* in-bed periods than either PD participants (Fisher's LSD: $p < 0.001$) or PTSD+PD participants (Fisher's LSD: $p < 0.01$). Mean KCGtime values are presented in Table 2. KCGtime also exhibited a main effect of diagnosis ($F(3,51) = 3.33$, $p < 0.05$, partial $\eta^2 = 0.164$). As before, PTSD participants exhibited raw KCGtimes that were significantly longer than both PD participants (Fisher's LSD, $p < 0.01$) and PTSD+PD participants (Fisher's LSD, $p < 0.001$), but not longer than controls'. The KCGtimes of the PTSD+PD group were also shorter than those of controls (Fisher's LSD, $p < 0.05$). KCGtime exhibited an effect of gender ($F(1,51) = 4.43$, $p < 0.05$, partial $\eta^2 = 0.080$), being on average 30 minutes shorter in women. When KCGtime was re-analyzed as a percentage of the ISP, there was no main effect of diagnosis ($F < 1$) but again an effect of gender ($F(1,51) = 8.30$, $p < 0.01$, partial $\eta^2 = 0.140$) with males exhibiting more KCGtime than females. In males, KCG-based HR and RSA were obtained from approximately 51% of the intended sleep period, in females, 41%. Typical total sleep HR and RSA samples were approximately 40 hours.

Sleep HR exhibited a main effect of diagnosis ($F(3,51) = 2.89$, $p < 0.05$, partial $\eta^2 = 0.145$). PTSD participants' sleep HRs were higher than controls' (Fisher's LSD: $p < 0.05$) as were those of PTSD+PD participants' (Fisher's LSD: $p < 0.05$). HRs of controls and PD participants did not differ (See Fig 1.). There was no effect of gender and no diagnosis $\times$ gender interaction (F 's < 1). Sleep RSA magnitude also exhibited a main effect of diagnosis ($F(3,51) = 3.55$, $p < 0.05$, partial $\eta^2 = 0.173$). PD participants exhibited higher sleep RSA magnitude than PTSD (Fisher's LSD: $p < 0.001$) and PTSD+PD participants (Fisher's LSD: $p < 0.05$; See Fig. 1.). Covariation for age, AHI, BMI, PLMAI, and KCG time had no impact on the between-group results for sleep HR and RSA.

Neither BDI scores, BAI scores, nor CAPS severity scores (in PTSD and PTSD+PD participants) were correlated with sleep HR or RSA magnitude. BAI scores were inversely correlated with KCGtime over all participants ($r(57) = -0.341$, $p < 0.01$) and after excluding controls ($r(43) = -0.341$, $p < 0.05$). Sleep HRs correlated positively ($r(56) = 0.332$, $p < 0.05$) and sleep RSA magnitudes negatively ($r(56) = -0.333$, $p < 0.05$) with PSQI total scores (See Fig. 2). Sleep HR was also correlated with the PSQI PTSD Addendum score ($r(56) = 0.406$, $p < 0.01$). This pattern was preserved when controls were excluded. The relation between PSQI scores and RSA magnitudes was re-evaluated controlling for age, BAI and BDI scores. The intent of this analysis was to isolate variance in PSQI not associated with aging and waking dysphoria. Significant increases in PSQI variance explained were evident at all steps. In the final model, only BAI and sleep RSA magnitude retained beta weights significantly different from zero (BAI: $\beta = 0.413$, $t = 3.05$, $p < 0.01$; RSA: $\beta = -0.284$, $t = 2.39$, $p < 0.05$). The model accounted for 37% of variance in PSQI scores in the sample composed of both anxiety-disordered participants and controls (adjusted $R^2 = 0.371$, $F_{\text{regression}}(4,53) = 9.48$, $p < 0.001$). Restricting the analyses to anxiety-disordered participants produced a similar result (BAI: $\beta = 0.339$, $t = 2.28$, $p < 0.05$; RSA: $\beta = -0.405$, $t = 2.74$, $p < 0.01$; adjusted $R^2 = 0.247$, $F_{\text{regression}}(4,40) = 4.61$, $p < 0.01$). The waking disturbance score of the PSQI (not incorporated into the PSQI score, itself) was positively correlated with sleep HR ($r(56) = 0.337$, $p < 0.01$) and negatively correlated with KCGtime ($r(56) = -0.324$, $p < 0.05$). The latter correlations remained significant after controlling for age, BDI, BAI, and PSQI total score in the whole sample (HR: $r(52) = 0.381$, $p < 0.01$; KCGtime: $r(52) = -0.275$, $p < 0.05$), and in the anxiety-disordered participants (HR: $r(39) = 0.407$, $p < 0.01$; KCGtime: $r(39) = -0.349$, $p < 0.05$).

Discussion

Consistent with the findings of both Beckham and Buckley, sleep HR in PTSD and PTSD+PD participants was significantly elevated relative to controls. For PTSD participants, the magnitude of HR elevation was very similar to observations made by those investigators. For PTSD+PD participants, the elevation was over 8 BPM. The pattern of results among PTSD, PD and controls closely approximated the observations of Blechert et al (31) who found baseline HR elevation in PTSD but not PD in a community sample. Sleep RSA magnitude presented a corresponding pattern, with PTSD and PTSD+PD participants exhibiting lower magnitudes; although here the significant contrasts were with PD participants. In waking, Blechert et al observed RSA power to be lower in PTSD than in PD, and not different between PD and controls; however, in that study, PTSD participants also exhibited significantly lower RSA power than controls. The association of PTSD with elevated sleep HR provides an intriguing context for further tests of prazosin to treat sleep disturbances in this disorder. While prazosin reduces nightmares and improves subjective sleep in PTSD (32,33), it does not, at least when administered to hypertensive men, lower heart rate (34).

An important limitation of this study is the absence of a subgroup with histories of trauma that did not manifest PTSD. Hence, trauma *per se* cannot be excluded as the cause of excess autonomic activation in the PTSD and PTSD+PD groups. We also note that multiple factors favored low symptom severity among the anxiety-disordered persons studied. These include the intensity of the study, the use of a community sample, and the exclusion of additional Axis I psychopathology. Commensurate with this are a preponderance of single-incident trauma histories among PTSD participants and a low rate of psychotropic use overall (26%). That such a low-severity sample manifested significant associations between PTSD and altered sleep cardiac autonomic status while chronic severe combat-related PTSD inpatient males studied in our laboratory have not is intriguing. Extended in-home studies of combat veterans may resolve this apparent inconsistency.

We suspect that contrasting persons with PD, PTSD and PTSD+PD promoted the creation of diagnostically “clean” groups. From this perspective, it is noteworthy that, when considered simultaneously in terms of sleep cardiac autonomic status and actigraphic sleep/wake findings, these groups are not metrically related. PD participants appeared essentially normal in both domains. Findings in comorbid participants, though broadly consistent with the exaggeration of subjective sleep complaints reported by Leskin (35), could not be extrapolated from those with PTSD and PD. In a similar vein, the actigraphic sleep profiles of PD, PTSD, and PTSD +PD groups fell into distinct risk categories in relation to metabolic syndrome (13). PD participants exhibited no evidence of elevated risk from sleep restriction or tonic sympathetic activation. PTSD+PD participants exhibited elevated risk due to both sleep restriction and sympathetic activation. PTSD-only patients presented contradictory indications, combining tonically elevated sympathetic tone with extended sleep periods. Careful attention to distinctions among PD, PTSD and PTSD+PD phenotypes may prove productive in future studies of associations between metabolic syndrome and psychiatric status. Finally, PTSD only participants spent more time in bed, and more time immobile, yet still slept “worse” to the extent that autonomic quiescence during sleep has positive health implications. If this paradoxical pattern can be replicated, we may then ask whether it represents an adaptive strategy for achieving the biological goals of sleep while retaining the capacity for rapid awakening.

Actigraphy infers sleep from immobility; however, immobility can be measured with varying precision, and may mean different things in different diagnostic groups (21,36). Analyses are now underway in our laboratory to evaluate the relative capacities of the different immobility indices obtainable from mattress actigraphy to predict PSG-defined sleep. Among these,

KCGtime is a very strict index because it excludes any epoch containing movement sufficient to introduce artifact into the KCG time series. The observed relationships between KCGtime and the waking disturbance subscore of the PSQI provides preliminary evidence that quantifying in-bed time devoid of even the smallest and briefest of movements may prove a productive extension of the actigraphic measurement of sleep.

Acknowledgments

This research was funded by grant number MH64724 from the National Institute of Mental Health to J.I. Sheikh, M.D. The authors wish to thank European Sleep Works, Berkeley, CA, who donated expertise and materials towards fabrication of the mattress toppers. We are also grateful for the administrative support provided by the Research Service of the Veterans Affairs Palo Alto Health Care System, and the Sponsored Projects Office, Stanford University.

Dr. Woodward is the inventor of the SensorBed. The patent (US 6,485,441) is co-assigned to the Department of Veterans Affairs and to Leland Stanford University. The SensorBed has not been licensed for commercialization. In the event of such licensure, Dr. Woodward could benefit indirectly in his official capacity as a federal employee as one-third of any royalties would be paid into the general fund of the VA Palo Alto Health Care System and, from there, could be assigned to Dr. Woodward for research-related expenditures. No royalty earnings would accrue to Dr. Woodward were he to separate from the Department of Veterans Affairs; however, he could profit financially as a private citizen through personally licensing and commercializing the SensorBed in the event that profits were realized.

References

1. Inoue T, Iseki K, Iseki C, Ohya Y, Kinjo K, Takishita S. Association between heart rate and multiple risk factor syndrome. *Circ J* 2008;72:454–7. [PubMed: 18296845]
2. Oberlinner C, Humpert PM, Nawroth PP, Zober A, Morcos M. Metabolic syndrome in a large chemical company: prevalence in a screened worksite sample. *Acta Diabetol* 2008;45:31–5. [PubMed: 17882349]
3. Tomiyama H, Yamada J, Koji Y, Yambe M, Motobe K, Shiina K, et al. Heart rate elevation precedes the development of metabolic syndrome in Japanese men: a prospective study. *Hypertens Res* 2007;30:417–26. [PubMed: 17587754]
4. Blanchard EB. Elevated basal level of cardiovascular responses in Vietnam veterans with PTSD: A health problem in the making? *J Anxiety Disord* 1990;4:233–237.
5. Keane TM, Kolb LC, Kaloupek DG, Orr SP, Blanchard EB, Thomas RG, et al. Utility of psychophysiological measurement in the diagnosis of posttraumatic stress disorder: results from a Department of Veterans Affairs Cooperative Study. *J Consult Clin Psychol* 1998;66:914–23. [PubMed: 9874904]
6. Buckley TC, Kaloupek DG. A meta-analytic examination of basal cardiovascular activity in posttraumatic stress disorder. *Psychosom Med* 2001;63:585–94. [PubMed: 11485112]
7. Shalev AY, Sahar T, Freedman S, Peri T, Glick N, Brandes D, et al. A prospective study of heart rate response following trauma and the subsequent development of posttraumatic stress disorder. *Arch Gen Psychiatry* 1998;55:553–9. [PubMed: 9633675]
8. Kuhn E, Blanchard EB, Fuse T, Hickling EJ, Broderick J. Heart rate of motor vehicle accident survivors in the emergency department, peritraumatic psychological reactions, ASD, and PTSD severity: a 6-month prospective study. *J Trauma Stress* 2006;19:735–40. [PubMed: 17075910]
9. Zatzick DF, Russo J, Pitman RK, Rivara F, Jurkovich G, Roy-Byrne P. Reevaluating the association between emergency department heart rate and the development of posttraumatic stress disorder: A public health approach. *Biol Psychiatry* 2005;57:91–5. [PubMed: 15607305]
10. Murburg, MM.; McFall, ME.; Veith, RC. Basal sympathoadrenal function in patients with PTSD and depression. In: Murburg, MM., editor. *Progress in psychiatry*; no 42. Vol. 1. Vol. 42. Washington, DC: American Psychiatric Press; 1994. p. xxxip. 371
11. Orr SP, Meyerhoff JL, Edwards JV, Pitman RK. Heart rate and blood pressure resting levels and responses to generic stressors in Vietnam veterans with posttraumatic stress disorder. *J Trauma Stress* 1998;11:155–64. [PubMed: 9479684]

12. Woodward SH, Kaloupek D, Schaer M, Martinez C, Eliez S. Right anterior cingulate cortical volume covaries with respiratory sinus arrhythmia magnitude in combat veterans. *J Rehabil R D* 2008;43:451–464.
13. Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep and sleep loss. *Sleep Med* 2008;9 Suppl 1:S23–8. [PubMed: 18929315]
14. Woodward SH, Murburg MM, Bliwise DL. PTSD-related hyperarousal assessed during sleep. *Physiol Behav* 2000;70:197–203. [PubMed: 10978496]
15. Blake, DD.; Weathers, FW.; Nagy, LM.; Kaloupek, DG.; Charney, DS.; Keane, TM. Clinician-Administered PTSD Scale for DSM-IV: Current and Lifetime Version. Behavioral Science Division; Boston VA: Medical Center/Neurosciences Division; West Haven VA Medical Center: 1997.
16. Woodward SH, Arsenault NJ, Murray C, Bliwise DL. Laboratory sleep correlates of nightmare complaint in PTSD inpatients. *Biol Psychiatry* 2000;48:1081–7. [PubMed: 11094141]
17. Beckham JC, Feldman ME, Barefoot JC, Fairbank JA, Helms MJ, Haney TL, et al. Ambulatory cardiovascular activity in Vietnam combat veterans with and without posttraumatic stress disorder. *J Consult Clin Psychol* 2000;68:269–76. [PubMed: 10780127]
18. Buckley TC, Holohan D, Greif JL, Bedard M, Suvak M. Twenty-four-hour ambulatory assessment of heart rate and blood pressure in chronic PTSD and non-PTSD veterans. *J Trauma Stress* 2004;17:163–71. [PubMed: 15141790]
19. Dimsdale JE, Coy TV, Ancoli-Israel S, Clausen J, Berry CC. The effect of blood pressure cuff inflation on sleep. A polysomnographic examination. *Am J Hypertens* 1993;6:888–91. [PubMed: 8267947]
20. Morgan BJ, Crabtree DC, Puleo DS, Badr MS, Toiber F, Skatrud JB. Neurocirculatory consequences of abrupt change in sleep state in humans. *J Appl Physiol* 1996;80:1627–36. [PubMed: 8727549]
21. Woodward SH, Arsenault NJ, Voelker K, Nguyen T, Lynch J, Leskin G, et al. Estimating heart rate and RSA from the mattress-recorded kinetocardiogram. *Psychophysiology* 2007;44:635–8. [PubMed: 17437553]
22. Sheikh JI, Woodward SH, Leskin GA. Sleep in post-traumatic stress disorder and panic: convergence and divergence. *Depress Anxiety* 2003;18:187–97. [PubMed: 14661188]
23. Maislin G, Pack AI, Kribbs NB, Smith PL, Schwartz AR, Kline LR, et al. A survey screen for prediction of apnea. *Sleep* 1995;18:158–166. [PubMed: 7610311]
24. Spitzer, RL.; Williams, JBW.; Gibbon, M.; First, MB. Structured Clinical Interview for DSM-IV. New York: Biometrics Research Department, New York State Psychiatric Institute; 1996.
25. Blake, D.; Weathers, F.; Nagy, L.; Kaloupek, D.; Klauminzer, G.; Charney, D., et al. Clinician-Administered PTSD Scale (CAPS) Instructional Manual: National Center for PTSD. 2000.
26. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998;59 Suppl 20:22–33. [PubMed: 9881538]quiz 34–57
27. Beck, AT.; Steer, RA.; Brown, GK. Manual for the Beck Depression Inventory-II. San Antonio, TX: Psychological Corporation; 1996.
28. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 1988;56:893–897. [PubMed: 3204199]
29. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: A new instrument for psychiatric practice and research. *Psychiatry Res* 1989;28:193–213. [PubMed: 2748771]
30. Woodward SH, Leskin GA, Sheikh JI. Movement during sleep: associations with posttraumatic stress disorder, nightmares, and comorbid panic disorder. *Sleep* 2002;25:681–8. [PubMed: 12224848]
31. Blechert J, Michael T, Grossman P, Lajtman M, Wilhelm FH. Autonomic and respiratory characteristics of posttraumatic stress disorder and panic disorder. *Psychosom Med* 2007;69:935–43. [PubMed: 17991823]
32. Raskind MA, Peskind ER, Kanter ED, Petrie EC, Radant A, Thompson CE, et al. Reduction of nightmares and other PTSD symptoms in combat veterans by prazosin: a placebo-controlled study. *Am J Psychiatry* 2003;160:371–3. [PubMed: 12562588]

33. Taylor FB, Martin P, Thompson C, Williams J, Mellman TA, Gross C, et al. Prazosin effects on objective sleep measures and clinical symptoms in civilian trauma posttraumatic stress disorder: a placebo-controlled study. *Biol Psychiatry* 2008;63:629–32. [PubMed: 17868655]
34. Materson BJ, Williams DW, Reda DJ, Cushman WC. Response to six classes of antihypertensive medications by body mass index in a randomized controlled trial. *J Clin Hypertens (Greenwich)* 2003;5:197–201. [PubMed: 12826782]
35. Leskin GA, Woodward SH, Young HE, Sheikh JI. Effects of comorbid diagnoses on sleep disturbance in PTSD. *J Psychiatr Res* 2002;36:449. [PubMed: 12393315]
36. Jean-Louis G, Mendlowicz MV, Gillin JC, Rapaport MH, Kelsoe JR, Zizi F, et al. Sleep estimation from wrist activity in patients with major depression. *Physiol Behav* 2000;70:49–53. [PubMed: 10978477]

Fig. 1.
Bar graphs plotting least-squares means and standard deviations of all-night heart rates and respiratory sinus arrhythmia magnitudes for each participant group.

Fig. 2. Scatterplot of PSQI total score and all-night respiratory sinus arrhythmia magnitudes across all participants.

Table 1
Means and standard deviations of indices of medical sleep disorder and psychiatric severity across groups.

n	CONTROL			PANIC DISORDER			PTSD			PTSD+PANIC DISORDER			Effect Size of Diagnosis (partial h^2) where $\alpha < .05$
	13			13			22			11			
	M	SD		M	SD		M	SD		M	SD		
Age	39.6	10.8		42.1	11.0		42.2	12.8		42.0	13.9		
BMI	22.1	4.8		25.5	4.9		26.2	5.7		26.6	6.2		
MAPI	0.1	0.1		0.2	0.1		0.2	0.2		0.2	0.2		
AHI	3.0	2.6		2.6	2.5		2.5	1.8		2.7	2.0		
PLMAI	0.7	3.1		2.7	3.2		0.9	3.7		1.7	4.1		
Education	16.1	3.3		14.7	3.5		14.7	4.3		13.5	4.3		
Work Hrs/Wk	28.5	18.5		38.8	20.1		27.6	24.4		22.9	24.2		
BDI	0.4	8.4		10.2	8.6		14.2	10.9		20.4	10.8		0.381
BAI	1.2	9.8		15.8	10.1		10.8	12.8		26.7	12.7		0.393
PSQI	8.0	2.2		9.4	2.3		11.5	2.9		11.4	2.9		0.273
PSQI PTSD Addendum	7.5	3.2		9.9	3.3		13.3	4.1		16.2	4.1		0.454

Least-squares means (controlled for gender) and standard deviations of mattress actigraphic indices of nocturnal quiescence and autonomic status.

		CONTROL				PANIC DISORDER		PTSD		PTSD+PANIC DISORDER		Effect Size of Diagnosis (partial h^2) where $\alpha < .05$	
		13		13		13		22		11			
n		M	SD	M	SD	M	SD	M	SD	M	SD		
actigraphy nights	6.7		4.3	12.0	4.4	14.6	5.1	10.9	5.5				0.346
ISP minutes	472.5		61.5	436.9	58.6	504.6	68.1	435.1	83.5				0.194
KCG minutes	224.4		49.1	195.4	50.3	240.5	58.4	179.9	63.4				0.156
HR	61.2		7.7	61.3	8.0	66.6	9.2	69.8	10.0				0.148
RSA magnitude	0.25		0.08	0.31	0.09	0.21	0.10	0.21	0.11				0.177