Published in final edited form as:

Br J Ophthalmol. 2008 August ; 92(8): 1150–1151. doi:10.1136/bjo.2007.122549.

A microstructural retinal analysis of membrano-proliferative glomerulonephritis type II

C Gerth¹, R J Zawadzki², C Licht³, J S Werner², and E Héon^{1,4}

¹ Department of Ophthalmology and Vision Sciences, The Hospital for Sick Children and The University of Toronto, Canada

² Department of Ophthalmology & Vision Science and Section of Neurobiology, Physiology & Behavior, University of California, Davis, USA

³ Department of Nephrology, The Hospital for Sick Children, Toronto, Canada

⁴ Program of Genetics and Genomic Biology, The Hospital for Sick Children, Toronto, Canada

Membranoproliferative glomerulonephritis type II (MPGN type II)/dense deposit disease (DDD) is characterised by electron deposits within the lamina densa of the glomerular basement membrane. Systemic manifestation may include lipodystrophy and a retinopathy characterised by drusenoid deposits.^{1–3} MPGN type II is caused by overactivation of the alternative complement pathway (C3 Nephritic Factor, Factor H deficiency or functional defects) in the majority of cases.⁴ Recently, Factor H H402Y has been identified as a major-risk haplotype within the alternative complement pathway.⁵ In vivo microstructural identification and localisation of these retinal deposits in humans are shown for the first time.

CASE REPORT

A 54-year-old woman who presented with a history of blurred vision in her right eye, nyctalopia and photophobia for 5 years, was referred with the possible diagnosis of fundus flavimaculatus/RP. The patient's past medical history revealed lipodystrophy (most remarkable in the face, fig 1A) diagnosed at age 11. She was diagnosed as having glomerulonephritis at age 21, for which she had renal transplants at ages 27 and 41. The patient also required insertion of a pacemaker at the age of 52 years. The diagnosis of MPGN type II was confirmed at the time of her first renal transplant. At the age of 54 years, her visual function assessment showed reduced visual acuity (right eye: 0.20 logMAR; left eye: 0.22 logMAR), reduced contrast sensitivity (either eye: 1.05 log) but normal colour vision when tested with HRR plates. Rod- and cone-mediated function was reduced significantly in the recorded full-field electroretinogram (ISCEV standard⁶) (rod-isolated b-wave: 20%; rod-cone a- and b-wave: 23% and 26%; cone a- and b-wave: 23% and 44%; 30 Hz flicker response: 34% of the mean control) when compared with age-matched control data. Fundus examination of both eyes demonstrated small yellowish partially confluent drusenoid deposits, which were irregularly distributed within the macular area and the posterior pole involving the peripapillary area (fig 1B,C).

Correspondence to: E Héon, University of Toronto and Hospital for Sick Children, Department of Ophthalmology and Vision Sciences, 555 University Avenue, Toronto, ON, M5G 1X8, Canada; eheon@attglobal.net.

Competing interests: None.

Patient consent: Obtained

High-speed (1000 A-scans/frame; 9 frames/s) high-resolution (axial 4 μm ; lateral: 10–15 μm) Fourier-domain optical coherence tomography (Fd-OCT)⁷ constructed at UC Davis, combined with a hand-held scanner (Biotigen, Triangle Park, NC), was used for the acquisition of retinal layer images. Microstructural Fd-OCT analysis revealed an intact inner retinal structure compared with an age-matched control subject.⁸ Deposits were observed within the RPE/Bruchs membrane complex, extending through the outer and inner segment layer up to the outer limiting membrane (fig 2).

Comment

Microstructural in vivo retinal high-resolution imaging identified the localisation and the extent of retinal deposits affecting the RPE/Bruch membrane complex. The mild impact on vision function, characteristic of MPGN type II, is supported by the integrity of retinal lamination including inner and outer segments, which is in contrast to other hereditary macular dystrophy where the photoreceptor structure is disrupted.⁹

Acknowledgments

Funding: This study was supported by the NEI (grant 014743) and an RPB Jules and Doris Stein Professorship (JSW).

References

1. Duvall-Young J, Short CD, Raines MF, et al. Fundus changes in mesangiocapillary glomerulonephritis type II: clinical and fluorescein angiographic findings. *Br J Ophthalmol* 1989;73:900–6. [PubMed: 2605144]
2. Duvall-Young J, MacDonald MK, McKechnie NM. Fundus changes in (type II) mesangiocapillary glomerulonephritis simulating drusen: a histopathological report. *Br J Ophthalmol* 1989;73:297–302. [PubMed: 2713310]
3. Leys A, Proesmans W, Van Damme-Lombaerts R, et al. Specific eye fundus lesions in type II membranoproliferative glomerulonephritis. *Pediatr Nephrol* 1991;5:189–92. [PubMed: 2031833]
4. Licht C, Schlotzer-Schrehardt U, Kirschfink M, et al. MPGN II—genetically determined by defective complement regulation? *Pediatr Nephrol* 2007;22:2–9. [PubMed: 17024390]
5. Hageman GS, Anderson DH, Johnson LV, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A* 2005;102:7227–32. [PubMed: 15870199]
6. Marmor MF, Holder GE, Seeliger MW, et al. Standard for clinical electroretinography (2004 update). *Doc Ophthalmol* 2004;108:107–14. [PubMed: 15455793]
7. Wojtkowski M, Leitgeb R, Kowalczyk A, et al. In vivo human retinal imaging by Fourier domain optical coherence tomography. *J Biomed Opt* 2002;7:457–63. [PubMed: 12175297]
8. Alam S, Zawadzki RJ, Choi S, et al. Clinical application of rapid serial Fourier-domain optical coherence tomography for macular imaging. *Ophthalmology* 2006;113:1425–31. [PubMed: 16766031]
9. Gerth C, Zawadzki RJ, Choi SS, et al. Visualization of lipofuscin accumulation in Stargardt macular dystrophy by high-resolution Fourier-domain optical coherence tomography. *Arch Ophthalmol* 2007;125:575. [PubMed: 17420386]

Figure 1.

Face (A) photograph illustrates extended lipodystrophy typical of MPGN type II. Fundus photographs of the right (B) and left eye (C) show drusenoid deposits involving the posterior pole and the peripapillary area.

Figure 2.

Horizontal Fourier-domain OCT 5 mm scans through the macula from the patient's right (A) and left eye (B): well-demarcated deposits within the RPE/Bruchs membrane extending through the outer and inner segment layer up to the outer limiting membrane. Retinal lamination is preserved. Retinal lamination depicted in the macular scan of an age-matched control subject shown for comparison (C; foveal magnification: D). CL, connecting cilia; GCL, ganglion cell layer; ILM/NFL, internal limiting membrane/nerve fibre layer; INL, inner nuclear layer; IPL, inner plexiform layer; ISL, inner segment layer; OLM, outer limiting membrane; ONL, outer nuclear layer; OPL, outer plexiform layer; OSL, outer segment layer; RPE/BM, retinal pigment epithelium/Bruch membrane; VM, Verhoeff membrane.