CORRESPONDENCE

Lyme Disease—Current State of Knowledge

by Prof. Dr. med. Roland Nau, Prof. Dr. med. Hans-Jürgen Christen,
Prof. Dr. med. Dr. rer. nat. Helmut Eiffert in volume 5/2009

Different Symptoms

There are four different species of human pathogen *Borrelia* in Europe (in central Europe, mainly *Borrelia afzelii* and *Borrelia garinii*), whereas the United States has only one (*Borrelia burgdorferi*). The vectors (ticks) on the two continents also differ. The result is that manifestations of disease or constellations of symptoms of borreliosis in Germany/Europe occasionally significantly differ from the US (1). Acrodermatitis chronica atrophicans, comparatively common in these parts, is almost unknown in the US, as is borreliolymphocytoma. Erythema migrans, arthritis, neuroborreliosis, and other manifestations often take a different clinical course in the US as they do in Europe. Because of these differences—some of them very pronounced—different preventive approaches (2) have been developed: In the US, a vaccine that was actually successful and effective against *B. burgdorferi* was taken off the market in 2001 because of side effects and lacking economic viability. Currently, antibiotic treatment with a single dose of 200 mg doxycycline, orally administered within 72 hours after the tick bite, seems promising in the US—in contrast to Germany. In terms of differential diagnoses and therapeutic strategy, the undifferentiated term "Lyme borreliosis" for all tickborne borrelioses, which was adopted perhaps slightly too uncritically from the Anglo-American literature, is confusing. For this reason, in 2003, I proposed a differentiation of currently (still) unified terminology into distinct variants of the infectious disease (3), such as has been the case for the different forms of hepatitis for a long time. But more is required: an interdisciplinary consensus (guideline) for borreliosis is needed in order to provide more detailed definitions of the occurring clinical entities of this infection with its many facets, which is often the subject of controversial discussions or used as an exclusion diagnosis, and to put this on a more differentiated footing.

DOI: 10.3238/arztebl.2009.0524a

REFERENCES

1. Hengge U, Tannapfel A, Tyring SK, Erbel R, Arendt G, Ruzicka T: Lyme borreliosis. *Lancet Infect Dis* 2003; 3: 489–500.
2. Hayes EB, Piesman J: How we can prevent Lyme disease? *N Engl J Med* 2003; 348: 2424–30.
3. Haufs MG: Preventing Lyme disease. *N Engl J Med* 2003; 349: 1192.
4. Nau R, Christen H, Eiffert H: Lyme disease—current state of knowledge. *Dtsch Arztebl Int* 2009; 106: 72–81.

Dr. med. Dr. rer. nat. Michael G. Haufs
Gluckweg 31, 48147 Münster, Germany
michaelhaufs@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Veterinary Medical Research

Since this is an anthroponosis that can affect grazing animals, especially horses, a look at research results from veterinary medicine is worth while, as this discipline has come to similar evaluations of clinical course and laboratory correlations.

I missed any comment about the effectiveness of insect repellents, because "wearing appropriate clothing" in the middle of summer is wishful thinking, especially for children. Searching the bends of the joints, the interdigital spaces especially on the feet, the gluteal folds, and the region behind the earlobes is essential.

DOI: 10.3238/arztebl.2009.0524b

REFERENCES

1. Nau R, Christen H, Eiffert H: Lyme disease—current state of knowledge. *Arztebl Int* 2009; 106: 72–81.

Marianne Temmesfeld

Schillerstr. 52
45894 Gelsenkirchen-Buer, Germany
familie-temmesfeld@web.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Higher Dosage at Later Stages

The authors report that seroconversion can be expected in 3 to 6% of cases after a tick bite, and that clinically manifest disease develops in only 0.3 to 1.4% of cases. How common is spontaneous cure at stages I and II? Do the authors have data/numbers on this that they could share?

In the skin biopsies of patients with erythema migrans who were treated in Slovenia, *borrelia* species (*B. burgdorferi sensu lato*) were found in the cultures of 19 of 1148 (1.7%) patients after antibiotic treatment had been completed (1). In five of the patients the authors specify the antibiotic treatment that was given: one patient received 2 g ceftriaxone i.v. for 14 days; two patients received cefuroxime (2×500 mg) for 14 and 10 days, respectively; one patient received amoxicillin orally (3×500 mg) for 14 days; and one patient received azithromycin orally for four days (1000 mg on the first day, followed by 1×500 mg). Persistence of the causative strain has been observed but seems rare.

Ceftriaxone at a dosage of 2 g/d is probably sufficient at the acute stage. At later stages, a higher dosage (2×2 g ceftriaxone) is obviously more favorable. As Schefold et al reported in 2003, in chronic disease a higher dosage of ceftriaxone—namely 2×2 g—should be given (3). If ceftriaxone is contraindicated, cefotaxime (3×2 g) can be given as an alternative. DOI: 10.3238/arztebl.2009.0524c

REFERENCES

1. Hunfeld KP, Ruzic-Sabljic E, Norris DE, Kraiczy P, Strle F: In vitro susceptibility testing of *Borrelia burgdorferi* sensu lato isolates cultured from patients with erythema migrans before and after antimicrobial chemotherapy. *Antimicrob Agents Chemother* 2005; 49: 1294–301.
2. Scheffold N, Bergler-Klein J, Sucker C, Cyran J: Kardiovaskuläre Manifestationsformen der Lyme-Borreliose: Therapeutische Situation im chronischen Stadium problematisch. *Dtsch Arztebl* 2003; 100(14): A 912–20.
3. Nau R, Christen H, Eiffert H: Lyme disease—current state of knowledge. *Dtsch Arztebl* 2009; 106: 72–81.

Prof. Dr. med. Pramod M. Shah

Gutzkowstr. 69
60594 Frankfurt am Main, Germany
infektiologie@online.de

Conflict of interest statement

The author has received honoraria for speaking and advisory activities from Sanofi-Aventis, Tanner-Cilag, and Pohl-Boskamp.

In Reply:

We welcome the interest expressed in our review article. Repellents, as discussed by Temmesfeld, are effective to only a limited degree. In most cases, the protective effect of repellents against ticks is clearly shorter than that against mosquitoes, so that the available substances have to be re-applied at relatively short intervals. It is of particular importance that bends of joints, interdigital spaces, gluteal folds, and the retroauricular region should be inspected thoroughly after any stay in an area populated by ticks.

As shown by Hunfeld's working group and mentioned by Shah, both in Frankfurt, persistence of culturable *Borrelia* species in the human organism may occur even after adequate antibiotic therapy for Lyme borreliosis (1). However, after guideline conform treatment this is a rare event, and it is unclear to what proportion the host organism will succeed in completely eliminating the *Borrelia* species—weakened by the preceding antibiotic treatment—on its own account without repeated treatment with antibiotics. With respect to reinfection by means of a repeated tick bite, we would not withhold a second course of antibiotic treatment, preferably on the basis of available sensitivity tests (2, 3) with ceftriaxone i.v. from any patient with clear clinical disease activity (for example, progressive neurological symptoms plus >6 months of persistent pleocytosis in the cerebrospinal fluid or persistent swollen joints plus positive borrelia polymerase chain reaction in fluid aspirated during puncture of a joint).

We are not in favor of the dosage increase of ceftriaxone to 4 g/d i.v., because after 3 weeks' treatment with this

dose the risk of precipitation of calcium-ceftriaxone salt in the gallbladder is high. The minimal inhibitory concentrations of different strains of *Borrelia burgdorferi* for ceftriaxone are 0.03 to 0.06 g/l (2, 3). Even after a single dose of ceftriaxone, ventricular CSF in patients without meningeal inflammation reached maximum concentrations of 0.18 to 1.04 mg/l (median 0.43 mg/l) (4). This means that they are one order of magnitude above the minimum inhibitory concentration. The elimination half life of ceftriaxone from CSF is about 17 hours (4), so that effective concentrations of ceftriaxone can be expected for the entire duration of treatment.

We think that the differentiation proposed by Haufs—of the variable European disease spectrum caused by different subspecies of *B burgdorferi* sensu lato versus the more uniform disease picture of the North American Lyme borreliosis (caused by *B burgdorferi* sensu stricto)—makes sense and is desirable. However, we are not in a position, nor do we wish to, decide here whether the term "non-Lyme" borreliosis is suitable for this or whether the disease should be termed "Eurasian" borreliosis.

DOI: 10.3238/arztebl.2009.0525

REFERENCES

1. Hunfeld KP, Ruzic-Sabljic E, Norris DE, Kraiczy P, Strle F: In vitro susceptibility testing of *Borrelia burgdorferi* sensu lato isolates cultured from patients with erythema migrans before and after antimicrobial chemotherapy. *Antimicrob Agents Chemother* 2005; 49: 1294–301.
2. Dever LL, Jorgensen JH, Barbour AG: In vitro antimicrobial susceptibility testing of *Borrelia burgdorferi*: a microdilution MIC method and time-kill studies. *J Clin Microbiol* 1992; 30: 2692–7.
3. Mursic VP, Wilske B, Schierz G, Holmberger M, Süß E: In vitro and in vivo susceptibility of *Borrelia burgdorferi*. *Eur J Clin Microbiol* 1987; 6: 424–6.
4. Nau R, Prange HW, Muth P et al.: Passage of cefotaxime and ceftriaxone into cerebrospinal fluid of patients with uninfamed meninges. *Antimicrob Agents Chemother* 1993; 37: 1518–24.
5. Nau R, Christen H, Eiffert H: Lyme disease—current state of knowledge. *Dtsch Arztebl* 2009; 106: 72–81.

Prof. Dr. med. Roland Nau

Prof. Dr. med. Hans-Jürgen Christen

Prof. Dr. med. Dr. rer. nat. Helmut Eiffert

Geriatrisches Zentrum
Evangelisches Krankenhaus Göttingen-Weende
Abteilung Neurologie und Medizinische Mikrobiologie
Universität Göttingen
Kinderkrankenhaus auf der Bult, Hannover
An der Lutter 24
37075 Göttingen, Germany
rnau@gwdg.de

Conflict of interest statement

The authors declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.