Published in final edited form as:

Free Radic Res. 2009 June ; 43(6): 613–621. doi:10.1080/10715760902942808.

Increasing expression of H- or L-ferritin protects cortical astrocytes from hemin toxicity

ZHI LI, JING CHEN-ROETLING, and RAYMOND F. REGAN*

Abstract

Iron toxicity may contribute to oxidative injury in cells surrounding an intracerebral hematoma. Cells detoxify iron by sequestering it in ferritin, a 24-mer heteropolymer constructed of H and L subunits. The relative antioxidant efficacy of H and L-ferritin has not been defined, and was tested in this study using an established model of hemin toxicity. Consistent with prior observations, cultures treated with 30 μ M hemin sustained loss of approximately half of cells by six hours, as measured by LDH and MTT assays, and a 14-fold increase in protein carbonyls. Increasing expression of either ferritin by adenoviral gene transfer prior to hemin treatment had a similar protective effect. Quenching of calcein fluorescence, a marker of the labile iron pool, in hemin-treated cultures was also equally reduced by either subunit. These results suggest that over-expression of either H or L ferritin protects astrocytes from hemin, and may be beneficial after CNS hemorrhage.

Keywords

Cell culture; free radical; hemoglobin toxicity; intracerebral hemorrhage; iron chelation; oxidative stress; stroke

1. Introduction

Iron toxicity may contribute to cell injury in tissue surrounding an intracerebral hemorrhage (ICH) [1]. The putative source of this iron is extravascular hemoglobin, which tends to release its heme moieties after undergoing auto-oxidation [2]. Their subsequent breakdown by the heme oxygenase enzymes releases equimolar iron, and likely accounts for the increase in nonheme iron observed in adjacent neurons and glial cells [3]. Nonheme iron is increased within 1-3 days of experimental ICH produced by direct blood injection in rat and rabbit models, and persists for at least three months in the former [1,4]. Some of this iron may be sequestered in ferritin, which is rapidly induced in peri-hematoma tissue [5]. However, it is unclear if this ferritin provides cells with any protection from heme toxicity. At best, it appears to be either insufficient or too late, since redox-active iron produces oxidative injury after experimental ICH despite upregulation of ferritin expression [6,7].

Mammalian ferritin is a 24-mer heteropolymer constructed of H and L subunits, with considerable variability in subunit composition in different cell populations. The antioxidant efficacy of H-rich and L-rich ferritin heteropolymers has been directly compared only in HeLa cells to date. Cozzi et al. reported that iron availability was negatively regulated by increasing expression of H-ferritin by gene transfer, but that L-ferritin had no effect per se [8]. The vulnerability of these cells to hydrogen peroxide (H_2O_2), which is an iron-dependent injury [9], was also inversely related to H-ferritin but not L-ferritin levels. In contrast, Orino et al.

*Corresponding Author, Department of Emergency Medicine, Thomas Jefferson University, 1020 Sansom Street, Thompson Building Room 239, Philadelphia, PA 19107, Telephone: 215-955-2695; FAX: 215-923-6225, E-mail: Raymond.Regan@jefferson.edu

reported that over-expression of either H or L-ferritin reduced oxidative stress equally in the same cell line after H₂O₂ treatment [10].

Differences in the antioxidant efficacy of H and L-ferritin have been attributed to the ferroxidase activity of the former, which is essential for rapid iron uptake by the heteropolymer and is lacking in the L-subunit [11]. Consistent with a critical role of H-ferritin expression in cellular iron homeostasis, homozygous H-ferritin knockout mouse embryos die at 3-9 days of development [12]. However, in vitro evidence suggests that increasing expression of L-ferritin may offer certain advantages to cells subjected to iron-loading conditions, due to the greater iron storage capacity, solubility, and stability of ferritins containing over 70-80% L-subunits [13]. These characteristics may account for the predominance of L-rich ferritin in cells that store iron, such as hepatocytes. The relative efficacy of H and L-ferritin in protecting CNS cells from injury produced by supraphysiologic iron concentrations has not yet been defined.

Although iron chelators are protective in some experimental ICH models [6,7], therapy with currently-available chelators may be limited by myriad toxic effects in humans, particularly if administered in the absence of systemic iron overload or at the high doses required for benefit in rodents [14-16]. An alternate or perhaps complementary approach is to increase the iron-sequestering capacity of cells adjacent to a hematoma via gene transfer, using vectors administered by stereotactic injection. Toward that end, we have constructed adenoviruses encoding murine H and L-ferritin genes driven by the human CMV promoter. In the present study, we compared the effect of increasing H or L-ferritin expression in an established astrocyte model of hemin toxicity.

2. Materials and Methods

Primary astrocyte cultures

All astrocyte cultures were prepared from 1-3 day postnatal C57BL/6 X 129/Sv mice that were bred in our animal facility. Mice were euthanized for culture preparation by deep isoflurane anesthesia followed by decapitation, via a protocol approved by the local Institutional Animal Care and Use Committee. Cortices were dissected free and incubated for 30 min in 0.09% trypsin. Tissue was dissociated by trituration through a Pasteur pipette with a narrowed (flame-polished) tip. The cell suspension was plated at a density of 1 hemisphere per 24-well plate, in medium consisting of minimal essential medium (MEM, Gibco/Invitrogen, Grand Island, NY) supplemented with 10% fetal bovine serum (Hyclone, Logan, UT), 10% equine serum (Hyclone), 10 ng/mL mouse epidermal growth factor (Sigma, St. Louis, MO), 23 mM glucose and 2 mM glutamine. Cultures were maintained at 37°C in a humidified 5% CO₂ atmosphere. Approximately two-thirds of the culture medium was replaced at five days in vitro and then twice weekly, using growth medium containing MEM, 23 mM glucose, 2 mM glutamine, and 10% equine serum.

H-ferritin heterozygous knockout (FTH^{+/-}) cultures were prepared using a similar procedure, except that cortices from each mouse were dissociated and plated separately. FTH^{+/-} mice were then distinguished from wild-type littermates by PCR-based genotyping, using genomic DNA extracted from residual brain tissue, and the following primer pair specific for the knockout gene:

5'-ATA CTT TCT CGG CAG GAG CA-3'

5'-TCA GAG CCA CAT CAT CTC GGT C-3'

Adenovirus preparation and astrocyte transfection

The open reading frames of the murine H-ferritin and L-ferritin genes were excised by digestion with EcoRI on the 5' end and NotI on the 3' end from pCMVSPORT6-mFTH1 (ATCC, GeneBank #BC012314) or pCMVSPORT6-LFER (ATCC, GeneBank# BC106146). The 0.9Kb (H-ferritin) and ~1Kb(L-ferritin) fragments were introduced to adenoviral shuttle vector pDUAL-CCM (Vector Biolabs, Philadelphia, PA) creating pDUAL-CMV- mFTH1 and pDUAL-CMV- mLFer. These constructs were sequenced and transfected to primary cultured wild type astrocytes (5 days in vitro, 1µg DNA per well) using Lipofectamine plus reagent in serum-free medium (OptiMEM, Invitrogen) for expression evaluation via immunoblotting. Sequence analysis demonstrated promoter - insert orientation and insert identity to mouse ferritin heavy chain or light chain sequences, using the Chromas sequences analyzing software and NCBI- The Basic Local Alignment Search Tool (BLAST).

SwaI endonuclease was used to release the insert from shuttle pDUAL-CMV-mFTH1 or pDUAL- CMV- mLFer and ligate the insert directly into the viral plasmid vector (pAd -VEC). For viral packaging in HEK293 cells, pAd-VEC- CMV-mFTH1 and pAd-VEC- CMV- mLFer were transfected in linear form (digested with Pac-I), to produce Ad-HF and Ad-LF. After propagation and harvesting, titer was quantified by cytopathic effect assay.

Adenoviral infection of astrocytes was accomplished in medium similar to feeding medium, except that it contained 3.3% equine serum. In a prior study using this culture system, treatment with 100 MOI (multiplicity of infection) of adenovirus serotype 5 in this medium resulted in transfection of approximately 80% of astrocytes [17].

Immunoblotting

Culture medium was aspirated, and cultures were then washed once with 1 ml MEM. After medium aspiration, 100µl ice-cold lysis buffer (210 mM mannitol, 70 mM sucrose, 5 mM HEPES, 1 mM EDTA, 0.1 % sodium dodecyl sulfate, 0.1 % Triton X-100) was added. Cells were collected, sonicated on ice, and centrifuged. Protein concentration of the supernatant was quantified by the BCA method (Pierce Biotechnology, Rockford, IL). Samples (20 µg in 30 µl) were diluted with 10 µl loading buffer (Tris-Cl 240 mmol/L, β-mercaptoethanol 20%, sodium dodecyl sulfate 8%, glycerol 40%, and bromophenol blue 0.2%) and heated to 95°C for 5 minutes. Proteins were separated on 12% SDS-PAGE gels (Ready Gel, Bio-Rad, Hercules, CA), and were then transferred to a polyvinylidene difluoride (PVDF) membrane (Imobilon-P, Millipore, Billerica, MA). After washing, nonspecific sites were blocked with 5% non-fat dry milk in a buffer containing 20 mM Tris, 500 mM NaCl, and 0.1% Tween 20 (pH 7.5) for 1 h at room temperature. For experiments depicted in Fig. 1, membranes were incubated at 4°C overnight with goat anti-H-ferritin or goat anti-L-ferritin (Santa Cruz Biotechnology, Santa Cruz, CA, USA, Product# SC-14416 and C-14420, 1:200 dilution). For experiments using FTH+/- and paired wild-type cultures (Fig. 7), rabbit anti-H-ferritin was a gift from Dr. James Connor (Pennsylvania State University, Hershey, PA, USA, 1:10,000 dilution), and anti-L-ferritin was a gift from Dr. Paolo Arosio (Università di Brescia, Brescia, Italy, 1:2000). After washing, membranes were treated with appropriate horseradish peroxidase-conjugated secondary antibody for one hour at room temperature. Immunoreactive proteins were visualized using Super Signal West Femto Reagent (Pierce, Rockford, IL) and Kodak Gel Logic 2200.

Cytotoxicity experiments

Experiments were conducted after 21 days in vitro. Confluent cultures were washed free of growth medium and were then exposed to hemin in serum-free medium consisting of MEM with 10 mM glucose (MEM10). In all experiments, cell injury was estimated at the end of the exposure period by visualizing cultures using phase contrast microscopy. Cell viability was

then quantified using the MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) and lactate dehydrogenase (LDH) release assays, as previously described in detail [18].

Cell protein oxidation was quantified by derivatized carbonyl assay, using the Oxyblot™ kit (Chemicon, Inc., Temecula, CA) and following the manufacturer's instructions. Briefly, proteins in cell lysates were protected from in vitro oxidation by adding 2-mercaptoethanol to a final concentration of 1%. 2,4-dinitrophenylhydrazine was then added to convert carbonyl groups to 2,4-dinitrophenylhydrazone (DNP) derivatives. Proteins were separated on 12% polyacrylamide gels and were transferred to a PVDF membrane filter. After incubation with rabbit anti-DNP primary antibody (1:150) followed by goat anti-rabbit HRP-conjugated secondary antibody (1:300), immunoreactive proteins were detected as described above. Oxidized protein bands densities were analyzed using Kodak 1D software.

Assessment of the labile iron pool

Changes in cell chelatable iron were detected by quantifying calcein fluorescence quenching in hemin-treated cultures [19]. After medium exchange to remove extracellular hemin, cultures were loaded with calcein by incubation for 15 minutes with 0.1 μ M calcein-AM, its cell-permeable ester. Cultures were then examined using an inverted microscope with epifluorescence attachment and an FITC filter. Digital 100X images in the center of each well (1 field/culture) were rapidly captured, and fluorescence intensity was quantified using Scanalytics IPLab software. Membrane integrity, essential for calcein retention, was verified by staining with propidium iodide, which is excluded by intact cells. For these experiments, hemin exposure was conducted in 3.3% equine serum, which attenuates the toxicity of hemin, and was terminated by medium exchange at three or five hours. Fluorescence of hemin-treated cultures was expressed as a percentage of that in sister cultures from the same plating treated with the same adenovirus, but not with hemin.

Data analyses

All data were analyzed using one-way analysis of variance (ANOVA) followed by the Bonferroni multiple comparisons test, using Graphpad (San Diego, CA) Prism.

3. Results

Treatment with Ad-HF and Ad-LF increases ferritin expression

We have previously demonstrated that transgene expression using adenovirus serotype 5 in this culture system is optimized at a dose of 100 MOI for 24h [17]. These observations were confirmed using ad-HF and ad-LF (Fig. 1). In cultures subjected to medium exchange only, baseline H and L-ferritin levels were minimal when membranes were probed with isoform-specific antibodies purchased from Santa Cruz Biotechnology. Over a range of 10-200 MOI, maximal ferritin expression was observed at 100 MOI (Fig. 1) for each construct. Treatment with an empty control adenovirus (Ad-Null) at this dose did not increase endogenous H or L-ferritin expression.

Protective effect of H or L-ferritin over-expression

Consistent with prior observations in this system [17], cultures pretreated with 100 MOI Ad-Null for 24h and then treated with 30 μ M hemin sustained widespread cell injury, which was morphologically apparent within 6h (Fig. 2). At this time point, $44.0 \pm 1.5\%$ of culture LDH was released into the culture medium (Fig. 3A). In cultures pretreated with either Ad-HF or Ad-LF, less injury was detected, with release of $22.0 \pm 1.7\%$ and $23.3 \pm 2.0\%$ of culture LDH,

respectively. A similar but somewhat weaker effect was observed when injury was assessed by the ability of cells to reduce MTT to formazan (Fig. 3B).

Protein carbonylation is a sensitive marker of hemin-mediated oxidative injury in this culture system [18]. Protein carbonyl levels were increased by 14-fold over baseline in cultures pretreated with 100 MOI Ad-Null for 24h followed by 30 μ M hemin for 6 h. Carbonyls were reduced by approximately 40% in cultures pretreated with either Ad-HF or Ad-LF (Fig. 4).

Effect of ferritin over-expression on the labile iron pool

In cultures pretreated with Ad-Null, hemin treatment for 3-5 hours reduced calcein fluorescence by over 90% (Figs. 5, 6). Most of this quenching was prevented in cultures pretreated with either Ad-HF or Ad-LF. Fluorescence differences in cultures pretreated with Ad-HF and Ad-LF were not statistically significant. Quenching was completely reversed by treating cultures with the cell-permeable iron chelator 1,10-phenanthroline (100 μ M) for 15 minutes. The fluorescence intensity in cultures treated with hemin for five hours followed by 1,10-phenanthroline for 15 min exceeded that of control cultures treated with 1,10-phenanthroline after exposure to the experimental medium (MEM10) only.

Effect of decreased H:L ferritin ratio on cell vulnerability to hemin

Homozygous H-ferritin gene knockout is lethal in early gestation, but heterozygotes (FTH^{+/-}) survive and are fertile [20]. These mice express less H-ferritin in the CNS but compensate by increasing L-ferritin expression. In order to investigate the effect of altering the H:L ferritin ratio on astrocyte vulnerability to hemin, cultures were prepared from FTH^{+/-} mice and their wild-type littermates. The antibodies used in these experiments were more sensitive than the commercially-available antibodies used in experiments depicted in Fig. 1, and detected H and L-ferritin expression in control cultures subjected to medium exchange only. In confluent FTH^{+/-} cultures, H-ferritin expression was reduced to almost half of that in wild-type cultures at baseline or after ferritin induction by hemoglobin (Fig. 7), which slowly delivers heme to astrocytes and is nontoxic to this cell population at low micromolar concentrations [21]. L-ferritin expression was increased by 2.2-fold at baseline and remained significantly increased with induction. The vulnerability of FTH^{+/-} and wild-type cells to hemin was not significantly different despite the decreased H:L ferritin ratio of the former (Fig. 7).

4. Discussion

At low micromolar concentrations, hemin produces an iron-dependent oxidative injury in cultured cells [22,23]. The protective effect of ferritin over-expression is consistent with prior observations that pretreatment with horse spleen ferritin, which is taken up by endocytosis, attenuated hemin toxicity in endothelial cells and astrocytes [22,24]. However, data comparing the efficacy of H and L-ferritin against heme-mediated oxidative injury have not been previously reported. The present results demonstrate that in primary cultured astrocytes, increasing expression of either subunit has a comparable effect on the labile iron pool and cell injury after hemin treatment. The hypothesis that both H and L-subunits are protective in this model is further supported by observations that altering the H:L ferritin ratio by heterozygous H-ferritin gene knockout, which is associated with a compensatory increase in L-ferritin, had no effect on astrocyte vulnerability to hemin.

The present results are consistent with those of Orino et al. [10], who reported that over-expression of either H or L-ferritin attenuated cellular reactive oxygen species levels after H₂O₂ treatment. However, they disagree with data of Cozzi et al. suggesting that cell iron availability and vulnerability to oxidant challenge were primarily regulated by H-ferritin levels only [8]. These disparate results are likely reconciled by differences in endogenous H-ferritin

expression in these models. Iron uptake by ferritin heteropolymers is optimal when the H-ferritin content is 20-30%, suggesting that additional ferroxidase activity provides no benefit [13,25]. Furthermore, L-rich heteropolymers are more soluble and incorporate up to fourfold more iron than H-homopolymers [13]. Astrocytes in this culture system rapidly upregulate H-ferritin after exposure to heme, in contrast with the very weak expression observed in neurons [26]. Since heme breakdown by the heme oxygenases releases ferrous iron [27], and since ferritin heteropolymers incorporate only ferric iron in their mineral core, ferroxidase activity is likely critical for rapid iron uptake. The present results suggest that adequate ferroxidase activity is provided by endogenous H-ferritin in cultured astrocytes, even in heterozygous knockouts.

The effect of ferritin gene transfer on cell viability as measured by the MTT assay was less than that observed with the LDH release assay. Reduction of MTT to its formazan product is catalyzed in part by mitochondrial enzymes [28]. Heteropolymers constructed of H and L-ferritin are the primary cytoplasmic iron regulators. However, a unique ferritin encoded by a separate nuclear gene predominates in mitochondria [29], and was not directly altered in the current series of experiments, although indirect alterations as a consequence of the experimental conditions cannot be excluded. The effect of specifically increasing expression of mitochondrial ferritin on heme-mediated astrocyte injury seems a worthy topic for future investigation. It is noteworthy that when over-expressed in HeLa cells, it reduced the levels of reactive oxygen species after treatment with H₂O₂ or antimycin A [30].

The protection provided by H or L-ferritin gene transfer was also less than predicted from its effect on the labile iron pool, as measured by calcein fluorescence quenching. Two factors may contribute to this discrepancy. First, heme is lipid soluble and accumulates in membranes, where it may catalyze free radical reactions in a cellular compartment that is inaccessible to hydrophilic ferritin and calcein [31]. Second, the calcein method may underestimate the labile iron pool. Tenopoulou et al. have recently reported that calcein does not measure lysosomal iron, due to its limited ability to penetrate lysosomal membranes and chelate iron in an acidic environment [32]. Lysosomal iron mediates most of the cell death produced by H₂O₂ in vitro [33]. Although its role in heme-mediated injury has not been defined, any contribution may not be detected by changes in calcein fluorescence.

Ferritin expression is minimal at baseline in this culture system, but it is induced within 6 hours by heme treatment [26]. While that induction likely contributes to the resistance of astrocytes to the more-slowly developing toxicity of hemoglobin, it apparently occurs too late to prevent widespread cell death per se in this heme toxicity model. This observation is consistent with the hypothesis that endogenous ferritin levels may be suboptimal in cells adjacent to an intracerebral hematoma, which has a heme content in the high micromolar range [34], and that therapies aiming to upregulate its expression prior to a heme-mediated oxidative insult may be beneficial. In support of this hypothesis, we have recently reported that neurons lacking iron regulatory protein-2 (IRP2), which represses H and L-ferritin translation by binding to an iron responsive element in the 5' regions of their mRNA, are highly resistant to iron-dependent oxidative injury in vitro and to experimental intracerebral hemorrhage in vivo [26,35]. It remains to be determined if any treatment administered after the onset of hemorrhage would alter ferritin levels quickly enough to increase cell resistance to heme. It is noteworthy, however, that iron-mediated injury is delayed until approximately 2-3 days after experimental ICH, which likely reflects the time required for erythrocyte lysis and hemoglobin oxidation [36]. The therapeutic window for the component of injury produced by heme breakdown may therefore be sufficient for delivery of viral vectors to astrocytes adjacent to a hematoma.

The long-term consequences of iron accumulation in the ferritin of astrocytes after hemorrhagic injury is undefined, and may best be addressed in vivo, due to the shorter lifespan of cells in

primary cultures. Studies using genetically modified mice suggest that sustained upregulation of ferritin per se may be associated with toxicity, although precise mechanisms have yet to be delineated. In aging ceruloplasmin knockout mice, ferritin-bound iron accumulation in astrocytes is associated with reduced immunoreactivity to the astrocyte markers glial fibrillary acidic protein and S100 β , consistent with cell loss [37]. IRP2 knockout mice may develop a late-onset movement disorder, in which iron deposition in white matter tracts and nuclei precede neurodegeneration [38]. The toxicity of sustained ferritin over-expression may be mitigated by the use of adenoviral vectors, which tend to produce transient gene expression [39]. However, any benefit may then be negated by a direct cytotoxic or inflammatory response [40]. Further investigation is needed to determine if ferritin gene transfer is feasible in vivo, and if it also protects neurons and other vulnerable cell populations.

5. Acknowledgements

This study was supported by a grant from the National Institutes of Health (NS042273) to R.F.R.. The authors thank Dr. James Connor for providing H-ferritin heterozygous knockout mice and anti-H-ferritin, and Dr. Paolo Arosio for anti-L-ferritin.

6. References

- [1]. Hua Y, Nakamura T, Keep RF, Wu J, Schallert T, Hoff JT, Xi G. Long-term effects of experimental intracerebral hemorrhage: the role of iron. *J Neurosurg* 2006;104:305–12. [PubMed: 16509506]
- [2]. Bunn HF, Jandl JH. Exchange of heme along hemoglobins and between hemoglobin and albumin. *J. Biol. Chem* 1967;243:465–475. [PubMed: 4966113]
- [3]. Koeppen AH, Dickson AC, Smith J. Heme oxygenase in experimental intracerebral hemorrhage: the benefit of tin-mesoporphyrin. *J. Neuropathol. Exp. Neurol* 2004;63:587–597. [PubMed: 15217087]
- [4]. Koeppen AH, Dickson AC, McEvoy JA. The cellular reactions to experimental intracerebral hemorrhage. *J Neurol Sci* 1995;134(Suppl):102–112. [PubMed: 8847540]
- [5]. Wu J, Hua Y, Keep RF, Nakamura T, Hoff JT, Xi G. Iron and iron-handling proteins in the brain after intracerebral hemorrhage. *Stroke* 2003;34:2964–2969. [PubMed: 14615611]
- [6]. Wan S, Hua Y, Keep RF, Hoff JT, Xi G. Deferoxamine reduces CSF free iron levels following intracerebral hemorrhage. *Acta Neurochir Suppl* 2006;96:199–202. [PubMed: 16671454]
- [7]. Masuda T, Hida H, Kanda Y, Aihara N, Ohta K, Yamada K, Nishino H. Oral administration of metal chelator ameliorates motor dysfunction after a small hemorrhage near the internal capsule in rat. *J Neurosci Res* 2007;85:213–22. [PubMed: 17061255]
- [8]. Cozzi A, Corsi B, Levi S, Santambrogio P, Biasiotto G, Arosio P. Analysis of the biologic functions of H- and L-ferritins in HeLa cells by transfection with siRNAs and cDNAs: evidence for a proliferative role of L-ferritin. *Blood* 2004;103:2377–83. [PubMed: 14615379]
- [9]. Starke PE, Farber JL. Ferric iron and superoxide ions are required for the killing of cultured hepatocytes by hydrogen peroxide. Evidence for the participation of hydroxyl radicals formed by an iron-catalyzed Haber-Weiss reaction. *J Biol Chem* 1985;260:10099–104. [PubMed: 2991275]
- [10]. Orino K, Lehman L, Tsuji Y, Ayaki H, Torti SV, Torti FM. Ferritin and the response to oxidative stress. *Biochem J* 2001;357:241–7. [PubMed: 11415455]
- [11]. Arosio P, Levi S. Ferritin, iron homeostasis, and oxidative damage. *Free Radic Biol Med* 2002;33:457–63. [PubMed: 12160928]
- [12]. Ferreira C, Bucchini D, Martin ME, Levi S, Arosio P, Grandchamp B, Beaumont C. Early embryonic lethality of H ferritin gene deletion in mice. *J Biol Chem* 2000;275:3021–4. [PubMed: 10652280]
- [13]. Levi S, Santambrogio P, Cozzi A, Rovida E, Corsi B, Tamborini E, Spada S, Albertini A, Arosio P. The role of the L-chain in ferritin iron incorporation. Studies in homo and heteropolymers. *J. Mol. Bio.* 1994;238:649–654. [PubMed: 8182740]
- [14]. Porter JB, Huehns ER. The toxic effects of desferrioxamine. *Baillieres Clin Haematol* 1989;2:459–74. [PubMed: 2660937]
- [15]. Kushner JP, Porter JP, Olivieri NF. Secondary Iron Overload. *Hematology* 2001;2001:47–61. [PubMed: 11722978]

- [16]. Chaston TB, Richardson DR. Iron chelators for the treatment of iron overload disease: relationship between structure, redox activity, and toxicity. *Am J Hematol* 2003;73:200–10. [PubMed: 12827659]
- [17]. Teng ZP, Chen J, Chau LY, Galunic N, Regan RF. Adenoviral transfer of the heme oxygenase-1 gene protects cortical astrocytes from heme-mediated oxidative injury. *Neurobiol. Dis* 2004;17:179–187. [PubMed: 15474356]
- [18]. Chen J, Regan RF. Increasing expression of heme oxygenase-1 by proteasome inhibition protects astrocytes from heme-mediated oxidative injury. *Curr Neurovasc Res* 2005;2:189–96. [PubMed: 16181113]
- [19]. Kakhlon O, Cabantchik ZI. The labile iron pool: characterization, measurement, and participation in cellular processes(1). *Free Radic Biol Med* 2002;33:1037–46. [PubMed: 12374615]
- [20]. Thompson K, Menzies S, Muckenthaler M, Torti FM, Wood T, Torti SV, Hentze MW, Beard J, Connor J. Mouse brains deficient in H-ferritin have normal iron concentration but a protein profile of iron deficiency and increased evidence of oxidative stress. *J Neurosci Res* 2003;71:46–63. [PubMed: 12478613]
- [21]. Chen-Roetling J, Regan RF. Effect of heme oxygenase-1 on the vulnerability of astrocytes and neurons to hemoglobin. *Biochem Biophys Res Commun* 2006;350:233–7. [PubMed: 16999934]
- [22]. Balla G, Jacob HS, Balla J, Rosenberg M, Nath K, Apple F, Eaton JW, Vercellotti GM. Ferritin: A cytoprotective strategem of endothelium. *J. Biol. Chem* 1992;267:18148–18153. [PubMed: 1517245]
- [23]. Goldstein L, Teng ZP, Zeserson E, Patel M, Regan RF. Hemin induces an iron-dependent, oxidative injury on human neuron-like cells. *J. Neurosci. Res* 2003;73:113–121. [PubMed: 12815715]
- [24]. Regan RF, Kumar N, Gao F, Guo YP. Ferritin induction protects cortical astrocytes from heme-mediated oxidative injury. *Neuroscience* 2002;113:985–994. [PubMed: 12182902]
- [25]. Santambrogio P, Levi S, Cozzi A, Rovida E, Albertini A, Arosio P. Production and characterization of recombinant heteropolymers of human ferritin H and L chains. *J Biol Chem* 1993;268:12744–8. [PubMed: 8509409]
- [26]. Regan RF, Chen M, Li Z, Zhang X, Benvenisti-Zarom L, Chen-Roetling J. Neurons lacking iron regulatory protein-2 are highly resistant to the toxicity of hemoglobin. *Neurobiol Dis* 2008;31:242–249. [PubMed: 18571425]
- [27]. Yoshida T, Migita CT. Mechanism of heme degradation by heme oxygenase. *J Inorg Biochem* 2000;82:33–41. [PubMed: 11132636]
- [28]. Bernas T, Dobrucki J. Mitochondrial and nonmitochondrial reduction of MTT: interaction of MTT with TMRE, JC-1, and NAO mitochondrial fluorescent probes. *Cytometry* 2002;47:236–42. [PubMed: 11933013]
- [29]. Santambrogio P, Biasiotto G, Sanvito F, Olivieri S, Arosio P, Levi S. Mitochondrial Ferritin Expression in Adult Mouse Tissues. *J. Histochem. Cytochem* 2007;55:1129–1137. [PubMed: 17625226]
- [30]. Campanella A, Rovelli E, Santambrogio P, Cozzi A, Taroni F, Levi S. Mitochondrial ferritin limits oxidative damage regulating mitochondrial iron availability: hypothesis for a protective role in Friedreich ataxia. *Hum Mol Genet* 2009;18:1–11. [PubMed: 18815198]
- [31]. Hebbel RP, Eaton JW. Pathobiology of heme interaction with the erythrocyte membrane. *Sem. Hematol* 1989;26:136–149.
- [32]. Tenopoulou M, Kurz T, Doulias PT, Galaris D, Brunk UT. Does the calcein-AM method assay the total cellular ‘labile iron pool’ or only a fraction of it? *Biochem J* 2007;403:261–6. [PubMed: 17233627]
- [33]. Starke PE, Gilbertson JD, Farber JL. Lysosomal origin of the ferric iron required for cell killing by hydrogen peroxide. *Biochem Biophys Res Commun* 1985;133:371–9. [PubMed: 4084283]
- [34]. Letarte PB, Lieberman K, Nagatani K, Haworth RA, Odell GB, Duff TA. Hemin: levels in experimental subarachnoid hematoma and effects on dissociated vascular smooth muscle cells. *J. Neurosurg* 1993;79:252–255. [PubMed: 8331409]
- [35]. Chen, M.; Chen-Roetling, J.; Regan, RF. Neuroscience Meeting Planner. Society for Neuroscience, Online; Washington, DC: 2008. Reduced oxidative injury after experimental intracerebral hemorrhage in iron regulatory protein-2 knockout mice. 757.12

- [36]. Xi GH, Keep RF, Hoff JT. Erythrocytes and delayed brain edema formation following intracerebral hemorrhage in rats. *J. Neurosurg* 1998;89:991–996. [PubMed: 9833826]
- [37]. Jeong SY, David S. Age-related changes in iron homeostasis and cell death in the cerebellum of ceruloplasmin-deficient mice. *J Neurosci* 2006;26:9810–9. [PubMed: 16988052]
- [38]. LaVaute T, Smith S, Cooperman S, Iwai K, Land W, Meyron-Holtz E, Drake SK, Miller G, Abu-Asab M, Tsokos M, Switzer R 3rd, Grinberg A, Love P, Tresser N, Rouault TA. Targeted deletion of the gene encoding iron regulatory protein-2 causes misregulation of iron metabolism and neurodegenerative disease in mice. *Nat Genet* 2001;27:209–14. [PubMed: 11175792]
- [39]. Gonzalez SC, McMenamin MM, Charlton HM, Goodman J, Lantos T, Simpson C, Wood MJ. Readministration of adenoviral gene delivery to dopamine neurons. *Neuroreport* 2007;18:1609–14. [PubMed: 17885611]
- [40]. McMenamin MM, Lantos T, Carter EE, Hamilton L, Charlton HM, Gonzalez SC, Wood MJ. Neuropathological consequences of delivering an adenoviral vector in the rat brain. *J Gene Med* 2004;6:740–50. [PubMed: 15241781]

A **B** **Fig. 1.**

Dose-response effect of adenoviruses encoding murine H (Ad-HF) or L-ferritin (Ad-LF) in primary astrocyte cultures. Representative immunoblots from astrocyte cultures treated with 10-200 MOI (multiplicity of infection) Ad-HF (A), Ad-LF (B), or control empty adenovirus (Null) for 24 hours, stained with goat antibody specific for H or L-ferritin (Santa Cruz Biotechnology). Control cultures were subjected to medium exchange (sham-wash, SW) only. M denotes protein molecular weight marker. Membranes were also stained with rabbit anti- β -actin (1:1250, Sigma) as a gel loading control. Entire membranes are shown, which were cut to allow separate incubation with different secondary antibodies.

Fig. 2.

Ad-HF and Ad-LF protect astrocytes from hemin. Phase-contrast photomicrographs of astrocyte cultures, 6 hours after: (A) medium change only; (B) exposure to 30 μ M hemin after 24 hour incubation with 100 MOI Ad-Null; (C) exposure to 30 μ M hemin after 24 hour incubation with 100 MOI Ad-HF; (D) exposure to 30 μ M hemin after 24 hour incubation with Ad-LF. Scale bar = 100 μ m.

Fig. 3.

Quantification of the protective effect of Ad-HF and Ad-LF. A) Mean culture percentage LDH release ($\pm$ SEM), 6 hours after treatment with 30 μ M hemin, preceded by 24 hour incubation with 100 MOI Ad-Null, Ad-HF or Ad-LF. Values are scaled to those in control sister cultures treated with 0.1% Triton X-100 (= 100), which releases 100% of culture LDH. The LDH values in sister cultures subjected to sham-wash only were subtracted from all values. B) Quantification of cell viability using the MTT assay at 3 hours. Values are scaled to those in control sister cultures subjected to sham-wash only (=100). *** P <0.001, * P < 0.05 v. mean value in cultures pretreated with Ad-Null, n = 14-24/condition, Bonferroni multiple comparisons test.

Fig. 4.

Astrocyte protein oxidation after hemin treatment is reduced by pretreatment with Ad-HF or Ad-LF. Upper figure is a representative immunoblot of lysates from astrocyte cultures treated with sham wash (SW) only or with 30 μ M hemin for 6 hours, preceded by 24 hour incubation with 100 MOI Ad-Null, Ad-HF or Ad-LF. Bars represent mean carbonyl signal intensities ($\pm$ SEM, $n = 5$ /condition). *** $P < 0.001$ v. mean value in cultures pretreated with Ad-Null, Bonferroni multiple comparisons test.

Fig. 5. Calcein fluorescence quenching after hemin treatment is attenuated by pretreatment with Ad-HF or Ad-LF. A) Phase contrast photomicrograph of culture treated with 100 MOI Ad-Null for 24 hours; B) Same culture as A, demonstrating baseline calcein fluorescence without hemin treatment; C) Sister culture, treated with 100 MOI Ad-Null for 24 hours followed by 30 μ M hemin for 3 hours; fluorescence intensity is markedly diminished; D) Same culture as C, 15 minutes after treatment with 100 μ M 1,10-phenanthroline, demonstrating reversal of fluorescence quenching with a cell-permeable iron chelator; E, F) Cultures pretreated with 100 MOI Ad-HF (E) or Ad-LF (F) for 24 hours prior to 3 hour hemin treatment; fluorescence quenching is reduced compared with culture pretreated with Ad-Null.

Fig. 6.

Quantification of calcein fluorescence quenching by hemin after treatment with viral vectors. Cultures were treated with 100 MOI Ad-Null, Ad-HF, or Ad-LF for 24 hours. Half were then treated with 30 μ M hemin for 3 or 5 hours. Mean calcein fluorescence ($\pm$ SEM, $n = 5-8$ /condition) is expressed as a percentage of that in control cultures not treated with hemin. Cultures pretreated with Ad-Null were subsequently treated with 100 μ M 1,10-phenanthroline (Phe) to reverse quenching. *** $P < 0.001$ v. corresponding value in cultures pretreated with Ad-Null.

Fig. 7.

Heterozygous H-ferritin gene deletion has no effect on astrocyte vulnerability to hemin. A) Cell injury, as assessed by LDH release and MTT assays, in wild-type and H-ferritin heterozygous knockout (FTH ^{+/-}) cultures produced by exposure to 30 μM hemin as in Fig. 3. Mean culture LDH values (±SEM, n= 28-32/condition) are scaled to those in sister cultures treated with 0.1% Triton X-100 (=100), after subtraction of mean values in sham-washed (SW) controls, to yield the value associated with hemin toxicity. Values for MTT reduction to formazan (n=24/condition) are scaled to those in sister cultures subjected to sham-wash only (=100). B) Expression of H-ferritin and L-ferritin in astrocyte cultures prepared from wild-type (WT) and FTH ^{+/-} mice, at indicated intervals after treatment with 10 μM hemoglobin, using non-commercial antibodies described in Methods. C) Bars represent mean ferritin band densities (±SEM, n= 4 condition). **P<0.01, ***P<0.001 v. corresponding wild-type condition, Bonferroni multiple comparisons test.