Published in final edited form as:

J Org Chem. 2009 August 7; 74(15): 5516–5522. doi:10.1021/jo900767x.

A Divergent Synthesis of the Δ^{13} -9-Isofurans

Douglass F. Taber^{*}, Peiming Gu, and Rui Li

Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19716

Abstract

A stereodivergent total synthesis of the Δ^{13} -9-isofurans has been developed. The four core substituted tetrahydrofurans were prepared by the Sharpless asymmetric epoxidation and Sharpless asymmetric dihydroxylation followed by cascade cyclization. The relative configuration at C-8 was inverted by oxidation followed by immediate L-selectride reduction. The relative configuration of the C-15 diastereomers were assigned by (*S*)-Binol/LAH/EtOH reduction of the corresponding enone. This synthesis of the Δ^{13} -9-isofurans will provide sufficient material for further investigation of their biological activity.

Introduction

The isofurans (Fig. 1), represented by SC- Δ^{13} -9-IsoF **1aaa**, are produced by human metabolism of arachidonic acid.^{1,2} Previously,³ we prepared the two diastereomeric isofurans that we judged were the closest in structure to the enzymatically produced human hormone PGF_{2 α} . One of the diastereomers was found to have significant bioactivity. On re-preparation, however, no bioactivity was observed. Suspecting that the initially observed bioactivity might have been due to a minor contaminating diastereomer, we have developed a general strategy for the preparation of each of the 32 of the enantiomerically-pure diastereomers of the Δ^{13} -9-isofurans. As the previous route to the isofurans had suffered from poor regioselectivity in the dihydroxylation step, we have devised an improved and much more robust route to the isofurans.

Taberdf@udel.edu.

Supporting Information Available. General experimental procedures, experimental procedures and spectra for all new compounds, and details of the x-ray structural analysis of **11a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Results and Discussion

Our interest was to develop a stereodivergent route to the Δ^{13} -9-isofurans, such that each of the enantiomerically-pure (ep) diastereomers could be prepared by branching from advanced intermediates. The eight substituted tetrahydrofurans **2a–2h** (Scheme 1) would each lead to four of the final thirty-two (ep) diastereomers. The four intermediates **2a–2d** were to be prepared from the previously unknown monotrityl ether **3** by Sharpless asymmetric epoxidation (SAE)⁴ followed by Sharpless asymmetric dihydroxylation (SAD).^{3,5} We planned to prepare the tetrahydrofurans **2e–2h** from the intermediates **2a–2d** by Mitsunobu esterification^{3,6} followed by hydrolysis.

For this approach to the isofurans to be successful, there were three central issues to be addressed (Eq. 1):

1. Would the asymmetric dihydroxylation of the epoxy sulfonate derived from **3** show matched vs unmatched diastereoselectivity?
2. Could the stereocenters at C-8 and C-11 be inverted without elimination?
3. Could the anticipated C-15 allylic alcohol diastereomers be separated?

(1)

Preparation of the monotrityl ether **3**

We prepared **3** (Scheme 2) from methyl 4-bromocrotonate **5**. Reduction by $\text{LiAlH}_4/\text{AlCl}_3$ ⁷ afforded the allylic alcohol **6**. This alcohol was coupled⁸ with propargyl alcohol to give a mixture (2:1) of the linear product **7a** and the branched product **7b**, that were not easily separable. The mixture was submitted to LiAlH_4 reduction to give the diols **8a** and **8b**. After protection with TrCl , the monoether **3** was readily purified.

Preparation of the Four Core Tetrahydrofurans

The monoether **3** was first carried on (Scheme 3, Table 1) to the epoxy alcohol **9a** by Sharpless asymmetric epoxidation with the adjunct D-diethyl tartrate (D-DET) (Scheme 3). The derived benzenesulfonate **10a** was subjected to the Sharpless asymmetric dihydroxylation with AD-mix followed by direct treatment with K_2CO_3 in MeOH ^{3,9} to give the cyclized product. Using L-DET in the Sharpless asymmetric epoxidation, when AD-mix α was employed (entry 1), it gave a 3:1 mixture of diastereomers, that were not readily separable by silica gel chromatography. We were pleased to observe that the two diastereomers were readily separable after silylation of the secondary hydroxyl group, leading to **2a** and **2b**. The structure of **2a** was confirmed by X-ray analysis of the crystalline free alcohol (**11a**, Scheme 4). Alternatively, when AD-mix β was used (entry 2), primarily, (>9:1) the single diastereomer **2b** was found. Similarly, using L-DET in the Sharpless asymmetric epoxidation and AD-mix α in the dihydroxylation (entry 4) delivered primarily (>9:1) the diastereomer **2d**, whereas L-DET followed by AD-mix β (entry 3) gave a 2:1 mixture of diastereomers of **2c** and **2d**. Note that

the major diastereomers prepared this way showed >99% ee (chiral HPLC), while the minor diastereomers (entry 1, entry 3) were about 67% ee and 53% ee, respectively (chiral HPLC). Only the high ee diastereomers were carried on in the synthesis.

Inversion of the C-11 Stereochemistry

To achieve stereodivergence, it was necessary to invert the alcohols at C-11 (isofuran numbering, Scheme 4). Deprotection of the silyl group of **2a** with TBAF gave the free alcohol **11a**. The secondary alcohol at C-11 participated efficiently in Mitsunobu esterification⁶ to give, after methanolysis, the inverted alcohol **11e**. Protection of **11e** with TBS gave the tetrahydrofuran diastereomer **2e**. Similarly, **2b** – **2d** were converted to **2f** – **2h**.

Construction of the Upper Side Chain

Lewis-acid BF₃·OEt₂ assisted¹⁰ opening (Scheme 5) of the epoxide **2a** with the lithium anion derived from the commercially available 4-pentynenitrile at low temperature gave efficient conversion to the alkyne **12a**. Partial hydrogenation using P-2 Ni catalyst¹¹ afforded the *cis*-alkene **13aa**. Protection of **13aa** with TBS resulted in the trityl ether **4aa**.

Inversion of the C-8 Stereochemistry

When Mitsunobu inversion was attempted with alcohol **13aa** (Scheme 6), predominantly dehydration was observed. Fortunately, oxidation¹² followed by immediate L-Selectride reduction of the alcohol **13aa** gave predominantly the inverted alcohol **13ab**. Protection of **13ab** with TBS gave the diastereomer **4ab**.

Construction of the Lower Side Chain

The next step in the synthesis was the removal of the trityl group from **4aa** (Scheme 7). Formic acid¹³ worked well with **4aa**, but destroyed **4ab**. Deprotection of **4aa** with 3 eq of Et₂AlCl at –50 °C¹⁴ gave the desired alcohol **14aa** accompanied by some unidentified impurities. Fortunately, deprotection of **4aa** and related compounds with 5 eq of Et₂AlCl at –78 °C for 10 min gave clean conversion to the primary alcohol. Dess-Martin oxidation¹² of the alcohol **14aa** afforded the aldehyde. Horner-Emmons condensation of the aldehyde with commercial dimethyl 2-oxo-heptylphosphonate **15** gave the enone **16aa**. Luche reduction¹⁵ of the enone gave the allylic alcohols **17aaa** and **17aab** in a ratio of about 1:1. We were pleased to observe that these two diastereomers were readily separable by silica gel chromatography.

Assignment of the C-15 Absolute Configuration

The enone **16aa** (Scheme 7) was also reduced with LiAlH₄/(*S*)-Binol/EtOH at –100 °C, that was expected¹⁶ to give predominantly **17aaa**. The major diastereomer from the reduction was contaminated with the (*S*)-Binol. However, this material was found to be congruent with the upper TLC spot from the Luche reduction by ¹³C NMR (δ 72.4 vs 72.4, lower spot δ 72.6). This allowed us to assign the absolute configurations at C-15 of the two diastereomeric alcohols **17aaa** (*S*) and **17aab** (*R*).

Synthesis of the Isofurans

Desilylation of **17aaa** with TBAF gave the free alcohol (Scheme 8). The nitrile group was hydrolyzed¹⁷ with 15% aqueous NaOH in EtOH at 85 °C, and the desired carboxylic acid **1aaa** was isolated in excellent yield. Isofuran **1aab** was prepared in the same way. The ¹³C NMR of the two free acids appeared to be complicated, so the **1aab** was converted to its methyl ester **18aab**. Both **1aaa** and **1aab** and the methyl ester **18aab** were found to be congruent (¹H, ¹³C NMR, [α]_D) to the substances that we had previously^{3a} prepared.

Using this approach, we have prepared all thirty-two of the enantiomerically pure diastereomers of Δ^{13} -9-IsoF. The full experimental details for the preparation of a representative six more of those diastereomers are included in the supporting information, along with physical characteristics and spectroscopic data for each of the other 31 enantiomerically-pure diastereomers of the Δ^{13} -9-isofurans.

Conclusion

We have developed a general route to the enantiomerically-pure diastereomers of Δ^{13} -9-IsoF. The four key intermediates **2a–2d** were obtained by Sharpless asymmetric epoxidation and Sharpless dihydroxylation, followed by cascade cyclization. Mitsunobu inversion of each of these four key intermediates (**2a–2b**) gave the other four diastereomers (**2e–2h**). This approach has made each of the thirty-two ep diastereomers of Δ^{13} -9-IsoF, previously known only in microgram quantities as mixtures, available in sufficient quantity to assess the individual physiological activity of each. We believe that the robust synthetic approach outlined here will allow ready access both to the other regioisomeric isofurans and to the neurofurans.^{18,19}

Experimental Section

((2R, 3R)-3-((E)-4-trityloxybut-2-enyl) oxiran-2-yl) methanol **9a**

To a stirred solution of diethyl D-tartrate (680 mg, 3.3 mmol) in CH_2Cl_2 (150 mL) was added $\text{Ti}(\text{O}^i\text{Pr})_4$ (924 mg, 3.3 mmol) at -30 to -20 °C. The reaction mixture was stirred for 30 min at this temperature, then a solution of **3** (1.11 g, 3.0 mmol) in CH_2Cl_2 (18 mL) was added over a period of 15 min. After an additional 30 mins at -30 to -20 °C, $t\text{BuOOH}$ (4.7 M in CH_2Cl_2 , 1.7 mL, 8.0 mmol) was added dropwise over a period of 5 min. The reaction mixture was stirred for 3 h at -30 to -20 °C. Aqueous L-(+)-tartaric acid (10%, 6 mL) was added, then the reaction mixture was stirred at -20 °C for 25 min, allowed to warm to rt, and stirred for 40 min. Aqueous NaOH (1 N, 18 mL) was added at 0 °C, and the resulting reaction mixture was stirred for 1.5 h. The reaction mixture was then partitioned between CH_2Cl_2 and water. The organic extract was dried (Na_2SO_4) and concentrated. The residue was chromatographed to afford the epoxide **9a** as a colorless oil (879 mg, 76% yield from monotrityl ether **3**), TLC R_f (MTBE/petroleum ether = 3/7) = 0.28; $[\alpha]_D^{+13.8}$ (c 1.0, CH_2Cl_2); ^1H NMR δ 7.44–7.46 (m, 6 H), 7.20–7.30 (m, 9 H), 5.67–5.75 (m, 2 H), 3.87–3.91 (m, 1 H), 3.58–3.63 (m, 3 H), 3.02–3.05 (m, 1 H), 2.93–2.95 (m, 1 H), 2.34–2.37 (m, 2 H), 2.03 (t, 1 H, J = 6.0 Hz); ^{13}C NMR δ u^{20} : 144.2, 86.9, 64.6, 61.6, 34.4 d: 130.1, 128.7, 127.9, 127.0, 126.0, 58.1, 55.0; IR (film) 3415, 2862, 1447, 1049, 751 cm^{-1} ; HRMS: calcd. $\text{C}_{26}\text{H}_{26}\text{O}_3\text{Na}$ 409.1774, found 409.1775.

((2R, 3R)-3-((E)-4-trityloxybut-2-enyl)oxiran-2-yl) methyl benzenesulfonate **10a**

To a stirring solution of **9a** (879 mg, 2.28 mmol) in CH_2Cl_2 (20 mL) was added DMAP (20 mg, cat) and Et_3N (806 mg, 7.98 mmol) at 0 °C. Then a solution of benzenesulfonyl chloride (1.01 g, 5.70 mmol) in CH_2Cl_2 (4 mL) was added. The reaction mixture was stirred for 30 min at 0 °C, then partitioned between CH_2Cl_2 and water. The organic extract was dried (Na_2SO_4) and concentrated. The residue was chromatographed to give the desired epoxide **10a** (1.14 g, 95% yield from epoxide **9a**), TLC R_f (MTBE/petroleum ether = 2/8) = 0.28; $[\alpha]_D^{+14.7}$ (c 1.6, CH_2Cl_2); ^1H NMR δ 7.90–7.93 (m, 2 H), 7.62–7.64 (m, 1 H), 7.52–7.58 (m, 2 H), 7.42–7.45 (m, 6 H), 7.21–7.32 (m, 9 H), 5.66–5.68 (m, 2 H), 4.24 (dd, 1 H, J = 11.2 Hz and 3.6 Hz), 4.01 (dd, 1 H, J = 11.2 Hz and 6.0 Hz), 3.57 (d, 2 H, J = 3.6 Hz), 2.92–2.99 (m, 1 H), 2.87–2.88 (m, 1 H), 2.32–2.33 (m, 2 H); ^{13}C NMR δ u : 144.1, 135.8, 86.9, 70.1, 64.5, 34.0 d: 134.0, 130.5, 129.3, 128.6, 127.9, 127.8, 127.0, 125.2, 55.7, 54.0; IR (film) 3058, 1447, 1185, 863, 753 cm^{-1} ; HRMS: calcd. $\text{C}_{32}\text{H}_{30}\text{O}_5\text{SNa}$ 549.1731, found 549.1713.

((2S,3S,5S)-tetrahydro-5-((S)-oxiran-2-yl)-2-(trityloxymethyl)furan-3-yloxy)(tert-butyl)dimethylsilane 2a

To a stirred solution of **10a** (770 mg, 1.46 mmol) in a mixed ^tBuOH and H₂O solution (v/v 1:1, 3 mL) was added AD-mix-α (2.13 g). After 10 min at room temperature, methanesulfonamide (141 mg, 1.48 mmol) was added. The reaction mixture was stirred for 60 h. NaHSO₃ (2.18 g, 21.0 mmol) was added and the mixture was stirred for 60 min. The reaction mixture was partitioned between EtOAc and water. The organic extract was dried (Na₂SO₄) and concentrated to afford the diol crude (871 mg).

To a stirred solution of the crude diol (499 mg) in MeOH (7.5 mL) at 0 °C was added powdered K₂CO₃ (307 mg, 2.23 mmol). After an additional 2.5 h at 0 °C, the reaction mixture was partitioned between EtOAc and water. The organic extract was dried (Na₂SO₄) and concentrated. The residue was chromatographed to afford the mixture of alcohol **11a** and **11b** (244 mg, two steps, 72% yield from **10a**).

To a stirred solution of the mixture of alcohols (360 mg, 0.90 mmol) in CH₂Cl₂ (7 mL) was added TBSCl (405 mg, 2.69 mmol), imidazole (350 mg, 5.15 mmol) and DMAP (6 mg, cat) at room temperature. After an additional 40 min, the reaction mixture was partitioned between CH₂Cl₂ and water. The organic extract was dried (Na₂SO₄) and concentrated. The residue was chromatographed to afford **2b** (110 mg, 24% yield from mixture of **11a** and **11b**, 17% yield for three steps from **10a**), TLC *R_f* (MTBE/petroleum ether = 1/9) = 0.46; [α]_D −23.8 (c 1.0, CH₂Cl₂). Further elution gave **2a** as a colorless oil (316 mg, 68% yield from the mixture of **11a** and **11b**, 49% yield for three steps from **10a**), TLC *R_f* (MTBE/petroleum ether = 1/9) = 0.31; [α]_D +42.2 (c 1.0, CH₂Cl₂); ¹H NMR δ 7.46–7.49 (m, 6 H), 7.19–7.29 (m, 9 H), 4.29–4.31 (m, 1 H), 4.05–4.09 (m, 1 H), 3.79–3.83 (m, 1 H), 3.43 (dd, 1 H, *J* = 6.8 Hz and 10.0 Hz), 3.13 (dd, 1H, *J* = 4.0 Hz and 10.0 Hz), 3.04–3.10 (m, 1 H), 2.78 (dd, 1 H, *J* = 4.0 Hz and 5.2 Hz), 2.70 (dd, 1 H, *J* = 2.4 Hz and 5.2 Hz), 2.23–.30 (m, 1 H), 1.95 (ddd, 1H, *J* = 2.0 Hz, 3.6 Hz and 14.6 Hz), 0.70 (s, 9 H), −0.03 (s, 3 H), −0.18 (s, 3 H); ¹³C NMR δ u: 144.1, 86.9, 64.2, 46.9, 38.9, 17.8 d: 128.8, 127.7, 126.9, 83.2, 78.6, 72.5, 54.0, 25.7, −4.8, −5.4; IR (film) 2930, 2857, 1254, 1074, 835 cm^{−1}; HRMS: calcd. C₃₂H₄₀O₄SiNa 539.2594, found 539.2572.

(2S,3S,5S)-2-((trityloxy)methyl)-tetrahydro-5-((S)-oxiran-2-yl)furan-3-ol 11a

To a stirred solution of **2a** (1.57 g, 3.3 mmol) in THF (20 mL) at room temperature was added a Bu₄NF solution in THF (1M, 6.6 mL, 6.6 mmol) at 0 °C. After an additional 5 h, the mixture was partitioned between CH₂Cl₂ and water. The organic extract was dried (Na₂SO₄) and concentrated. The residue was chromatographed to afford **11a** as a solid (1.15 g, 96% yield from **2a**), A 50 mg portion was crystallized from 1.0 mL of 1:5 CH₂Cl₂/hexanes, m.p. = 142 °C; TLC *R_f* (MTBE/petroleum ether = 3/7) = 0.33; [α]_D +13.3 (c 0.60, CH₂Cl₂); ¹H NMR δ 7.44–7.48 (m, 6 H), 7.19–7.30 (m, 9 H), 4.32 (d, 1 H, *J* = 5.2 Hz), 4.06–4.08 (m, 1 H), 3.94 (dd, 1 H, *J* = 2.0 Hz and 5.2 Hz), 3.45 (dd, 1 H, *J* = 5.6 Hz and 9.6 Hz), 3.38 (dd, 1 H, *J* = 5.6 Hz and 9.6 Hz), 3.17–3.20 (m, 1 H), 2.77–2.82 (m, 2 H), 2.60 (dd, 1 H, *J* = 2.8 Hz and 4.8 Hz), 2.18–2.26 (m, 1 H), 1.75–1.79 (m, 1 H); ¹³C NMR δ u: 143.8, 87.1, 62.8, 45.7, 36.0 d: 128.8, 128.0, 127.2, 82.3, 77.8, 72.1, 53.7; IR (film) 3458, 1265, 1072, 756 cm^{−1}; HRMS: calcd. C₂₆H₂₆O₄Na 425.1729, found 425.1733.

Alkyne 12a

To a stirred solution of 5-hexynenitrile (1.07 g, 11.44 mmol) in THF (10 mL) was added dropwise n-BuLi (2.28 M in hexane, 4.1 mL, 9.36 mmol) at −78 °C over a period of 5 min. After an additional 40 min at −78 °C, a solution of **2a** (1.00 g, 1.94 mmol) in THF (15 mL) was added, followed by the addition of a solution of BF₃·OEt₂ (986 mg, 6.95 mmol) in THF (3 mL) over a period of 3 min. The reaction mixture was stirred for 40 min at −78 °C. The reaction mixture was quenched with saturated aqueous NH₄Cl. The mixture was partitioned

between CH₂Cl₂ and water. The organic extract was dried (Na₂SO₄) and concentrated. The residue was chromatographed to afford **12a** (1.08 g, 91% yield from **2a**), TLC *R_f* (MTBE/petroleum ether = 3/7) = 0.30; [α]_D +27.0 (c 1.0, CH₂Cl₂); ¹H NMR δ 7.45–7.48 (m, 6 H), 7.20–7.30 (m, 9 H), 4.26–4.27 (m, 1 H), 4.19–4.21 (m, 1 H), 3.97–4.01 (m, 1 H), 3.87–3.93 (m, 1 H), 3.42 (dd, 1 H, *J* = 6.8 Hz and 10.0 Hz), 3.12 (dd, 1 H, *J* = 4.0 Hz and 10.0 Hz), 2.73 (d, 1 H, *J* = 2.8 Hz), 2.43–2.49 (m, 3 H), 2.33–2.37 (m, 3 H), 2.12–2.17 (m, 1 H), 1.93 (ddd, 1H, *J* = 0.8 Hz, 4.4 Hz and 14.6 Hz), 1.81–1.88 (m, 2 H), 0.71 (bs, 9H), –0.03 (s, 3 H), –0.19 (s, 3 H); ¹³C NMR δ u: 144.1, 119.3, 86.9, 79.5, 78.3, 64.0, 35.1, 24.8, 24.0, 18.0, 17.9, 16.2 d: 128.8, 127.8, 127.0, 82.6, 80.0, 72.6, 71.1, 25.7, –4.9, –5.4; IR (film) 3454, 2932, 2048, 1256, 1072, 705 cm^{–1}; HRMS: calcd. C₃₈H₄₇NO₄SiNa 632.3172, found 632.3177.

Alkene 13aa

To a stirred solution of Ni(OAc)₄·4H₂O (259 mg, 1.04 mmol) in EtOH (8 mL) was added NaBH₄ solution in EtOH (1 M, 1.0 mL, 1.0 mmol). The black mixture was evacuated and backfilled with H₂ three times. Ethylenediamine (70 uL, 0.95 mmol) was added, followed by alkyne **12aa** (317 mg, 0.52 mmol) in EtOH (5 mL). The reaction mixture was stirred for 2.5 h at rt under H₂. The reaction mixture was diluted in MTBE and filtered through a pad of silical gel. The filtrate was concentrated and the residue was chromatographed to afford the desired alkene **13aa** as a colorless oil (301 mg, 95% yield from the alkyne **12aa**), TLC *R_f* (MTBE/petroleum ether = 3/7) = 0.37; [α]_D +24.2 (c 1.0, CH₂Cl₂); ¹H NMR δ 7.64–7.66 (m, 6 H), 7.38–7.48 (m, 9 H), 5.75–5.81 (m, 1 H), 5.61–5.65 (m, 1 H), 4.43–4.46 (m, 1 H), 4.22–4.26 (m, 1 H), 4.12–4.14 (m, 1 H), 4.04–4.08 (m, 1 H), 3.61 (dd, 1 H, *J* = 6.4 Hz and 10.0 Hz), 3.32 (dd, 1 H, *J* = 4.0 Hz and 10.0 Hz), 2.83 (s, 1 H), 2.50 (t, 2 H, *J* = 7.2 Hz), 2.29–2.41 (m, 1 H), 2.11–2.15 (m, 1 H), 1.89–1.93 (m, 2 H), 0.90 (s, 9 H), 0.16 (s, 3 H), 0.00 (s, 3 H); ¹³C NMR δ u: 144.1, 119.7, 86.9, 63.9, 34.7, 31.8, 26.1, 25.3, 17.9, 16.5 d: 130.0, 128.8, 127.9, 127.8, 127.0, 82.4, 80.6, 72.6, 71.8, 25.7, –4.9, –5.5; IR (film) 3440, 2933, 2245, 2048, 1448, 1072, 774, 704 cm^{–1}; HRMS: calcd. C₃₈H₄₉NO₄SiNa 634.3329, found 634.3329.

Alkene 4aa

To a stirred solution of **13aa** (348.3 mg, 0.57 mmol) in CH₂Cl₂ (8 mL) was added TBSCl (430 mg, 2.85 mmol), imidazole (371 mg, 5.70 mmol) and DMAP (5 mg, cat) at room temperature. After an additional 17 h, the reaction mixture was partitioned between CH₂Cl₂ and water. The organic extract was dried (Na₂SO₄) and concentrated. The residue was chromatographed to afford **4aa** (349 mg, 84% yield from **13aa**), TLC *R_f* (MTBE/petroleum ether = 1/9) = 0.50; [α]_D +14.2 (c 1.0, CH₂Cl₂); ¹H NMR δ 7.70–7.72 (m, 6 H), 7.43–7.53 (m, 9 H), 5.87–5.91 (m, 1 H), 5.65–5.68 (m, 1 H), 4.45–4.48 (m, 1 H), 4.13–4.17 (m, 1 H), 4.03–4.06 (m, 1 H), 3.89–3.94 (m, 1 H), 3.57–3.60 (m, 1 H), 3.29 (dd, 1 H, *J* = 2.4 Hz and 10.0 Hz), 2.78–2.85 (m, 1 H), 2.42–2.50 (m, 6 H), 1.92–1.97 (m, 3 H), 1.10 (s, 9 H), 0.92 (s, 9 H), 0.31 (s, 3 H), 0.27 (s, 3 H), 0.14 (s, 3 H), 0.00 (s, 3 H); ¹³C NMR δ u: 144.3, 119.9, 86.6, 64.7, 38.8, 32.3, 26.4, 25.5, 18.1, 17.9, 16.4 d: 128.9, 128.9, 127.7, 126.9, 82.5, 79.6, 74.4, 73.0, 25.9, 25.8, –4.0, –4.3, –4.8, –5.4; IR (film) 3432, 2932, 2264, 1253, 1078, 834 cm^{–1}; HRMS: calcd. C₄₄H₆₃NO₄Si₂Na 748.4193, found 748.4192.

Alcohol 14aa

To a stirred solution of **4aa** (20 mg, 0.0275 mmol) in CH₂Cl₂ (4 mL) was added diethyl aluminum chloride (1N in hexane, 0.15 mL, 0.15 mmol) at –78 °C. After an additional 10 min, the reaction mixture was quenched with water and filtered through a pad of silica gel, washing with methanol. The filtrate was concentrated and the residue was chromatographed to give the desired alcohol **14aa** (11.2 mg, 84% yield from the ether **4aa**), TLC *R_f* (MTBE/petroleum ether = 3/7) = 0.32; [α]_D –5.2 (c 1.0, CH₂Cl₂); ¹H NMR δ 5.58–5.61 (m, 1 H), 5.38–5.41 (m, 1 H), 4.50 (dd, 1 H, *J* = 6.0 Hz and 13.2 Hz), 3.79–3.83 (m, 2 H), 3.72–3.75 (m, 2 H), 3.63–3.68 (m,

1 H), 2.32–2.40 (m, 4 H), 2.17–2.25 (m, 4 H), 1.78–1.83 (m, 1 H), 1.70–1.76 (m, 2 H), 0.88 (s, 18 H), 0.07 (s, 12 H); ^{13}C NMR δ u: 119.7, 62.7, 38.1, 32.5, 26.5, 25.3, 18.1, 18.0, 16.5 d: 128.8, 127.7, 81.1, 79.4, 73.8(2), 25.9, 25.8, –4.2, –4.4, –4.6, –5.2; IR (film) 2932, 2256, 1254, 1079, 835 cm^{-1} ; HRMS: calcd. $\text{C}_{25}\text{H}_{49}\text{NO}_4\text{Si}_2\text{Na}$ 506.3098, found 506.3097.

Enone 16aa

To a stirred solution of **14aa** (64 mg, 0.13 mmol) in CH_2Cl_2 (1.5 mL) was added Dess-Martin periodinane (111 mg, 0.26 mmol). After an additional 40 min at rt, the reaction mixture was directly poured on the silica gel and chromatographed to afford the crude aldehyde (53 mg).

To a stirred suspension of NaH (26 mg, 0.66 mmol) in THF (5 mL) was added dimethyl 2-oxoheptylphosphonate **15** (244 mg, 1.10 mmol) at 0°C . After an additional 1 hr at 0°C , a solution of the crude aldehyde (53 mg) in THF (1 mL) was added. After an additional 40 min at 0°C , the mixture was partitioned between CH_2Cl_2 and saturated aqueous NH_4Cl . The organic extract was dried (Na_2SO_4) and concentrated. The residue was chromatographed to afford **16aa** (64 mg, 84% yield from **14aa**), TLC R_f (MTBE/petroleum ether = 3/7) = 0.73; $[\alpha]_D +5.9$ (c 1.0, CH_2Cl_2); ^1H NMR δ 6.79 (dd, 1 H, $J = 4.2$ Hz and 16.0 Hz), 6.26 (dd, 1 H, $J = 1.6$ Hz and 16.0 Hz), 5.595.64 (m, 1 H), 5.39–5.45 (m, 1 H), 4.42–4.46 (m, 1 H), 4.28 (dt, 1 H, $J = 1.6$ Hz and 5.6 Hz), 3.86 (q, 1 H, $J = 5.2$ Hz), 3.74 (q, 1 H, $J = 7.2$ Hz), 2.54–2.58 (m, 2 H), 2.43–2.50 (m, 1 H), 2.19–2.35 (m, 6 H), 1.82–1.89 (m, 1 H), 1.70–1.77 (m, 2 H), 1.59–1.66 (m, 2 H), 1.25–1.37 (m, 4 H), 0.86–0.90 (m, 21 H), 0.09 (s, 6 H), 0.05 (s, 3 H), 0.02 (s, 3 H); ^{13}C NMR δ u: 200.5, 119.7, 39.9, 38.3, 32.4, 31.5, 26.3, 25.3, 23.8, 22.5, 18.1, 18.0, 16.5 d: 142.7, 130.8, 128.9, 127.6, 82.3, 80.1, 74.3, 73.8, 25.9, 25.8, 13.9, –4.1, –4.3, –4.7, –5.0; IR (film) 3442, 2929, 2250, 1641, 833, 775 cm^{-1} ; HRMS: calcd. $\text{C}_{32}\text{H}_{59}\text{NO}_4\text{Si}_2\text{Na}$ 600.3880, found 600.3876.

Alcohol 17aaa and 17aab

To a stirred solution of **16aa** (118 mg, 0.20 mmol) in MeOH (6 mL) was added $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (74 mg, 0.20 mmol). After 5 min, NaBH_4 (12 mg) was added. The mixture was stirred for an additional 30 min at rt, then the reaction mixture was filtered through a pad of silica gel, washing with MTBE, the filtrate was concentrated and the residue was chromatographed to give the desired alcohol **17aaa** (45 mg, 38% yield from the enone **16aa**), TLC R_f ($\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ /petroleum ether = 1/3/6) = 0.23; $[\alpha]_D +4.5$ (c 1.0, CH_2Cl_2); ^1H NMR δ 5.75–5.82 (m, 2 H), 5.58–5.62 (m, 1 H), 5.36–5.43 (m, 1 H), 4.28–4.31 (m, 1 H), 4.06–4.15 (m, 2 H), 3.84 (q, 1 H, $J = 4.8$ Hz), 3.66 (q, 1 H, $J = 7.2$ Hz), 2.65 (d, 1 H, $J = 9.6$ Hz), 2.39–2.47 (m, 1 H), 2.30–2.34 (m, 2 H), 2.17–2.27 (m, 4 H), 1.83–1.87 (m, 1 H), 1.68–1.76 (m, 2 H), 1.50–1.56 (m, 4 H), 1.26–1.43 (m, 4 H), 0.87–0.90 (m, 21 H), 0.03–0.08 (m, 12 H); ^{13}C NMR δ u: 119.8, 38.3, 37.1, 32.3, 31.8, 26.4, 25.3, 25.2, 22.6, 18.2, 18.1, 16.5 d: 136.3, 128.7, 127.9, 127.6, 83.5, 79.7, 74.3, 74.0, 72.4, 25.9 (3), 25.9 (3), 14.1, –4.1, –4.3, –4.6, –4.9; IR (film) 2931, 2250, 1464, 1252, 1073, 834 cm^{-1} ; HRMS: calcd. $\text{C}_{32}\text{H}_{61}\text{NO}_4\text{Si}_2\text{Na}$ 602.4033, found 602.4025. Further elution gave **17aab** (41 mg, 35% yield from the enone **16aa**), TLC R_f ($\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ /petroleum ether = 1/3/6) = 0.33; $[\alpha]_D +6.9$ (c 1.0, CH_2Cl_2); ^1H NMR δ 5.68–5.81 (m, 2 H), 5.58–5.64 (m, 1 H), 5.37–5.42 (m, 1 H), 4.25–4.29 (m, 1 H), 4.03–4.13 (m, 2 H), 3.82–3.86 (m, 1 H), 3.64–3.70 (m, 1 H), 2.65 (d, 1 H, $J = 9.6$ Hz), 2.42–2.48 (m, 1 H), 2.31–2.35 (m, 2 H), 2.18–2.25 (m, 4 H), 1.81–1.86 (m, 1 H), 1.60–1.75 (m, 4 H), 1.49–1.55 (m, 2 H), 1.37–1.44 (m, 1 H), 1.26–1.36 (m, 3 H), 0.88–0.90 (m, 21 H), 0.01–0.09 (m, 12 H); ^{13}C NMR δ u: 119.9, 38.6, 37.1, 32.2, 31.8, 26.4, 25.3, 25.2, 22.6, 18.2, 18.1, 16.5 d: 136.8, 128.7, 128.0, 127.8, 83.9, 79.7, 74.1, 74.0, 72.6, 25.90 (3), 25.87 (3), 14.1, –4.1, –4.3, –4.6, –4.9; IR (film) 3449, 2932, 2250, 1254, 1074, 835 cm^{-1} ; HRMS: calcd. $\text{C}_{32}\text{H}_{61}\text{NO}_4\text{Si}_2\text{Na}$ 602.4037, found 602.4023.

Isofuran 1aaa

To a stirred solution of **17aaa** (45 mg, 0.078 mmol) in THF (0.8 ml) was added a 1M solution of Bu₄NF (1.0 M in THF, 1.0 mL). After an additional 1.5 hr at rt, the reaction mixture was partitioned between CH₂Cl₂ and H₂O. The organic extract was concentrated to afford the crude triol. The crude triol was dissolved in EtOH (1.0 mL), then 15% aqueous NaOH (2.0 mL) was added. After an additional 16 h at 85 °C, the reaction mixture was cooled to rt and acidified with NaH₂PO₄ (0.5 M, 3 mL) and AcOH (3 mL). The mixture was partitioned between CH₂Cl₂ and water. The organic extract was concentrated and the residue was chromatographed to give **1aaa** (26 mg, 89% yield from **17aaa**), TLC R_f (Et₂O/0.5 M aqueous NaH₂PO₄/HOAc = 90/9/1) = 0.35; [α]_D +25.0 (c 0.5, CH₂Cl₂) [α]_D +35.4 (c 0.58, CH₃OH); ¹H NMR δ 5.98 (s, 3 H), 5.81–5.82 (m, 2 H), 5.44–5.49 (m, 2 H), 4.08–4.16 (m, 4 H), 3.86–3.87 (m, 1 H), 2.35 (s, 1 H), 2.02–2.27 (m, 7 H), 1.70 (s, 2 H), 1.22–1.53 (m, 9 H), 0.86–0.91 (m, 3 H); ¹³C NMR δ u: 36.7, 34.1, 31.8, 26.4, 25.2, 24.4, 22.6 d: 137.5, 131.6, 126.0, 125.9, 84.0, 80.4, 73.1, 72.0, 71.8, 30.3, 14.1; the carbonyl carbon was not observed; IR (film) 3334, 2925, 1708, 1018 cm⁻¹; HRMS: calcd C₂₀H₃₄O₆Na 393.2253, found 393.2246.

Isofuran 1aab

The reaction was performed with the alcohol **17aab** (36 mg, 0.062 mmol) in the same manner as described for the preparation of isofuran **1aaa** to give isofuran **1aab** (24 mg, 92% yield from **17aab**), TLC R_f (Et₂O/0.5 M aqueous NaH₂PO₄/HOAc = 90/9/1) = 0.38; [α]_D +15.6 (c 0.48, CH₂Cl₂) [α]_D +23.1 (c 0.50, CH₃OH); ¹H NMR δ 5.71–5.88 (m, 3 H), 5.43–5.48 (m, 3 H), 4.05–4.20 (m, 4 H), 3.89 (s, 1 H), 2.36 (s, 2 H), 2.04–2.27 (m, 6 H), 1.70–1.73 (m, 2 H), 1.43–1.57 (m, 3 H), 1.25–1.36 (m, 7 H), 0.86–0.89 (m, 3 H); ¹³C NMR δ u: 178.1, 36.5, 34.2, 32.0, 31.8, 26.4, 25.2, 24.4, 22.6 d: 136.5, 131.3, 126.2, 126.2, 83.1, 80.3, 72.7, 72.3, 71.8, 30.3, 14.1; the carbonyl carbon was not initially observed, but when **1aab** was prepared by saponification of the ester **18aab**, the carbonyl was evident; IR (film) 3447, 3034, 2926, 1708, 1240, 1018 cm⁻¹; HRMS: calcd. C₂₀H₃₄O₆Na 393.2253, found 393.2246.

Ester 18aab

To a stirred solution of isofuran **1aab** (8 mg, 0.021 mmol) in MeOH/benzene (1:4, 0.5 mL) was added trimethylsilyl diazomethane (2.0 M in Et₂O, 50 uL). The mixture was stirred at room temperature for 20 min. The mixture was concentrated and the residue was chromatographed to afford the ester **18aab** (8 mg, 96% yield from the isofuran **1aab**), TLC R_f (acetone/CH₂Cl₂ = 4/1) = 0.38; [α]_D +23.8 (c 0.38, CH₂Cl₂) [α]_D +23.1 (c 0.55, CH₃OH); ¹H NMR δ 5.81–5.87 (m, 1 H), 5.73 (dd, *J* = 15.6 Hz and 5.2 Hz, 1 H), 5.44–5.51 (m, 2 H), 4.88 (s, 1 H), 4.79 (s, 1 H), 4.18–4.20 (m, 1 H), 4.11–4.16 (m, 2 H), 4.02–4.08 (m, 1 H), 3.85–3.90 (m, 1 H), 3.70–3.85 (m, 1 H), 3.67 (s, 3 H), 2.32 (t, 2 H, *J* = 7.2 Hz), 2.03–2.29 (m, 5 H), 1.66–1.74 (m, 3 H), 1.41–1.60 (m, 2 H), 1.21–1.38 (m, 6 H), 0.87 (t, 3 H, *J* = 6.4 Hz); ¹³C NMR δ u: 174.0, 36.5, 34.0, 33.4, 32.2, 31.7, 26.7, 25.3, 24.7, 22.6 d: 136.4, 131.2, 126.5, 126.2, 83.1, 80.2, 72.8, 72.2, 71.8, 51.5, 14.1. HRMS: calcd. C₂₁H₃₆O₆Na 407.2404, found 407.2410.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Institutes of Health (GM42056) for support of this work. We thank Dr. John Dykins for mass spectrometric measurements, supported by the NSF (0541775), and Dr. Glenn Yap for X-ray analysis.

References

1. (a) Fessell JP, Porter NA, Moore KP, Sheller JR, Roberts LJ II. *Proc Nat Acad Sci USA* 2002;99:16713. [PubMed: 12482927] For a recent review, see: (b) John U, Galano JM, Durand T. *Angew Chem Int Ed* 2008;47:5894.
2. For a detailed account of isofuran nomenclature, see: Taber DF, Morrow JD, Roberts LJ. *Prostaglandins Other Lipid Mediators* 2004;73:47. [PubMed: 15165030]
3. (a) Taber DF, Pan Y, Zhao X. *J Org Chem* 2004;69:7234. [PubMed: 15471474] (b) Taber DF, Zhang Z. *J Org Chem* 2006;71:926. [PubMed: 16438503]
4. Hanson RM, Sharpless KB. *J Org Chem* 1986;51:1922.
5. (a) Sharpless KB, Amberg W, Bennani YL, Crispino GA, Hartung J, Jeong KS, Kwong HL, Morikawa K, Wang ZM, Xu D, Zhang XL. *J Org Chem* 1992;57:2768. (b) Xu D, Crispino GA, Sharpless KB. *J Am Chem Soc* 1992;114:7570. (c) Kolb HC, VanNieuwenhze MS, Sharpless KB. *Chem Rev* 1994;94:2483.
6. (a) Mitsunobu O. *Synthesis* 1981;1: and references cited therein (b) Martin SF, Dodge JA. *Tetrahedron Lett* 1991;32:3017.
7. (a) Jorgenson MJ. *Tetrahedron Lett* 1962:559. (b) Brown HC, Hess HM. *J Org Chem* 1969;34:2206.
8. Taber D, Green JH, Zhang W, Song R. *J Org Chem* 2000;65:5436. [PubMed: 10993381]
9. For the development of this strategy, see (a) Taber DF, Bhamidipati RS, Thomas ML. *J Org Chem* 1994;59:3442. For a related acid-catalyzed cyclization, see: (b) Sivakumar M, Borhan B. *Tetrahedron Lett* 2003;44:5547. For a recent application, see (c) Aho JE, Salomäki E, Rissanen K, Pihko PM. *Org Lett* 2008;10:4179. [PubMed: 18763802]
10. Sinha SC, Keinan E. *J Org Chem* 1999;64:7067.
11. Brown CA, Ahuja VK. *J Chem Soc, Chem Comm* 1973:553.
12. (a) Dess DB, Martin JC. *J Org Chem* 1983;48:4155. (b) Ireland RE, Liu L. *J Org Chem* 1993;58:2899.
13. Bessodes M, Komiotis D, Antonakis K. *Tetrahedron Lett* 1986;27:579.
14. Koster H, Sinha ND. *Tetrahedron Lett* 1982;23:2641.
15. Germal AL, Luche JL. *J Am Chem Soc* 1981;103:5454.
16. Noyori R, Tomino I, Yamada M, Nishizawa M. *J Am Chem Soc* 1984;106:6717.
17. (a) Magatti CV, Kaminski JJ, Rothberg I. *J Org Chem* 1991;56:3102. (b) Taber DF, Hoerrner RS. *J Org Chem* 1992;57:441.
18. Browne SE, Roberts LJ II, Dennery PA, Doctrow SR, Beal MF, Barlow C, Levine R. *Free Rad Biol Med* 2004;36:938. [PubMed: 15019978]
19. (a) Durand T, Guy A, Vidal J-P, Viala J, Rossi J-C. *Tetrahedron Lett* 2000;41:3859. (b) Quan LG, Cha JK. *J Am Chem Soc* 2002;124:12424. [PubMed: 12381179] (c) Quan LG, Cha JK. *Chem Phys Lip* 2004;128:3. (d) Kim S, Lawson JA, Pratico D, FitzGerald GA, Rokach J. *Tetrahedron Lett* 2002;43:2801. (e) Durand T, Guy A, Herry O, et al. *Chem Phys Lip* 2004;128:14. (f) Rokach J, Kim S, Bellone S, et al. *Chem Phys Lip* 2004;128:35.
20. ^{13}C multiplicities were determined with the aid of a JVERT pulse sequence, differentiating the signals for methyl and methine carbons as “d” and for methylene and quaternary carbons as “u”.

Fig. 1.
There are 32 ep diastereoisomers of SC- Δ^{13} -9-IsoF.

SCHEME 1.

SCHEME 2.

SCHEME 3.

SCHEME 5.

SCHEME 7.

SCHEME 8.

Table 1

Tetrahydrofurans by sequential SAE and SAD

entry	SAE	SAD	Products	Ratio	yield ^a
1	D-DET	AD-mix α	2a/2b	3/1	66%
2	D-DET	AD-mix β	2b	>9/1	57%
3	L-DET	AD-mix β	2c/2d	2.2/1	54%
4	L-DET	AD-mix α	2d	>9/1	62%

^a Combined yield of the two diastereomers.