Published in final edited form as:

J Org Chem. 2008 December 19; 73(24): 9594–9600. doi:10.1021/jo801159k.

Synthetic Studies on Et-743. Assembly of the Pentacyclic Core and a Formal Total Synthesis

Dan Fishlock^a and Robert M. Williams^{a,b,*}

^aDepartment of Chemistry, Colorado State University Fort Collins, Colorado 80523, U.S.A.

^bUniversity of Colorado Cancer Center, Aurora, Colorado 80045, U.S.A.

Abstract

A formal total synthesis of the potent anticancer agent Et-743 is described. The tetrahydroisoquinoline core is stereoselectively constructed using a novel radical cyclization of a glyoxalimine. Further elaboration of this core rapidly accessed the pentacyclic core of Et-743, but a mixture of regioisomers was obtained in the key Pictet-Spengler ring closure. A known advanced intermediate in the synthesis of Et-743 was intercepted, constituting a formal synthesis of the molecule.

Introduction

Members of the tetrahydroisoquinoline family of alkaloids display a wide range of biological properties such as anti-tumor and antimicrobial activities.ⁱ Of particular significance within this family is Ecteinascidin 743 (Et-743, **1**; Figure 1), which has been demonstrated to possess extremely potent cytotoxic activity with *in vitro* IC₅₀ values in the 0.1 ~ 1 ng/mL range in several cell lines (as a measure of RNA, DNA and protein synthesis inhibition).ⁱⁱ Et-743 is currently in phase II/III clinical trials for the treatment of ovarian, endometrial, and breast cancers, and several sarcoma lines.ⁱⁱⁱ The scarcity of the natural product from marine sources renders Et-743 an important target for synthesis. Corey and co-workers reported the first total synthesis of Et-743 in thirty-six steps with an overall yield of 0.72%.^{iva} A second-generation synthesis improved the overall yield to 2.04%, but still required thirty-six steps.^{ivb} Fukuyama and co-workers achieved a total synthesis of Et-743 in fifty steps and 0.56% overall yield.^v More recently, Zhu and co-workers reported a thirty-one step synthesis in 1.7% overall yield.^{vi} Most recently, Danishefsky and co-workers reported a formal total synthesis^{vii} via a pentacyclic compound that intercepted a late-stage intermediate of Fukuyama's route.⁵ Despite the advancements in the state-of-the-art in total synthetic approaches to Et-743, the clinical supply of this complex drug is semi-synthetically derived from natural cyanosafrafin B, obtained by fermentation as reported by PharmaMar.^{viii}

Our laboratory has been developing methodology for the assembly of tetrahydroisoquinoline natural products and has reported syntheses of *D,L*-quinocarcinamide,^{ix} (–)-tetrazomine,^x (–)-renieramycin G,¹¹ (–)-jorumycin^{xi} and cribrostatin 4 (renieramycin H).^{xii} As a part of this program we have targeted Et-743 by a convergent route that envisioned coupling of a suitably functionalized tyrosine derivative^{xiii} with the complete tetrahydroisoquinoline core (Scheme 1). We have successfully deployed this strategy, with the present objective of construction of pentacycle **A**, in the synthesis of (–)-renieramycin G and (–)-jorumycin.^{xi,xii}

We have previously reported a concise and highly diastereoselective synthesis of the tetrahydroisoquinoline core of Et-743 (**E**).^{xiv} This was achieved *via* an intramolecular 6-*endo* radical closure on a glyoxalimine, and the desired 1, 3 *cis* diastereomer was obtained exclusively. The synthesis of a tetrahydroisoquinoline such as **E** can be problematic because of the acid sensitivity of the benzylic hydroxyl, particularly because it is *ortho*- to the phenolic hydroxyl of the aromatic ring and thus has a high propensity for *ortho*-quinonemethide formation. Herein, we report a formal total synthesis of Et-743 as part of our ongoing efforts to devise a practical and scaleable synthesis of this potent antitumor antibiotic that would be amenable to the construction of analogs with anticipated potent cytotoxic activity.

Results and Discussion

The synthesis began with Borchardt's catechol **3**^{xv} that was regioselectively brominated to generate **4** (92% yield) (Scheme 2). Conversion of catechol **4** to the methylenedioxy aldehyde **5** was accomplished using bromochloromethane in a sealed vessel (69% yield). Baeyer-Villiger oxidation using *m*-CPBA provided bromophenol **6** as an off-white solid following hydrolysis of the resulting formate intermediate (73% yield). Stereoselective aldol condensation of the titanium phenolate of **6** with (*R*)-Garner's aldehyde (**7**)^{xvi} using a modification of Casiraghi's method^{xvii} provided the *anti*-product **8** followed by allyl protection of the phenolic oxygen delivering **9** (65% yield, two steps). Subsequent hydrolysis of the oxazolidine and formation of the *trans*-acetone (84% yield, two steps) provided **10** as an oil that cleanly underwent *N*-Boc deprotection using Ohfuné's protocol^{xviii} (76% yield) to afford free amine **11** as a stable crystalline solid. From **11** the glyoxalimine intermediate **13** (see Scheme 3) was readily obtained by condensation with ethyl glyoxalate. Following isolation by filtration through Celite and concentration, the radical ring closure commenced with slow addition of Bu₃SnH and AIBN *via* syringe pump to a refluxing dilute solution of the glyoxalimine (**13**). Concentration and KF/silica chromatography^{xix} of the crude reaction mixture provided solid **12** as a single diastereomer (58% yield, two steps). The relative stereochemistry of **12** was secured ¹H NMR data and corroborated by X-ray crystallography. Examination of the crude ¹H NMR revealed the formation of a single diastereomer in the radical closure and exclusive 6-*endo* regioselectivity. In addition to **12** and tin impurities visible in the ¹H NMR spectrum, an aromatic proton arising from hydride quenching of the aryl radical revealed a ~6.6:1 ratio of **12** to reduced substrate. Slower addition rates (over 18 or 36 h) did not improve the isolated yield of **12**.

The diastereoselectivity of this reaction stands apart from numerous Pictet-Spengler cyclizations on related substrates that provide tetrahydroisoquinolines exclusively as the 1, 3-*trans*-diastereomers.^{xi,xx,xxi} We qualitatively rationalize the *cis*-diastereoselectivity of this radical process using the Beckwith-Houk chair-like transition state model for intramolecular radical ring closures (Figure 2).^{xxii} The lowest-energy chair conformation (**A**) adopted by the *trans*-acetone of the substrate (**13**) results in both the glyoxalimine and aryl substituent being in an equatorial disposition. In this conformation, 1, 3-diaxial steric effects and allylic strain interactions are minimized in the ring-forming transition state. To further examine the stereocontrol imparted by the acetone ring, the *cis*-acetone substrate **14** was prepared (using Casiraghi's method from the magnesium phenolate of **6**).^{xvii} Substrate **14** resulted in a 1:1 mixture of 1, 3-*trans* and 1, 3-*cis*-tetrahydroisoquinolines (**15** and **16**, both are known compounds),²⁰ that suggests the energy difference between transition state conformations **B** and **C** (axial aryl group *versus* axial glyoxalimine) is negligible.

As shown in Scheme 3, reduction of the tetrahydroisoquinoline ester (**12**)¹⁴ with LAH, followed by immediate protection as the benzyl ether (**17**) proceeded cleanly in 77% yield over two steps. The substituted tyrosine amino acid component (**18**) has been previously reported by us, utilizing the oxazinone template technology developed in our laboratory, that was

benzylated with the advanced aromatic side chain.^{xiii} Thus, acylation of the tetrahydroisoquinoline (**17**) was achieved *via* the *N*-Fmoc-protected amino acid chloride (**18**) to give amide **19a** without epimerization. The use of the *N*-Boc free acid with a variety of coupling agents (DCC, HOBt, HATU) all resulted in very sluggish reactions with poor isolated yields, as did the attempted use of the *N*-Boc acid fluoride.

Treatment of **19a** with piperidine provided the free amine, which was not isolated in favor of immediate evaporation of excess base and solvent, and subsequent Boc protection of the crude material. Isolation following chromatography provided compound **19b** in 90% yield. Removal of the acetonide from **19b** was accomplished using the extremely mild, albeit slow, method of stirring with Dowex 50W-X8 cationic resin in methanol. Complete deprotection took 8–12 hours, but the yield was quantitative following simple filtration and concentration. Instead of providing the usual diol product, this substrate incorporated methanol at the benzylic position thus providing the methyl ether as a ~ 1:1 mixture of diastereomers. Not unexpectedly, the benzylic stereogenic center loses stereochemical integrity since the methanol is incorporated *via* the incipient *ortho*-quinonemethide species arising from the acidic deprotection conditions.

Alternatively, we found that the use of water/dichloromethane with cationic resin on **19b** could provide the corresponding free diol but oxidation of the primary alcohol (in the presence of the free benzylic alcohol) could not, in our hands, be cleanly accomplished. The methyl ether was thus a fortuitous selective protection of the benzylic alcohol, ultimately simplifying the subsequent manipulations.

Facile deprotection of the *O*-TBS-protected phenol using TBAF was followed by oxidation of the primary alcohol using Swern conditions in high yield. This oxidation product (**20**) existed as an equilibrium mixture of the aldehyde and the corresponding hemi-aminal species (illustrated) as observed by ¹H NMR, which was otherwise additionally complicated by amide and carbamate rotamers. The attempted oxidation using either Dess-Martin periodinane or TPAP/NMO both failed, leading to extensive decomposition. Following filtration of crude **20** through a plug of silica gel, this substance was immediately subjected to the Pictet-Spengler conditions.

The objective at this stage was to achieve the Pictet-Spengler reaction *via* *N*-Boc deprotection, iminium ion formation and electrophilic aromatic substitution to provide the desired pentacyclic core of Et-743. This meant that the aromatic substitution must occur *ortho*- to the free phenol, and the benzylic methyl ether must survive these conditions. Unfortunately, it had already been demonstrated above that the electron-rich aromatic ring of the tetrahydroisoquinoline component was highly sensitive to protic conditions, leading to *ortho*-quinonemethide formation.

Indeed, when substrate **20** was treated with trifluoroacetic acid in methylene chloride it cleanly underwent the expected pentacycle formation furnishing **21** + **22** as a ~0.72:1 *ortho*:*para* mixture of regioisomers in 72% combined yield. As anticipated, the benzylic methoxy group was eliminated under these conditions, presumably *via* the incipient *ortho*-quinonemethide species that forms under these conditions. In a fruitless effort to circumvent the vexing olefin formation, pentacycle formation with TFA in dry methanol resulted in extensive decomposition of the substrate.

As part of these synthetic investigations, the intermediate **23** was prepared (in parallel with the *O*-benzyl protected synthesis) bearing an *O*-allyl protected hydroxymethyl at C1 of the THIQ core. This substrate was used to examine the regioselectivity of the pentacycle-forming ring closure and was utilized to acquire detailed ¹H NMR data, while the *O*-benzyl material **21** was carried forward in the synthesis. One interesting observation was the behavior of compound **25** containing the *O*-Boc carbonate-protected phenolic oxygen. Treatment of **25** under the same

reaction conditions provided the pentacycles **26** + **27** in a 2:3 ratio of *ortho:para* regioisomers. The *O*-Boc carbonate would presumably be deprotected quickly under these conditions to reveal the free phenol-containing reactive species, thus resulting in a comparable regioselectivity as observed with substrate **20** (beginning with a free phenol on the aryl nucleophile moiety). Notably, however, when substrate **25** was treated with K₂CO₃/MeOH, the *O*-Boc carbonate was selectively removed (**28**) with apparent olefin formation prior to the Pictet-Spengler reaction and pentacycle formation. Treatment of **28** with TFA in dichloromethane produced the pentacycles **26** + **27** in a 1:3 ratio of *ortho:para* regioisomers, supporting the hypothesis that some regioselectivity in the closure might arise from an intramolecular H-bond with a heteroatom at the benzylic position.^{xic}

In their synthesis of renieramycin H, the Zhu group has interestingly reported control of Pictet-Spengler regioselectivity in a related system by variation of acid concentration.^{xxiii} It was found in that case that lowering the concentration of methanesulfonic acid to 0.1% in CH₂Cl₂ could invert the *ortho:para* selectivity from 3.4:1 to 2:3. Furthermore, the use of acetonitrile as the solvent instead of dichloromethane favored the undesired isomer, giving *ortho:para* selectivity of 1:10. Our attempt to reproduce the Zhu conditions on substrate **24** using 0.1% methanesulfonic acid in CH₂Cl₂ did not affect the regioselectivity of this reaction. The substrate was consumed to provide some material that appeared to still contain the *N*-Boc protecting group, but the ¹H NMR of the crude product was prohibitively complex. Subsequent treatment of this reaction crude with a TFA/anisole/CH₂Cl₂ mixture provided the pentacycles **26** + **27** with ~1:1 regioselectivity. The same ratio is obtained if the TFA/anisole conditions are used directly on substrate **24**.

In order to redeem the synthetic utility of the olefinic products (**21** or **26**), our attention was captured by the recent formal synthesis of Et-743 reported by the Danishefsky group^{vii} in which the olefin (**29**, Scheme 5) underwent facile oxidation using DMDO, and immediate hydride reduction delivering the benzylic alcohol **30**. With the availability of this methodology in the literature, our efforts were briefly re-directed to convert our synthetic pentacycle **21** into compound **29** which would constitute a formal total synthesis of Et-743 by relay through the Danishefsky^{vii} and then Fukuyama^v syntheses, respectively.

In the event, the desired pentacycle **21** (Scheme 3) was *N*-protected as the trichloroethyl carbamate (Troc), and the phenolic residue was protected as the corresponding *O*-benzyl ether in 85% yield for the two steps (Scheme 5). Removal of the *O*-allyl group under standard conditions followed by re-protection as the corresponding MOM ether provided compound **29** (56% yield for the two steps). Compound **29** perfectly matched Danishefsky's substrate by ¹H, ¹³C NMR and optical rotation, confirming the structure of compound **29**.

Since Danishefsky has previously converted^{vii} compound **29** into a late-stage intermediate in Fukuyama's total synthesis^v (namely, compound **30**, Scheme 5), this two-stage relay of our synthetic **21** thus constitutes a formal total synthesis of Et-743 and provides firm structural corroboration of our synthetic material and methods.

While the present formal synthesis reveals that our glyoxalimine radical cyclization technology^{xiv} holds considerable potential for the efficient total synthesis of Et-743 and congeners, we are currently endeavoring to improve the regioselectivity of the key pentacycle formation (**20** to **21**) as well as refining the overall synthetic efficiency of our approach. These objectives are currently under study in our laboratory and will be reported in due course.

Experimental Section

For General Methods and considerations, see Supporting Information.

Compound 19

The Fmoc-amino acid (410 mg, 0.727 mmol, 1.2 eq) was dissolved in dry toluene and concentrated (x2), and then dried under high vacuum. This oil was dissolved in dry CH₂Cl₂ (4 mL) to which was added oxalyl chloride (1 mL) at room temperature, followed by dry DMF (20 μ L). After stirring for 20 min the solution was concentrated and re-concentrated from dry toluene (x2) and then dried under high vacuum. This acid chloride **18** was dissolved in CH₂Cl₂ (4 mL) and cooled to 0°C. THIQ(OBn) **17** (275 mg, 0.61 mmol, 1 eq) was dissolved in dry CH₂Cl₂ (2 mL) and 2,6-lutidine (77 μ L 0.67 mmol, 1.1 eq). This solution was transferred into the acid chloride solution slowly drop-wise, and the resulting mixture warmed to rt and stirred 7h (TLC showed consumption of the THIQ(OBn) starting material). The reaction was quenched with sat. NH₄Cl (aq) and then extracted to EtOAc (x3). The combined organic fractions were dried (Na₂SO₄), filtered and concentrated to provide a crude orange oil. Purification by flash chromatography (hexanes:EtOAc 5:1, silica gel) gave the peptide **19a** as a pale yellow oil (426 mg, 70%). R_f = 0.34 (3:1 Hexanes:EtOAc, UV, CAM) [α]_D²⁵ -22.8 (c 1.14, CH₂Cl₂). IR (thin film) 3289, 2929, 2858, 1717, 1634 cm⁻¹. ¹H and ¹³C NMR spectra are extremely complex due to amide and carbamate rotamers. See the rt (CDCl₃) and 373K (DMSO-*d*₆) ¹H spectra and rt (CDCl₃) ¹³C spectra in the Supporting Information. HRMS(ESI/APCI+) *m/z* calcd. For C₅₈H₆₈N₂O₁₁NaSi (M+Na)⁺ 1019.4485; found 1019.4499.

Compound 19b

Fmoc (OBn) peptide **19a** (146 mg, 0.146 mmol) was dissolved in a 20% v/v solution of Et₂NH in CH₂Cl₂ [CH₂Cl₂ (2.5 mL) and diethylamine (0.6 mL)]. After stirring for 6h the solution was concentrated, and then re-concentrated from toluene and dried under high vacuum. The crude material was dissolved in EtOH:CH₂Cl₂ (2:0.5 mL) to which was added Boc₂O (370 mg, 10 eq). After stirring for 12 h the reaction was concentrated and immediately purified by flash chromatography (9:1 to 5:1 hexanes:EtOAc, silica gel) to provide **19b** as a clear colourless oil (115 mg, 90% over 2 steps). R_f = 0.43 (3:1 Hexanes:EtOAc, UV, CAM). [α]_D²⁵ -26.6 (c 1.0, CH₂Cl₂). IR (thin film) 3319, 2930, 2858, 1711, 1646 cm⁻¹. ¹H and ¹³C NMR spectra are extremely complex due to amide and carbamate rotamers. See the ¹H spectra (CDCl₃, rt) and (DMSO-*d*₆, 373K) and ¹³C spectrum (CDCl₃, rt) in the Supporting Information. HRMS(ESI/APCI+) *m/z* calcd. For C₄₈H₆₆N₂O₁₁NaSi (M+Na)⁺ 897.4328; found 897.4310.

Compounds 21 and 22

Boc (OBn) peptide **19b** (115 mg, 0.132 mmol) was dissolved in dry MeOH (5 mL) and Dowex 50W-X8 cationic resin (100 mg) was added (the resin was first rinsed with dry methanol and dried under a stream of argon). After 65 h the reaction was complete by TLC and a single streak was observed (during the course of the reaction two streaks initially arise due to a mixture of diol and methyl ether/alcohol products). The reaction was filtered through a plug of Celite, eluting with dry MeOH, and the filtrate combined to provide the methyl ether as clear, colourless oil (100 mg, 90% yield). R_f = 0 to 0.35 streak (3:1 Hexanes:EtOAc, UV, CAM). HRMS(FAB+) *m/z* calcd. For C₄₆H₆₅N₂O₁₁Si (M+H)⁺ 849.4358; found 849.4354. The methyl ether (100 mg) was dissolved in THF (3 mL) and TBAF (1M in THF, 125 μ L, 1.06 eq) was added in one portion. After 20 minutes the reaction was concentrated by rotary evaporation and passed through a silica plug (eluting with 3:1 to 1:1 hexanes:EtOAc) to provide the free phenol as a clear, colourless oil (82 mg, 95% yield). R_f = 0 to 0.43 streak (3:1 Hexanes:EtOAc, UV, CAM). HRMS(FAB+) *m/z* calcd. For C₄₀H₅₁N₂O₁₁ (M+H)⁺ 735.3493; found 735.3490. Oxalyl chloride (15 μ L 1.5 eq) was added carefully to a solution of DMSO (25 μ L, 3.2 eq) in CH₂Cl₂ (1 mL) previously cooled to -78°C. A solution of the above alcohol (82 mg, 0.11 mmol) in CH₂Cl₂ (2 mL) was added drop-wise, and the resulting mixture stirred at -78°C for 40 min. The reaction was quenched with Et₃N (125 μ L, 8 eq) and then allowed to warm to rt. The reaction was diluted with CH₂Cl₂ and washed with brine, and then the combined organic

fractions dried (Na_2SO_4), filtered and concentrated. The crude material was passed through a silica gel plug (eluting with hexanes:EtOAc 1:1) to provide a yellow oil/foam (82 mg, quant.), of hemi-aminal **20** which was used without further purification. $R_f = 0.5$ (Hexanes:EtOAc 1:1, UV, CAM). Hemi-aminal **20** (232 mg, 0.32 mmol) was dissolved in CH_2Cl_2 (3 mL) to which was added TFA (3 mL) and anisole (0.350 mL) at rt. The reaction was stirred for 14 hours and then concentrated to remove TFA, then re-dissolved in CH_2Cl_2 and washed with saturated aq. NaHCO_3 . The organic fraction was dried (Na_2SO_4), filtered and concentrated. Crude ^1H NMR shows 0.72:1 *ortho* (**21**) to *para* (**22**) regioisomers. Purification by PTLC (2% MeOH in EtOAc) provided the *ortho* (63 mg) and *para* products (69 mg) for a combined yield of 72%. Data for **21**: $R_f = 0.61$ (EtOAc:MeOH 95:5, UV, CAM). $[\alpha]^{25}_{\text{D}} -18.0$ (c 1.0, CH_2Cl_2). IR (thin film) 3295, 2932, 1672, 1632, 1455, 1428 cm^{-1} . ^1H NMR (400 MHz, CDCl_3) δ 7.14–7.30 (m, 3H), 6.98 (s, 1H), 6.97 (s, 1H), 6.24 (s, 1H), 6.19 (s, 1H), 6.12 (dddd, $J = 16.0, 11.0, 5.4, 5.4$ Hz, 1H), 6.05 (dd, $J = 7.2, 5.0$ Hz, 1H), 5.85 (br s, 1H), 5.82 (br s, 1H), 5.78 (v br s, 1H), 5.45 (app dd, $J = 17.1, 1.1$ Hz, 1H), 5.29 (app dd, $J = 10.3, 0.8$ Hz, 1H), 4.9 (s, 1H), 4.30 (app d of AB quartet, $J = 12.3, 5.4$ Hz, 2H), 4.03 (d, $J = 6.1$ Hz, 1H), 3.87 (AB quartet, $J = 12.1$ Hz, 2H), 3.63 (s, 3H), 2.95–3.2 (m, 5H), 2.11 (s, 3H), 2.09 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 168.8, 147.6, 145.6 (x2), 143.4, 139.7, 138.7, 134.5, 133.9, 129.4, 128.8, 128.0 (x2), 127.1, 126.8 (x2), 122.5, 119.3, 117.7, 117.5, 113.0, 108.7, 101.5, 100.2, 75.3, 72.6, 70.0, 60.8, 54.4, 50.0, 46.9, 33.4, 15.9, 9.4. HRMS(ESI/APCI+) m/z calcd. For $\text{C}_{34}\text{H}_{35}\text{N}_2\text{O}_7$ ($\text{M}+\text{H}$) $^+$ 583.2439; found 583.2441. **Data for 22**: $R_f = 0.5$ (EtOAc:MeOH 95:5, UV, CAM). $[\alpha]^{25}_{\text{D}} +47.8$ (c 1.45, CH_2Cl_2). IR (thin film) 3298, 2931, 1671, 1631, 1430, 1409 cm^{-1} . ^1H NMR (400 MHz, CDCl_3) δ 7.11–7.22 (m, 3H), 6.94 (s, 1H), 6.92 (s, 1H), 6.41 (s, 1H), 6.11 (dddd, $J = 16.1, 10.6, 5.5, 5.5$ Hz, 1H), 6.08 (s, 1H), 6.03 (dd, $J = 6.6, 5.1$ Hz, 1H), 5.86 (br s, 1H), 5.83 (br s, 1H), 5.45 (app dd, $J = 17.1, 1.1$ Hz, 1H), 5.30 (app dd, $J = 10.4, 0.8$ Hz, 1H), 4.65 (s, 1H), 4.36 (app d of A of AB quartet, $J = 12.5, 5.5$ Hz, 1H), 4.24 (app d of B of AB quartet, $J = 12.5, 5.5$ Hz, 2H), 4.01 (d, $J = 6.0$ Hz, 1H), 3.91 (AB quartet, $J = 12.2$ Hz, 1H), 3.56 (s, 3H), 2.95–3.24 (m, 5H), 2.27 (s, 3H), 2.12 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) δ 168.7, 148.0, 147.6, 145.9, 144.3, 139.7, 138.4, 134.6, 133.7, 129.6, 128.7, 128.1 (x2), 127.2, 126.9 (x2), 124.9, 117.8, 117.1, 113.8, 113.0, 108.8, 101.5, 100.5, 75.4, 72.8, 70.1, 61.0, 54.3, 52.6, 46.8, 35.4, 12.0, 9.4. HRMS (ESI/APCI+) m/z calcd. For $\text{C}_{34}\text{H}_{35}\text{N}_2\text{O}_7$ ($\text{M}+\text{H}$) $^+$ 583.2439; found 583.2429.

Preparation of compound 29

The desired *ortho*-regioisomer **21** (55 mg, 0.095 mmol) was dissolved in CH_2Cl_2 (2 mL) and pyridine (11 μL , 0.14 mmol, 1.5 eq) at 0°C . TrocCl (13.5 μL , 0.1 mmol, 1.0 eq) was added and the reaction maintained at 0°C for 2 h, and then diluted with CH_2Cl_2 and washed with saturated aq. NH_4Cl . The organic layer was dried (Na_2SO_4) and filtered, then concentrated. The crude oil was passed through a plug of silica gel eluting with EtOAc, and then concentrated and dried under vacuum. The resulting oil was dissolved in CH_2Cl_2 (600 μL) and MeOH (200 μL) and K_2CO_3 (52 mg, 0.38 mmol, 4 eq) were added followed by benzyl bromide (22 μL , 0.19 mmol, 2 eq) and a catalytic amount of tetrabutylammonium iodide. The resulting mixture was stirred at rt for 13.5 h then filtered through a pad of Celite, rinsing with CH_2Cl_2 . Flash chromatography (5:1 hexanes:EtOAc) provided the *N*-Troc/*O*-benzyl product as a pale yellow oil (68 mg, 85% over 2 steps). $R_f = 0.46$ (Hexanes:EtOAc 3:1, UV, CAM). $[\alpha]^{25}_{\text{D}} +58.1$ (c 1.7, CH_2Cl_2). IR (thin film) 2927, 1724, 1681, 1434, 1371 cm^{-1} . ^1H NMR (400 MHz, CDCl_3) Mixture of carbamate rotamers: δ : 7.30–7.56 (m, 3H), 7.14–7.25 (m, 4H), 6.92–7.00 (m, 2H), 6.45 (d, $J = 9.9$ Hz, 1H), 6.22 (d, $J = 4.1$ Hz, 1H), 6.12 ($J = 16.1$ Hz, 1H), 6.01–6.08 (m, 1H), 5.79–5.90 (m, 3H), 4.98–5.29 (m, 5H), 4.85 (d, $J = 12.0, 2.8$ Hz, 1H), 4.60 (d, $J = 11.9, 6.5$ Hz, 1H), 3.97–4.11 (m, 3H), 2.85 (app d, $J = 12.1$ Hz, 1H), 3.70 (s, 3H), 3.04–3.30 (m, 4H), 2.10 (s, 3H), 2.07 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3) Mixture of carbamate rotamers: δ : 166.1/166.0, 151.5/151.4, 149.9, 148.9, 148.4, 148.3/148.2, 146.2, 139.6, 138.5, 137.6/137.5, 133.8/133.7, 132.6/132.5, 131.3/131.2, 128.9 (x2), 128.3, 128.1 (x2), 128.0, 127.2, 126.8, 126.6/126.5, 125.4/125.0, 117.6, 117.3, 116.9, 113.3/113.3, 108.6/108.4, 103.3,

102.9, 101.6, 95.3/95.2, 75.4/75.3, 75.2/75.0, 74.6/74.4, 72.6, 69.9/69.8, 60.5, 54.4/53.7, 50.9/50.1, 47.3/47.2, 32.8/32.4, 16.0, 9.5. HRMS(ESI/APCI+) m/z calcd. For $C_{44}H_{42}N_2O_9Cl_3$ (M+H)⁺ 847.1950; found 847.1949.

The allyl protected pentacycle obtained above (20 mg, 0.024 mmol) was dissolved in CH_2Cl_2 (400 μ L), and pyrrolidine (6 μ L, 3 eq) was added, followed by $Pd(PPh_3)_4$ (2 mg, 0.002 mmol) under Ar. After 16 h the reaction was still not complete, so additional portions of pyrrolidine and palladium catalyst were added. After stirring an additional 4 h (20 hours total) the dark green reaction was applied directly to flash chromatography (silica gel, hexanes:EtOAc 3:1). The pure fractions were combined to provide the phenol as yellow oil (11 mg 56%), used without characterization. R_f = 0.26 (Hexanes:EtOAc 3:1, UV, CAM). Phenol (11 mg, 0.014 mmol) was dissolved in CH_2Cl_2 (200 μ L) to which was added iPr_2NEt (12 μ L, 0.07 mmol, 5 eq) and MOMBr (3.3 μ L, 0.042 mmol, 3 eq). The mixture was stirred for 30 min at rt and then quenched with water and extracted with CH_2Cl_2 (x3). The combined organic fractions were dried (Na_2SO_4), filtered and concentrated. Flash chromatography (hexanes:EtOAc 3:1) provided the protected pentacycle **29** (11.5 mg, quant.). R_f = 0.41 (Hexanes:EtOAc 3:1, UV, CAM). $[\alpha]^{25}_D +45.4$ (c 0.8, $CHCl_3$) [Lit. +50 (c 1.0, $CHCl_3$)] IR (thin film) 2932, 1723, 1681, 1654, 1432, 1371 cm^{-1} . 1H and ^{13}C NMR spectra perfectly match the data provided by the Danishefsky group for this intermediate in their formal synthesis (copies of their spectra included below).⁷ 1H NMR (400 MHz, $CDCl_3$) Mixture of carbamate rotamers: δ : 7.56–7.31 (m, 5H), 7.13–7.23 (m, 3H), 6.96 (app br d, J = 6.9 Hz, 2H), 6.46 (d, J = 9.4 Hz, 1H), 6.01–6.15 (m, 3H), 5.86 (app d, J = 3.0 Hz, 2H), 5.82 (br s, 1H), 4.97–5.19 (m, 4H), 4.86 (d, J = 11.9 Hz, 1H), 4.79 (A of AB quart, J = 12.0 Hz, 1H), 4.68 (B of AB quart, J = 11.9 Hz, 1H), 4.49–4.60 (m, 2H), 4.43 (app d, J = 6.1 Hz, 1H), 4.01 (d of A of AB quart, J = 11.8, 4.4 Hz, 1H), 3.85 (B of AB quart, J = 12.1 Hz, 1H), 3.71 (app d, J = 10.6 Hz, 3H), 3.38 (rotomeric s, 3H), 3.03–3.29 (m, 5H), 2.11 (s, 6H). ^{13}C NMR (100 MHz, $CDCl_3$) Mixture of carbamate rotamers: δ : 166.0/165.9, 151.6/151.4, 149.9, 148.7/148.2, 147.3, 146.2/146.1, 139.8, 138.4/138.4, 137.8/137.7, 132.6/132.5, 131.1/131.1, 128.8, 128.2, 127.9, 127.2, 126.8, 126.6/126.5, 125.2/125.0, 117.0/116.8, 113.7/113.6, 108.5/108.4, 103.3/102.7, 101.6, 100.4/100.4, 95.3/95.2, 75.4/75.3, 74.4/74.0, 72.6/72.6, 69.9/69.9, 60.4, 57.6/57.5, 54.4/53.7, 50.8/50.1, 47.4/47.3, 32.7/32.3, 16.0, 9.9. HRMS(ESI/APPI+) m/z calcd. For $C_{43}H_{42}N_2O_{10}Cl_3$ (M+H)⁺ 851.1900; found 851.1897.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This paper is fondly dedicated to the memory of the late Professor A.I. Meyers. We gratefully acknowledge financial support from the National Institutes of Health (Grant CA85419). We are grateful to Prof. Samuel J. Danishefsky of the Memorial Sloan-Kettering Cancer Research Institute for kindly providing characterization data for compound **29**.

References

- i. Scott JD, Williams RM. Chem. Rev 2002;102:1669–1730. [PubMed: 11996547]
- ii. (a) Rinehart KL, Holt TG, Fregeau NL, Keifer PA, Wilson GR, Perun TJ, Sakai R, Thompson AG, Stroh JG, Shield LS, Seigler DS. J. Nat. Prod 1990;53:771–792. [PubMed: 2095373] (b) Rinehart KL, Holt TG, Fregeau NL, Stroh JG, Keifer PA, Sun F, Li LH, Martin DG. J. Org. Chem 1990;55:4512–4515. (c) Wright AE, Forleo DA, Gunawardana GP, Gunasekera SP, Koehn FE, McConnell OJ. J. Org. Chem 1990;55:4508–4512. (d) Guan Y, Sakai R, Rinehart KL, Wang AH-J. J. Biomol. Struct. Dyn 1993;10:793–818. [PubMed: 8318161] (e) Aune GJ, Furuta T, Pommier

- Y. *Anti-Cancer Drugs* 2002;13:545–555. [PubMed: 12172500] (f) Rinehart KL. *Med. Drug Rev* 2000;1–27.
- iii. Zelek, L.; Yovine, A.; Etienne, B.; Jimeno, J.; Taamma, A.; Martín, C.; Spielmann, M.; Cvitkovic, E.; Misset, JL. Ecteinacidin-743 in taxane / Anthracycline pretreated advanced / metastatic breast cancer patients: preliminary results with the 24 hour continuous infusion Q3 week schedule. presented at the American Society of Clinical Oncology, 36th Annual Meeting; New Orleans. 2000 May 20–23. Abstract number 592Delalogue, S.; Yovine, A.; Taamma, A.; Cottu, P.; Riofrio, M.; Raymond, E.; Brain, E.; Marty, M.; Jimeno, J.; Cvitkovic, E.; Misset, JL. Preliminary evidence of activity of Ecteinacidin-743 (ET-743) in heavily pretreated patients with bone and soft tissue sarcomas. New Orleans. presented at the American Society of Clinical Oncology, 36th Annual Meeting; 2000 May 20–23. Abstract number 2181Le Cesne, A.; Judson, I.; Blay, JY.; Radford, J.; an Oosterom, A.; Lorigan, P.; Rodenhuis, E.; Donato Di Paoula, E.; Van Glabbeke, M.; Jimeno, J.; Verweij, J. Phase II of ET-743 in advance soft tissue sarcoma in adult: a STBSG-EORTC trial. New Orleans. presented at the American Society of Clinical Oncology, 36th Annual Meeting; 2000 May 20–23. Abstract number 2181 (c) Aune GJ, Furuta T, Pommier Y. *Anti-Cancer Drugs* 2002;13:545–555. [PubMed: 12172500]
- iv. (a) Corey EJ, Gin DY, Kania RS. *J. Am. Chem. Soc* 1996;118:9202–9203. (b) Martinez EJ, Corey E. *J. Org. Lett* 2000;2:993–996.
- v. Endo A, Yanagisawa A, Abe M, Tohma S, Kan T, Fukuyama T. *J. Am. Chem. Soc* 2002;124:6552–6554. [PubMed: 12047173]
- vi. Chen J, Chen X, Bois-Choussy M, Zhu J. *J. Am. Chem. Soc* 2005;128:87–89. [PubMed: 16390134]
- vii. Zheng S, Chan C, Furuuchi T, Wright BJD, Zhou B, Guo J, Danishefsky S. *J. Angew. Chem. Int. Ed* 2006;45:1754–1759.
- viii. Cuevas C, Pérez M, Martín MJ, Chicharro JL, Fernández-Rivas C, Flores M, Francesch A, Gallego P, Zarzuelo M, de la Calle F, García J, Polanco C, Rodríguez I, Manzanares I. *Org. Lett* 2000;2:2545–2548. [PubMed: 10956543]
- ix. Flanagan ME, Williams RM. *J. Org. Chem* 1995;60:6791–6797.
- x. (a) Scott JD, Williams RM. *Angew. Chem. Int. Ed* 2001;40:1463–1465. (b) Scott JD, Williams RM. *J. Am. Chem. Soc* 2002;124:2951–2956. [PubMed: 11902886]
- xi. (a) Lane JW, Chen Y, Williams RM. *J. Am. Chem. Soc* 2005;127:12684–12690. [PubMed: 16144418] (b) Lane JW, Estevez A, Mortara K, Callan O, Spencer JR, Williams RM. *Bioorg. Med. Chem. Lett* 2006;16:3180–3183. [PubMed: 16632360] (c) Vincent G, Lane JW, Williams RM. *Tetrahedron Lett* 2007;48:3719–3722. [PubMed: 19578531] (d) Jin W, Metobo S, Williams RM. *Org. Lett* 2003;5:2095–2098. [PubMed: 12790537]
- xii. (a) Vincent G, Williams RM. *Angew. Chem. Int. Ed. Engl* 2007;46:1517–1520. [PubMed: 17387658] (b) Vincent G, Chen Y, Lane JW, Williams RM. *Heterocycles* 2007;72:385–398.
- xiii. Jin W, Williams RM. *Tetrahedron Lett* 2003;44:4635–4639.
- xiv. Fishlock D, Williams RM. *Org. Lett* 2006;8:3299–3301. [PubMed: 16836390]
- xv. Shinhababu AK, Ghosh AK, Borchardt RT. *J. Med. Chem* 1985;28:1273–1279. [PubMed: 4032430] Prepared from 2,3-dimethoxytoluene according to:
- xvi. Garner P, Park JM. *Org. Synth* 1992;70:18–28.(R)-Garner's aldehyde was synthesized from D-serine according to:
- xvii. Casiraghi G, Cornia M, Rassu GJ. *Org. Chem* 1988;53:4919–4922. In our case, sonication was not required as described in the original paper
- xviii. Sakaitani M, Ohfune YJ. *Org. Chem* 1990;55:870–876.
- xix. Harrowven DC, Guy IL. *Chem. Commun* 2004:1968–1969. Effective removal of tin impurities from Bu₃SnH mediated reactions:
- xx. (a) De Paolis M, Chiaroni A, Zhu J. *Chem. Commun* 2003:2896–2897. (b) Chen X, Chen J, De Paolis M, Zhu J. *J. Org. Chem* 2005;70:4397–4408. [PubMed: 15903318]
- xxi. (a) Herberich B, Kinugawa M, Vazquez A, Williams RM. *Tetrahedron Lett* 2001;42:543–546. (b) Jin W, Metobo S, Williams RM. *Org. Lett* 2003;5:2095–2098. [PubMed: 12790537]
- xxii. (a) Beckwith ALJ, Schiesser CH. *Tetrahedron* 1985;41:3925–3941. (b) Spellmeyer DC, Houk KN. *J. Org. Chem* 1987;52:959–974.
- xxiii. Chen X, Zhu J. *Angew. Chem. Int. Ed* 2007;46:3962–3965.

1, ecteinascidin 743

Figure 1.
Ecteinascidin 743 (**1**).

Figure 2. Transition state models to rationalize the observed 1,3 relative stereochemistry in the tetrahydroisoquinoline radical ring closure.

Scheme 1.
Synthetic plan.

Scheme 2.
Tetrahydroisoquinoline core of Et-743.

Scheme 3.
Pentacycle construction.

Scheme 4.
Pictet-Spengler regioselectivity.

Scheme 5.
Formal synthesis of Et-743 via **21** to **29** and Danishefsky to Fukuyama relay.