Published in final edited form as:

Endocr Relat Cancer. 2008 December ; 15(4): 841–849. doi:10.1677/ERC-08-0084.

Androgen receptor and growth factor signaling cross-talk in prostate cancer cells

Meng-Lei Zhu and Natasha Kyprianou

Departments of Urology and Toxicology, University of Kentucky College of Medicine, University of Kentucky Medical Center, Combs Research Building Room 306, Lexington, Kentucky 40536, USA

Abstract

Androgens promote the growth and differentiation of prostate cells through ligand activation of the androgen receptor (AR). Sensitization of the androgenic response by multifunctional growth factor signaling pathways is one of the mechanisms via which AR contributes to the emergence of androgen-independent prostate tumors. The ability of AR to cross-talk with key growth factor signaling events toward the regulation of cell cycle, apoptosis, and differentiation outcomes in prostate cancer cells is established. In this paper, we review the functional interaction between AR and an array of growth factor signal transduction events (including epidermal growth factor; fibroblast growth factor; IGF1; vascular endothelial growth factor; transforming growth factor- β) in prostate tumors. The significance of this derailed cross-talk between androgens and key signaling networks in prostate cancer progression and its value as a therapeutic forum targeting androgen-independent metastatic prostate cancer is discussed.

Introduction

Prostate cancer development and growth is dependent on androgens and can be suppressed by androgen ablation monotherapy. Due to the emergence of androgen-independent prostate tumor growth however, prostate cancer recurs as androgen-independent, highly metastatic advanced disease (Wang *et al.* 2007).

Androgen functions through an axis involving testicular synthesis of testosterone, conversion by 5 reductase to the active metabolite 5 dihydrotestosterone (DHT), and its binding to androgen receptor (AR) to induce transcriptional activation of target genes (Siiteri & Wilson 1974, Imperato-McGinley *et al.* 1985, Heinlein & Chang 2002). In the adult prostate, androgens promote survival of epithelial cells, the primary step to malignant transformation to prostate adenocarcinoma (De Marzo *et al.* 1998). Androgen-induced prostate epithelial cell proliferation is regulated by an indirect pathway involving paracrine mediators produced by stromal cells, such as insulin-like growth factor (IGF), fibroblast growth factor (FGF), and epidermal growth factor (EGF; Cunha & Donjacour 1989, Byrne *et al.* 1996). The absence of a link between elevated serum testosterone, DHT, or adrenal androgens and prostate cancer risk suggests that androgens are not sufficient to promote prostate carcinogenesis (Roberts & Essenhight 1986, Hsing 2001). The current evidence on the cross-talk between AR/androgen axis and signaling effectors of growth factors, as the contributing mechanism to prostate tumor initiation and progression, is discussed in this review.

Correspondence should be addressed to N Kyprianou; Email: nkypr2@uky.edu.

Declaration of interest

The authors hereby declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

AR connects with EGF

EGF and its membrane receptor, the epidermal growth factor-1 receptor (EGFR), are involved in the pathogenesis of different tumors, including prostate cancer (Russell *et al.* 1998). Both the ligand and its signaling receptor partner are frequently up-regulated in advanced stages of prostate cancer (Di Lorenzo *et al.* 2002). Targeting EGFR with monoclonal antibodies or with tyrosine kinase inhibitors suppresses growth and invasion of androgen-dependent and -independent prostate cancer cells *in vitro* (Bonaccorsi *et al.* 2004b, Festuccia *et al.* 2005). The involvement of EGFR in proliferation and invasion of cancer cells have been supported by other evidence (Wells *et al.* 2002). EGFR also participates in the formation of plasma membrane structures (lamellipodia) that mediate migration through the basal membrane (Rabinovitz *et al.* 2001). Significantly, elevated EGFR enhances the invasion potential of mammary tumors by increasing cell motility, without affecting tumor growth (Xue *et al.* 2006), pointing the key role exerted by the EGF/EGFR system in invasion and metastasis. Moreover, the robust evidence on the interaction between EGF/EGFR and androgen signaling provides proof of principle that engagement of multi-crossed signals is crucial for the acquisition and the maintenance of androgen sensitivity (Leotoing *et al.* 2007). Expression of the androgen-regulated prostate specific antigen, (*KLK3*) gene, is induced by the administration of interleukin-6 (IL6), which activates EGFR (Hobisch *et al.* 1998, Ueda *et al.* 2002). This evidence initially pointed to the contribution of EGFR in dictating AR outcomes in prostate cancer cells. ERBB2, a lead member of the EGFR family of receptor tyrosine kinases, was shown to be overexpressed in prostate cancer during progression to androgen-independent metastatic disease (Heinlein & Chang 2004). The mechanistic basis for important correlative cross-talk between AR and Erb2 has been provided by other reports indicating that modulation of AR signaling activity by the HER-2/neu tyrosine kinase promotes androgen-independent prostate tumor growth *in vitro* and *in vivo* (Craft *et al.* 1999, Yeh *et al.* 1999). More recent evidence further supports the signaling interaction by indicating that the loss of ERBB2 by siRNA impaired prostate cancer cell growth via targeting AR activity (Mellinghoff *et al.* 2004). Taken together, these lines of evidence converge to the recognition of the ERBB2 kinase activity being required for optimal transcriptional activity of AR in prostate cancer cells (Mellinghoff *et al.* 2004, Liu *et al.* 2005).

Androgens can post-transcriptionally control protein expression by regulating the binding of endogenous HuR to the AU-rich 3'UTRs, e.g. *EGF* mRNA (Myers *et al.* 1999, Topping *et al.* 2003). The ability of androgens to regulate the expression of androgen response element (ARE)-binding proteins that bind to these instability elements, supports an additional mechanistic involvement (by androgens) in the post-transcriptional control of EGF (Simons & Toomre 2000, DiNitto *et al.* 2003, Kuhajda 2006). In a 'reversal-of-action' mode, EGF reduces AR expression and blocks androgen-dependent transcription in differentiated cells, while it activates the AR promoter (Culig *et al.* 1994). This mechanistic EGF-AR interplay is an important contributor to prostate tumor progression, but it is not exclusive to EGF, as AR activity can be modulated by other growth factors (Orio *et al.* 2002).

AR interacts with the mitogen-activated protein kinase (MAPK)/extracellular signaling-regulated kinase kinase-1 (MEKK1) and the EGFR (Abreu-Martin *et al.* 1999, Bonaccorsi *et al.* 2004a; Fig. 1). Androgen-activated AR activates MAPK (Peterziel *et al.* 1999) and in a 'functional-symmetry', EGF-activated MAPK signaling cascade interferes with AR function, modulating the androgen response. MAPK extracellular kinase (MEK) inhibition reverses the EGF-mediated AR down-regulation in differentiated cells, thus suggesting the existence of an inverse correlation between EGF and androgen signaling in non-tumor epithelial cells (Leotoing *et al.* 2007). Additional key signal transducers in this dynamic, include transducer activator of transcription 3 (STAT3), most probably required for AR activation by IL6 toward promoting metastatic progression of prostate cancer (Abdulghani *et al.* 2008). Increased levels

of Stat3 have been shown to lead to Stat3–AR complex formation in response to EGF and IL6 (as shown on Fig. 1). Moreover, Stat3 increases the EGF-induced transcriptional activation of AR, while androgen pre-treatment increases Stat3 levels in an IL6 autocrine-/paracrine-dependent manner suggesting an intracellular feedback loop (Aaronson *et al.* 2007). AR can also affect clathrin-mediated endocytosis pathway of EGFR, an essential step in its signaling integrity. The significance of engaging such a robust cross-signaling by prostate cancer cells toward determining their survival and response to the microenvironment is established by growing evidence (Bonaccorsi *et al.* 2007).

The recently identified active integration of AR and EGFR signaling within the lipid raft microdomains in target cells provides an intriguing topological twist to this cross-talk. Thus, considering that the serine–threonine kinase AKT1 is a convergence point of the two hormonal stimuli and AR is localized in lipid raft membranes where it is stabilized by androgens (Freeman *et al.* 2007), one could easily argue that the newly found membrane ‘domain’ harboring AR is responsible for the non-genomic signaling by AR. The emerging concept that AKT1 is sensitive to manipulations in cholesterol levels, gains direct support from biochemical analysis verifying that a subpopulation of AKT1 molecules resides within lipid raft micro-domains (Bauer *et al.* 2003, Zhuang *et al.* 2005). Distinct changes in phosphorylation state of AKT1 in response to androgen occur quickly but temporally independent in the raft and non-raft compartment, implicating processing of dissimilar signals. Interestingly, EGF triggers AKT1 phosphorylation via more rapid kinetics than those induced by androgens; this was recently documented by studies on the sensitivity of EGFR family proteins to disruptions in cholesterol synthesis and homeostasis, supporting the functional significance of EGF signal transduction through lipid rafts (Freeman *et al.* 2007).

AR and IGF interactions

Signaling by IGF1 is of major mechanistic and biological significance (Burfeind *et al.* 1996, Pollak *et al.* 1998, Wolk *et al.* 1998, Nickerson *et al.* 2001). In a scenario, fostering AR reactivation in a low-androgen environment (Grossmann *et al.* 2001), insulin resistance, and hyperinsulinemia correlates with an increased incidence of prostate cancer (Fan *et al.* 2007). High IGF1 levels in the serum correlate with an increased risk of prostate cancer (Pollak *et al.* 1998, Wolk *et al.* 1998), whereas IGF1 enhances AR transactivation under low/absent androgen levels (Culig *et al.* 1994, Orio *et al.* 2002) and promotes prostate tumor cell proliferation (Burfeind *et al.* 1996).

Endogenous AR expression as well as AR transcriptional activity is regulated by insulin via activation of the phosphatidylinositol 3-kinase (PI3K) transduction pathway (Manin *et al.* 1992, 2000, 2002). FOXO1, as a downstream molecule becomes phosphorylated and inactivated by PI3K/AKT kinase in response to IGF1 or insulin, and subsequently suppresses ligand-mediated AR transactivation (Fig. 1). FOXO1 is recruited by liganded AR to the AR promoters and interacts directly with the C terminus of AR in a ligand-dependent manner disrupting ligand-induced AR nuclear compartmentalization. This FOXO1 interference with AR–DNA interactions suppresses androgen-induced AR activity resulting in prostate tumor cell growth suppression (Fan *et al.* 2007).

An intracrine positive feedback between IGF1 and AR signaling has been implicated in prostate cancer cells. Liganded AR up-regulates IGF1 receptor expression in HepG2 and LNCaP cells, presumably resulting in higher IGF1 signaling in prostate cancer cells (Wu *et al.* 2007). Two AREs within the IGF1 upstream promoter activate IGF1 expression (Wu *et al.* 2007). In addition, androgens can control IGF signaling via modulation of IGF-binding proteins (IGFBPs) in prostate epithelial cells, while both androgens and IGF1 up-regulate *IGFBP5* mRNA in androgen-responsive human fibroblasts (Yoshizawa & Ogikubo 2006). IGFBP5

initially binds IGFs with high affinity, principally by an N-terminal motif, and inhibits IGF activity by preventing IGF interaction with the type 1 receptor (Kalus *et al.* 1998). Taken together, this evidence supports a 'higher-level' interaction between AR and the IGF signaling, via recruitment of direct pathways toward activation, transcriptional regulation, and protein post-translational changes, all critical to tumor cell survival.

AR and TGF β interactions: cell death and survival partners

Transforming growth factor- β (TGFB) is a ubiquitous cytokine that plays a critical role in numerous pathways regulating cellular and tissue homeostasis. The TGFB superfamily members regulate proliferation, growth arrest, differentiation, and apoptosis of prostatic stromal and epithelial cells, as well as the formation of osteoblastic metastases. TGFB is over-expressed in advanced prostate cancer and exerts diverse functions in stromal cells via both SMAD-dependent and SMAD-independent signaling pathways (Coffey *et al.* 1986, Roberts *et al.* 1986, Derynck & Zhang 2003, Zhu & Kyprianou 2005). Recently, cofilin and prohibitin, two novel signaling effectors of TGFB1, that serve as potential intracellular effectors of its apoptotic action were identified in human prostate cancer cells (Zhu *et al.* 2006). Cancer cells become refractory to the growth inhibitory activity of TGFB due to the loss or mutation of transmembrane receptors or intracellular TGFB signaling effectors during tumor initiation (Akhurst & Derynck 2001).

During prostate tumor progression to metastatic disease, TGFB1 ligand overexpression results in prooncogenic rather than growth suppressive effect. In human prostate cancer cells, TGFB signaling proceeds via ligand binding and subsequent phosphorylation of TGFBR2 receptor to the TGFBR1 kinase to SMAD activation (Zhu & Kyprianou 2005). Interaction of SMAD4, (alone or together with SMAD3), with the AR in the DNA-binding and ligand-binding domains, may result in the modulation of DHT-induced AR transactivation (Zhu *et al.* 2008). Interestingly, in the human prostate cancer cell lines PC3 and LNCaP, addition of SMAD3 enhances AR transactivation, while co-transfection of SMAD3 and SMAD4 actually repress AR transactivation (Kang *et al.* 2002). A protein-protein interaction between AR and SMAD3 has been documented both *in vitro* and *in vivo* via the transcription activation domain of AR and the MH2 of SMAD3; AR repression by SMAD3 is mediated through the MH2 domain (Hayes *et al.* 2001). In PC-3 prostate cancer cells, AR expression reduces the TGFB1/SMAD transcriptional activity and the growth effects of TGFB1 (in the absence of DHT), thus preventing TGFB1-induced growth inhibition and apoptosis. Furthermore, TGFB1 suppresses the E2F transcriptional activity of AR activation by DHT, an event that is associated with a reduced c-Myc expression. An ARE sequence in the TGFB promoter may provide a mechanistic basis for TGFB promoter activity toward DHT in both Huh7 and PC3/AR-expressing cells. A direct interaction between AR and TGFB1 has been causally implicated in other human tumors including hepatocarcinogenesis (Yoon *et al.* 2006). Androgens can inhibit TGFB1-induced transcriptional activity in prostate cancer cells (Chipuk *et al.* 2002), an interaction that is regulated by AR-associated protein 55 (ARA55/Hic-5; LIM protein superfamily). Overexpression of ARA55 inhibits TGFB-mediated up-regulation of SMAD transcriptional activity in rat prostate epithelial cells, as well as human prostate cells, via an interaction between ARA55 and SMAD3 mediated through the MH2 domain of SMAD3 and the C terminus of ARA55 (Wang *et al.* 2005).

The involvement of AR in the apoptosis outcomes of TGFB signaling in prostate cancer cells is supported by work from this laboratory. Treatment of TGFB receptor II overexpressing LNCaP TGFBR2 cells with TGFB in the presence of DHT, both cell cycle arrest and apoptosis induction are significantly enhanced over TGFB alone, through caspase-1 activation and targeting of BCL-2 (Bruckheimer & Kyprianou 2001). Enforced BCL2 expression significantly inhibits the combined TGFB and DHT apoptotic effect in prostate cancer cells

(Bruckheimer & Kyprianou 2002). An androgenic contribution, with TGFB enhancement, on the epithelial-mesenchymal transition (EMT) provides an attractive mechanistic possibility in view of the assigned role of EMT during cancer metastasis (Zavadil & Bottinger 2005), with E-cadherin being considered as a potential target for such a dynamic duo.

AR and FGF interactions

The FGF family is a large family of proteins with broad spectrum of functions, including cell migration, differentiation, and angiogenesis (Ornitz & Itoh 2001). Changes in the expression of FGFs and/or their receptors are involved in prostate tumor progression toward androgen-independent disease. The estrogen receptor (ER) can regulate the synthesis of FGF2 and FGF7 in prostate cells, while stromal ER can mediate the synthesis of stromally derived growth factors, both in coordination with AR activation. AR signaling can directly dictate dramatic changes in the expression pattern of FGFs in both prostate tumor epithelial cells and stromal cells, primarily via changes in FGF1, FGF2, FGF8, and FGF10 (Saric & Shain 1998, Nakano *et al.* 1999, Rosini *et al.* 2002). Via a positive feedback, AR is up-regulated by paracrine FGF10 and synergizes with cell-autonomous activated AKT in prostate cancer cells (Memarzadeh *et al.* 2007). Moreover, in response to FGFs, AR facilitates FGF-induced survival of prostate cancer cells, possibly through BCL2 induction and down-regulation of AR, allowing the escape of selected clones from androgenic control (Rosini *et al.* 2002, Gonzalez-Herrera *et al.* 2006).

AR and vascular endothelial growth factor (VEGF) interactions

VEGF, originally known as vascular permeability factor, is a well-characterized angiogenic cytokine, responsible for endothelial cell proliferation, migration, and vessel assembly (Fong *et al.* 1995). Its value as a diagnostic tool as well as a therapeutic target for advanced metastatic prostate cancer has been examined at the molecular and translational level.

The 'hypoxia-response' signaling system up-regulates the expression of a network of effectors that increase the propensity of tumor cells for survival, even in this adverse environment (Anastasiadis *et al.* 2003). Expression of VEGF is transcriptionally induced by hypoxia-inducible factor (HIF1A) in response to oxygen changes in the microenvironment (Delongchamps *et al.* 2006). Androgen-stimulated growth of the glandular ventral prostate is preceded by increased VEGF synthesis, endothelial cell proliferation, vascular growth, and increased blood flow (Joseph *et al.* 1997, Franck-Lissbrant *et al.* 1998). The role of VEGF in androgen-mediated prostate vascularity was further supported by additional studies (Lissbrant *et al.* 2004). In prostate cancer, the effect of androgens on angiogenesis is mediated via their ability to regulate VEGF through activation of HIF1A in androgen-sensitive tumors (Boddy *et al.* 2005). The significant correlation between HIF1A and HIF2A expression and with AR and VEGF expression (Boddy *et al.* 2005, Banham *et al.* 2007) provides firm support for such a control system. The driving mechanism involves the direct up-regulation of VEGF-C in response to androgen depletion in prostate cancer cells (Rinaldo *et al.* 2007), via activation of the small GTPase, RalA; VEGF-C can increase the AR co-activator BAG-1L expression that facilitates AR transactivation. Under conditions of low-androgen levels, the intracellular reactive oxygen species induce RalA activation and VEGF-C synthesis (Rinaldo *et al.* 2007).

AR and growth factor interplay in the stroma

The stroma is a lead component of the prostate microenvironment contributing to tumor heterogeneity and growth dynamics. Stroma-derived fibroblasts play an active role in carcinogenesis in addition to structurally supporting the epithelial cell growth (Chung *et al.* 1989, 1991, Camps *et al.* 1990, Cunha *et al.* 1996). Studies in the early 1990s established that human prostate-derived stromal cells stimulate growth of prostate cancer cells *in vitro* and *in*

vivo (Gleave *et al.* 1991). This evidence widely popularized the belief that disturbance in the epithelial–stromal interactions is most critical in the pathogenesis of prostate cancer (Hayward *et al.* 1998). Androgenic control during normal growth and differentiation of the prostate gland is regulated via nuclear AR in both stromal and epithelial cells (Sar *et al.* 1990). The close association between low-AR levels in the stroma adjacent to malignant epithelium, with a poor clinical outcome in prostate cancer patients is of high translational value (Henshall *et al.* 2001). Androgens increase VEGF transcription and active VEGF secretion from prostatic stroma, thus indirectly enhancing prostate cancer growth and angiogenesis (Levine *et al.* 1998). DHT and FGF2 can synergistically stimulate prostate stromal cell proliferation (Niu *et al.* 2001), while androgen depletion rapidly reduces stroma IGF1 synthesis and its action in the prostate epithelium. Close rules of compartmentalization become ‘loose’ here: although IGF1 is principally produced in the stroma and IGF-R1 in the epithelium, both are under androgenic regulation as stroma *IGF1* mRNA is significantly decreased after castration, correlating with epithelial cell apoptosis (Ohlson *et al.* 2007).

TGFB1 is also regulator of stromal cell proliferation and differentiation, depending on the specific stromal cell type, microenvironment, and contributing activities of other growth factors (Sporn & Roberts 1992). A distinct in its complexity cross-talk between androgens and TGFB1 signaling in prostate stromal cells affects AR localization, cell proliferation, and myodifferentiation, thus defining its mechanistic contribution to the reactive stroma. AR and TGFB1 levels significantly correlate in the stromal component of prostatic intraepithelial neoplasia (Cardillo *et al.* 2000). Induction of rat PS-1 prostate stromal cell proliferation by androgens can be antagonized by TGFB1. Furthermore, TGFB1 triggers a cytoplasmic translocation of nuclear AR during myodifferentiation in the prostate stroma (Gerdes *et al.* 1998, 2004), while androgens enhance TGFB1-mediated proliferation of prostatic smooth muscle cells PSMC1 (Salm *et al.* 2000).

During prostate cancer progression the androgen axis engages the growth factor network to an active cross-talk toward conferring a survival and invasion advantage of prostate cancer cells. The current evidence dissecting this signaling interaction between the AR and growth factors is discussed in this review. Androgens can modify prostate cancer cell response to growth factor signals from growth inhibitory to tumor promoting during the metastatic process. A better understanding of such cross-talk between the AR axis and critical growth factor signaling in the context of the tumor microenvironment, may identify a mechanism underlying the emergence of androgen-independent prostate cancer, and provide new opportunities for therapeutic targeting of aggressive prostate tumors.

Acknowledgments

The authors acknowledge the assistance of Lorie Howard during the submission process.

Funding

This work was supported by an NIH/National Institutes of Diabetes, Digestive and Kidney Diseases R01 grant (DK-53525-08).

References

- Aaronson DS, Muller M, Neves SR, Chung WC, Jayaram G, Iyengar R, Ram PT. An androgen-IL-6-Stat3 autocrine loop re-routes EGF signal in prostate cancer cells. *Molecular and Cellular Endocrinology* 2007;270:50–56. [PubMed: 17374439]
- Abdulghani J, Gu L, Dagvadorj A, Lutz J, Leiby B, Bonuccelli G, Lisanti MP, Zellweger T, Alanen K, Mirtti T, et al. Stat3 promotes metastatic progression of prostate cancer. *American Journal of Pathology* 2008;172:1717–1728. [PubMed: 18483213]

- Abreu-Martin MT, Chari A, Palladino AA, Craft NA, Sawyers CL. Mitogen-activated protein kinase kinase 1 activates androgen receptor-dependent transcription and apoptosis in prostate cancer. *Molecular and Cellular Biology* 1999;19:5143–5154. [PubMed: 10373563]
- Akhurst RJ, Derynck R. TGF-beta signaling in cancer – a double-edged sword. *Trends in Cell Biology* 2001;11:S44–S51. [PubMed: 11684442]
- Anastasiadis AG, Bemis DL, Stisser BC, Salomon L, Ghafar MA, Buttyan R. Tumor cell hypoxia and the hypoxia-response signaling system as a target for prostate cancer therapy. *Current Drug Targets* 2003;4:191–196. [PubMed: 12643469]
- Banham AH, Boddy J, Launchbury R, Han C, Turley H, Malone PR, Harris AL, Fox SB. Expression of the forkhead transcription factor FOXP1 is associated both with hypoxia inducible factors (HIFs) and the androgen receptor in prostate cancer but is not directly regulated by androgens or hypoxia. *Prostate* 2007;67:1091–1098. [PubMed: 17477366]
- Bauer B, Jenny M, Fresser F, Uberall F, Baier G. AKT1/PKBalpha is recruited to lipid rafts and activated downstream of PKC isotypes in CD3-induced T cell signaling. *FEBS Letters* 2003;541:155–162. [PubMed: 12706837]
- Boddy JL, Fox SB, Han C, Campo L, Turley H, Kanga S, Malone PR, Harris AL. The androgen receptor is significantly associated with vascular endothelial growth factor and hypoxia sensing via hypoxia-inducible factors HIF-1a, HIF-2a, and the prolyl hydroxylases in human prostate cancer. *Clinical Cancer Research* 2005;11:7658–7663. [PubMed: 16278385]
- Bonaccorsi L, Carloni V, Muratori M, Formigli L, Zecchi S, Forti G, Baldi E. EGF receptor (EGFR) signaling promoting invasion is disrupted in androgen-sensitive prostate cancer cells by an interaction between EGFR and androgen receptor (AR). *International Journal of Cancer* 2004a;112:78–86.
- Bonaccorsi L, Marchiani S, Muratori M, Forti G, Baldi E. Gefitinib ('IRESSA', ZD1839) inhibits EGF-induced invasion in prostate cancer cells by suppressing PI3 K/AKT activation. *Journal of Cancer Research and Clinical Oncology* 2004b;130:604–614. [PubMed: 15258753]
- Bonaccorsi L, Nosi D, Muratori M, Formigli L, Forti G, Baldi E. Altered endocytosis of epidermal growth factor receptor in androgen receptor positive prostate cancer cell lines. *Journal of Molecular Endocrinology* 2007;38:51–66. [PubMed: 17242169]
- Bruckheimer EM, Kyprianou N. Dihydrotestosterone enhances transforming growth factor-beta-induced apoptosis in hormone-sensitive prostate cancer cells. *Endocrinology* 2001;142:2419–2426. [PubMed: 11356690]
- Bruckheimer EM, Kyprianou N. BCL-2 antagonizes the combined apoptotic effect of transforming growth factor-beta and dihydrotestosterone in prostate cancer cells. *Prostate* 2002;53:133–142. [PubMed: 12242728]
- Burfeind P, Chernicky CL, Rininsland F, Ilan J. Antisense RNA to the type I insulin-like growth factor receptor suppresses tumor growth and prevents invasion by rat prostate cancer cells *in vivo*. *PNAS* 1996;93:7263–7268. [PubMed: 8692980]
- Byrne RL, Leung H, Neal DE. Peptide growth factors in the prostate as mediators of stromal epithelial interaction. *British Journal of Urology* 1996;77:627–633. [PubMed: 8689101]
- Camps JL, Chang SM, Hsu TC, Freeman MR, Hong SJ, Zhau HE, von Eschenbach AC, Chung LW. Fibroblast-mediated acceleration of human epithelial tumor growth *in vivo*. *PNAS* 1990;87:75–79. [PubMed: 2296606]
- Cardillo MR, Petrangeli E, Perracchio L, Salvatori L, Ravenna L, Di Silverio F. Transforming growth factor-beta expression in prostate neoplasia. *Analytical and Quantitative Cytology and Histology* 2000;22:1–10. [PubMed: 10696454]
- Chipuk JE, Cornelius SC, Pultz NJ, Jorgensen JS, Bonham MJ, Kim SJ, Danielpour D. The androgen receptor represses transforming growth factor-beta signaling through interaction with Smad3. *Journal of Biological Chemistry* 2002;277:1240–1248. [PubMed: 11707452]
- Chung LW, Chang SM, Bell C, Zhau HE, Ro JY, von Eschenbach AC. Co-inoculation of tumorigenic rat prostate mesenchymal cells with non-tumorigenic epithelial cells results in the development of carcinosarcoma in syngeneic and athymic animals. *International Journal of Cancer* 1989;43:1179–1187.

- Chung LW, Gleave ME, Hsieh JT, Hong SJ, Zhau HE. Reciprocal mesenchymal–epithelial interaction affecting prostate tumour growth and hormonal responsiveness. *Cancer Surveys* 1991;11:91–121. [PubMed: 1726790]
- Coffey RJ Jr, Shipley GD, Moses HL. Production of transforming growth factors by human colon cancer lines. *Cancer Research* 1986;46:1164–1169. [PubMed: 3002610]
- Craft N, Shostak Y, Carey M, Sawyers CL. A mechanism for hormone-independent prostate cancer through modulation of androgen receptor signaling by the HER-2/neu tyrosine kinase. *Nature Medicine* 1999;5:280–285.
- Culig Z, Hobisch A, Cronauer MV, Radmayr C, Trapman J, Hittmair A, Bartsch G, Klocker H. Androgen receptor activation in prostatic tumor cell lines by insulin-like growth factor-I, keratinocyte growth factor, and epidermal growth factor. *Cancer Research* 1994;54:5474–5478. [PubMed: 7522959]
- Cunha GR, Donjacour AA. Mesenchymal–epithelial interactions in the growth and development of the prostate. *Cancer Treatment and Research* 1989;46:159–175. [PubMed: 2577188]
- Cunha GR, Hayward SW, Dahiya R, Foster BA. Smooth muscle–epithelial interactions in normal and neoplastic prostatic development. *Acta Anatomica* 1996;155:63–72. [PubMed: 8811117]
- Delongchamps NB, Peyromaure M, Dinh-Xuan AT. Role of vascular endothelial growth factor in prostate cancer. *Urology* 2006;68:244–248. [PubMed: 16904427]
- Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature* 2003;425:577–584. [PubMed: 14534577]
- DiNitto JP, Cronin TC, Lambright DG. Membrane recognition and targeting by lipid-binding domains. *Science's STKE* 2003;2003:re16.
- Fan W, Yanase T, Morinaga H, Okabe T, Nomura M, Daitoku H, Fukamizu A, Kato S, Takayanagi R, Nawata H. Insulin-like growth factor 1/insulin signaling activates androgen signaling through direct interactions of Foxo1 with androgen receptor. *Journal of Biological Chemistry* 2007;282:7329–7338. [PubMed: 17202144]
- Festuccia C, Muzi P, Millimaggi D, Biordi L, Gravina GL, Specia S, Angelucci A, Dolo V, Vicentini C, Bologna M. Molecular aspects of gefitinib antiproliferative and pro-apoptotic effects in PTEN-positive and PTEN-negative prostate cancer cell lines. *Endocrine-Related Cancer* 2005;12:983–998. [PubMed: 16322337]
- Fong GH, Rossant J, Gertsenstein M, Breitman ML. Role of the Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. *Nature* 1995;376:66–70. [PubMed: 7596436]
- Franck-Lissbrant I, Haggstrom S, Damber JE, Bergh A. Testosterone stimulates angiogenesis and vascular regrowth in the ventral prostate in castrated adult rats. *Endocrinology* 1998;139:451–456. [PubMed: 9449610]
- Freeman MR, Cinar B, Kim J, Mukhopadhyay NK, Di Vizio D, Adam RM, Solomon KR. Transit of hormonal and EGF receptor-dependent signals through cholesterol-rich membranes. *Steroids* 2007;72:210–217. [PubMed: 17173942]
- Gerdes MJ, Dang TD, Larsen M, Rowley DR. Transforming growth factor-beta1 induces nuclear to cytoplasmic distribution of androgen receptor and inhibits androgen response in prostate smooth muscle cells. *Endocrinology* 1998;139:3569–3577. [PubMed: 9681509]
- Gerdes MJ, Larsen M, Dang TD, Ressler SJ, Tuxhorn JA, Rowley DR. Regulation of rat prostate stromal cell myodifferentiation by androgen and TGF-beta1. *Prostate* 2004;58:299–307. [PubMed: 14743470]
- Gleave M, Hsieh JT, Gao CA, von Eschenbach AC, Chung LW. Acceleration of human prostate cancer growth *in vivo* by factors produced by prostate and bone fibroblasts. *Cancer Research* 1991;51:3753–3761. [PubMed: 1712249]
- Gonzalez-Herrera IG, Prado-Lourenco L, Pileur F, Conte C, Morin A, Cabon F, Prats H, Vagner S, Bayard F, Audigier S, et al. Testosterone regulates FGF-2 expression during testis maturation by an IRES-dependent translational mechanism. *FASEB Journal* 2006;20:476–478. [PubMed: 16423876]
- Grossmann ME, Huang H, Tindall DJ. Androgen receptor signaling in androgen-refractory prostate cancer. *Journal of the National Cancer Institute* 2001;93:1687–1697. [PubMed: 11717329]
- Hayes SA, Zarnegar M, Sharma M, Yang F, Peehl DM, ten Dijke P, Sun Z. SMAD3 represses androgen receptor-mediated transcription. *Cancer Research* 2001;61:2112–2118. [PubMed: 11280774]

- Hayward SW, Grossfeld GD, Tlsty TD, Cunha GR. Genetic and epigenetic influences in prostatic carcinogenesis (review). *International Journal of Oncology* 1998;13:35–47. [PubMed: 9625801]
- Heinlein CA, Chang C. The roles of androgen receptors and androgen-binding proteins in nongenomic androgen actions. *Molecular Endocrinology* 2002;16:2181–2187. [PubMed: 12351684]
- Heinlein CA, Chang C. Androgen receptor in prostate cancer. *Endocrine Reviews* 2004;25:276–308. [PubMed: 15082523]
- Henshall SM, Quinn DI, Lee CS, Head DR, Golovsky D, Brenner PC, Delprado W, Stricker PD, Grygiel JJ, Sutherland RL. Altered expression of androgen receptor in the malignant epithelium and adjacent stroma is associated with early relapse in prostate cancer. *Cancer Research* 2001;61:423–427. [PubMed: 11212224]
- Hobisch A, Eder IE, Putz T, Horninger W, Bartsch G, Klocker H, Culig Z. Interleukin-6 regulates prostate-specific protein expression in prostate carcinoma cells by activation of the androgen receptor. *Cancer Research* 1998;58:4640–4645. [PubMed: 9788616]
- Hsing AW. Hormones and prostate cancer: what's next? *Epidemiologic Reviews* 2001;23:42–58. [PubMed: 11588854]
- Imperato-McGinley J, Binienda Z, Arthur A, Mininberg DT, Vaughan ED Jr, Quimby FW. The development of a male pseudohermaphroditic rat using an inhibitor of the enzyme 5 alpha-reductase. *Endocrinology* 1985;116:807–812. [PubMed: 3967629]
- Joseph IB, Nelson JB, Denmeade SR, Isaacs JT. Androgens regulate vascular endothelial growth factor content in normal and malignant prostatic tissue. *Clinical Cancer Research* 1997;3:2507–2511. [PubMed: 9815654]
- Kalus W, Zweckstetter M, Renner C, Sanchez Y, Georgescu J, Grol M, Demuth D, Schumacher R, Dony C, Lang K, et al. Structure of the IGF-binding domain of the insulin-like growth factor-binding protein-5 (IGFBP-5): implications for IGF and IGF-I receptor interactions. *EMBO Journal* 1998;17:6558–6572. [PubMed: 9822601]
- Kang HY, Huang KE, Chang SY, Ma WL, Lin WJ, Chang C. Differential modulation of androgen receptor-mediated transactivation by Smad3 and tumor suppressor Smad4. *Journal of Biological Chemistry* 2002;277:43749–43756. [PubMed: 12226080]
- Kuhajda FP. Fatty acid synthase and cancer: new application of an old pathway. *Cancer Research* 2006;66:5977–5980. [PubMed: 16778164]
- Leotoing L, Manin M, Monte D, Baron S, Communal Y, Lours C, Veyssiere G, Morel L, Beaudoin C. Crosstalk between androgen receptor and epidermal growth factor receptor-signalling pathways: a molecular switch for epithelial cell differentiation. *Journal of Molecular Endocrinology* 2007;39:151–162. [PubMed: 17693613]
- Levine AC, Liu XH, Greenberg PD, Eliashvili M, Schiff JD, Aaronson SA, Holland JF, Kirschenbaum A. Androgens induce the expression of vascular endothelial growth factor in human fetal prostatic fibroblasts. *Endocrinology* 1998;139:4672–4678. [PubMed: 9794479]
- Lissbrant IF, Hammarsten P, Lissbrant E, Ferrara N, Rudolphsson SH, Bergh A. Neutralizing VEGF bioactivity with a soluble chimeric VEGF-receptor protein flt(1–3)IgG inhibits testosterone-stimulated prostate growth in castrated mice. *Prostate* 2004;58:57–65. [PubMed: 14673953]
- Liu Y, Majumder S, McCall W, Sartor CI, Mohler JL, Gregory CW, Earp HS, Whang YE. Inhibition of HER-2/neu kinase impairs androgen receptor recruitment to the androgen responsive enhancer. *Cancer Research* 2005;65:3404–3409. [PubMed: 15833875]
- Di Lorenzo G, Tortora G, D'Armiento FP, De Rosa G, Staibano S, Autorino R, D'Armiento M, De Laurentiis M, De Placido S, Catalano G, et al. Expression of epidermal growth factor receptor correlates with disease relapse and progression to androgen-independence in human prostate cancer. *Clinical Cancer Research* 2002;8:3438–3444. [PubMed: 12429632]
- Manin M, Veyssiere G, Cheyvialle D, Chevalier M, Lecher P, Jean C. *In vitro* androgenic induction of a major protein in epithelial cell subcultures from mouse vas deferens. *Endocrinology* 1992;131:2378–2386. [PubMed: 1385102]
- Manin M, Martinez A, Van Der Schueren B, Reynaert I, Jean C. Acquisition of androgen-mediated expression of mouse vas deferens protein (MVDP) gene in cultured epithelial cells and in vas deferens during postnatal development. *Journal of Andrology* 2000;21:641–650. [PubMed: 10975410]

- Manin M, Baron S, Goossens K, Beaudoin C, Jean C, Veyssiere G, Verhoeven G, Morel L. Androgen receptor expression is regulated by the phosphoinositide 3-kinase/Akt pathway in normal and tumoral epithelial cells. *Biochemical Journal* 2002;366:729–736. [PubMed: 11971763]
- De Marzo AM, Nelson WG, Meeker AK, Coffey DS. Stem cell features of benign and malignant prostate epithelial cells. *Journal of Urology* 1998;160:2381–2392. [PubMed: 9817389]
- Mellinghoff IK, Vivanco I, Kwon A, Tran C, Wongvipat J, Sawyers CL. HER2/neu kinase-dependent modulation of androgen receptor function through effects on DNA binding and stability. *Cancer Cell* 2004;6:517–527. [PubMed: 15542435]
- Memarzadeh S, Xin L, Mulholland DJ, Mansukhani A, Wu H, Teitell MA, Witte ON. Enhanced paracrine FGF10 expression promotes formation of multifocal prostate adenocarcinoma and an increase in epithelial androgen receptor. *Cancer Cell* 2007;12:572–585. [PubMed: 18068633]
- Myers RB, Oelschlager D, Manne U, Coan PN, Weiss H, Grizzle WE. Androgenic regulation of growth factor and growth factor receptor expression in the CWR22 model of prostatic adenocarcinoma. *International Journal of Cancer* 1999;82:424–429.
- Nakano K, Fukabori Y, Itoh N, Lu W, Kan M, McKeenhan WL, Yamanaka H. Androgen-stimulated human prostate epithelial growth mediated by stromal-derived fibroblast growth factor-10. *Endocrine Journal* 1999;46:405–413. [PubMed: 10503993]
- Nickerson T, Chang F, Lorimer D, Smeekens SP, Sawyers CL, Pollak M. *In vivo* progression of LAPC-9 and LNCaP prostate cancer models to androgen independence is associated with increased expression of insulin-like growth factor I (IGF-I) and IGF-I receptor (IGF-IR). *Cancer Research* 2001;61:6276–6280. [PubMed: 11507082]
- Niu Y, Xu Y, Zhang J, Bai J, Yang H, Ma T. Proliferation and differentiation of prostatic stromal cells. *BJU International* 2001;87:386–393. [PubMed: 11251536]
- Ohlson N, Bergh A, Stattin P, Wikstrom P. Castration-induced epithelial cell death in human prostate tissue is related to locally reduced IGF-1 levels. *Prostate* 2007;67:32–40. [PubMed: 16998819]
- Orio F Jr, Terouanne B, Georget V, Lumbroso S, Avances C, Siatka C, Sultan C. Potential action of IGF-1 and EGF on androgen receptor nuclear transfer and transactivation in normal and cancer human prostate cell lines. *Molecular and Cellular Endocrinology* 2002;198:105–114. [PubMed: 12573820]
- Ornitz DM, Itoh N. Fibroblast growth factors. *Genome Biology* 2001;2:REVIEWS3005. [PubMed: 11276432]
- Peterziel H, Mink S, Schonert A, Becker M, Klocker H, Cato AC. Rapid signalling by androgen receptor in prostate cancer cells. *Oncogene* 1999;18:6322–6329. [PubMed: 10597231]
- Pollak M, Beamer W, Zhang JC. Insulin-like growth factors and prostate cancer. *Cancer and Metastasis Reviews* 1998;17:383–390. [PubMed: 10453282]
- Rabinovitz I, Gipson IK, Mercurio AM. Traction forces mediated by alpha6beta4 integrin: implications for basement membrane organization and tumor invasion. *Molecular Biology of the Cell* 2001;12:4030–4043. [PubMed: 11739798]
- Rinaldo F, Li J, Wang E, Muders M, Datta K. RalA regulates vascular endothelial growth factor-C (VEGF-C) synthesis in prostate cancer cells during androgen ablation. *Oncogene* 2007;26:1731–1738. [PubMed: 16964283]
- Roberts JT, Essenhigh DM. Adenocarcinoma of prostate in 40-year-old body-builder. *Lancet* 1986;2:742. [PubMed: 2876202]
- Roberts AB, Sporn MB, Assoian RK, Smith JM, Roche NS, Wakefield LM, Heine UI, Liotta LA, Falanga V, Kehr JH, et al. Transforming growth factor type beta: rapid induction of fibrosis and angiogenesis *in vivo* and stimulation of collagen formation *in vitro*. *PNAS* 1986;83:4167–4171. [PubMed: 2424019]
- Rosini P, Bonaccorsi L, Baldi E, Chiasserini C, Forti G, De Chiara G, Lucibello M, Mongiat M, Iozzo RV, Garaci E, et al. Androgen receptor expression induces FGF2, FGF-binding protein production, and FGF2 release in prostate carcinoma cells: role of FGF2 in growth, survival, and androgen receptor down-modulation. *Prostate* 2002;53:310–321. [PubMed: 12430142]
- Russell PJ, Bennett S, Stricker P. Growth factor involvement in progression of prostate cancer. *Clinical Chemistry* 1998;44:705–723. [PubMed: 9554481]

- Salm SN, Koikawa Y, Ogilvie V, Tsujimura A, Coetzee S, Moscatelli D, Moore E, Lepor H, Shapiro E, Sun TT, et al. Generation of active TGF-beta by prostatic cell cocultures using novel basal and luminal prostatic epithelial cell lines. *Journal of Cellular Physiology* 2000;184:70–79. [PubMed: 10825235]
- Sar M, Lubahn DB, French FS, Wilson EM. Immunohistochemical localization of the androgen receptor in rat and human tissues. *Endocrinology* 1990;127:3180–3186. [PubMed: 1701137]
- Saric T, Shain SA. Androgen regulation of prostate cancer cell FGF-1, FGF-2, and FGF-8: preferential down-regulation of FGF-2 transcripts. *Growth Factors* 1998;16:69–87. [PubMed: 9777371]
- Siiteri PK, Wilson JD. Testosterone formation and metabolism during male sexual differentiation in the human embryo. *Journal of Clinical Endocrinology and Metabolism* 1974;38:113–125. [PubMed: 4809636]
- Simons K, Toomre D. Lipid rafts and signal transduction. *Nature Reviews. Molecular Cell Biology* 2000;1:31–39. [PubMed: 11413487]
- Sporn MB, Roberts AB. Transforming growth factor-beta: recent progress and new challenges. *Journal of Cell Biology* 1992;119:1017–1021. [PubMed: 1332976]
- Torring N, Dagnaes-Hansen F, Sorensen BS, Nexø E, Hynes NE. ErbB1 and prostate cancer: ErbB1 activity is essential for androgen-induced proliferation and protection from the apoptotic effects of LY294002. *Prostate* 2003;56:142–149. [PubMed: 12746839]
- Ueda T, Mawji NR, Bruchovsky N, Sadar MD. Ligand-independent activation of the androgen receptor by interleukin-6 and the role of steroid receptor coactivator-1 in prostate cancer cells. *Journal of Biological Chemistry* 2002;277:38087–38094. [PubMed: 12163482]
- Wang H, Song K, Sponseller TL, Danielpour D. Novel function of androgen receptor-associated protein 55/Hic-5 as a negative regulator of Smad3 signaling. *Journal of Biological Chemistry* 2005;280:5154–5162. [PubMed: 15561701]
- Wang X, Yin L, Rao P, Stein R, Harsch KM, Lee Z, Heston WD. Targeted treatment of prostate cancer. *Journal of Cellular Biochemistry* 2007;102:571–579. [PubMed: 17685433]
- Wells A, Kassir J, Solava J, Turner T, Lauffenburger DA. Growth factor-induced cell motility in tumor invasion. *Acta Oncologica* 2002;41:124–130. [PubMed: 12102155]
- Wolk A, Mantzoros CS, Andersson SO, Bergstrom R, Signorello LB, Lagiou P, Adami HO, Trichopoulos D. Insulin-like growth factor 1 and prostate cancer risk: a population-based, case-control study. *Journal of the National Cancer Institute* 1998;90:911–915. [PubMed: 9637140]
- Wu Y, Zhao W, Zhao J, Pan J, Wu Q, Zhang Y, Bauman WA, Cardozo CP. Identification of androgen response elements in the insulin-like growth factor I upstream promoter. *Endocrinology* 2007;148:2984–2993. [PubMed: 17363459]
- Xue C, Wyckoff J, Liang F, Sidani M, Violini S, Tsai KL, Zhang ZY, Sahai E, Condeelis J, Segall JE. Epidermal growth factor receptor overexpression results in increased tumor cell motility *in vivo* coordinately with enhanced intravasation and metastasis. *Cancer Research* 2006;66:192–197. [PubMed: 16397232]
- Yeh S, Lin HK, Kang HY, Thin TH, Lin MF, Chang C. From HER2/Neu signal cascade to androgen receptor and its coactivators: a novel pathway by induction of androgen target genes through MAP kinase in prostate cancer cells. *PNAS* 1999;96:5458–5463. [PubMed: 10318905]
- Yoon G, Kim JY, Choi YK, Won YS, Lim IK. Direct activation of TGF-beta1 transcription by androgen and androgen receptor complex in Huh7 human hepatoma cells and its tumor in nude mice. *Journal of Cellular Biochemistry* 2006;97:393–411. [PubMed: 16187311]
- Yoshizawa A, Ogikubo S. IGF binding protein-5 synthesis is regulated by testosterone through transcriptional mechanisms in androgen responsive cells. *Endocrine Journal* 2006;53:811–818. [PubMed: 17001112]
- Zavadil J, Bottinger EP. TGF-beta and epithelial-to-mesenchymal transitions. *Oncogene* 2005;24:5764–5774. [PubMed: 16123809]
- Zhu B, Kyprianou N. Transforming growth factor beta and prostate cancer. *Cancer Treatment and Research* 2005;126:157–173. [PubMed: 16209066]
- Zhu B, Fukada K, Zhu H, Kyprianou N. Prohibitin and cofilin are intracellular effectors of transforming growth factor beta signaling in human prostate cancer cells. *Cancer Research* 2006;66:8640–8647. [PubMed: 16951178]

- Zhu ML, Partin JV, Bruckheimer EM, Strup SE, Kyprianou N. TGF-beta signaling and androgen receptor status determine apoptotic cross-talk in human prostate cancer cells. *Prostate* 2008;68:287–295. [PubMed: 18163430]
- Zhuang L, Kim J, Adam RM, Solomon KR, Freeman MR. Cholesterol targeting alters lipid raft composition and cell survival in prostate cancer cells and xenografts. *Journal of Clinical Investigation* 2005;115:959–968. [PubMed: 15776112]

Figure 1.

Growth factors cross-talk with AR in prostate cancer cells. IGF, FGF, VEGF, and TGFβ secreted by the prostate stromal cells activate their receptors and interact with AR signal axis. In prostate epithelial cells, the androgenic signal engages secreted VEGF and TGFβ which affects the prostate tumor microenvironment by inducing angiogenesis and stromal cell growth and differentiation. EGF signaling encounters AR signal in a tight control of multiple pathways. Growth factor signaling may proceed via AR signal and regulate the downstream effectors of AR regulating key cellular processes including proliferation, differentiation, apoptosis, and survival of prostate cancer cells.