Children are often undertreated for pain

Janice Hopkins Tanne *New York*

"Children in pain rarely seem to get the attention they need. There is a substantive gap between what we know and what we do," said Dr Richard Howard, from the anaesthesiology department at Great Ormond Street Children's Hospital, London, speaking at the American Medical Association's briefing in New York to launch JAMA's theme issue on pain.

"Infants used to get much less treatment than adults for the same conditions," he said, because doctors feared side effects in children. Although improvements had been made in the past 20 years—as doctors learnt more about the immature nervous system—there were few good evidence based studies, he said.

In his paper reviewing the current status of pain management in children, he said that children were not good at reporting pain, even when using visual scales ranging from no pain to worst pain (*JAMA* 2003;290:2464-9). Doctors are now better at using indirect measures to gauge pain and the effectiveness of treatment, and self controlled analgesia and

epidural anaesthesia have been adapted for use in children.

Neuropathic pain resulting from damage to the nervous system is more common than previously thought, but little is known about treating it, said Dr Howard. He said that the nervous system is not fully developed at birth and grows through different stages in infancy and that infants and children may perceive pain differently at the different stages.

Little is known about the long term effect of brief pain—as from needles used in immunisation—compared with pain after surgery, he said. For minor pain, such as that from needles, he suggested using local anaesthetic creams or behavioural methods to reduce children's anxiety.

A Dutch team, led by Dr Dick Tibboel of Erasmus Medical Center-Sophia Children's Hospital in Rotterdam, reported in another journal published by the American Medical Association that 151 premature babies in a neonatal intensive care unit had 14 painful procedures every day during their first two weeks of life, many without analgesia

Doctors estimate that 26 out of 31 procedures used in neonatal intensive care are painful, but many babies do not receive analgesia

(*Archives of Pediatric and Adolescent Medicine* 2003;157:1058-64).

The clinicians estimated that 26 of 31 procedures in a checklist of all procedures would be painful. Each day less than 35% of the babies received analgesia before a painful procedure, and nearly 40% did not receive any analgesia during their time in the unit. The team said painful procedures should be reduced and analgesia improved.

However, another study by Dr Tibboel's team found no analgesic effect from routine continuous intravenous morphine in premature babies on ventilator support, nor did they find a beneficial neurological

effect (*JAMA* 2003;290:2419-27).

There is confusion over the use of morphine in such babies. The team studied 150 preterm infants in neonatal intensive care units. The babies were randomised to placebo or morphine (100 micrograms per kg and 10 micrograms per hour, for up to seven days).

The pre-emptive morphine infusion did reduce intraventricular bleeding, but the authors suggest that morphine infusions should not be used in premature infants on ventilator support until more is known about the long term consequences. □

Painkillers before knee surgery improve outcomes

Janice Hopkins Tanne *New York*

Patients who had pre-emptive treatment of pain with a cyclooxygenase-2 (COX 2) inhibitor before knee replacement surgery had less pain and vomiting after the surgery, needed fewer opioids, slept better, had better knee flexion, and needed less physical therapy. The randomised controlled trial was reported by Dr Asokumar Buvanendran and colleagues at Rush-Presbyterian-St Luke's Medical Center, Chicago (*JAMA* 2003;290:2411-8).

Knee replacement is among the most painful of operations, Dr Buvanendran said. The 70 patients, aged 44 to 77 years,

were assigned to placebo treatment or to treatment with 50 mg of oral rofecoxib 24 hours before surgery, another 50 mg an hour or two before surgery, 50 mg daily for five days after surgery, and 25 mg daily for the next eight days. All patients were given epidural anaesthesia and self controlled analgesia after surgery.

Range of motion is an important outcome measure after knee replacement. The degree of knee flexion needed for climbing stairs is 83°, for descending stairs 90°, and for tying shoelaces 106°. At discharge, patients in the rofecoxib group had 84.2° of active flexion and patients in the placebo group had 73.2°. One month after surgery the figures were 109.3° and 100.8° respectively.

Dr Buvanendran said that the improvement meant an increase in satisfaction among patients, a quicker return to work or usual activities, and a reduction in the costs of physical therapy. □

Treatment for depression improves pain in elderly patients

Janice Hopkins Tanne *New York*

Treating elderly patients for their depression can reduce their pain and disability from arthritis, finds a randomised controlled trial of 1801 elderly patients with depression in five western US states (*JAMA* 2003; 290:2428-34).

Of the patients, who were aged over 60 years, 1001 (56%) had arthritis as well as depression. They were randomly assigned to usual care or to treatment for depression with antidepressants or six to eight sessions of psychotherapy.

At follow up at three, six, and 12 months all the patients who

were treated for depression said they had less functional impairment, better health, and better quality of life.

Dr Elizabeth Lin, of the Center for Health Studies at the Group Health Cooperative, Seattle, who coordinated the study, said that before entering the study "patients had significant pain, but few were taking pain medications or antidepressants."

"Depression treatment was doubly effective," she said. "Most patients had other chronic diseases, and depression leads to less compliance with medication regimens. Better depression care reduced depression, reduced pain severity, and increased the quality of life."

Treatment for depression probably also reduced healthcare costs, she said, because depressed people use health care 50% more than people without depression. □