Published in final edited form as:

Inorg Chem. 2009 September 7; 48(17): 8054–8056. doi:10.1021/ic900983v.

Water-Soluble Macrocycles Synthesized via the Weak-Link Approach

Michael J. Wiester and Chad A. Mirkin*

Department of Chemistry and the International Institute for Nanotechnology, Northwestern University, Evanston, Illinois 60208-3113

Abstract

We report a general, high-yielding method for the synthesis of water-soluble complexes, which is based upon the weak-link approach to supramolecular coordination chemistry. Specifically, we have utilized oligomeric ethylene glycol functional groups appended to the aryl groups of the diphenylphosphine moieties to achieve solubility. Small molecules or halide ions can be used to expand these complexes into larger more flexible macrocyclic structures. The realization of this approach should allow for the preparation of allosteric biomimetic structures which can be used in aqueous media.

For decades, chemists have been attempting to mimic the sophisticated structures and exquisite reactivity of compounds commonly found in Nature. To effectively realize the properties of such systems, and enzymes in particular, chemists need high yielding synthetic strategies that allow one to construct highly functional molecules: 1) with tailorable recognition and catalytic properties, 2) that are dynamic and conformationally addressable, and 3) that exhibit solubility in aqueous media.¹ Coordination chemistry provides a platform for synthesizing biomimetic supramolecular structures, whereby it is possible to assemble structures in a convergent manner and typically in high yields and with excellent selectivity.^{2,3} In this regard, structures that mimic the reactivity of enzymes have been prepared by a variety of different researchers.⁴ Some of these structures are highly charged and soluble in aqueous media, creating a hydrophobic reaction pocket where, in some instances, new or unusual reactivity is observed.⁵ For example, the Fujita group has shown that a cage-like complex in the shape of an octahedron can accelerate the Diels-Alder cyclization between anthracene and *N*-cyclohexylmaleimide, where the maleimide reacts at the terminal anthracene ring instead of the central one, as is typically seen.^{5a} In a second example, Raymond, Bergman, and coworkers have demonstrated that a reaction that is normally acid catalyzed, the hydrolysis of orthoformates, proceeds in the cavity of a tetrahedron-shaped complex in basic solutions.^{5b} It should be noted that both of these examples use rigid, non-conformationally addressable structures and there are relatively few coordination chemistry based synthetic strategies, which allow one to construct molecules that exhibit all of the aforementioned properties.

There are many powerful methods for constructing large biomimetic systems. Two of the most popular and extensively developed are the directional-bonding approach² and the weak-link approach (WLA).^{3,6} The former yields large rigid structures with well defined cavities while the latter yields two types of structures that can be chemically interconverted through small molecule reactions that occur at metal hinge sites (Figure 1). Therefore, the WLA allows one

*To whom correspondence should be addressed. chadnano@northwestern.edu..

Supporting Information Available: Experimental procedures for compounds 1-11. This material is available free of charge via the Internet at <http://pubs.acs.org>.

to fulfill two of the requirements for a biomimetic structure in that it yields complexes that can have tailorable pockets for both stoichiometric and catalytic reactions, and structures that can be chemically toggled between active and inactive conformations, much in the way a natural allosteric system works. Indeed, this approach has been used to prepare a vast array of supramolecular structures, including mimics of single and multieffector allosteric enzymes, ELISA, and PCR.⁷ However, thus far all structures made by the WLA are only soluble in organic solvents limiting studies of such systems to environments that are not relevant to the structures one is trying to mimic. Herein, we present a general, high-yielding synthetic methodology for the synthesis of a new type of water soluble hemilabile ligand that can be used in the WLA to rapidly prepare water-soluble macrocyclic coordination complexes from d⁸ transition metal centers.

There are a variety of ways of making ligands water soluble. They include the incorporation of charged groups, such as metal ions or sulfonate, carboxylate, or ammonium salts, or functional groups that can form hydrogen bonds with water, such as ethylene glycol.^{8,9} To minimize the challenge of ligand purification, we targeted hemilabile structures with non-ionic oligomeric ethylene glycol functional groups appended to aryl moieties. This approach has many benefits. First, all of the intermediates en route to the target ligand can be purified by column chromatography. Second, the metal complexes formed from such ligands typically are soluble in water as well as polar organic solvents such as CH₂Cl₂, THF, and MeOH. Third, the oligomeric ethylene glycol functionality does not compete with the ligating properties of the hemilabile ligand with respect to the transition metal centers of interest.

Water soluble hemilabile ligand **5** was synthesized in five steps in 61% yield starting with commercially available p-bromophenol and p-toluene sulfonic acid 2-[2-(2-methoxy-ethoxy)-ethoxy]-ethyl ester (Scheme 1). These starting reagents were converted to 1-bromo-4-(2-[2-(2-methoxy-ethoxy)-ethoxy]-ethoxy)-benzene **1** by refluxing in a DMF solution of K₂CO₃. Diethylphosphoramidous dichloride was used to prepare the Lewis acid (BH₃) complexed diarylphosphine with oligomeric ethylene glycol groups, **2**, in a one-pot procedure. The borane protecting group allows for purification under ambient conditions without the formation of phosphine oxide. The protecting group can be removed under mild conditions using piperazine.¹⁰ To achieve this deprotection, we used polymer-bound piperazine, which can be readily removed from the reaction mixture, after the reaction, by filtration. The deprotection reaction can be followed by ³¹P{¹H} NMR spectroscopy, with the resonance corresponding to **2** at -2.8 ppm decreasing with a concomitant increase in the intensity of the resonance for BH₃-free **3** at -44.4 ppm.¹⁰ Compound **3** was converted into the chloroethyl diarylphosphine **4** by lithiation followed by reaction with 1,2-dichloroethane. Like 1-chloro-2-diphenylphosphinoethane, compound **4** will react with benzenedithiol to form the targeted ligand **5**. Compounds **1-5** have been characterized by ¹H, ³¹P{¹H}, ¹³C{¹H}, and ¹¹B{¹H} NMR spectroscopy and high resolution mass spectrometry (see Supporting Information).

To probe the reactivity of ligand **5**, we synthesized a series of transition metal complexes **6**, **7**, and **8** from it and a variety of metal ions (Rh^I, Cu^I, and Pd^{II}, respectively, Scheme 2). These complexes were formed by adding one equivalent of **5** in CH₂Cl₂ to a CH₂Cl₂ solution of the appropriate metal ion precursor. All of the resulting binuclear complexes were isolated as viscous oils and were characterized by ¹H and ³¹P{¹H} NMR spectroscopy and high resolution mass spectrometry. The ³¹P{¹H} NMR spectrum of each complex in CD₂Cl₂ exhibits a single highly diagnostic resonance in the δ 61-64 range for **6** and **8** and near δ 0 for **7**, assigned to the corresponding highly symmetric rigid condensed macrocycle with a *cis*-phosphine coordination environment. Each of these resonances are within 2 ppm of model complexes without oligomeric ethylene glycol groups.^{11,12} All other data are fully consistent with the proposed structures **6-8**.

The most extensively characterized macrocycles made via the WLA are Rh^I complexes, and ³¹P{¹H} NMR spectroscopy is one of the most utilized tools for the characterization of these complexes. For example, complex **6** contains a doublet at δ 63.7 with a diagnostic J_{Rh-P} coupling constant of 161 Hz, which is indicative of a phosphine-thioether ligand chelated to the rhodium with the phosphines bound in a cis fashion.¹¹ The HRMS spectrum of **6** exhibits the expected M²⁺ ion peak at m/z 1494.6748. When CO is added to a solution of complex **6**, complex **9** is formed immediately and in quantitative yield, as seen in the ³¹P{¹H} NMR spectrum. Complex **9** exhibits the expected upfield shift in its ³¹P{¹H} NMR spectrum,^{>11} a doublet at 23.3 ppm with a J_{Rh-P} coupling constant of 110 Hz, which indicates the thioether ligand is no longer chelated to the rhodium and the complex contains a *trans*-P-Rh-P coordination environment.¹¹ Significantly, the opened macrocycle **9** is also soluble in aqueous media, despite its neutral charge, because of the presence of the oligomeric ethylene glycol moieties. All of the synthesized complexes exhibit ³¹P{¹H} NMR resonances that correspond well with analogous complexes without oligomeric ethylene glycol units (see Supporting Information).^{11,12}

Interestingly, complexes **6-8** exhibit solubility in water where small molecules or halide ions can be used to expand them into larger more flexible macrocyclic structures such as **9-11**. The type of small molecules and halide used for this ring expansion depend upon the transition metal center. The Pd^{II} complex **8** can be opened by the addition of CN⁻, the Cu^I complex **7** by the addition of a nitrogen donor such as pyridine, and the Rh^I complex **6** by the addition of Cl⁻ and CO (1 atm).⁶ All of the open complexes, except **10**, are uncharged and yet highly soluble in water (e.g. **11** in Scheme 2, inset).

In conclusion, we have described the synthesis of a new water-soluble hemilabile ligand **5**, suitable for use in the WLA. Using a variety of transition metal precursors, we have synthesized a series of condensed macrocyclic structures, which can be opened into flexible water-soluble macrocycles through ligand displacement reactions. These are the first water-soluble complexes made by the WLA, and their realization points towards routes to biomimetic structures that can function as novel catalysts and amplified chemical detection systems.⁷ Significantly, this strategy of appending oligomeric ethylene glycol groups to hemilabile phosphine containing ligands should be general enough to confer water solubility to many ligands previously used in the WLA.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We acknowledge the NSF, AFOSR, ARO, and DDRE for financial support of this research. C.A.M. is grateful for a NIH Director's Pioneer Award.

REFERENCES

- (1) (a). Breslow R. Acc. Chem. Res 1995;28:146–153. (b) Vriezema DM, Aragones MC, Elemans JAAW, Cornelissen JJLM, Rowan AE, Nolte RJM. Chem. Rev 2005;105:1445–1490. [PubMed: 15826017] (c) Das S, Brudvig GW, Crabtree RH. Chem. Commun 2008:413–424.
- (2). Fujita M, Tominaga M, Hori A, Therrien B. Acc. Chem. Res 2005;38:369–378. [PubMed: 15835883]
- (3) (a). Holliday BJ, Mirkin CA. Angew. Chem., Int. Ed 2001;40:2022–2043. (b) Oliveri CG, Ulmann PA, Wiester MJ, Mirkin CA. Acc. Chem. Res 2008;41:1618–1629. [PubMed: 18642933]
- (4) (a). Collman JP, Decreau RA. Chem. Commun 2008:5065–5076. (b) Que L, Tolman WB. Nature 2008;455:333–340. [PubMed: 18800132] (c) Hooley RJ, Rebek J Jr. Chem. Biol 2009;16:255–264. [PubMed: 19318207] (d) Yoshizawa M, Klosterman Jeremy K, Fujita M. Angew. Chem., Int. Ed

- 2009;48:3418–3438. (e) Amijs CHM, van Klink GPM, van Koten G. Dalton Trans 2006:308–327. [PubMed: 16365645] (f) Wurthner F, You C-C, Saha-Moller CR. Chem. Soc. Rev 2004;33:133–146. [PubMed: 15026818]
- (5) (a). Yoshizawa M, Tamura M, Fujita M. Science 2006;312:251–254. [PubMed: 16614218] (b) Pluth MD, Bergman RG, Raymond KN. Science 2007;316:85–88. [PubMed: 17412953]
- (6). Gianneschi NC, Masar MS III, Mirkin CA. Acc. Chem. Res 2005;38:825–837. [PubMed: 16285706]
- (7) (a). Gianneschi NC, Cho S-H, Nguyen ST, Mirkin CA. Angew. Chem., Int. Ed 2004;43:5503–5507. (b) Yoon HJ, Heo J, Mirkin CA. J. Am. Chem. Soc 2007;129:14182–14183. [PubMed: 17973396] (c) Oliveri CG, Gianneschi NC, Nguyen ST, Mirkin CA, Stern CL, Wawrzak Z, Pink M. J. Am. Chem. Soc 2006;128:16286–16296. [PubMed: 17165783] (d) Gianneschi NC, Bertin PA, Nguyen ST, Mirkin CA, Zakharov LN, Rheingold AL. J. Am. Chem. Soc 2003;125:10508–10509. [PubMed: 12940719] (e) Kuwabara J, Stern CL, Mirkin CA. J. Am. Chem. Soc 2007;129:10074–10075. [PubMed: 17665915] (f) Heo J, Mirkin CA. Angew. Chem., Int. Ed 2006;45:941–944.
- (8) (a). Kusakawa T, Nakai T, Okano T, Fujita M. Chem. Lett 2003;32:284–285. (b) Ito H, Kusakawa T, Fujita M. Chem. Lett 2000:598–599. (c) Yamaguchi T, Fujita M. Angew. Chem., Int. Ed 2008;47:2067–2069. (d) Fiedler D, Leung DH, Bergman RG, Raymond KN. J. Am. Chem. Soc 2004;126:3674–3675. [PubMed: 15038695] (e) Fiedler D, Pagliero D, Brumaghim JL, Bergman RG, Raymond KN. Inorg. Chem 2004;43:846–848. [PubMed: 14753801] (f) Leung DH, Fiedler D, Bergman RG, Raymond KN. Angew. Chem., Int. Ed 2004;43:963–966. (g) Brauer DJ, Hingst M, Kottsieper KW, Liek C, Nickel T, Tepper M, Stelzer O, Sheldrick WS. J. Organomet. Chem. 2002;645:14–26.
- (9) (a). Hong SH, Grubbs RH. J. Am. Chem. Soc 2006;128:3508–3509. [PubMed: 16536510] (b) Wu J, Li J, Kolb U, Mullen K. Chem. Commun 2006:48–50. (c) Kahn A, Muller S, Hecht S. Chem. Commun 2005:584–586.
- (10). Sayalero S, Pericàs M. Synlett 2006:2585–2588.
- (11) (a). Sanger AR. Can. J. Chem 1983;61:2214–2219. (b) Eisenberg AH, Dixon FM, Mirkin CA, Stern CL, Incarvito CD, Rheingold AL. Organometallics 2001;20:2052–2058. (c) Masar MS III, Mirkin CA, Stern CL, Zakharov LN, Rheingold AL. Inorg. Chem 2004;43:4693–4701. [PubMed: 15257598]
- (12). Masar MS III, Ovchinnikov MV, Mirkin CA, Zakharov LN, Rheingold AL. Inorg. Chem 2003;42:6851–6858. [PubMed: 14552636]

Figure 1.
General strategy of the WLA.

Scheme 1.

Synthesis of Water-Soluble Ligand **5**^a.

a (i) K_2CO_3 , DMF, reflux, 90%; (ii) a) $n\text{-BuLi}$, $\text{Cl}_2\text{P}(\text{NEt}_2)$, THF, -78°C ; b) HCl , THF, -78°C ; c) LiBH_4 , THF, 0°C , 77%; (iii) polymer-bound piperazine, THF, 60°C , >99%; (iv) $n\text{-BuLi}$, $\text{ClCH}_2\text{CH}_2\text{Cl}$, THF, -78°C , 95%; (v) 1,4-benzenedithiol, K_2CO_3 , CH_3CN , reflux, 95%.

^a (i) for **6**: [Rh(cyclooctene)₂Cl]₂, CH₂Cl₂; for **7**: Cu(CH₃CN)₄PF₆, CH₂Cl₂; for **8**: Pd(CH₃CN)₄(BF₄)₂, Acetone, CH₂Cl₂. (ii) for **9**: CO (1 atm), D₂O; for **10**: pyridine, D₂O; for **11**: KCN, D₂O. Inset: A photograph of a solution of **11** in D₂O.

Scheme 2.

Synthesis of Water-Soluble Complexes^a.

a (i) for **6**: [Rh(cyclooctene)₂Cl]₂, CH₂Cl₂; for **7**: Cu(CH₃CN)₄PF₆, CH₂Cl₂; for **8**: Pd(CH₃CN)₄(BF₄)₂, Acetone, CH₂Cl₂. (ii) for **9**: CO (1 atm), D₂O; for **10**: pyridine, D₂O; for **11**: KCN, D₂O. Inset: A photograph of a solution of **11** in D₂O.