Retention of Motility of *Treponema pallidum* (Nichols Virulent Strain) in an Anaerobic Cell Culture System and in a Cell-Free System

P. L. SANDOK,* H. M. JENKIN, S. R. GRAVES, AND S. T. KNIGHT

The Hormel Institute, University of Minnesota, Austin, Minnesota 55912

Received for publication 11 July 1975

Optimum parameters for retention of motility of *Treponema pallidum* (Nichols virulent strain) were found by anaerobic co-incubation of the treponeme with rat glial cells and anaerobic incubation in spent medium obtained from glial cells originally grown aerobically.

The successful in vitro cultivation of virulent *Treponema pallidum* will probably follow the discovery of environmental parameters that extend its survival time during incubation. To date, the short-term survival of the virulent treponeme has been accomplished in various complex artificial media (3, 7, 8) and in attempts to grow it in cell cultures (2, 9, 10). However, since the microorganism has yet to be cultivated in vitro, the investigation of cell culture systems for this purpose remains a viable alternative to studies based solely on the use of artificial media. In addition, recent work by Sykes and Miller (5), Sykes et al. (6), Ovcinnikov and Delektorskij (4), and Fitzgerald et al. (2) on the detection of intracellular treponemes suggests that a utilizable cell culture system for treponemal co-incubation could provide valuable insights into the mechanism(s) of host-parasite interaction (10). To date there are no reports of a universally acceptable system for the co-incubation or cultivation of *T. pallidum* in cell cultures. Consequently, we have sought to develop such a system.

Modified Eagle medium (prereduced Eagle minimal essential medium [1] plus 10% newborn calf serum, PRMEM₁₀) was prepared as described in Table 1. This basal medium was diluted with unreduced Eagle medium plus 10% newborn calf serum (MEM₁₀) to obtain a gradual variation in redox potential from -408 to -232 mV E_{ca1} (Tables 2 and 3). Although rat glial cells (kindly supplied by W. Bondareff) exhibited no growth under anaerobic conditions after 48 h at 33.5 C (Table 2), cytopathological effects were minimal or undetectable. The cells could be repressed in aerobic medium after at least 96 h of anaerobic cultivation in various dilutions of PRMEM₁₀.

The glial cell cultures co-incubated with *T.*

pallidum were maintained under deoxygenated N₂ in sealed (air-tight) Leighton tubes containing removable cover slips for subsequent dark-field microscopic examination. New Zealand rabbits (6 to 7 lb [2,721 to 3,175 g]) inoculated with 10⁷ to 3 × 10⁷ microorganisms/testes were killed 8 to 9 days later after observing the initiation of an orchitis. When freshly harvested virulent treponemes were inoculated into anaerobic glial cell cultures (1 to 5 treponemes/glial cell) and incubated 24 h at 33.5 C, motile treponemes were detected, both in the supernatant fluid (at about 1/5 to 1/10 the initial concentration) and attached to the cells (Table 3). The treponemes could not be dislodged by agitation with a Vortex tube mixer, and they appeared to have become elongated, compared with unattached nonmotile microorganisms, without any evidence for division. In our earlier experience media having a redox potential of -250 ± 50 mV E_{ca1} appeared to be optimal for this system (Tables 2 and 3). The longevity of motility of *T. pallidum* has been extended well beyond the time reported by Fitzgerald et al. (2).

In another series of experiments, unreduced MEM₁₀ was used to cultivate approximately 3.0 × 10⁷ rat glial cells in 32-oz (0.946-liter) bottles for 48 h at 33.5 C. The supernatant fluid was removed and diluted in PRMEM₁₀ (pH 7.4 to 7.6) under deoxygenated N₂ or 50% N₂ plus 50% H₂. The controls consisted of fresh MEM₁₀ diluted in PRMEM₁₀, as described above. Freshly harvested treponemes (0.3 ml or approximately 10⁷ to 3 × 10⁷) were inoculated into each tube containing 10 ml of medium. Percentage of motility was estimated at 24-h intervals by counting 50 to 100 random fields or about 30 to 70 treponemes from triplicate tubes under dark-field microscopy using a 40× objective.

TABLE 1. Preparation and components of PRMEM₁₀

Procedure	Solution designation	Components	Final concn (per liter)
1. Prereduce the salt solution ^a by heating in autoclave for 8 min at 121 C, at slow exhaust. Cool to room temperature under deoxygenated N ₂ .	Salt solution	NaCl	5,440 mg
		KCl	320 mg
		MgSO ₄ ·7H ₂ O	140 mg
		KH ₂ PO ₄	48 mg
		Na ₂ HPO ₄	48 mg
		Glucose	800 mg
		CaCl ₂ ·2H ₂ O	112 mg
		Resazurin	1 mg
2. Add the reducing agents in powder form under N ₂ . Adjust pH to 7.8 to 7.9 and sterilize in stoppered tubes or flask at 121 C for 25 min. ^b Cool to room temperature.	Reducing agents	Reduced glutathione	1,200 mg
		L-Cysteine	800 mg
		Sodium thioglycolate	700 mg
3. Mix appropriate volumes of the serum solution ^c with the reduced salts solution to obtain a final concentration of 10% serum plus 1× concentration of all the other nutrients listed.	Serum solution	50× MEM essential amino acids	20 ml
		100× MEM vitamin mixture	10 ml
		200 mM glutamine	10 ml
		1.4% NaHCO ₃	20 ml
		0.2 N NaOH	20 to 30 ml
		Newborn calf serum	100 ml
4. Mix appropriate proportion of unreduced MEM ₁₀ with reduced MEM ₁₀ (Tables 2 and 3) under deoxygenated N ₂ to obtain media at various redox potentials.			

^a The sodium, potassium, and magnesium salts, the sodium and potassium phosphate buffers containing glucose, the calcium chloride, and the resazurin solutions were prepared separately at 10× concentration prior to mixing the final solution.

^b The pH was adjusted with about 0.5 ml of 10 N NaOH, and the stoppered tubes and Pyrex flasks were sterilized in an autoclave press (Bellco, Vineland, N.J.).

^c The serum solution was adjusted with 0.2 N NaOH to pH 7.8 prior to mixing with the reduced salts to obtain a final pH of 7.4 to 7.6.

Results from our earlier preliminary work using the protocol above suggested that the optimal conditions for the long-term retention of treponemal motility in spent medium were: (i) a redox potential of -250 ± 50 mV E_{ca} (i.e., the medium mixture of one part of PRMEM₁₀ to nine parts of spent medium); (ii) a pH range of 7.3 to 7.6; (iii) a deoxygenated atmosphere of 50% N₂ plus 50% H₂; and (iv) an incubation temperature of 33 ± 1 C. The data in Table 4 are typical of results obtained from a minimum of six independent experiments performed under optimal conditions. The estimated time at which motility drops to 50% for PRMEM₁₀ diluted with the supernatant fluid from glial cells is about the same under both N₂ and N₂ plus H₂ (i.e., 50 to 60 h) (Table 4). However, the estimated time to reach 10% motility under N₂ plus H₂ is longer than that obtained under N₂ alone. The 10 and 50% motilities obtained

for the controls (i.e., mixtures of fresh MEM₁₀ and PRMEM₁₀) are considerably shorter (i.e., 15 to 20 h). Thus, the supernatant fluid appears to contain some factor(s) not available in fresh MEM₁₀ that enhances the retention of treponemal motility.

In summary, we have presented evidence that the retention of motility of *T. pallidum* in cells and cell-free supernatant fluid is greatly enhanced under anaerobic conditions in appropriately reduced media. This evidence is in sharp contrast to previous reports using tissue culture systems for the co-incubation of *T. pallidum*, in which no attempt was made to control the redox potential (9, 10) and/or strict anaerobiosis was not maintained during incubation (2, 9). We are now in a position to perform a series of metabolic experiments which shall be less affected by rapid decreases in *T. pallidum* viability in cell-free and in cell cul-

TABLE 2. Rat glial cell cultivation under anaerobic conditions using various dilutions of PRMEM₁₀

Mixture of PRMEM ₁₀ /MEM ₁₀ (%)	<i>E</i> _{cal} ^a (mV) at pH 7.6	Cell count ^b (× 10 ⁵)
100/0	-408	1.61 ± 0.43
70/30	-380	2.37 ± 0.32
30/70	-334	3.40 ± 0.32
10/90	-232	2.07 ± 0.28
0/100	+48	3.87 ± 0.43
Aerobic control	+48	5.23 ± 0.87

^a Redox potentials were obtained at room temperature with a saturated KCl combination calomel-platinum electrode (Orion model 96-78, Orion Research, Cambridge, Mass.).

^b Initial inoculum was 2.2×10^5 rat glial cells per Leighton tube in triplicate samples. The cells were cultivated 48 h at 33.5 C under deoxygenated N₂ except for the aerobic control, which was cultivated in air. Counts were performed on a Coulter counter after the cells were removed from the glass surfaces by a solution containing 0.05% trypsin and 0.05% ethylenediaminetetraacetic acid.

TABLE 3. Motility retention of virulent *T. pallidum* co-incubated with rat glial cells^a

Mixture of PRMEM ₁₀ /MEM ₁₀ ^b (%)	Motility (%) ^c	
	Supernatant fluids	Attached treponemes
100/0	0	0
70/30	45 ± 17	92.8
30/70	77 ± 8	95.0
10/90	81 ± 10	100.0
0/100	0	0

^a The cultures were incubated for 24 h under deoxygenated N₂ at 33.5 C.

^b The redox potentials for each mixture of PRMEM₁₀/MEM₁₀ are given in Table 2.

^c Percentage of motility was determined by counting 50 to 100 treponemes from Leighton tubes with cover slips containing approximately 2.2×10^5 rat glial cells in 1 ml of medium inoculated with 3×10^5 to 10×10^5 treponemes in 0.05 ml. Cover slips with attached glial cells were placed face down on glass slides and inspected under dark field microscopy using a 40× objective lens.

ture systems. In addition, a useful by-product of the present report is that it may be possible to ship fresh (unfrozen) human isolates in the cell-free medium from many parts of the world for immediate laboratory use.

We thank D. L. Harris, College of Veterinary Medicine, Iowa State University, Ames, for his helpful suggestions during these experiments. We wish to thank Jeff Robbins and Gregg Jorgenson for their expert technical assistance.

TABLE 4. Motility retention of virulent *T. pallidum* in reduced fresh Eagle medium and in reduced supernatant fluids from rat glial cells^a diluted with PRMEM₁₀

Culture medium ^b	M ₅₀ ^c		M ₁₀ ^c	
	N ₂	N ₂ + H ₂	N ₂	N ₂ + H ₂
Fresh MEM ₁₀	16.5	18.0	41.5	48.0
Supernatant fluids	51.5	57.8	71.5	119.0

^a Rat glial cells (10^7 to 3×10^7) were cultivated for 48 h at 33.5 C aerobically in MEM₁₀, and then the supernatant fluid was collected and mixed with PRMEM₁₀, as described in the text and in footnote b below.

^b Nine parts of supernatant fluid and nine parts of fresh MEM₁₀ were each diluted with one part of PRMEM₁₀ by volume to obtain a final *E*_{cal} of -250 mV at pH 7.6 at room temperature.

^c The estimation of the time in which the motility of the treponemes dropped to 50% (M₅₀) and to 10% (M₁₀), respectively, was obtained from triplicate samples of a plot of percentage of motility versus time. The cultures were incubated at 33.5 C.

This investigation was supported by Public Health Service contract NO1-AI-42537, S. A. 1 from the National Institute of Allergy and Infectious Diseases and by The Hormel Foundation, Austin, Minn.

LITERATURE CITED

1. Eagle, H. 1959. Amino acid metabolism in mammalian cell cultures. *Science* 130:432-437.
2. Fitzgerald, T. J., J. N. Miller, and J. A. Sykes. 1975. *Treponema pallidum* (Nichols strain) in tissue cultures: cellular attachment, entry, and survival. *Infect. Immun.* 11:1133-1140.
3. Nelson, R. A., and J. A. Diesendruck. 1951. Studies on treponemal immobilizing antibodies in syphilis. I. Technique of measurement of factors influencing immobilization. *J. Immunol.* 66:667-685.
4. Ovcinnikov, N. M., and V. V. Delektorskij. 1975. *Treponema pallidum* in nerve fibers. *Br. J. Vener. Dis.* 51:10-14.
5. Sykes, J. A., and J. N. Miller. 1971. Intracellular location of *Treponema pallidum* (Nichols strain) in the rabbit testes. *Infect. Immun.* 4:307-314.
6. Sykes, J. A., J. N. Miller, and A. J. Kalan. 1974. *Treponema pallidum* within cells of a primary chancre from a human female. *Br. J. Vener. Dis.* 50:40-44.
7. Weber, M. M. 1960. Factors influencing the *in vitro* survival of *Treponema pallidum*. *Am. J. Hyg.* 71:410-417.
8. Willcox, R. R., and J. Guthe. 1966. *Treponema pallidum*. A bibliographical review of the morphology, culture and survival of *T. pallidum* and associated organisms. W.H.O. Monogr. Ser. 35(Suppl.):35-90.
9. Wright, M. I. 1962. Exploratory studies in tissue culture of *T. pallidum*, p. 884-887. In D. M. Pillsbury and C. S. Livingood (ed.), *Proc. Int. Congr. Dermatol.*, 12th, vol. 2. Excerpta Medica Foundation, Amsterdam.
10. World Health Organization. 1970. Treponematoses research, p. 1-91. Technical Report Series no. 455. World Health Organization, Geneva.