Published in final edited form as:

Pers Individ Dif. 2009 December 1; 47(8): 812–816. doi:10.1016/j.paid.2009.06.019.

Mapping Gray's BIS and BAS Constructs onto Factor 1 and Factor 2 of Hare's Psychopathy Checklist - Revised

John F. Wallace, Melanie B. Malterer, and Joseph P. Newman

University of Wisconsin - Madison

Abstract

Reinforcement Sensitivity Theory (RST; Gray, 1987; Gray & McNaughton, 2000) has proven to be a valuable tool for understanding psychopathy (e.g., Fowles, 1980, 1988; Newman & Malterer, 2009; Poythress, Edens, Landfield, Lilienfeld, Skeem, & Douglas, 2008). Recent research has linked two RST constructs, the Behavioral Inhibition System (BIS) and the Behavioral Activation System (BAS), to individuals with primary psychopathy and secondary psychopathy (Lykken, 1995; Newman, MacCoun, Vaughn, & Sadeh, 2005): Primary psychopaths manifest low BIS reactivity and secondary psychopaths manifest high BAS reactivity. In the present study, we examine the relationships between the BIS/BAS constructs and Factors 1 and 2 of the Psychopathy Checklist – Revised (PCL-R) in a sample of 472 incarcerated male offenders. Paralleling their relationships with primary and secondary psychopathy, the BIS/BAS constructs were differentially related to the two PCL-R factors. Specifically, the influence of the BIS was found to be more prominent than the influence of the BAS for Factor 1, and the influence of the BAS was more prominent than that of the BIS for Factor 2.

Keywords

Psychopathy; Reinforcement Sensitivity Theory; BIS; BAS; Psychopathy Checklist – Revised; PCL-R Factors

Introduction

Psychopathic individuals are known for their exceptionally poor judgment, penchant for acting in response to relatively low levels of motivation (i.e., on a whim), inability to learn from negative consequences, extreme egocentricity, and lack of remorse for misdeeds (Cleckley, 1976). Hence, these individuals are prone to engage in impulsive, maladaptive, and antisocial acts, which cause them to experience substantial marital, academic, and employment

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

¹Although the BAS subscales were not a focus of this study, we provide the relevant zero-order correlations involving the BAS subscales for those having an interest in this issue: PCL-R Factor 2 was significantly associated with BAS drive ($r = .216, p < .001$) and BAS fun seeking ($r = .246, p < .001$), but not BAS reward reactivity ($r = .057, p = .217$). BAS drive was also significantly associated with PCL-R Factor 1 ($r = .124, p < .01$), but this correlation was smaller than that with PCL-R Factor 2.

²Although the results obtained by Newman et al. (2005) and those of the present study are similar, the data do not indicate that the primary/secondary psychopathy distinction and the PCL-R factors reference identical constructs (e.g., Newman, 2009).

³Results obtained using other measures of the psychopathy construct (e.g., Uzieblo, Verschuere, & Crombez, 2007), particularly those that employ different assessment methods such as self-report questionnaires, may not be entirely comparable to those obtained using the PCL-R (which uses a semi-structured interview and a review of participants' institutional files to assign designations to study participants).

disruption. Furthermore, although the prevalence of psychopathy in the general population is only 1%, estimates of the prevalence among incarcerated offenders range from 15% to 25% (Hare, 1996), and psychopathic individuals re-offend two to five times more frequently than do nonpsychopathic individuals (Hemphill, Hare, & Wong, 1998; Quinsey, Rice, & Harris 1995; Serin, 1996, Walters, 2003).

The use of Gray's reinforcement sensitivity theory (RST; 1975, 1987; Gray & Smith, 1969) to explain psychopathic behavior dates back nearly 30 years (e.g., Fowles, 1980). Briefly, the most recent formulation of RST (Gray & McNaughton, 2000) postulates three interacting brain systems. The Fight-Flight-Freeze System (FFFS) is activated in the presence of both conditioned and unconditioned aversive stimuli. Its activation is associated with the emotion of fear and with behaviors that diminish the perceived threat (e.g., escape or avoidance). The Behavioral Activation System (BAS) is activated in the presence of both conditioned and unconditioned appetitive stimuli (e.g., opportunities for reward). Increased BAS activity is associated with positive emotional states (e.g., optimism, hopefulness) and appetitive (e.g., approach) behavior, as well as with aggression (e.g. Carver 2004; Harmon-Jones & Peterson, 2008). Finally, the Behavioral Inhibition System (BIS) is activated when conflicts occur between concurrent goals (e.g., the classic approach-avoidance conflict); specifically, when both the BAS and the FFFS are activated (e.g., Corr, 2006). Increased activity of the BIS is associated with the emotion of anxiety, and causes the inhibition of ongoing behaviors and the initiation of information processing to resolve the conflict.

Fowles (1980) proposed one of the initial explanations of psychopathy that made use of the BIS and BAS constructs, attributing psychopathy to a weak or hypo-reactive BIS (which leads to lower than normal levels of BIS activity in response to stimuli to which the BIS is sensitive). More recently, Lykken (1995) expanded upon Fowles' use of RST by focusing on the distinction between primary and secondary psychopathy (e.g., Blackburn, 1979; Hare, 1970; Skeem, Johansson, Andershed, Kerr, & Loudon, 2007). Primary psychopaths are considered to be true or prototypical psychopaths (Cleckley, 1976). They are characterized by relatively low levels of anxiety, and their maladaptive, antisocial actions are considered to be the result, at least in part, of the failure to experience anxiety even when the likelihood of adverse consequences is high (e.g., Cleckley, 1976). On the other hand, secondary psychopaths experience higher-than-normal levels of negative emotions (e.g., anxiety), and their antisocial actions are thought to occur in response to intra-psychic conflicts or emotional distress (e.g., Cleckley, 1976).

Lykken (1995) proposed that primary psychopathy is associated with weak BIS reactivity (but normal BAS reactivity) and, in consequence, primary psychopaths are prone to experience smaller than normal reactions to BIS inputs. Conversely, secondary psychopathy is associated with strong BAS reactivity (but normal BIS reactivity); that is, secondary psychopaths are hyperreactive to opportunities for reward. To test Lykken's hypotheses, Newman et al. (2005) administered two measures of the BIS and BAS constructs--the BIS/BAS Scales (Carver & White, 1994) and the Sensitivity to Punishment and Sensitivity to Reward Questionnaire (Torrubia, Ávila, Moltó, & Caseras, 2001)--to a sample of incarcerated male offenders (and these same measures also were employed in the present study--see below). Consistent with Lykken's hypotheses, for individuals with primary psychopathy, the influence of low BIS reactivity predominated, whereas for those with secondary psychopathy, the influence of high BAS reactivity was most prominent.

Another way of conceptualizing the psychopathy construct (and thus the relationships between psychopathy and RST) involves the Psychopathy Checklist - Revised (PCL-R; Hare, 2003), which is the principal instrument used to identify psychopathic individuals in correctional settings. Of particular relevance to the present manuscript, factor-analytic studies have

identified two broad PCL-R factors with distinct personality, behavioral, physiological, and social correlates (e.g., Harpur, Hare, & Hakstian, 1989; Patrick, Bradley, & Lang, 1993; Ross et al., 2007). One of these factors (Factor 1), is hypothesized to reflect psychopathy's distinguishing characteristics, such as low anxiety and deficient emotion processing (e.g., Patrick, 1994; Patrick et al., 1993), whereas the second factor (Factor 2) relates to more general externalizing tendencies that are shared by persons having diagnoses such as Conduct Disorder and Antisocial Personality Disorder (Patrick, Hicks, Krueger, & Lang, 2005). Psychopathic individuals usually score above the mean on both PCL-R Factor 1 and Factor 2, although, for a particular psychopathic individual, the score on one factor may be substantially higher than the score on the other.

In this study, we examined the relationships among PCL-R Factors 1 and 2 and the BIS and BAS constructs in a large sample of incarcerated male offenders to determine whether the RST constructs are differentially related to the two principal factors derived from the PCL-R. In past research, inmates with high PCL-R scores (i.e., psychopathic individuals) displayed both lower BIS and higher BAS scores than non-psychopathic control participants (Book and Quinsey, 2004). However, paralleling results for primary and secondary psychopathy, such findings may mask the unique contributions of the two PCL-R Factors. To the extent that Factor 1, like primary psychopathy, is associated with low anxiety, we predicted that Factor 1 would be associated predominantly with low scores on measures of the BIS construct. Conversely, to the extent that Factor 2, like secondary psychopathy, is associated with general externalizing tendencies, we predicted that Factor 2 would be associated predominantly with high scores on measures of the BAS construct.

In addition to the two-factor model upon which we based our predictions, a four-factor model for the PCL-R has been proposed (see, for example, Neumann, Hare, & Newman, 2007). Hence, although comparing and contrasting the two- and four-factor models was not an aim of this study, we conducted post-hoc analyses to determine whether the two-factor results (see below) were moderated by underlying correlations involving the four-factor model.

Method

Participants

The sample consisted of 472 adult male offenders (13.8% African-American, 85.4% Caucasian) incarcerated in the Wisconsin prison system, ranging in age from 18 – 45 years of age ($M = 31.66$ years, $SD = 7.38$). Researchers selected potential participants from a comprehensive prison roster and screened out anyone older than 45 years, with diagnoses of Bipolar Disorder or Schizophrenia, or receiving psychotropic medication. Only participants for whom all relevant data points were available were included in this study. Hence, all correlations and analyses reported herein pertain to the same 472 participants.

Measures

PCL-R—The PCL-R (Hare, 1991, 2003) consists of 20 items indicative of psychopathic traits as delineated by Cleckley (1976). Each PCL-R item is scored on a 3-point scale (0=clearly not present; 1=may be present; 2 = clearly present), and the scores are summed to yield a total score (range 0 – 40). Scores are based on information derived from both a semi-structured interview and a review of institutional file information. The reliability and construct validity of the PCL-R are well established (see Hare, 1985; 1996; Hare, Harpur, Hakstian, Forth, Hart, & Newman, 1990). Early work with the Psychopathy Checklist revealed a replicable two-factor structure (Harpur et al., 1989) with Factor 1 assessing interpersonal and affective characteristics (e.g., pathological lying, shallow affect) and Factor 2 assessing impulsive and antisocial

behavior. The internal consistency of the PCL-R Factors, as indexed by Cronbach's alpha, is reported in Table 1.

Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ)—

The SPSRQ (Torrubia, Ávila, Moltó, & Caseras, 2001) is a 48-item self-report measure consisting of two scales: Sensitivity to Punishment (SP; 24 items) and Sensitivity to Reward (SR; 24 items). Each item is answered dichotomously (*yes* or *no*). The SPSRQ has shown good reliability and construct validity (Ávila & Parcet, 2000, 2001; Caseras, Ávila, & Torrubia, 2003). The SP scale measures individual differences in BIS functioning, including a) behavioral inhibition in response to novelty or aversive consequences, and b) cognitive processes or worry in response to failure or punishment cues. The SR scale measures individual differences in BAS functioning, including sensation seeking and approach behaviors in situations involving possible rewards. See Table 1 for the internal consistency of the SP/SR Scale scores.

BIS/BAS Scales (BIS/BAS)—The BIS/BAS (Carver & White, 1994) is a 20-item self-report questionnaire consisting of four scales: BIS (7 items), BAS—Reward Responsiveness (5 items), BAS—Drive (4 items), and BAS—Fun Seeking (4 items). Each item is rated on a 4-point scale ranging from 1 (*strongly agree*) to 4 (*strongly disagree*). The BIS scale measures sensitivity to punishment or aversive stimuli. The three BAS scales (Reward Responsiveness, Drive, and Fun Seeking) measure sensitivity or positive responses to anticipated rewards, persistent pursuit and motivation to achieve desired goals, and willingness to approach new reward stimuli and to do so on the spur of a moment, respectively. Because previous research (e.g., Jorm, Christensen, Henderson, Jacomb, Korten, & Rodgers, 1999) suggests that the three BAS scales strongly load on a second-order (BAS) factor, we used the total BAS score for our analyses. Although the use of the BAS subscales, rather than the BAS Total Score, recently has been advocated (e.g., Heym, Ferguson, & Lawrence, 2008), in our sample the internal consistency of the BAS (i.e., the items comprising the BAS Total Score) exceeded not only that of the BIS, but also those of both the SPSPQ scales (SP and SR). See Table 1 for the internal consistency of the BIS/BAS Scale scores.

We should also note here that the use of the BIS/BAS with samples of incarcerated offenders, as well as its validity as a measure of the RST constructs, have both been questioned (e.g., Poythress, Edens, Landfield, Lilienfeld, Skeem, & Douglas, 2008; Poythress, Skeem, Lilienfeld, Douglas, & Edens, 2009) and defended (e.g., Newman & Malterer, 2009). In any event, as described below, neither the BIS/BAS nor the SPSRQ is used as a stand-alone measure of the RST construct in this study.

Procedure

All participants included in these analyses were informed of the components of the study, potential risks and benefits, and procedures for safeguarding information. Consent information was presented to participants both orally and in written form, and they were informed that their participation in the study would have no impact on their correctional status.

After providing informed consent, each participant was interviewed and assigned a PCL-R score based on information obtained during the interview and a post-interview review of his prison files (see Hare, 2003). In addition, each participant completed the BIS/BAS (Carver & White, 1994) and SPSRQ (Torrubia et al., 2001)—the same two measures of the BAS and BIS constructs that were used in the Newman et al. (2005) study. Hence, each participant generated two scores reflecting his status with respect to the BAS construct (BAS and SR) and two scores pertaining to the BIS construct (BIS and SP).

Results

Because the correlations between the corresponding components of the BIS/BAS and SPSRQ measures are moderate to high (see Table 1), and we had no *a priori* grounds to consider one measure or the other as a better operationalization of the RST constructs, we created composite variables using participants' scores from both the BIS/BAS and SPSRQ. Specifically, participants' raw scores on each scale were converted to z-scores. The two z-scores reflecting BIS functioning (i.e., BIS and SP) were then averaged, creating the composite BIS variable – BIS/SP. Likewise, the two z-scores pertaining to the BAS (i.e., BAS and SR) were averaged to create the BAS composite – BAS/SR. Using those two composite variables, we performed two correlational analyses, with the BIS composite as the criterion variable for one analysis and the BAS composite as the criterion variable for the other, and with PCL-R Factor 1 and Factor 2 as predictor variables in both analyses. Zero-order correlations for these and other variables are reported in Table 1.

For the BIS analysis, consistent with prediction, BIS/SP and PCL-R Factor 1 were significantly associated ($r = -.242, p < .001$), and this correlation was larger than the zero-order correlation between BIS/SP and Factor 2 ($r = -.173, p < .01$). To determine whether this difference was statistically significant, the correlations were transformed to z-scores and compared using the procedure described by Papoulis (1990). All subsequent comparisons of correlation magnitudes were computed using this method. In this case, the correlations of BIS/SP with Factor 1 and Factor 2 did not, in fact, differ significantly in magnitude.

To provide a more specific assessment of the association of the BIS construct with PCL-R Factor 1, we examined whether the association between BIS/SP and Factor 1 scores remained significant after removing the contribution of Factor 2. Even when the variance associated with PCL-R Factor 2 was removed using partial correlation, the association between BIS/SP and Factor 1 remained significant ($r_{\text{partial}} = -.175, p < .001$). Conversely, the previously significant correlation of BIS/SP with Factor 2 was rendered non-significant when controlling for Factor 1 ($r_{\text{partial}} = -.034, p > .6$). The difference between these two partial correlations (i.e., $-.175$ and $-.034$) was statistically significant, $p < .05$, indicating that the BIS composite was more strongly associated with PCL-R Factor 1 than with Factor 2.

For the BAS analysis, as predicted, BAS/SR was significantly associated with PCL-R Factor 2 ($r = .291, p < .001$). Moreover, this association was significantly larger in magnitude ($p < .01$) than the corresponding correlation with PCL-R Factor 1 ($r = .130, p < .05$)¹. As above, we also examined whether the association between BAS/SR and PCL-R Factor 2 scores retained its statistical significance after removing the contribution of PCL-R Factor 1 to this association. Even when the variance associated with PCL-R Factor 1 was removed using partial correlation analysis, the association of the BAS composite with Factor 2 remained statistically significant ($r_{\text{partial}} = .269, p < .001$). On the other hand, when we repeated this analysis partialling the contribution of Factor 2 from Factor 1, the previously significant association between the BAS/SR and PCL-R Factor 1 was no longer statistically significant ($r_{\text{partial}} = -.061, p > .3$). The difference between these two partial correlations (.269 and $-.061$) was statistically significant, $p < .001$, indicating that the BAS composite was more strongly associated with PCL-R Factor 2 than with Factor 1.

Supplementary Analyses

The PCL-R total score (PCL-R Total) was significantly associated with both BIS/SP ($r = -.229, p < .001$) and BAS/SR ($r = .245, p < .001$). To address the extent to which the association between psychopathy and the BIS/BAS measures was due to the unique contributions of the PCL-R factors, we next investigated the association between PCL-R Total and the BIS/BAS constructs while partialling the effects of the PCL-R factors.

We first examined whether the association between PCL-R Total and BIS/SP scores remained significant after removing the contribution of PCL-R Factor 1 scores to this association. Once the variance associated with PCL-R Factor 1 was removed using partial correlation analysis, the previously significant association between the PCL-R total scores and BIS/SP was no longer significant ($r_{\text{partial}} = -.033, p > .4$). On the other hand, after removing the variance due to PCL-R Factor 2 scores, the association between PCL-R Total and BIS/SP scores remained significant ($r_{\text{partial}} = -.165, p < .001$).

Parallel analyses were performed to test whether the BAS/SR association with PCL-R Total was due primarily to the specific contribution of PCL-R Factor 2. After the variance associated with PCL-R Factor 2 scores was partialled, the previously significant association between PCL-R Total and BAS/SR was no longer significant ($r_{\text{partial}} = -.037, p > .4$). Conversely, after the variance associated with PCL-R Factor 1 was removed, the association between the PCL-R Total and BAS/SR remained significant ($r_{\text{partial}} = .273, p < .001$).

These analyses suggest that the association of psychopathy as indexed by PCL-R total scores with both low BIS reactivity and high BAS reactivity (e.g., Book and Quinsey, 2004) may be due largely to the specific contributions of Factors 1 and 2: The association of psychopathy with low BIS reactivity is due largely to the contribution of Factor 1, and the association of psychopathy with high BAS reactivity is due largely to the contribution of Factor 2.

Finally, a four-factor model for the PCL-R also has been described (e.g., Neumann, Hare, & Newman, 2007), two factors of which (the Interpersonal and Affective factors) are associated with PCL-R Factor 1, and two of which (the Lifestyle and Antisocial factors) are associated with PCL-R Factor 2. Hence, we examined whether 1) the results pertaining to Factor 1 were moderated by underlying correlations involving the Interpersonal and Affective factors, and 2) the results pertaining to Factor 2 were moderated by underlying correlations involving the Lifestyle and Antisocial factors.

None of the differences between the relevant pairs of correlations approached statistical significance. The zero-order correlations of BIS/SP with the Interpersonal and Affective factors were $-.202$ and $-.237$, respectively ($p > 0.5$), and the correlations with the Lifestyle and Antisocial factors were $-.155$ and $-.160$, respectively ($p > 0.9$). The correlations of BAS/SR with the Interpersonal and Affective factors were $.143$ and $.078$, respectively ($p > 0.3$), and the correlations with the Lifestyle and Antisocial factors were $.289$ and $.227$, respectively ($p > 0.3$). Because none of the differences between any of these four pairs of correlations approached statistical significance, the results pertaining to Factors 1 and 2 reported above do not appear to be moderated by differential associations of the BIS and BAS measures with the constituent factors of the four factor model of the PCL-R.

Discussion

The present study examined the associations between the BIS and BAS constructs and PCL-R Factors 1 and 2. As predicted, Factor 1 was significantly associated with the BIS composite and Factor 2 was significantly associated with the BAS composite. Moreover, supporting the specificity of the hypothesized relationships, the partial correlations between Factor 1 and BAS, and between Factor 2 and BIS, were low and non-significant ($r'_{\text{partial}} = -.061$ and $-.034$, respectively) after controlling for the effects of the other factor. Thus, just as primary psychopathy is associated predominantly with a weak BIS and secondary psychopathy is associated predominantly with a strong BAS (Newman et al., 2005), the influence of the BIS is significantly more prominent than the influence of the BAS for PCL-R Factor 1, and the influence of the BAS is significantly more prominent than that of the BIS for Factor 2.² That

is, Factor 1 is associated with low BIS reactivity and Factor 2 is associated with high BAS reactivity.³

These findings also complement earlier reports (e.g., Patrick, 2007; Patrick et al., 2005) that PCL-R Factor 1 is associated with low anxiety and other affective deficits (e.g., weak fear-potentiated startle), which plausibly reflect, at least in part, individual differences in BIS functioning. Conversely, PCL-R Factor 2 is associated with impulsivity and disinhibited approach behavior, which plausibly reflect individual differences in BAS-mediated processes.

Finally, we conducted supplementary analyses to examine whether the results involving the two-factor model were moderated by underlying correlations involving the four-factor model. Because these analyses were of a post-hoc nature, we are hesitant to make general statements regarding the relative validities of the two models. Nonetheless, at least for the present data set, the four-factor model does not appear to add appreciably to the explanatory power of the two-factor variant.

We conclude by concurring with Lilienfeld (1994) and others (e.g., Frick, Lilienfeld, Ellis, Loney, & Silverthorn, 1999; Harpur et al., 1989; Patrick et al., 2005; Sellbom & Verona, 2007) that the explication of the independent contributions of psychopathy's major factors advances the understanding of psychopathy as a whole. Given the prominence of RST in theoretical formulations of psychopathy, as well as the burgeoning literature that links the RST constructs to diverse normal and psychopathological processes (e.g., Corr, 2008), continuing to clarify the associations between the PCL-R factors and the BIS and BAS is likely to be of considerable relevance. In addition, although the BAS and the BIS traditionally have been more prominent than the FFFS as explanatory mechanisms for understanding psychopathy, some of the observations associated with psychopathy (e.g., decreased fear-related startle response) might be conceptualized as reflecting deficient FFFS functioning. Hence, examining the role that the FFFS plays mediating psychopaths' deficits may well be a fruitful area for future study.

References

- Ávila C, Parcet MA. The role of Gray's impulsivity in anxiety-mediated differences in resistance to extinction. *European Journal of Personality* 2000;14:185–198.
- Ávila C, Parcet MA. Personality and inhibitory deficits in the stop-signal task: The mediating role of Gray's anxiety and impulsivity. *Personality and Individual Differences* 2001;29:975–986.
- Blackburn R. Cortical and autonomic arousal in primary and secondary psychopaths. *Psychophysiology* 1979;16:143–150. [PubMed: 424497]
- Book AS, Quinsey VL. Psychopaths: Cheaters or warrior-hawks? *Personality and Individual Differences* 2004;36:33–45.
- Carver CS. Negative affects deriving from the behavioral approach system. *Emotion* 2004;4:3–22. [PubMed: 15053723]
- Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social Psychology* 1994;67:319–333.
- Caseras X, Ávila C, Torrubia R. The measurement of individual differences in behavioural inhibition and behavioural activation Systems: A comparison of personality scales. *Personality and Individual Differences* 2003;34:999–1013.
- Cleckley, H. *The mask of sanity*. 5th ed.. St. Louis, MO: Mosby; 1976.
- Corr, PJ. *Understanding biological psychology*. Malden, MA: Blackwell; 2006.
- Corr, PJ. *Reinforcement Sensitivity Theory of Personality*. New York, NY: Cambridge University Press; 2008.
- Fowles DC. The three arousal model: Implications of Gray's two-factor learning theory for heart rate, electrodermal activity, and psychopathy. *Psychophysiology* 1980;17:87–104. [PubMed: 6103567]

- Fowles DC. Psychophysiology and psychopathy: A motivational approach. *Psychophysiology* 1988;25:373–391. [PubMed: 3051073]
- Frick PJ, Lilienfeld SO, Ellis M, Loney B, Siverthorn P. The association between anxiety and psychopathy dimensions in children. *Journal of Abnormal Child Psychology* 1999;27:383–392. [PubMed: 10582839]
- Gray, JA. The psychology of fear and stress. New York: Cambridge University Press; 1987.
- Gray, JA. Elements of a Two-Process Theory of Learning. London: Academic Press; 1975.
- Gray, JA.; McNaughton, N. The Neuropsychology of Anxiety: an Enquiry into the Functions of the Septo-hippocampal System. 2nd ed.. New York: Oxford University Press; 2000.
- Gray, JA.; Smith, PT. An arousal-decision model for partial reinforcement and discrimination learning. In: Gilbert, R.; Sutherland, NS., editors. *Animal Discrimination Learning*. New York: Academic Press; 1969. p. 243–272.
- Hare, RD. Psychopathy: Theory and Research. New York: Wiley; 1970.
- Hare RD. A comparison of procedures for the assessment of psychopathy. *Journal of Consulting and Clinical Psychology* 1985;53:7–16. [PubMed: 3980831]
- Hare, RD. The Hare Psychopathy Checklist-Revised (PCL-R). Toronto, Ontario: Multi-Health Systems; 1991.
- Hare RD. Psychopathy: A clinical construct whose time has come. *Criminal Justice and behavior* 1996;23:25–54.
- Hare, RD. Manual for the Hare Psychopathy Checklist—Revised. 2nd ed.. Toronto, Ontario, Canada: Multi-Health Systems; 2003.
- Hare RD, Harpur TJ, Hakstian AR, Forth AE, Hart SD, Newman JP. The Revised Psychopathy Checklist: Reliability and factor structure. *Psychological Assessment* 1990;2:338–341.
- Harmon-Jones E, Peterson CK. Effect of trait and state approach motivation on aggressive inclinations. *Journal of Research in Personality* 2008;42:1381–1385.
- Harpur TJ, Hare RD, Hakstian AR. Two-factor conceptualization of psychopathy: Construct validity and assessment implications. *Psychological Assessment: A Journal of Consulting and Clinical Psychology* 1989;1:6–17.
- Hemphill JF, Hare RD, Wong S. Psychopathy and recidivism: A review. *Legal and Criminological Psychology* 1998;3:139–170.
- Heym N, Ferguson E, Lawrence C. An evaluation of the relationship between Gray's revised RST and Eysenck's PEN: Distinguishing BIS and FFFS in Carver and White's BIS/BAS scales. *Personality and Individual Differences* 2008;45:709–715.
- Jorm AF, Christensen H, Henderson AS, Jacomb PA, Korten AE, Rodgers B. Using the BIS/BAS scales to measure behavioral inhibition and behavioral activation: Factor structure, validity and norms in a large community sample. *Personality and Individual Differences* 1999;26:49–58.
- Lilienfeld SO. Conceptual problems in the assessment of psychopathy. *Clinical Psychology Review* 1994;14:17–38.
- Lilienfeld SO, Andrews BP. Development and preliminary validation of a self report measure of psychopathic personality traits in noncriminal populations. *Journal of Personality Assessment* 1996;66:488–524. [PubMed: 8667144]
- Lykken, DT. The antisocial personalities. Mahwah, NJ: Erlbaum; 1995.
- Neumann CS, Hare RD, Newman JP. The super-ordinate nature of the Psychopathy Checklist - Revised. *Journal of Personality Disorders* 2007;21:102–117. [PubMed: 17492916]
- Newman JP. Relationships between primary/secondary psychopathy and PCL-R Factors 1 and 2. Unpublished raw data. 2009
- Newman JP, MacCoon DG, Vaughn LJ, Sadeh N. Validating a distinction between primary and secondary psychopathy with measures of Gray's BIS and BAS constructs. *Journal of Abnormal Psychology* 2005;114:319–323. [PubMed: 15869363]
- Newman JP, Malterer MB. Problems with the BIS/BAS scales or Lykken's model of primary psychopathy? A reply to Poythress et al(2008). *Personality and Individual Differences* 2009;46:673–677. [PubMed: 20161337]
- Papoulis, A. Probability and Statistics. Englewood Cliffs, NJ: Prentice-Hall; 1990.

- Patrick CJ. Emotion and psychopathy: startling new insights. *Psychophysiology* 1994;31:319–330. [PubMed: 10690912]
- Patrick, CJ. Getting to the heart of psychopathy. In: Hervé, H.; Yuille, JC., editors. *The psychopath: Theory, research, and social implications*. Hillsdale, NJ: Lawrence Erlbaum Associates; 2007. p. 207–252.
- Patrick CJ, Bradley MM, Lang PJ. Emotion in the criminal psychopath: Startle reflex modulation. *Journal of Abnormal Psychology* 1993;102:82–92. [PubMed: 8436703]
- Patrick CJ, Hicks BM, Krueger RF, Lang AR. Relations between psychopathy facets and externalizing in a criminal offender sample. *Journal of Personality Disorders* 2005;19:339–356. [PubMed: 16178678]
- Poythress NG, Edens JF, Landfield K, Lilienfeld SO, Skeem JL, Douglas KS. A critique of Carver and White's (1994) behavioral inhibition scale (BIS) for investigating Lykken's (1995) theory of primary psychopathy. *Personality and Individual Differences* 2008;45:269–275. [PubMed: 19727321]
- Poythress NG, Skeem JL, Lilienfeld SO, Douglas KS, Edens JF. Carver and White's (1994) BIS scale and Lykken's (1995) primary psychopath: A reply to Newman and Malterer. *Personality and Individual Differences* 2009;46:678–681.
- Quinsey VL, Rice ME, Harris GT. Actuarial prediction of sexual recidivism. *Journal of Interpersonal Violence* 1995;10:85–105.
- Ross SR, Moltó J, Poy R, Segarra P, Pastor MC, Montañés S. Gray's model and psychopathy: BIS but not BAS differentiates primary from secondary psychopathy in noninstitutionalized young adults. *Personality and Individual Differences* 2007;43:1644–1655.
- Sellbom M, Verona E. Neuropsychological Correlates of Psychopathic Traits in a Non-Incarcerated Sample. *Journal of Research in Personality* 2007;41:276–294.
- Serin RC. Violent recidivism in criminal psychopaths. *Law and Human behavior* 1996;20:207–217.
- Skeem J, Johansson P, Andershed H, Kerr M, Louden JE. Two subtypes of psychopathic violent offenders that parallel primary and secondary variants. *Journal of Abnormal Psychology* 2007;116:395–409. [PubMed: 17516770]
- Torrubia R, Ávila C, Moltó J, Caseras X. The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and impulsivity dimensions. *Personality and Individual Differences* 2001;31:837–862.
- Uzieblo K, Verschuere B, Crombez G. The Psychopathic Personality Inventory: Construct validity of the two-factor structure. *Personality and Individual Differences* 2007;43:657–667.
- Walters GD. Predicting institutional adjustment and recidivism with the psychopathy checklist factor scores: a meta-analysis. *Law and Human behavior* 2003;27:541–558. [PubMed: 14593797]

Table 1

Inter-correlations among scales and scale reliabilities

	1	2	3	4	5	6	7	8
PCL-R F1	.829							
PCL-R F2	.605**	.775						
BIS	-.210**	-.181**	.744					
BAS	.068	.211**	.110*	.855				
SP	-.214**	-.123**	.539**	.071	.839			
SR	.162**	.304**	.075	.570**	.072	.820		
BIS/SP	-.242**	-.173**	.876**	.103*	.878**	.084	.861	
BAS/SR	.130**	.291**	.104*	.886**	.081	.886**	.105*	.887

Note. Values in the **diagonal** represent the internal consistency (Cronbach's alpha) of each scale. PCL-R = Psychopathy Checklist – Revised; BIS = Behavioral Inhibition System Scale; BAS = Behavioral Activation System Scale; SP = Sensitivity to Punishment Scale; SR = Sensitivity to Reward Scale.

N = 473.
* p<.05.
** p< .01.