Published in final edited form as:

Curr Probl Cancer. 2009 ; 33(1): 7–64. doi:10.1016/j.currprobcancer.2009.01.001.

Multiple Myeloma

S. Vincent Rajkumar, M.D.

Division of Hematology, Mayo Clinic, Rochester, MN USA

INTRODUCTION

Multiple myeloma is a malignant plasma cell disorder that accounts for approximately 10% of all hematologic cancers.^{1,2} It is felt to evolve from an asymptomatic pre-malignant stage of clonal plasma cell proliferation termed monoclonal gammopathy of undetermined significance (MGUS). MGUS is present in over 3% of the population above the age of 50, and progresses to myeloma or related malignancy at a rate of 1% per year.^{3,4} In some patients, an intermediate asymptomatic but more advanced pre-malignant stage referred to as smoldering multiple myeloma (SMM) can be recognized clinically.

The annual incidence of myeloma, age-adjusted to the 2000 United States population, is 4.3 per 100,000.⁵ Almost 20,000 new cases and 11,000 new deaths are estimated to occur in the United States in 2007.⁶ Multiple myeloma is twice as common in African-Americans compared to Caucasians, and slightly more common in males than females.

The median age at diagnosis is 66 years.⁷ The most common presenting symptoms are fatigue and bone pain.⁷ Osteolytic bone lesions and/or compression fractures are the hallmark of the disease, and cause significant morbidity. Anemia occurs in 70% of patients at diagnosis and is the primary cause of fatigue. Renal dysfunction occurs in 50% and hypercalcemia in 25% of patients.

HISTORICAL PERSPECTIVE

Initial Description of the disease

Although multiple myeloma has most likely been present for thousands of years, the first well-documented case was a patient, Sarah Newbury, described by Solly in 1844⁸. Later, the case of Thomas Alexander McBean, led to the discovery of the Bence Jones protein in the late 1840s.⁹⁻¹² Multiple myeloma is sometimes referred to as Kahler's disease following a case report by Professor Otto Kahler. The term "plasma cell" was introduced by Waldeyer in 1875,¹³ but it was Ramon y Cajal, the neuroanatomist, to first accurately describe the plasma cell. In 1895, Marschalko, published the best description of plasma cells which included blocked chromatin, eccentric position of the nucleus, a perinuclear pale area (hof) and a spherical or irregular cytoplasm. Wright¹⁴ was among the first to state that the tumor cells of myeloma consisted of plasma cells or immediate descendants of these cells. The first large case series of myeloma (n=412) was reported by Geschickter and Copeland.¹⁵

Corresponding author: S. Vincent Rajkumar, M.D., Mayo Clinic, 200 First Street SW, Rochester, MN 55905 USA, Phone: 507-538-0591, Fax: 507-266-4972, Email: rajkumar.vincent@mayo.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Monoclonal Proteins

Hyperproteinemia was first demonstrated in multiple myeloma in 1928 by Perlzweig et al.¹⁶ The term Bence Jones protein was first used by Fleischer in 1880.¹⁷ In 1922, Bayne-Jones and Wilson¹⁸ described the existence of two distinct groups of Bence Jones protein. The group of light chains were later characterized by Korngold and Lipari using the Ouchterlony test in 1956.¹⁹ As a tribute to Korngold and Lipari, the two classes of Bence Jones proteins were subsequently designated kappa and lambda. In 1937, Tiselius²⁰ separated serum globulins into three components: alpha, beta, and gamma. In 1939, Tiselius and Kabat²¹ demonstrated that the gamma globulin fraction consisted of antibodies. The tall, narrow-based, “church spire” peak on serum protein electrophoresis characteristic of multiple myeloma was recognized in 1939.²² Immunoelectrophoresis was described by Grabar and Williams in 1953,²³ and eleven years later, immunofixation was introduced by Wilson²⁴. Immunofixation is now the most commonly used approach for the identification of monoclonal proteins. The concept of monoclonal versus polyclonal gammopathies was enunciated by Jan Waldenström in 1961.²⁵

Alkylator and Corticosteroid Based therapy

Rhubarb pill and infusion of orange peel is the first documented treatment for myeloma, administered to Sarah Newbury in 1844. Mr. McBean was treated with phlebotomy, with the application of leeches used as “maintenance therapy.” Later, he was treated with steel and quinine.^{8,9,26} Urethane introduced by Alwall in 1947 was the standard of therapy for more than 15 years.²⁷ But urethane was discarded in 1966 when Holland et al.²⁸ randomized 83 patients with treated or untreated multiple myeloma to receive urethane or a placebo consisting of cherry and cola-flavored syrup and found no difference in the objective status or survival between the two treatment groups.

Melphalan (sarcolysin), a mainstay of therapy even today, was first introduced in 1958 by Blokhin et al.²⁹ He found benefit in three of six patients with multiple myeloma. Later, in 1962, Bergsagel et al.³⁰ reported significant improvement in 8 of 24 patients with multiple myeloma who were treated with melphalan.

In 1967, prednisone was reported by Maas et al to produce benefit in 8 of 10 patients with poor-risk myeloma.³¹ In a placebo-controlled double-blind trial, prednisone as a single agent produced significant decreases in serum globulin and an increase in hematocrit but no difference in survival when compared with a placebo.³² In a reanalysis of 2 CALGB myeloma treatment protocols, prednisone as a single agent produced a 44% objective response rate.³³

In 1969, Alexanian found in a randomized trial of 183 myeloma patients that survival was 6 months longer with melphalan and prednisone than melphalan alone, giving rise to the classic melphalan-prednisone (MP) regimen.³⁴ For the next 30 years, numerous chemotherapy combinations were compared to MP in an attempt to improve on overall survival, without avail. In a large meta-analysis of over 20 randomized trials comparing melphalan and prednisone with various combinations of therapeutic agents, response rates were significantly higher with combination chemotherapy (60%) compared with melphalan and prednisone (53%) ($P < 0.001$). However, there was no significant difference in overall survival.³⁵ It is only recently with the arrival of stem cell transplantation and novel agents as described below that improvements in overall have occurred in myeloma.

Stem Cell Transplantation

In 1957, Thomas et al. treated 6 human patients (1 of whom had multiple myeloma) with TBI or chemotherapy followed by an intravenous infusion of bone marrow cells.³⁶ McElwain & Powles reported the first successful cases of autologous bone marrow transplantation in myeloma.³⁷ Subsequently, Barlogie developed a series of treatment programs using autologous

transplantation (“total therapy I, II and III”) that played a major role in establishing high-dose therapy and stem cell rescue as standard therapy for myeloma.

DISEASE DEFINITION

The disease definition for myeloma has undergone multiple revisions over the years. The current definition requires 10% or more clonal plasma cells on bone marrow examination (or biopsy proven plasmacytoma), M protein in the serum and/or urine (except in patients with true non-secretory myeloma) and evidence of end-organ damage (hypercalcemia, renal insufficiency, anemia, or bone lesions) that is felt to be secondary to the underlying plasma cell disorder.^{38,39} Table 1 provides the diagnostic criteria for myeloma, as well as related clonal plasma cell disorders that must be entertained in the differential diagnosis and distinguished from myeloma.

PATHOGENESIS

Evolution of MGUS

The first step in myeloma pathogenesis is the establishment of MGUS, a precursor lesion that precedes almost all cases of myeloma. The origins of MGUS are likely linked to antigenic stimulation. Studies show that human myeloma cell lines and primary myeloma cells express a broad range of Toll-like receptors (TLRs) which normally help B cells recognize infectious agents and pathogen-associated molecular patterns (PAMP).⁴⁰⁻⁴² TLR-specific ligands cause increased myeloma cell proliferation, survival, and resistance to dexamethasone induced apoptosis. In addition, there is overexpression of CD126 (interleukin-6 receptor alpha-chain) in MGUS compared to normal plasma cells.^{43,44} Thus abnormal TLR expression and/or overexpression of IL-6 receptors in plasma cells may result in an enhanced response to infection and provide a sustained, autocrine IL-6 dependent, proliferative trigger for plasma cells which cause (or act in concert with) the certain cytogenetic events described below to establish the premalignant MGUS stage.

Approximately 50% of MGUS is associated with primary translocations in the clonal plasma cells involving the immunoglobulin heavy chain (IgH) locus on chromosome 14q32 (IgH translocated MGUS/SMM).^{1,45} The most common partner chromosome loci and genes dysregulated in these translocations are: 11q13 (CCND1 [cyclin D1 gene]), 4p16.3 (FGFR-3 and MMSET), 6p21 (CCND3 [cyclin D3 gene]), 16q23 (c-maf), and 20q11 (mafB)(Table 2).⁴⁶⁻⁴⁸ Most of the remaining cases of MGUS are associated with hyperdiploidy (IgH non-translocated MGUS).^{45,49-54}

Transformation of MGUS to Myeloma

The progression of MGUS to myeloma represents a simple, random, two-hit genetic model of malignancy.³ Several abnormalities such as Ras mutations, p16 methylation, abnormalities involving the myc family of oncogenes, secondary translocations, and p53 mutations have all been identified in clonal plasma cells in association with progression to the symptomatic stage.⁴⁶ The bone marrow microenvironment also undergoes marked changes with progression, including induction of angiogenesis,^{55,56} suppression of cell-mediated immunity,⁵⁷ and paracrine loops involving cytokines such as interleukin-6 and VEGF (vascular endothelial growth factor).⁵⁸

Lytic bone lesions characteristic of myeloma are caused by an imbalance between the activity of osteoclasts and osteoblasts. There is an increase in RANKL (receptor activator of nuclear factor κB ligand) expression by osteoblasts (and possibly plasma cells) accompanied by a reduction in the level of its decoy receptor, osteoprotegerin (OPG).^{59,60} This leads to an increase in RANKL/OPG ratio, which causes osteoclast activation and bone resorption. In

addition, increased levels of macrophage inflammatory protein-1 α (MIP-1 α), IL-3 and IL-6 produced by marrow stromal cells also contribute to the overactivity of osteoclasts. At the same time, increased levels of IL-3, IL-7 and dickkopf 1 (DKK1) inhibit osteoblast differentiation in myeloma.⁶¹

ADVANCES IN DIAGNOSIS

Myeloma is characterized by the presence of monoclonal immunoglobulins in the serum and/or urine. Monoclonal immunoglobulins are commonly referred to as monoclonal proteins, M proteins, or paraproteins. The presence of M proteins is indicative of a clonal plasma cell proliferative disorder such as myeloma, MGUS, Waldenstrom macroglobulinemia, etc; additional tests are required to distinguish between the various plasma cell disorders. M proteins are detected by serum protein electrophoresis (SPEP) in 82% of patients with myeloma, and by serum immunofixation (IFE) in 93%.⁷ The addition of urine protein electrophoresis and IFE (or the serum free light chain assay) increases the sensitivity of detecting monoclonal proteins in patients with myeloma to 97%.⁶² Currently only 1-2% of patients with myeloma will have no detectable M on any of these tests; these patients have true non-secretory myeloma.

Protein electrophoresis and immunofixation

Electrophoresis of serum and urine proteins (SPEP and UPEP, respectively) and immunofixation of these samples are the preferred methods of detection of monoclonal (M) proteins. UPEP requires a 24 hour urine collection for accuracy. M proteins on SPEP and UPEP appear as a localized band, and electrophoresis also allows measurement of their size.⁶³ After recognition of a possible M protein on electrophoresis, immunofixation is necessary for confirmation, and to determine the heavy- and light-chain class of the M protein. In addition, immunofixation is more sensitive than electrophoresis, and allows detection of smaller amounts of M protein and should therefore be performed whenever myeloma, amyloidosis, macroglobulinemia or a related disorder is suspected. Quantitation of serum immunoglobulins is performed with a rate nephelometer in patients in whom serum M proteins are detected as an added parameter that can be followed to assess response. Quantitative immunoglobulins are particularly useful in patients with myeloma and macroglobulinemia where at times the M protein size estimated on SPEP may be unreliable (eg., small beta migrating proteins, and IgA / IgM proteins that tend to polymerize)

Serum free light chain assay

The serum free light-chain (FLC) assay (Freelite™, The Binding Site Limited, Birmingham, U.K.) has been recently introduced into clinical practice.⁶⁴ This automated nephelometric assay allows quantitation of free kappa (κ) and lambda (λ) chains (i.e., light chains that are not bound to intact immunoglobulin) secreted by plasma cells. An abnormal kappa/lambda FLC ratio indicates an excess of one light chain type versus the other, and is interpreted as a surrogate for clonal expansion based on extensive testing in normal volunteers, and patients with myeloma, amyloidosis and renal dysfunction.^{64,65} The assay is performed on automated chemistry analyzers, is widely available, and is commonly used to monitor patients with oligo-secretory or non-secretory myeloma and primary amyloidosis, as well as patients with the light-chain only form of myeloma.⁶⁵⁻⁶⁷

The normal serum free-kappa level is 3.3 to 19.4 mg/L and the normal free-lambda level is 5.7 to 26.3 mg/L.⁶⁸ The normal ratio for FLC-kappa/lambda is 0.26-1.65. The normal reference range in the FLC assay reflects a higher serum level of free lambda light chains than would be expected given the usual kappa/lambda ratio of 2 for intact immunoglobulins. This occurs because the renal excretion of free kappa (which exists usually in a monomeric state) is much

faster than free lambda (which is usually in a dimeric state).^{64,65} Patients with a kappa/lambda FLC ratio <0.26 are typically defined as having monoclonal lambda free light chain and those with ratios >1.65 are defined as having a monoclonal kappa free light chain. If the FLC ratio is >1.65 , kappa is considered to be the “involved” FLC and lambda the “uninvolved” FLC, and vice versa if the ratio is less than 0.26.

In a recent study, 428 patients with a monoclonal gammopathy and urine M protein at initial diagnosis of plasma cell dyscrasia who had also undergone serum immunofixation and serum FLC quantitation within 30 days of diagnosis were studied.⁶² The results of SPEP, serum immunofixation, serum FLC, UPEP, and urine immunofixation were then analyzed to determine if all patients could be accurately identified as having a monoclonal plasma cell disorder had the urine studies not been done. All 3 serum studies were normal in only 2 patients (0.5% of the cohort), indicating that urine studies can be eliminated when screening for the presence of monoclonal plasma cell disorders by using the serum FLC assay in combination with the SPEP and immunofixation. However, urine studies are required if a monoclonal process is identified since it helps in monitoring disease progression and response to therapy over time.

Evaluation of bone disease

Plain radiographic examination of all bones including long bones (skeletal survey) is the preferred method of detecting lytic bone lesions in myeloma. CT and MRI scans are more sensitive than conventional radiography in detecting bone disease and are indicated when symptomatic areas show no abnormality on routine radiographs. Fluoro-deoxyglucose positron emission tomography (FDG-PET) and bone densitometry studies have shown promise in the evaluation of bone disease in myeloma and need further investigation.

Bone marrow studies

By definition, all patients with myeloma have 10% or more clonal bone marrow plasma cells. If a lower extent of involvement is detected, one is either dealing with an erroneous diagnosis or a sampling error due to patchy marrow involvement. Occasionally an entity called “multiple” solitary plasmacytomas has been described in which there are clearly multiple plasmacytomas on clinical and radiographic examination, but marrow involvement is either minimal or absent. The monoclonal nature of bone marrow plasma cells is established by the demonstration of an abnormal kappa/lambda ratio by immunohistochemistry or flow cytometry. Plasma cells in multiple myeloma typically stain positive for CD38, CD56, and CD138, and are usually negative for surface immunoglobulin and CD19. Bone marrow samples should be studied conventional karyotyping studies and/or fluorescent in-situ hybridization studies to detect specific abnormalities such as t(4;14), t(14;16), and 17p-.

CURRENT STAGING AND PROGNOSTIC EVALUATION

Staging

Since 1975, the Durie-Salmon staging system has been used to stratify patients with multiple myeloma.⁶⁹ However, this staging system has limitations especially in the categorization of bone lesions.^{70,71} Recently Greipp et al, developed an International Staging System (ISS) based on data from 11, 171 patients.⁷² The ISS overcomes the limitations of the Durie-Salmon staging, and divides patients into 3 distinct stages and prognostic groups based solely on the beta-2-microglobulin and albumin levels in the serum. Table 3 provides the details, and the advantages and disadvantages of these two staging systems.

Prognostic Factors

Once patients are staged, several independent prognostic factors are helpful in predicting outcome in myeloma (Table 4).^{45,70,73-75} A combination of two or more of the independent prognostic factors listed on Table 4 provides adequate information to estimate prognosis. For example, Facon et al have shown that patients undergoing stem cell transplantation can be grouped into three clear prognostic categories using 2 prognostic factors; median survival was 25 months if patients had a high beta-2 microglobulin and deletion of chromosome 13 by fluorescent in situ hybridization.⁷⁶ Patients with only one factor abnormal had a median survival of 47 months, while survival was in excess of 111 months in patients in whom both factors were normal.

Prognostic factors such as age, hemoglobin concentration, creatinine, calcium, albumin, immunoglobulin class subtype, and extent of bone marrow involvement have demonstrated some prognostic value.^{70,73,75} However, they are of limited value once stage and the independent prognostic factors discussed earlier are known.^{77,78} Newer risk factors such as angiogenesis shed light on biology, but clinical utility is unclear.

Risk-stratification

Using some of the most important clinically available prognostic factors discussed above, a risk-stratification method has been adopted by the Mayo Clinic, called mSMART (Mayo Stratification for Myeloma and Risk-adapted Therapy; www.msmaart.org). This model helps stratify patients into high-risk and standard-risk myeloma based on detection of deletion 13 or hypodiploidy on metaphase cytogenetic studies, deletion 17p- or immunoglobulin heavy chain (IgH) translocations t(4;14) or t(14;16) on molecular genetic studies, or plasma cell labeling index of 3% or higher on bone marrow immunofluorescent studies. Presence of any one or more of these high-risk factors (Table 5) classifies a patient as having high-risk MM.⁷⁹ The median survival of high-risk MM is only 2-3 years even with tandem stem cell transplantation, compared to over 6-7 years in patients with average-risk MM.²

RESPONSE ASSESSMENT

Several criteria have been used over the years to assess response to therapy in myeloma. In recent years, the European Group for Blood and Bone Marrow Transplant/ International Bone Marrow Transplant Registry/ American Bone Marrow Transplant Registry (EBMT/ IBMTR/ ABMTR criteria for the response and progression of MM treated by stem cell transplantation, commonly referred to as the Blade criteria or the EBMT criteria was adopted in many trials.⁸⁰ Other commonly used response criteria were those developed by the Chronic Leukemia-Myeloma Task Force, Southwest Oncology group (SWOG), and the Eastern Cooperative Oncology Group (ECOG) have been largely abandoned. In 2006, the International Myeloma Working Group (IMWG) recognized the need for uniformity and published uniform response criteria that are to be used in future clinical trials.⁸¹

The IMWG uniform response criteria are similar to the EBMT criteria with a few major exceptions: addition of free light chain response and progression criteria for patients without measurable disease (see Table 6 for definition of measurable disease), modification of the definition for disease progression for patients in complete response, addition of very good partial response and stringent response categories, elimination of the minor response category, elimination of the mandatory 6 week wait time to confirm response, and some additional clarifications and correction of errors. The IMWG criteria for response and progression are listed on Table 7.

NEW AGENTS IN THE TREATMENT OF MYELOMA

As described earlier, for decades, the mainstay of therapy has been alkylators and corticosteroids, specifically the oral regimen of melphalan and prednisone (MP). High dose therapy with autologous stem cell transplantation (ASCT) was subsequently shown to prolong survival compared to conventional chemotherapy, and adopted into standard practice. More recently, thalidomide,⁸² bortezomib,^{83,84} and lenalidomide,^{85,86} have emerged as effective agents in the treatment of myeloma (Table 8), and each has an interesting historical perspective.

In this section, the new agents for the treatment of myeloma will be reviewed. Subsequent sections will discuss in detail the current approach to initial therapy, stem cell transplantation, maintenance therapy, and treatment of relapsed refractory disease.

Thalidomide

Historical perspective—Chemie Grünenthal, a German pharmaceutical company introduced thalidomide(α -N-[phthalimido] glutarimide) into the market as a sedative on October 1, 1957.⁸⁷⁻⁸⁹ By 1960, it was sold in over 40 countries, and became popular both as a sedative and as treatment for morning sickness of pregnancy. It was marketed under various commercial names such as ‘Contergan’, ‘Distal’, ‘Softanon’, ‘Neurosedyn’, ‘Isomin’, ‘Kedavon’, ‘Telargan’, and ‘Sedalis.’⁹⁰

On November 18, 1961, Widukind Lenz, a German physician and geneticist determined that thalidomide was associated with severe teratogenic malformations.⁹⁰ In December 1961, independent confirmation came from William McBride, an Australian obstetrician.⁹¹ Fetal malformations occurred inevitably when the drug was taken in the first trimester between days 35-49 after the last menstrual period.^{87,88} By the end of 1961, thalidomide was taken off the market in most countries but almost 10,000 infants had already been affected. The United States was spared, because the drug had been denied approval by Dr. Frances Kelsey at the FDA who was concerned about the lack of safety data.⁹²

Shortly after the teratogenic properties of thalidomide were known, it was considered as a possible treatment against cancer.^{93,94} At least three clinical trials were undertaken in cancer,^{95,96} but no significant activity was seen. Although thalidomide was taken off the market and was unsuccessful against cancer, it nevertheless persisted as a therapeutic agent due to promising activity detected in leprosy (1964),^{97,98} Behcet’s Disease (1979),⁹⁹ graft versus host disease (1988),¹⁰⁰⁻¹⁰⁴ and human immunodeficiency virus (HIV) associated oral ulcers¹⁰⁵ and wasting (1989).¹⁰⁶ Under pressure from HIV activists and to discourage illegal drug distribution networks, the FDA approved thalidomide for the treatment of erythema nodosum leprosum in the United States in July 1998 with a built in safety system termed the System for Thalidomide Education and Prescribing Safety (STEPS) program.^{107,108}

Thalidomide is a chiral molecule and is dispensed as a racemic mixture of the (R)- (dextrorotatory) and (S)- (levorotatory) enantiomers.^{109,110} Although it is felt that the (R)- enantiomer is responsible for most of its sedative activities, and the (S)- enantiomer has a significant role in the teratogenic and anti-tumor properties, differences in enantio-selectivity are not clinically relevant because of rapid chiral interconversion *in vivo* at physiologic pH.¹¹¹ As a result of this rapid racemization, the use of a pure (R)- enantiomer therefore would not have prevented the thalidomide tragedy. Most of the metabolism is by non-enzymatic hydrolysis, and no major dose reductions are generally needed in liver or renal failure.

Initial results in myeloma—Based on the significant anti-angiogenic properties of thalidomide,^{112,113} and evidence of increased angiogenesis in myeloma, Singhal, Barlogie and colleagues treated patients with relapsed and refractory multiple myeloma with the drug in late

1997. Thirty-two percent of 84 patients responded, making thalidomide the first new drug with single-agent activity for myeloma in over three decades.^{82,114} These results were confirmed by many other centers, in all phases of the disease.¹¹⁵⁻¹²²

Studies show that response rates in relapsed disease are about 50% with the combination of thalidomide and steroids,¹²³ and increase to over 65% with an oral three-drug combination of thalidomide, steroids and cyclophosphamide (CTD).¹²⁴⁻¹²⁶ Several other combination chemotherapy regimens containing thalidomide have been developed including DT-PACE (dexamethasone, thalidomide, cisplatin, adriamycin, cyclophosphamide and etoposide), BLT-D (clarithromycin, low dose thalidomide, and dexamethasone), and MTD (melphalan, thalidomide and dexamethasone).¹²⁷

Dosage—Initially thalidomide was used in doses of up to 800 mg per day. However, most side-effects of thalidomide are dose-dependent, and there is no major improvement in clinical activity beyond a dose of 200 mg. Currently, thalidomide is usually given in a dosage of 50 to 200 mg daily; the dose is typically adjusted to the lowest dose that can achieve and maintain a response in order to minimize long-term toxicity. In most elderly patients (age >65) a dose of 50 to 100 mg is better tolerated.

Side-effects—The known toxicities of thalidomide include peripheral neuropathy, constipation, fatigue and sedation.¹²⁸ Therapy for myeloma has revealed new toxicities of thalidomide that were previously unrecognized. These include deep vein thrombosis,¹²⁹⁻¹³² toxic epidermal necrolysis,¹³³ and hypothyroidism.¹³⁴ The incidence of deep vein thrombosis (DVT) is only 1-3% in patients receiving thalidomide alone, but rises to approximately 10-15% in patients receiving thalidomide in combination with dexamethasone, and to about 25% in patients receiving the agent in combination with other cytotoxic chemotherapeutic agents, particularly doxorubicin.^{129-131,135,136} The management of thalidomide toxicity has been reviewed in detail by Ghobrial et al.¹²⁸

Bortezomib

Historical Perspective—The multi-catalytic ubiquitin-proteasome pathway is responsible for the degradation of eukaryotic cellular proteins.¹³⁷⁻¹⁴⁰ The orderly degradation of cellular proteins by the ubiquitin-proteasome pathway is critical for signal transduction, transcriptional regulation, response to stress, and control of receptor function.¹⁴¹⁻¹⁴⁴ The 26S proteasome (1500-2000 kDa) consists of a core 20S catalytic complex (approximately 700 kDa) and a 19S regulatory complex (Figure 1).^{137,141,144-148} Ubiquitin tagged proteins are recognized by the 19S regulatory complex, where the ubiquitin tags are removed proteins are fed into the inner catalytic compartments of the 20S proteasome cylinder for hydrolysis into small polypeptides ranging from 3 to 22 amino acids in length.^{149,150} 151 152 137,150,153

Inhibition of the proteasome inhibition leads to cellular apoptosis, with malignant, transformed and proliferating cells being more susceptible.^{148,154} 137,155-158 Numerous proteasome inhibitors were developed but none was suitable for clinical use.^{141,148,159,160} Subsequently, Adams and colleagues designed and developed several boronic acid derived compounds that inhibit the proteasome pathway in a highly specific manner.¹⁵⁴ Bortezomib (*N*-pyrazinecarbonyl-L-phenylalanine-L-leucine boronic acid; previously known as PS-341 or MLN-341), a boronic acid dipeptide was then selected for preclinical and clinical testing.^{83, 154} Bortezomib inhibits the proteasome pathway rapidly and in a reversible manner by binding directly with the 20S proteasome complex and blocking its enzymatic activity.

Preclinical studies demonstrated that bortezomib had potent cytotoxic and growth inhibitory effects across the panel of 60 cell lines from the NCI and in several murine models.^{139, 161-166} Bortezomib also showed striking activity against myeloma cells in several preclinical

models.¹⁶⁷⁻¹⁷⁰ Orlowski and colleagues conducted a phase 1 study in which responses were seen in several myeloma patients.

Initial results in myeloma—In the first phase II trial conducted in 202 patients with relapsed/refractory MM, approximately one-third responded to bortezomib therapy, with an average response duration of one year.⁸³ These results were confirmed in a randomized phase II trial in patients who failed to respond or relapsed after front-line therapy for myeloma.¹⁷¹ Although initial trials only allowed a maximum of 8 cycles of bortezomib, responding patients were allowed to receive continued treatment on a compassionate use protocol, data from which indicate that it is safe to give at least an additional 5-6 cycles of therapy, without undue toxicity.¹⁷¹ In a recent randomized trial, progression free survival was superior with bortezomib compared to dexamethasone alone in patients with relapsed, refractory myeloma.⁸⁴

Dosing—The starting dose of bortezomib is 1.3 mg/m² given twice weekly on days 1, 4, 8, and 11 every 21 days. Dosage may need to be decreased to 1.0 mg/m² or 0.7 mg/m² based on toxicity. Although no clear maintenance dose has been defined, after 4-6 cycles, the dosage often needs to be decreased if there is a necessity to continue the drug in order to minimize the risk of neuropathy. One strategy is to lengthen the cycle length from 21 days to 28 or even 35 days. Another option is to administer bortezomib once a week.

Side-effects—The most common toxicities attributable to bortezomib therapy are gastrointestinal side effects, cytopenias, fatigue, fever, and peripheral neuropathy. Peripheral neuropathy occurs in approximately 45% of patients, and is the most worrisome adverse effect. It can be painful. It is more frequent in patients who have received prior neurotoxic therapy and in those with pre-existing neuropathy. Neuropathy can be of grade 3 severity in about 10% of patients receiving therapy. Postural hypotension occurs in approximately 10% of patients.

Lenalidomide

Historical Perspective—Shortly after thalidomide was determined to be a teratogen in the 1960s, a number of analogs of thalidomide were synthesized to determine the mechanism of its teratogenicity. Similarly in the 1990s, a number of thalidomide analogs were synthesized in an attempt to increase efficacy and minimize toxicity. Lenalidomide, a 4-amino substituted analog of thalidomide formerly called CC-5013, is in a class of thalidomide analogs termed immunomodulatory drugs (IMiDs). It is safer and more effective than thalidomide in preclinical and clinical studies. In contrast to the clinical development of thalidomide, the development of lenalidomide can be considered an inevitable occurrence, commonly seen in various fields of medicine where pharmaceutical companies develop closely related and presumably more effective alternatives once an effective parent drug is identified.

Based on the preclinical studies conducted by the manufacturer and at the Dana-Farber Cancer Institute, lenalidomide was tested in phase I trials in relapsed refractory myeloma.¹⁷²

Initial Results in myeloma—Lenalidomide first showed promising activity in MM in 2 phase I trials.^{172,173} In a phase II trial conducted at Mayo Clinic, 31 of 34 patients (91%) with newly diagnosed myeloma achieved an objective response with lenalidomide plus dexamethasone.⁸⁵ Richardson et al reported on single-agent activity in a multi-center randomized phase II trial that enrolled 102 patients with relapsed/refractory myeloma.⁸⁶ Overall response rate with single-agent lenalidomide was 17%. Two large phase III trials demonstrated significantly superior time to progression and overall survival with lenalidomide plus dexamethasone compared to placebo plus dexamethasone in relapsed myeloma.^{174,175} Based on these two randomized trials, lenalidomide plus dexamethasone was approved by the FDA in June 2006 for the treatment of myeloma in patients who have failed one prior therapy.

Dosing—Typical dosing of lenalidomide for myeloma is 25 mg per day on days 1-21 of a 28 day cycle, with dose adjustments based on toxicity. With prolonged therapy dose adjustments are usually needed. Most patients who receive lenalidomide for over one year are usually on lower doses of 5-15 mg per day given in the usual schedule. The dosing of lenalidomide in renal failure is unclear, and caution must be exercised since the drug is renally excreted.

Side-effects—The most common side-effects of lenalidomide are cytopenias. Fatigue, dryness of skin, and rash are other common side effects. The incidence of DVT is low (2-3%) with single-agent lenalidomide or lenalidomide plus low-dose dexamethasone, but rises markedly when the agent is combined with high-dose dexamethasone (approximately 20%). The risk of DVT can be decreased to less than 10% by reducing the dose of dexamethasone to 40 mg once a week.

Liposomal-doxorubicin

For many years, doxorubicin has been used in myeloma as part of the VAD regimen and other combination regimens. However, the single-agent activity of doxorubicin is minimal. Recently, liposomal doxorubicin has shown activity in newly diagnosed and relapsed myeloma. In a randomized trial, bortezomib plus liposomal doxorubicin was associated with a superior time to progression, resulting in the approval of liposomal doxorubicin for the treatment of relapsed myeloma.¹⁷⁶ Further studies are needed to define the role of liposomal doxorubicin in myeloma.

Pomalidomide (CC-4047)

Pomalidomide (an analog of thalidomide and lenalidomide) has shown promising single-agent activity.¹⁷⁷ In combination with low-dose dexamethasone pomalidomide has shown a response rate of over 60% in relapsed myeloma in a trial conducted by the Mayo Clinic.¹⁷⁸ In this trial, pomalidomide was given orally 2 mg daily on days 1-28 of a 28-day cycle. Dexamethasone was given orally at a dose of 40 mg daily on days 1, 8, 15 and 22 of each cycle. Additional studies are ongoing. Pomalidomide is not approved by the FDA.

Carfilzomib

Carfilzomib, an irreversible proteasome inhibitor has shown significant single-agent activity in myeloma in two separate trials.^{179,180} The dose of carfilzomib was 20 mg/m² intravenously days 1, 2, 8, 9, 15 and 16 every 28 days, for up to 12 cycles. There is a suggestion that carfilzomib may have less neurotoxicity compared with bortezomib. It is also not approved by the FDA so far. Clinical trials with carfilzomib are ongoing.

Other new drugs

There is a continued search for other active agents based on advances in myeloma biology.¹⁸¹ Other agents which have raised some interest and are undergoing testing include heat shock protein inhibitors, histone deacetylase inhibitors, perifosine, and bendamustine.

INITIAL TREATMENT OF MYELOMA

The median survival of symptomatic myeloma is 4 years. For over 3 decades, the mainstay of therapy was oral chemotherapy with melphalan and prednisone (MP). High dose therapy with autologous stem cell transplantation (ASCT) prolongs survival compared to conventional chemotherapy, and is now routinely incorporated into the treatment strategy either early in the disease course or at the time of relapse in eligible patients. As discussed earlier, thalidomide,⁸² bortezomib,^{83,84} and lenalidomide,^{85,86} have emerged as effective agents in the treatment of myeloma and have greatly altered the way myeloma is treated today.

Despite many advances, there is still no evidence that early treatment of patients with asymptomatic (smoldering) multiple myeloma prolongs survival compared to therapy at the time of symptoms. Similarly the use of bisphosphonates does not alter the natural history of smoldering myeloma, and are not indicated.¹⁸² Therapy is therefore initiated only in patients defined as having multiple myeloma based on the criteria listed on Table 1; patients with smoldering myeloma should be observed closely without treatment. In all stages of myeloma, clinical trials are preferred over standard therapeutic approaches, since the disease is still incurable. Over the years several landmark trials have greatly altered the treatment of myeloma, details of which are listed on Table 9.

The choice of initial therapy depends on eligibility for stem cell transplantation and risk-stratification. The overall approach to treatment of symptomatic newly diagnosed multiple myeloma is outlined in Figure 1A and B. Eligibility for stem cell transplantation is determined by age, performance status, and coexisting comorbidities. Risk-Stratification is based on presence or absence of high-risk factors (Table 5)⁷⁹ Table 10 lists the most common regimens used in the treatment of newly diagnosed myeloma, including recommended doses and activity.

Transplant-Candidates

It is important to avoid protracted melphalan-based therapy in patients with newly diagnosed myeloma who are considered eligible for ASCT, since it can interfere with adequate stem cell mobilization, regardless of whether an early or delayed transplant is contemplated. Typically patients are treated with approximately 4 cycles of induction therapy prior to stem cell harvest. This includes patients who are transplant candidates but who wish to reserve ASCT as a delayed option for relapsed refractory disease. Such patients can resume induction therapy following stem cell collection until a plateau phase is reached, reserving ASCT for relapse.

Vincristine, doxorubicin, dexamethasone (VAD)—VAD was used for many years as pre-transplant induction therapy for patients considered candidates for ASCT. However, VAD requires an intravenous indwelling catheter, which predisposes patients to catheter related sepsis and thrombosis. Moreover, the neurotoxicity of vincristine can limit the future use of thalidomide and bortezomib, both of which also have neurotoxic potential. Most of the activity of VAD is from the high-dose dexamethasone component. Recently, Cavo and colleagues in a matched case-control study of 200 patients demonstrated that response rates with VAD were significantly lower compared to Thal/Dex; 76% versus 52%, respectively.¹⁸³ Preliminary results from a randomized trial from France confirm these findings.¹⁸⁴ As a result, VAD is not recommended anymore as initial therapy.

Pulsed Dexamethasone—Dexamethasone alone has also been used as induction therapy. It is typically used at a dose of 40 mg orally on days 1-4, 9-12, and 17-20 every 4-5 weeks. Objective response rates are approximately 45%,¹⁸⁵ significantly lower compared to newer induction regimens. In randomized trials the early mortality rate associated with dexamethasone is over 10% in the first 4 months of therapy, reflecting the toxicity and ineffectiveness of this regimen. Consequently, single-agent dexamethasone is also no longer recommended as initial therapy.

Thalidomide-Dexamethasone (Thal/Dex)—In the last few years, Thal/Dex has emerged as the most commonly used induction regimen for the treatment of newly diagnosed myeloma in the United States. The use of Thal/Dex was initially based on three phase II clinical trials.^{118,186,187} Response rates with Thal/Dex range 64-76% in these studies, comparable or better than those obtained with infusional VAD.

The Eastern Cooperative Oncology Group (ECOG) recently reported the results of a randomized trial comparing Thal/Dex to dexamethasone (Table 7).¹⁸⁵ Two hundred seven patients were studied. The best response within four cycles of therapy was significantly higher with Thal/Dex compared to dexamethasone alone; 63% versus 41%, respectively, $P=0.0017$. Adjusted response rates allowing for the use of serum M protein values alone in patients in whom a measurable urine protein at baseline was unavailable at follow-up were 72% with Thal/Dex versus 50% with dexamethasone alone. Stem cell harvest was successful in 90% of patients in each arm. DVT was more frequent with Thal/Dex (17% versus 3%). Overall, grade 3 or higher non-hematologic toxicities were seen in 67% of patients within four cycles with Thal/Dex and 43% with dexamethasone alone ($P < 0.001$). Early mortality (first 4 months) was 7% with Thal/Dex and 11% with dexamethasone alone. Based on this trial, the United States Food and Drug Administration (FDA) has granted accelerated approval for Thal/Dex for the treatment of newly diagnosed myeloma.

Confirmatory results are available from a separate randomized, double-blind, placebo-controlled study comparing Thal/Dex versus placebo/dexamethasone (Placebo/Dex) as primary therapy in 470 patients with newly diagnosed myeloma.¹⁸⁸ In this trial, the overall response rate was significantly higher with Thal/Dex compared with Placebo/Dex (63% v 46%), $P < 0.001$. Time to progression (TTP) was significantly longer with Thal/Dex compared with Placebo/Dex (median, 22.6 v 6.5 months, $P < 0.001$). As in the ECOG study, grade 4 adverse events were more frequent with Thal/Dex than with Placebo/Dex (30.3% v 22.8%).

A Dutch study compared thalidomide, Adriamycin, dexamethasone (TAD) with VAD as induction therapy.¹⁸⁹ After induction patients in both arms proceeded to stem cell transplantation. A total of 556 patients were randomized and progression free survival was superior with TAD compared with VAD, median 25 months to 33 months ($P < 0.001$). However no difference in overall survival was seen.

Patients receiving thalidomide in combination with high-dose steroids or chemotherapy need routine thromboprophylaxis with coumadin (target INR 2-3) or low-molecular weight heparin (equivalent of enoxaparin 40mg once daily). Aspirin can be used instead in patients receiving only low doses of dexamethasone (40mg once a week or lower) or prednisone in combination with thalidomide, provided no concomitant erythropoietic agents are used.

Lenalidomide-dexamethasone (Rev/Dex)—In a phase II trial conducted at the Mayo Clinic demonstrated remarkably high activity with the Rev/Dex regimen. Thirty-one of 34 patients (91%) achieved an objective response, including 2 (6%) achieving complete response (CR), and 11 (32%) meeting criteria for very good partial response (VGPR).⁸⁵ Lacy et al recently updated the results of this study. With longer follow-up, 56% of patients achieved very good partial response or better. In the subset of 21 patients receiving Rev/Dex as primary therapy without ASCT, 67% achieved VGPR or better.¹⁹⁰ Approximately fifty percent of patients experienced grade 3 or higher non-hematologic toxicity, similar to rates seen with dexamethasone alone.

ECOG recently reported preliminary findings of a randomized trial testing Rev/Dex as administered in the Mayo Phase II trial (and in the regulatory relapsed refractory myeloma studies) versus Rev/low-dose Dex (40 mg dexamethasone once weekly).¹⁹¹ Results so far show that toxicity rates are significantly higher with Rev/standard-dose Dex compared to Rev/low-dose Dex. Early (first 4 month) mortality rates were low in both arms, 5% and 0.5% respectively. The early mortality rate in the Rev/low-dose dexamethasone arm is probably the lowest reported in any large phase III newly diagnosed trial in which enrollment was not restricted by age or eligibility for stem cell transplantation; DVT rates are also low, making this one of the safest pre-transplant induction regimens for myeloma. Based on this Rev/low-

dose dexamethasone is currently the regimen of choice in the Mayo Stratification for Myeloma and Risk-adapted Therapy (mSMART) protocol for the treatment of standard-risk myeloma in patients who are candidates for ASCT outside the setting of a clinical trial.⁷⁹

Recommendations for thromboprophylaxis are similar to those discussed above with Thal/Dex; aspirin alone is probably sufficient for patients receiving lenalidomide plus low-dose dexamethasone.

Bortezomib-based regimens—In newly diagnosed myeloma, bortezomib has shown response rates of approximately 40% as a single-agent.¹⁹² Significantly higher response rates (approximately 70-90%) have been observed with bortezomib plus dexamethasone (Vel/Dex),^{193,194} bortezomib, thalidomide, dexamethasone (VTD) and other bortezomib-based combinations. The CR plus VGPR rate is approximately 25-30% with Vel/Dex in one study. No adverse effect on stem cell mobilization was noted.

Harousseau and colleagues recently reported preliminary results of a randomized trial comparing VAD versus Vel/Dex as pre-transplant induction therapy.¹⁹⁵ With over 400 patients enrolled, preliminary results show superior response rates and long-term outcome with Vel/Dex compared to VAD. DVT risk was low with bortezomib (<5%).

Bortezomib-based regimens are of particular value in patients with renal failure and in patients with high-risk myeloma (see below), and is not associated with high rates of thrombosis.

Other induction regimens—The role of other pre-transplant induction regimens, such as those containing doxorubicin or liposomal doxorubicin, need to be weighed in terms of the added side-effects that can affect quality of life, and should be considered investigational until future studies show that the addition of these agents improves long-term outcome compared to the regimens discussed above. One exception is patients presenting with very aggressive disease including plasma cell leukemia features or rapidly progressive disease with or without extramedullary features. In these patients, 2 cycles of the combination chemotherapy regimen VDT-PACE (bortezomib (Velcade), dexamethasone, thalidomide, cisplatin (platinum), doxorubicin (Adriamycin), cyclophosphamide, and etoposide) developed by Barlogie et al as part of total therapy III is highly effective in controlling the disease rapidly.¹⁹⁶

Initial Therapy in Standard-Risk Patients Eligible for Transplantation

Patients who are not transplant candidates are treated with standard alkylating agent therapy. For decades this has meant therapy with melphalan plus prednisone (MP).¹ Over the years, despite better response rates, no survival benefit has been reported with any of the more aggressive combination chemotherapy regimens compared to MP. Two recent randomized studies show that MPT (melphalan, prednisone, thalidomide) improves response and EFS compared to MP;^{197,198} an overall survival advantage has been observed in one of the two trials.¹⁹⁸ As a result, MPT has emerged as the current standard of care for patients not eligible for ASCT.

Melphalan, prednisone (MP)—MP has been used for decades in myeloma. The use of MP has decreased due to more active regimens as discussed below. However, it is still a regimen of value in frail, elderly patients who may not tolerate bortezomib or thalidomide well. The response rate with MP is approximately 50%, and complete responses occur in fewer than 5% of patients. Prior to the arrival of new agents such as thalidomide and bortezomib, despite better response rates no survival benefit was found with any combination chemotherapy regimen compared to MP.³⁵ The dosing of MP is given on Table 10.¹²² Blood counts are monitored 3 weeks after beginning therapy and the dose of melphalan is adjusted to produce mild cytopenia at mid-cycle.

Melphalan, prednisone, thalidomide (MPT)—Five randomized trials have compared MP to MPT.^{197,199-202} Only two show a clear overall survival benefit.^{199,200}

Facon et al¹⁹⁹ randomized 447 patients (ages 65-75) to MP versus MPT versus tandem autologous stem cell transplantation (ASCT) with reduced dose melphalan (Melphalan 100 mg/m²) (Table 9). Early (first 3 months) mortality rate was 8% with MP compared to 3% with MPT. Significantly higher response and PFS rates were observed with MPT compared with either MP or tandem ASCT groups. More importantly the trial demonstrated a significant survival advantage with MPT; median overall survival not reached at 52 months, 33 months, and 38 months, respectively (p<0.001).

Hulin et al²⁰⁰ confirmed a survival advantage with MPT compared with MP in a randomized trial in patients over the age of 75. After numerous attempts to improve on MP over the years with a variety of combination chemotherapy regimes, the results of these 3 randomized trials finally changed the standard of care for elderly patients.

Palumbo et al randomized patients to either standard dose MP for 6 months or to MPT for six months followed by maintenance thalidomide (Table 9).¹⁹⁷ Overall response rates were significantly higher with the MPT than the MP (76% versus 48%) as was the CR plus near CR rate (28% versus 7%). MPT also resulted in superior 2-year EFS rates (54% versus 27%, p=0.0006). No significant survival improvement has been seen even with longer follow up.²⁰³ Randomized trials led by Wijermans et al and Waage et al also do not show any significant survival benefit with MPT compared to MP.^{201,202}

MPT therapy is associated with greater toxicity than MP. Therefore not all elderly patients may be able to receive the regimen. Grade 3-4 adverse events occur in approximately 50% of patients treated with MPT, compared to 25% with MP.¹⁹⁷ As with Thal/Dex, there is a significant (20%) risk of DVT with MPT in the absence of thromboprophylaxis. However, this rate drops to approximately 3% with the use of thromboprophylaxis (eg., enoxaparin).¹⁹⁷

Melphalan, Prednisone, Bortezomib (MPV)—Mateos et al have reported the first results of the novel combination MPV in newly diagnosed myeloma in patients 65 years or older. Therapy was associated with a response rate of 89%, including 32% CR rate. Approximately half of the patients with CR also had no residual bone marrow plasma cells by immunophenotypic studies. In addition, MPV appeared to overcome the poor prognosis conferred by deletion 13 and IgH translocations. Event-free survival (EFS) and overall survival at 16 months were 83% and 90%, respectively. Most common grade 3 or higher adverse events were thrombocytopenia (51%), neutropenia (43%), peripheral neuropathy (17%), diarrhea (16%) and infection (16%). Herpes zoster infection occurred without prophylaxis in 13%; and was reduced to 7% in with prophylactic acyclovir. MPV appears very promising with high CR rates.

A recent randomized trial compared VMP to MP in 682 patients (Table 9).²⁰⁴ The time to progression among patients receiving VMP was 24 months versus 16.6 months with MP, P<0.001. Partial response or better was seen in 71% with VMP and 35% with MP; corresponding complete-response rates were 30% versus 4%, respectively (P<0.001). Overall survival was significantly better with VMP, p=0.008. Grade 3 events occurred more often with VMP versus MP (53% vs. 44%, P=0.02). Grade 3 neuropathy was seen in 13% with VMP. This study established the superiority of VMP over MP in myeloma.

Melphalan, Prednisone, Lenalidomide (MPR)—Palumbo and colleagues have studied the addition of lenalidomide to MP in newly diagnosed patients older than 65 years of age.²⁰⁵ Fifty-four patients were studied. At the maximum tolerated dose, the overall response rate

was 81%, with 48% of patients achieving at least very good partial response (VGPR) or better, and 24% of patients achieving CR. Major grade 3-4 adverse events were neutropenia, thrombocytopenia, anemia, rash and febrile neutropenia. All patients received prophylactic aspirin; DVT was uncommon, occurring in 3 patients 6% including two after aspirin discontinuation. Most patients needed growth factor support. Based on this trial, MPR appears to be a very effective oral regimen for the treatment of elderly patients with multiple myeloma who are not candidates for ASCT. An ECOG randomized trial is comparing MPR to MPT.

HEMATOPOIETIC STEM CELL TRANSPLANTATION

Autologous stem cell transplantation (ASCT)

Although not curative, ASCT improves complete response rates and prolongs median overall survival in myeloma by approximately 12 months.²⁰⁶⁻²⁰⁹ The mortality rate is 1-2%. The major studies that established the role of ASCT in myeloma are listed on Table 9. Melphalan, 200 mg/m² is the most widely used preparative (conditioning) regimen for ASCT. It is superior to the older regimen of melphalan 140 mg/m² and 8 Gy total body irradiation (TBI).²¹⁰ Studies are ongoing to determine if the conditioning regimen can be improved with the addition of radioactive compounds (Holmium (HO¹⁶⁶ DOTMP) or Samarium 153-SM-EDTMP)^{211,212} or bortezomib.

Three randomized trials show that survival is similar whether ASCT is done early (immediately following 4 cycles of induction therapy) or delayed (at the time of relapse as salvage therapy) (Table 9).²¹³⁻²¹⁵ In a Spanish PETHEMA group randomized trial, patients responding to induction therapy had similar overall and progression free survival with either ASCT or 8 additional courses of chemotherapy,²¹⁶ suggesting that the greatest benefit from ASCT may be in those with disease refractory to induction therapy.^{217,218}

There is little doubt that ASCT prolongs survival in myeloma, but its timing (early versus delayed) is controversial. Overall, given the inconvenience and side-effects of prolonged chemotherapy, insurance, and other issues we still favor early ASCT, especially for patients less than 65 years of age with adequate renal function.²⁰⁸ However, given effective new agents to treat myeloma, some patients and physicians may choose to delay the procedure. The need for early ASCT is an important question for future clinical trials.

Tandem transplantation—With tandem (double) ASCT, patients receive a second planned ASCT after recovery from the first procedure (Table 11).^{219,220} The recent IFM 94 randomized trial found significantly better event-free and overall survival in recipients of double versus single ASCT (Table 9).²²¹ A similar benefit was also demonstrated in a randomized trial conducted in Italy;²²² two other randomized trials are yet to show significant improvement in overall survival with tandem ASCT, but they have shorter follow up.^{223,224}

In both the French and Italian trials, the benefit of a second ASCT was restricted to patients failing to achieve a complete response or very good partial response (>90% reduction in M protein level) with the first procedure. Based on these results, we now routinely collect enough stem cells for two transplants in all eligible patients. Patients achieving a complete response or very good partial response with the first transplant are observed or offered clinical trials investigating maintenance therapy, reserving the second ASCT for relapse.²²⁵ Patients not achieving such a response are offered a second ASCT.

Allogeneic Transplantation

The advantages of allogeneic transplantation are lack of graft contamination with tumor cells and presence of a graft versus myeloma effect.^{226,227} However, only 5%-10% of patients are candidates because of age, availability of a HLA matched sibling donor, and adequate organ

function. Further, the high treatment related mortality (TRM), mainly related to graft versus host disease (GVHD) has made conventional allogeneic transplants unacceptable for most patients with myeloma.

Several recent trials have been conducted using non-myeloablative conditioning regimens (mini-allogeneic transplantation).²²⁸ Initial trials evaluating this approach included relapsed or refractory patients, which was thought to account at least in part for the poor outcomes. It also became apparent that this approach was less useful in patients who had significant residual tumor burden at the time of the non-myeloablative transplant, which led to the concept of a planned autologous SCT followed by a planned reduced intensity allogeneic stem cell transplant (RIC SCT) a couple of months later.²²⁹ The initial studies had treatment related mortalities approaching 25%, 3-year overall survival and progression-free survival rates were 41% and 21%, respectively.²²⁹ Adverse OS was associated with chemoresistant disease, more than 1 prior transplantation, and absence of chronic GVHD. Using the planned tandem autologous/nonmyeloablative allogeneic stem cell transplantation approach, outcomes have been better with treatment-related mortalities of 15% and 2 year overall survivals approaching 75%.²³⁰ However, there is also a high risk of acute and chronic GVHD, and the occurrence of GVHD appears necessary for disease control.

A French randomized trial addressed the role of allogeneic stem cell transplant in high risk patients as defined by the presence of deletion 13 by FISH and of beta-2 microglobulin greater than 3 mg/L.²³¹ Based on biological randomization, patients were allocated to either double autologous stem cell transplant or to autologous SCT followed by a RIC allogeneic SCT. Though patients did better than expected in both arms, with median overall survivals of 41 and 35 months, respectively, there was no significant difference between the 2 arms with a median follow-up of 24 months. The major criticisms of this study have been that the criteria used to select 'high-risk' disease did not select for the highest risk patients; and that the reduced intensity conditioning used may have been too immunosuppressive, thereby abrogating the graft versus myeloma effect.

In contrast to the French trial, a significant survival advantage was seen with ASCT followed by a RIC allogeneic SCT compared to tandem ASCT in a randomized trial conducted in Italy. On an intent to treat basis, the median overall and event-free survival were longer with autologous SCT followed by an RIC compared to tandem ASCT (80 months vs. 54 months, $p=0.01$; and 35 months vs. 29 months, $p=0.02$, respectively).²³² However, the sample size was modest ($n=162$), and additional confirmation is needed.

A subsequent trial investigating non-myeloablative transplants in myeloma strikes a note of caution. In this Spanish trial reported by Rosinol et al,²³³ 110 patients with myeloma failing to achieve near-complete remission (nCR) after first ASCT were assigned to a second ASCT (85 patients) or a reduced-intensity-conditioning allogeneic transplantation (allo-RIC; 25 patients), based on the human leukocyte antigen (HLA)-identical sibling donor availability. Although the CR rate was superior with allo-RIC (40% vs 11%, $P = 0.001$) transplantation-related mortality was higher (16% vs 5%, $P = .07$), and no benefit was seen with regards to overall survival. In addition, allo-RIC was associated with a 66% rate of chronic graft-versus-host disease. and no statistical difference in event-free survival and overall survival. At this time, mini-allogeneic transplantation remains investigational. It should only be considered in the context of clinical trials in standard-risk myeloma because current results with the tandem ASCT strategy described earlier yields 7-year survival rates in excess of 40%. In these patients the treatment related mortality and GVHD rates with non-myeloablative allogeneic transplantation are unacceptably high.

TREATMENT OF HIGH-RISK MYELOMA

Patients with high-risk myeloma tend to do poorly with median overall survival of approximately two years even with tandem ASCT. The main option for these patients is novel therapeutic strategies.^{2,79} Bortezomib containing regimens should be considered early in the disease course as primary therapy. In at least 3 separate studies, bortezomib appears to overcome the adverse effect of deletion 13.²³⁴⁻²³⁶ Due to the high risk of early relapse, stem cell transplantation is best reserved for relapse. Routine maintenance therapy could be considered (eg., lenalidomide plus low-dose dexamethasone) given the high risk of early relapse. Clearly clinical trials and new agents specifically designed for high-risk myeloma are needed.

MAINTENANCE THERAPY

Maintenance therapy with interferon- α is of limited value and is seldom used.²³⁷ Recent results from a large intergroup study showed no benefit with interferon as maintenance therapy.²¹⁵ A study by Berenson and colleagues suggests that prednisone may be useful for maintenance therapy.²³⁸ Progression-free survival (14 versus 5 months) and overall survival (37 versus 26 months) were significantly longer with 50 mg versus 10 mg of prednisone orally every other day. Since this comparison included only those who responded initially to steroid-based therapy and since patients did not receive ASCT, it is difficult to generalize these results to current practice. Clinical trials are currently evaluating thalidomide, dendritic cell vaccination, and other novel approaches as maintenance therapy.

A French randomized trial (IFM 99-02) randomized 597 patients (age <65) following tandem autologous stem cell transplantation to no maintenance, pamidronate, or pamidronate plus thalidomide.²³⁹ There was a significant improvement in event-free survival with maintenance thalidomide plus pamidronate; the 3-year event-free survival rate from the time of randomization was 36% in arm A, 37% in arm B, and 52% in arm C ($P < 0.009$). The corresponding 4-year overall survival rate was 77%, 74%, and 87%, respectively ($P < 0.04$). A similar overall survival benefit with thalidomide post-transplant has been noted in separate randomized trials conducted in Australia,²⁴⁰ Italy,²⁴¹ and Tunisia.²⁴² However, no significant overall survival benefit with maintenance thalidomide has been reported in a randomized trial conducted in Brazil,²⁴³ and in separate trials conducted by the Medical Research Council (MRC),²⁴⁴ and the University of Arkansas.²⁴⁵ Thus the results with thalidomide maintenance are conflicting. It also appears that most of the benefit is seen in patients who have failed to achieve a very good partial response with ASCT, and that thalidomide is functioning more as a fixed duration of consolidation therapy rather than maintenance. Further, it is not clear if all patients not receiving thalidomide maintenance had access to thalidomide at relapse in most trials. More data are needed before recommending thalidomide as maintenance particularly given the long term adverse effects of the agent. At present for standard risk patients, observation remains the standard following ASCT. Multiple studies addressing the use of lenalidomide post-stem cell transplantation are on-going.

TREATMENT OF RELAPSED MULTIPLE MYELOMA

Almost all patients with myeloma eventually relapse. If relapse occurs more than 6 months after stopping therapy, the initial chemotherapy regimen should be re-instituted. Patients who have cryopreserved stem cells early in the disease course can derive significant benefit from ASCT as salvage therapy.²⁴⁶

In general, patients who have indolent relapse can often be treated with single-agents or MP. These patients present with asymptomatic increases in serum and urine monoclonal protein levels, progressive anemia, or few small lytic bone lesions. In contrast, patients with more

aggressive relapse often require therapy with a combination of active agents. Given the non-curative nature of myeloma, patients with relapsed disease typically continue on one drug/regimen until relapse or toxicity and then try the next option.

The activity of various new agents in relapsed disease was discussed earlier. Besides thalidomide, lenalidomide, and bortezomib, high-dose pulse dexamethasone or intravenous methylprednisolone are reasonable options, particularly if the patient has had an initial response to steroids and was off high-dose steroids when the relapse occurred.^{247,248} Patients relapsing more than 6-12 months following an auto-SCT can respond to MP or MPT, since most would not have been exposed to this regimen as induction therapy.

Combination chemotherapy regimens such as VDT-PACE, VAD or other alkylator-based regimens can be effective in relapsed and refractory disease. Intravenous melphalan at a dose of 25mg/m² is another active regimen, but usually requires transfusion and growth factor support.

TREATMENT OF COMPLICATIONS

Hypercalcemia

Aggressive hydration with isotonic saline and corticosteroids are effective in most cases. A single dose of pamidronate, 60-90 mg intravenously over 2-4 hours,²⁴⁹ or zoledronic acid 4 mg intravenously over 15 minutes,²⁵⁰ will normalize the calcium levels within 24-72 hours in most patients

Skeletal Lesions

Surgical fixation of fractures or impending fractures of long bones may be needed. Local radiation should be limited to patients with disabling pain who have a well-defined focal process that has not responded to analgesics and/or chemotherapy. The administration of bisphosphonates significantly reduces the number of skeletal events (pathologic fracture, need for irradiation or surgery on bone and spinal cord compression).²⁵¹ In a randomized trial of 392 patients with at least one lytic lesion, skeletal events (pathologic fracture, radiation or surgery to bone, and spinal cord compression) after 9 months of therapy were significantly fewer with pamidronate compared to placebo, 24% versus 41%, respectively, $P < 0.001$.²⁵¹ More recently, zoledronic acid has been shown to have comparable efficacy to pamidronate.^{252,253} Either pamidronate (90 mg intravenously over at least 2 hours every 4 weeks) or zoledronic acid (4 mg intravenously over 15-30 minutes every 4 weeks) is recommended in patients with multiple myeloma who have one or more lytic lesions on skeletal roentgenograms.^{252,254}

The typical recommendation by initially was to continue bisphosphonates indefinitely at monthly intervals.²⁵⁴ However, by 2003, avascular osteonecrosis of jaw (ONJ) has been described as a new complication associated with their use²⁵⁵⁻²⁵⁹ The etiology of ONJ is unclear, but is likely multifactorial in origin. Although most patients who develop ONJ have had recent dental or oral surgical procedures (70%), the remainder develop spontaneous ONJ.²⁵⁹ Proposed mechanisms include that inhibition of osteoclast activity reduces bone turnover and remodeling and that bisphosphonates prevent release of bone-specific factors that promote bone formation.²⁶⁰

The incidence of ONJ ranges from 4.4% to 21% depending of they type and duration of bisphosphonate use.^{260,261} Risks increase with duration of therapy, and likely with use of zoledronic acid compared to pamidronate or other lower intensity bisphosphonates. A Mayo clinic consensus statement recommends pamidronate instead of zoledronic acid as the bisphosphonate of choice for long-term therapy.²⁶⁰ It also recommends 2 years of monthly

bisphosphonate therapy for patients with myelomatous bone disease, followed by either cessation of therapy in patient who are off active treatment for their myeloma or continuation of therapy every 3 months for those who are receiving myeloma therapy. These recommendations are bolstered by two recent observations. The first is that the rates of myeloma bone disease in the pre-bisphosphonate era was highest during the first 2 years after diagnosis.²⁶² The second supporting data element is the recent finding that pamidronate use after tandem transplant did not provide any significant reduction in skeletal events.²³⁹

Both vertebroplasty (injection of methylmethacrylate into a collapsed vertebral body) or kyphoplasty (introduction of an inflatable bone tamp into the vertebral body and after inflation the injection of methylmethacrylate into the cavity) have been successfully used to decrease pain and help restore height.²⁶³ Pain relief is generally rapid, and can be long-lasting.

Spinal cord compression from an extramedullary plasmacytoma should be suspected in patients with severe back pain, weakness or paresthesias of the lower extremities, or bladder or bowel dysfunction, or incontinence. The standard treatment for cord or cauda equina compression are corticosteroids (dexamethasone 10 mg to 40 mg IV followed by 4 mg orally/intravenously four times daily) and radiation therapy. On rare occasions, surgical decompression may be necessary.

Renal Insufficiency

Nonsteroidal anti-inflammatory agents can precipitate renal failure and should generally be avoided.^{264,265} Dehydration, infection, and radiographic contrast media may also contribute to acute renal failure. Maintenance of a high urinary output (3 L/day) is important in preventing renal failure in those with high levels of monoclonal light chains in the urine. Persons with acute or subacute renal failure due to light-chain cast nephropathy should be treated with Thal/Dex or dexamethasone to reduce the tumor mass as quickly as possible. A trial of plasmapheresis should be considered in these patients in an attempt to prevent irreversible renal damage.²⁶⁶

Anemia

Iron, folate, or B-12 deficiency may be responsible for the anemia as well and must be recognized and treated. Treatment of the underlying disease and renal failure often leads to improvement in the hemoglobin level. Erythropoietin (40,000 units subcutaneously weekly) or darbepoietin (200 micrograms subcutaneously every 2 weeks) are useful in patients with persistent symptomatic anemia,^{267,268} but should be restricted because of a higher risk of DVT when used in combination with thalidomide or lenalidomide. Blood transfusions are indicated for patients with symptomatic anemia who do not obtain benefit from other therapies.

Infections

Patients should receive pneumococcal and influenza vaccinations. Intravenously administered gamma globulin every 3-4 weeks is indicated if patients have recurrent serious infections associated with severe hypogammaglobulinemia. The use of prophylactic antibiotics in patients receiving chemotherapy for myeloma has not been settled. A small randomized placebo-controlled trial of trimethoprim-sulfamethoxazole in 57 patients with newly diagnosed myeloma demonstrated benefit with routine prophylaxis administered with the first two cycles of chemotherapy.²⁶⁹ Due to the high risk of infections, prophylactic antibiotics for 3-4 months should be considered for all patients starting therapy for newly diagnosed myeloma. Prophylaxis against *Pneumocystis carinii* pneumonia (PCP) should also be considered in all patients receiving high dose steroid therapy. However, due to the risk of serious skin toxicity, trimethoprim-sulfamethoxazole should be avoided in patients receiving Thal/Dex or Rev/Dex therapy. In these patients prophylaxis with other antibiotics (such as ciprofloxacin, levofloxacin

or cephalosporins) and alternative agents for PCP prophylaxis should be considered. A randomized trial comparing no prophylaxis, trimethoprim-sulfamethoxazole, and ciprofloxacin in patients with newly diagnosed myeloma receiving chemotherapy has just completed accrual, and results are awaited.

Hyperviscosity Syndrome

Infrequently patients with multiple myeloma develop hyperviscosity syndrome. Plasmapheresis promptly relieves the symptoms and should be done regardless of the viscosity level if the patient has signs or symptoms of hyperviscosity.²⁷⁰

Acknowledgments

Supported in part by Grants CA62242 and CA107476 from the National Cancer Institute, Bethesda, MD.

References

1. Kyle RA, Rajkumar SV. Multiple Myeloma. *N Engl J Med* 2004;351:1860–1873. [PubMed: 15509819]
2. Rajkumar SV, Kyle RA. Multiple Myeloma: Diagnosis and Treatment. *Mayo Clin Proc* 2005;80:1371–1382. [PubMed: 16212152]
3. Kyle RA, Therneau TM, Rajkumar SV, et al. A long-term study of prognosis of monoclonal gammopathy of undetermined significance. *N Engl J Med* 2002;346:564–569. [PubMed: 11856795]
4. Kyle RA, Therneau TM, Rajkumar SV, et al. Prevalence of Monoclonal Gammopathy of Undetermined Significance. *N Engl J Med* 2006;354:1362–1369. [PubMed: 16571879]
5. Kyle RA, Therneau TM, Rajkumar SV, Larson DR, Plevak MF, Melton LJ 3rd. Incidence of multiple myeloma in Olmsted County, Minnesota: Trend over 6 decades. *Cancer* 2004;101:2667–2674. [PubMed: 15481060]
6. Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer Statistics, 2007. *CA Cancer J Clin* 2007;57:43–66. [PubMed: 17237035]
7. Kyle RA, Gertz MA, Witzig TE, et al. Review of 1,027 patients with newly diagnosed multiple myeloma. *Mayo Clinic Proc* 2003;78:21–33.
8. Solly S. Remarks on the pathology of mollities ossium with cases. *Medica-Chirurgical Transactions of London* 1844;27:435–461.
9. Macintyre W. Case of mollities and fragilitas ossium, accompanied with urine strongly charged with animal matter. *Medica-Chirurgical Transactions of London* 1850;33:211–232.
10. Clapp JR. Some aspects of the first recorded case of multiple myeloma. *Lancet* 1967;2:1354–1356. [PubMed: 4170040]
11. Bence Jones H. Chemical pathology. *Lancet* 1847;2:88–92.
12. Bence Jones H. On the new substance occurring in the urine of a patient with mollities ossium. *Philos Trans R Soc Land* 1848;138:55–62.
13. Waldeyer W. Ueber bindegewebszellen. *Archiv fur microbiologie and anatomie* 1875;11:176–194.
14. Wright JH. A case of multiple myeloma. *Trans Assoc Am Phys* 1900;15:137–147.
15. Geschickter CF, Copeland MM. Multiple myeloma. *Arch surg* 1928;16:807–863.
16. Perlzweig WA, Delrue G, Geschichter C. Hyperproteinemia associated with multiple myelomas: report of an unusual case. *J Am Med Assoc* 1928;90:755–757.
17. Fleischer R XXIV. Ueber das Vorkommen des sogenannten Bence Jones' schen Eiweisskorpers im normalen Knochenmark. *Arch Pathol Anatom Physiol Klin Med* 1880;80:842–849.
18. Bayne-Jones S, Wilson DW. Immunological reactions of Bence-Jones proteins. II. Differences between Bence-Jones proteins from various sources. *Bulletin of the Johns Hopkins Hospital* 1922;33:119–125.
19. Korngold L, Lipari R. Multiple-myeloma proteins. III. The antigenic relationship of Bence Jones proteins to normal gamma-globulin and multiple-myeloma serum proteins. *Cancer* 1956;9:262–272. [PubMed: 13304841]

20. Tiselius A. A new apparatus for electrophoretic analysis of colloidal mixtures. *Transactions of the Faraday Society* 1937;33:524.
21. Tiselius A, Kabat EA. Electrophoretic study of immune sera and purified antibody preparation. *J Exp Med* 1939;69:119–131.
22. Longsworth LG, Shedlovsky T, MacInnes DA. Electrophoretic patterns of normal and pathological human blood, serum, and plasma. *Journal of Experimental Medicine* 1939;70:399–413.
23. Grabar P, Williams CA. Methode permettant l'étude conjuguee des proprietes electrophoretiques et immunochimiques d'un melange de proteines; application au serum sanguin. *Biochim Biophys Acta* 1953;10:193–194. [PubMed: 13041735]
24. Wilson AT. Direct Immunoelectrophoresis. *J Immunol* 1964;92:431–434. [PubMed: 14128988]
25. Waldenstrom J. Studies on conditions associated with disturbed gamma globulin formation (gammopathies). *Harvey Lectures* 1960/1961;56:211–231. [PubMed: 14004528]
26. Kyle RA. Multiple myeloma: an odyssey of discovery. *British Journal of Haematology* 2000;111:1035–1044. [PubMed: 11167737]
27. Alwall N. Urethane and stilbamidine in multiple myeloma: report on two cases. *Lancet* 1947;2:388–389.
28. Holland JR, Hosley H, Scharlau C, et al. A controlled trial of urethane treatment in multiple myeloma. *Blood* 1966;27:328–342. [PubMed: 5933438]
29. Blokhin N, Larionov L, Perevodchikova N, Chebotareva L, Merkulova N. Clinical experiences with sarcolysin in neoplastic diseases. *Ann N Y Acad Sci* 1958;68:1128–1132. [PubMed: 13627766]
30. Bergsagel DE, Sprague CC, Austin C, Griffith KM. Evaluation of new chemotherapeutic agents in the treatment of multiple myeloma. IV. L-Phenylalanine mustard (NSC-8806. *Cancer Chemother Rep* 1962;21:87–99. [PubMed: 13867794]
31. Salmon SE, Shadduck RK, Schilling A. Intermittent high-dose prednisone (NSC-10023) therapy for multiple myeloma. *Cancer Chemother Rep* 1967;51:179–187. [PubMed: 6043738]
32. Mass RE. A comparison of the effect of prednisone and a placebo in the treatment of multiple myeloma. *Cancer Chemother Rep* 1962;16:257–259. [PubMed: 14470881]
33. McIntyre OR, Pajak TF, Kyle RA, Cornwell GG 3rd, Leone L. Response rate and survival in myeloma patients receiving prednisone alone. *Medical & Pediatric Oncology* 1985;13:239–243. [PubMed: 4033539]
34. Alexanian R, Haut A, Khan AU, et al. Treatment for multiple myeloma. Combination chemotherapy with different melphalan dose regimens. *JAMA* 1969;208:1680–1685. [PubMed: 5818682]
35. Myeloma Trialists' Collaborative Group. Combination chemotherapy versus melphalan plus prednisone as treatment for multiple myeloma: An overview of 6,633 patients from 27 randomized trials. *J Clin Oncol* 1998;16:3832–3842. [PubMed: 9850028]
36. Thomas ED, Lochte HL Jr, Lu WC, Ferrebee JW. Intravenous infusion of bone marrow in patients receiving radiation and chemotherapy. *N Engl J Med* 1957;257:491–496. [PubMed: 13464965]
37. McElwain TJ, Powles RL. High-dose intravenous melphalan for plasma-cell leukaemia and myeloma. *Lancet* 1983;2:822–824. [PubMed: 6137651]
38. The International Myeloma Working Group. Criteria for the classification of monoclonal gammopathies, multiple myeloma and related disorders: a report of the International Myeloma Working Group. *Br J Haematol* 2003;121:749–757. [PubMed: 12780789]
39. Rajkumar SV, Dispenzieri A, Kyle RA. Monoclonal gammopathy of undetermined significance, Waldenstrom macroglobulinemia, AL amyloidosis, and related plasma cell disorders: diagnosis and treatment. *Mayo Clinic Proceedings* 2006;81:693–703. [PubMed: 16706268]
40. Jego G, Bataille R, Geffroy-Luseau A, Descamps G, Pellat-Deceunynck C. Pathogen-associated molecular patterns are growth and survival factors for human myeloma cells through Toll-like receptors. *Leukemia* 2006;20:1130–1137. [PubMed: 16628189]
41. Bohnhorst J, Rasmussen T, Moen SH, et al. Toll-like receptors mediate proliferation and survival of multiple myeloma cells. *Leukemia* 2006;20:1138–1144. [PubMed: 16617319]
42. Mantovani A, Garlanda C. Inflammation and multiple myeloma: the Toll connection. *Leukemia* 2006;20:937–938. [PubMed: 16721383]

43. Perez-Andres M, Almeida J, Martin-Ayuso M, et al. Clonal plasma cells from monoclonal gammopathy of undetermined significance, multiple myeloma and plasma cell leukemia show different expression profiles of molecules involved in the interaction with the immunological bone marrow microenvironment. *Leukemia* 2005;19:449–455. [PubMed: 15674420]
44. Rawstron AC, Fenton JA, Ashcroft J, et al. The interleukin-6 receptor alpha-chain (CD126) is expressed by neoplastic but not normal plasma cells. *Blood* 2000;96:3880–3886. [PubMed: 11090073]
45. Fonseca R, Barlogie B, Bataille R, et al. Genetics and cytogenetics of multiple myeloma: a workshop report. *Cancer Res* 2004;64:1546–1558. [PubMed: 14989251]
46. Kuehl WM, Bergsagel PL. Multiple myeloma: evolving genetic events and host interactions. *Nature Rev Cancer* 2002;2:175–187. [PubMed: 11990854]
47. Bergsagel PL, Kuehl WM. Chromosome translocations in multiple myeloma. *Oncogene* 2001;20:5611–5622. [PubMed: 11607813]
48. Fonseca R, Bailey RJ, Ahmann GJ, et al. Genomic abnormalities in monoclonal gammopathy of undetermined significance. *Blood* 2002;100:1417–1424. [PubMed: 12149226]
49. Drach J, Angerler J, Schuster J, et al. Interphase fluorescence in situ hybridization identifies chromosomal abnormalities in plasma cells from patients with monoclonal gammopathy of undetermined significance. *Blood* 1995;86:3915–3921. [PubMed: 7579361]
50. Fonseca R, Aguayo P, Ahmann GJ, et al. Translocations at 14q32 are common in patients with the monoclonal gammopathy of undetermined significance (MGUS) and involve several partner chromosomes. *Blood* 1999;94(Suppl 1):663a. [PubMed: 10397733]A 2943
51. Rajkumar SV. MGUS and Smoldering Multiple Myeloma: Update on Pathogenesis, Natural History, and Management. *Hematology (Am Soc Hematol Educ Program)* 2005;2005:340–345. [PubMed: 16304401]
52. Chng WJ, Van Wier SA, Ahmann GJ, et al. A validated FISH trisomy index demonstrates the hyperdiploid and nonhyperdiploid dichotomy in MGUS. *Blood* 2005;106:2156–2161.10.1182/blood-2005-02-0761 [PubMed: 15920009]
53. Magrangeas F, Lode L, Wuilleme S, Minvielle S, Avet-Loiseau H. Genetic heterogeneity in multiple myeloma. *Leukemia* 2005;19:191–194. [PubMed: 15538406]
54. Kaufmann H, Ackermann J, Baldia C, et al. Both IGH translocations and chromosome 13q deletions are early events in monoclonal gammopathy of undetermined significance and do not evolve during transition to multiple myeloma. *Leukemia* 2004;18:1879–1882. [PubMed: 15385925]
55. Rajkumar SV, Mesa RA, Fonseca R, et al. Bone marrow angiogenesis in 400 patients with monoclonal gammopathy of undetermined significance, multiple myeloma, and primary amyloidosis. *Clin Cancer Res* 2002;8:2210–2216. [PubMed: 12114422]
56. Kumar S, Fonseca R, Dispenzieri A, et al. Prognostic value of angiogenesis in solitary bone plasmacytoma. *Blood* 2003;101:1715–1717. [PubMed: 12393501]
57. Galea HR, Cogne M. GM-CSF and IL-12 production by malignant plasma cells promotes cell-mediated immune responses against monoclonal Ig determinants in a light chain myeloma model. *Clinical & Experimental Immunology* 2002;129:247–253. [PubMed: 12165080]
58. Vacca A, Ria R, Ribatti D, et al. A paracrine loop in the vascular endothelial growth factor pathway triggers tumor angiogenesis and growth in multiple myeloma. *Haematologica*. 2003
59. Roodman GD. Role of the bone marrow microenvironment in multiple myeloma. *Journal of Bone & Mineral Research* 2002;17:1921–1925. [PubMed: 12412796]
60. Roodman GD. Mechanisms of bone metastasis. *N Engl J Med* 2004;350:1655–1664. [PubMed: 15084698]see comment
61. Tian E, Zhan F, Walker R, et al. The role of the Wnt-signaling antagonist DKK1 in the development of osteolytic lesions in multiple myeloma. *N Engl J Med* 2003;349:2483–2494. [PubMed: 14695408] see comment
62. Katzmann JA, Dispenzieri A, Kyle R, et al. Elimination of the Need for Urine Studies in the Screening Algorithm for Monoclonal Gammopathies by Using Serum Immunofixation and Free Light Chain Assays. *Mayo Clin Proc* 2006;81:1575–1578. [PubMed: 17165636]
63. Kyle RA, Rajkumar SV. Monoclonal gammopathy of undetermined significance. *Br J Haematol* 2006;134:573–589. [PubMed: 16938117]

64. Katzmman JA, Clark RJ, Abraham RS, et al. Serum reference intervals and diagnostic ranges for free kappa and free lambda immunoglobulin light chains: relative sensitivity for detection of monoclonal light chains. *Clinical Chemistry* 2002;48:1437–1444. [PubMed: 12194920]
65. Bradwell AR, Carr-Smith HD, Mead GP, Harvey TC, Drayson MT. Serum test for assessment of patients with Bence Jones myeloma. *Lancet* 2003;361:489–491. [PubMed: 12583950]
66. Drayson M, Tang LX, Drew R, Mead GP, Carr-Smith H, Bradwell AR. Serum free light-chain measurements for identifying and monitoring patients with nonsecretory multiple myeloma. *Blood* 2001;97:2900–2902. [PubMed: 11313287]
67. Lachmann HJ, Gallimore R, Gillmore JD, et al. Outcome in systemic AL amyloidosis in relation to changes in concentration of circulating free immunoglobulin light chains following chemotherapy. *Br J Haematol* 2003;122:78–84. [PubMed: 12823348]
68. Abraham RS, Clark RJ, Bryant SC, et al. Correlation of serum immunoglobulin free light chain quantification with urinary Bence Jones protein in light chain myeloma. *Clin Chem* 2002;48:655–657. [PubMed: 11901068]
69. Durie BG, Salmon SE. A clinical staging system for multiple myeloma. Correlation of measured myeloma cell mass with presenting clinical features, response to treatment, and survival. *Cancer* 1975;36:842–854. [PubMed: 1182674]
70. Kyle RA. Prognostic factors in multiple myeloma. *Stem Cells* 1995;2:56–63. [PubMed: 8520513]
71. Rapoport BL, Falkson HC, Falkson G. Prognostic factors affecting the survival of patients with multiple myeloma. A retrospective analysis of 86 patients. *S-Afr-Med-J* 1990;79:65–67. [PubMed: 1989088]
72. Greipp PR, San Miguel JF, Durie BG, et al. International Staging System for Multiple Myeloma. *J Clin Oncol* 2005;23:3412–3420. [PubMed: 15809451]
73. Rajkumar SV, Greipp PR. Prognostic factors in multiple myeloma. *Hematology Oncology Clinics of North America* 1999;13:1295–1314.
74. Tricot G, Sawyer JR, Jagannath S, et al. Unique role of cytogenetics in the prognosis of patients with myeloma receiving high-dose therapy and autotransplants. *J Clin Oncol* 1997;15:2659–2666. [PubMed: 9215838]
75. Greipp PR. Prognosis in myeloma. *Mayo Clinic Proceedings* 1994;69:895–902. [PubMed: 8065197] [Review] [75 refs]
76. Facon T, Avet-Loiseau H, Guillemin G, et al. Chromosome 13 abnormalities identified by FISH analysis and serum beta2-microglobulin produce a powerful myeloma staging system for patients receiving high-dose therapy. *Blood* 2001;97:1566–1571. [PubMed: 11238092]
77. Greipp PR, Leong T, Bennett JM, et al. Plasmablastic morphology--an independent prognostic factor with clinical and laboratory correlates: Eastern Cooperative Oncology Group (ECOG) myeloma trial E9486 report by the ECOG Myeloma Laboratory Group. *Blood* 1998;91:2501–2507. [PubMed: 9516151]
78. Greipp PR, Lust JA, WM OF, Katzmman JA, Witzig TE, Kyle RA. Plasma cell labeling index and beta 2-microglobulin predict survival independent of thymidine kinase and C-reactive protein in multiple myeloma. *Blood* 1993;81:3382–3387. [PubMed: 8507875]see comments
79. Dispenzieri A, Rajkumar SV, Gertz MA, et al. Treatment of newly diagnosed multiple myeloma based on Mayo stratification of myeloma and risk-adapted therapy (mSMART): Consensus Statement. *Mayo Clin Proc* 2007;82:323–341. [PubMed: 17352369]
80. Blade J, Samson D, Reece D, et al. Criteria for evaluating disease response and progression in patients with multiple myeloma treated by high-dose therapy and haemopoietic stem cell transplantation. Myeloma Subcommittee of the EBMT. European Group for Blood and Marrow Transplant. *Br J Haematol* 1998;102:1115–1123. [PubMed: 9753033]
81. Durie BGM, Harousseau J-L, Miguel JS, et al. International uniform response criteria for multiple myeloma. *Leukemia* 2006;20:1467–1473. [PubMed: 16855634]
82. Singhal S, Mehta J, Desikan R, et al. Antitumor activity of thalidomide in refractory multiple myeloma. *N Engl J Med* 1999;341:1565–1571. [PubMed: 10564685]see comments
83. Richardson PG, Barlogie B, Berenson J, et al. A phase 2 study of bortezomib in relapsed, refractory myeloma. *N Engl J Med* 2003;348:2609–2617. [PubMed: 12826635]

84. Richardson PG, Sonneveld P, Schuster MW, et al. Bortezomib or high-dose dexamethasone for relapsed multiple myeloma. *N Engl J Med* 2005;352:2487–2498. [PubMed: 15958804]see comment
85. Rajkumar SV, Hayman SR, Lacy MQ, et al. Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma. *Blood* 2005;106:4050–4053. [PubMed: 16118317]
86. Richardson PG, Blood E, Mitsiades CS, et al. A randomized phase 2 study of lenalidomide therapy for patients with relapsed or relapsed and refractory multiple myeloma. *Blood* 2006;108:3458–3464.10.1182/blood-2006-04-015909 [PubMed: 16840727]
87. Lenz W. A short history of thalidomide embryopathy. *Teratology* 1988;38:203–215. [PubMed: 3067415]
88. Lenz W. A personal perspective on the thalidomide tragedy. *Teratology* 1992;46:417–418. [PubMed: 1462245]
89. Stephens, T.; Brynner, R. *Dark Remedy. The impact of thalidomide and its revival as a vital medicine.* Cambridge, MA: Perseus Publishing; 2001.
90. Lenz W. Thalidomide and congenital abnormalities. *Lancet* 1962;1:45.
91. McBride WG. Thalidomide and congenital abnormalities. *Lancet* 1961;2:1358.
92. Geraghty, K. Profile of a role model: Francis Oldham Kelsey, MD, PhD. Vol. 2003. American Medical Association; 2001. <http://www.ama-assn.org/ama/pub/printcat/5335.html>
93. Rogerson G. Thalidomide and congenital abnormalities. *Lancet* 1962;1:691.
94. Woodyatt PB. Thalidomide. *Lancet* 1962;1:750.
95. Grabstad H, Golbey R. Clinical experience with thalidomide in patients with cancer. *Clin Pharmacol Therap* 1965;6:298–302. [PubMed: 14296026]
96. Olson KB, Hall TC, Horton J, Khung CL, Hosley HF. Thalidomide (N-phthaloylglutamide) in the treatment of advanced cancer. *Clin Pharmacol Ther* 1965;6:292–297. [PubMed: 14296025]
97. Sheskin J. Thalidomide in the treatment of lepra reactions. *Clin Pharmacol Ther* 1965;6:303–306. [PubMed: 14296027]
98. Iyer CG, Languillon J, Ramanujam K, et al. WHO co-ordinated short-term double-blind trial with thalidomide in the treatment of acute lepra reactions in male lepromatous patients. *Bulletin of the World Health Organization* 1971;45:719–732. [PubMed: 4947831]
99. Mascaro JM, Lecha M, Torras H. Thalidomide in the treatment of recurrent, necrotic, and giant mucocutaneous aphthae and aphthosis. *Archives of Dermatology* 1979;115:636–637. [PubMed: 443847]
100. Lim SH, McWhannell A, Vora AJ, Boughton BJ. Successful treatment with thalidomide of acute graft-versus-host disease after bone-marrow transplantation. *Lancet* 1988;1:117. [PubMed: 2891954]
101. Saurat JH, Camenzind M, Helg C, Chapuis B. Thalidomide for graft-versus-host disease after bone marrow transplantation. *Lancet* 1988;1:359. [PubMed: 2893167]
102. Vogelsang GB, Santos GW, Colvin OM, Chen T. Thalidomide for graft-versus-host disease. *Lancet* 1988;1:827. [PubMed: 2895345]
103. Vogelsang GB, Wells MC, Santos GW, Chen TL, Hess AD. Combination low-dose thalidomide and cyclosporine prophylaxis for acute graft-versus-host disease in a rat mismatched model. *Transplantation Proceedings* 1988;20:226–228. [PubMed: 3284069]
104. Ringden O, Aschan J, Westerberg L. Thalidomide for severe acute graft-versus-host disease. *Lancet* 1988;2:568. [PubMed: 2900947]
105. Youle M, Clarbourn J, Farthing C, et al. Treatment of resistant aphthous ulceration with thalidomide in patients positive for HIV antibody. *BMJ* 1989;298:432. [PubMed: 2495052]
106. Reyes-Teran G, Sierra-Madero JG, Martinez del Cerro V, et al. Effects of thalidomide on HIV-associated wasting syndrome: a randomized, double-blind, placebo-controlled clinical trial. *Aids* 1996;10:1501–1507. [PubMed: 8931784]
107. Zeldis JB, Williams BA, Thomas SD, Elsayed ME. S.T.E.P.S.: a comprehensive program for controlling and monitoring access to thalidomide. *Clinical Therapeutics* 1999;21:319–330. [PubMed: 10211535]

108. Lary JM, Daniel KL, Erickson JD, Roberts HE, Moore CA. The return of thalidomide: can birth defects be prevented? *Drug Safety* 1999;21:161–169. [PubMed: 10487395]
109. Eriksson T, Bjorkman S, Hoglund P. Clinical pharmacology of thalidomide. *European Journal of Clinical Pharmacology* 2001;57:365–376. [PubMed: 11599654]
110. Stirling DI. Pharmacology of thalidomide. *Semin Hematol* 2000;37:5–14. [PubMed: 10676917]
111. Eriksson T, Bjorkman S, Roth B, Fyge A, Hoglund P. Stereospecific determination, chiral inversion in vitro and pharmacokinetics in humans of the enantiomers of thalidomide. *Chirality* 1995;7:44–52. [PubMed: 7702998]
112. D'Amato RJ, Loughnan MS, Flynn E, Folkman J. Thalidomide is an inhibitor of angiogenesis. *Proc Natl Acad Sci USA* 1994;91:4082–4085. [PubMed: 7513432]
113. Kenyon BM, Browne F, D'Amato RJ. Effects of thalidomide and related metabolites in a mouse corneal model of neovascularization. *Experimental Eye Research* 1997;64:971–978. [PubMed: 9301478]
114. Barlogie B, Spencer T, Tricot G, et al. Long term follow up of 169 patients receiving a phase II trial of single agent thalidomide for advanced and refractory multiple myeloma (MM). *Blood* 2000;96:514a. [PubMed: 10887113]A2213
115. Rajkumar SV, Fonseca R, Dispenzieri A, et al. Thalidomide in the treatment of relapsed multiple myeloma. *Mayo Clin Proc* 2000;75:897–902. [PubMed: 10994824]
116. Kumar S, Gertz MA, Dispenzieri A, et al. Response rate, durability of response, and survival after thalidomide therapy for relapsed multiple myeloma. *Mayo Clinic Proceedings* 2003;78:34–39. [PubMed: 12528875]comment
117. Rajkumar SV, Witzig TE. A review of angiogenesis and anti-angiogenic therapy with thalidomide in multiple myeloma. *Cancer Treat Rev* 2000;26:351–362. [PubMed: 11006136]
118. Rajkumar SV, Hayman S, Gertz MA, et al. Combination therapy with thalidomide plus dexamethasone for newly diagnosed myeloma. *J Clin Oncol* 2002;20:4319–4323. [PubMed: 12409330]
119. Rajkumar SV, Dispenzieri A, Fonseca R, et al. Thalidomide for previously untreated indolent or smoldering multiple myeloma. *Leukemia* 2001;15:1274–1276. [PubMed: 11480571]
120. Rajkumar SV, Gertz MA, Lacy MQ, et al. Thalidomide as initial therapy for early-stage myeloma. *Leukemia* 2003;17:775–779. [PubMed: 12682636]
121. Weber D, Rankin K, Gavino M, Delasalle K, Alexanian R. Thalidomide alone or with dexamethasone for previously untreated multiple myeloma. *J Clin Oncol* 2003;21:16–19. [PubMed: 12506164]
122. Rajkumar SV, Gertz MA, Kyle RA, Greipp PR. Current therapy for multiple myeloma. *Mayo Clin Proc* 2002;77:813–822. [PubMed: 12173715]
123. Anagnostopoulos A, Weber D, Rankin K, Delasalle K, Alexanian R. Thalidomide and dexamethasone for resistant multiple myeloma. *Br J Haematol* 2003;121:768–771. [PubMed: 12780791]
124. Dimopoulos MA, Hamilos G, Zomas A, et al. Pulsed cyclophosphamide, thalidomide and dexamethasone: an oral regimen for previously treated patients with multiple myeloma. *Hematology Journal* 2004;5:112–117. [PubMed: 15048060]
125. Garcia-Sanz R, Gonzalez-Fraile MI, Sierra M, Lopez C, Gonzalez M, San Miguel JF. The combination of thalidomide, cyclophosphamide and dexamethasone (ThaCyDex) is feasible and can be an option for relapsed/refractory multiple myeloma. *Hematology Journal* 2002;3:43–48. [PubMed: 11960395]
126. Kropff MH, Lang N, Bisping G, et al. Hyperfractionated cyclophosphamide in combination with pulsed dexamethasone and thalidomide (HyperCDT) in primary refractory or relapsed multiple myeloma. *Br J Haematol* 2003;122:607–616. [PubMed: 12899716]
127. Coleman M, Leonard J, Lyons L, et al. BLT-D (clarithromycin [Biaxin], low-dose thalidomide, and dexamethasone) for the treatment of myeloma and Waldenstrom's macroglobulinemia. *Leukemia & Lymphoma* 2002;43:1777–1782. [PubMed: 12685831]
128. Ghobrial IM, Rajkumar SV. Management of thalidomide toxicity. *J Support Oncol* 2003;1:194–205. [PubMed: 15334875]

129. Osman K, Comenzo R, Rajkumar SV. Deep vein thrombosis and thalidomide therapy for multiple myeloma. *N Engl J Med* 2001;344:1951–1952. [PubMed: 11419443]
130. Zangari M, Anaissie E, Barlogie B, et al. Increased risk of deep-vein thrombosis in patients with multiple myeloma receiving thalidomide and chemotherapy. *Blood* 2001;98:1614–1615. [PubMed: 11520815]
131. Zangari M, Siegel E, Barlogie B, et al. Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy. *Blood* 2002;100:1168–1171. [PubMed: 12149193]
132. Zangari M, Saghafifar F, Anaissie E, et al. Activated protein C resistance in the absence of factor V Leiden mutation is a common finding in multiple myeloma and is associated with an increased risk of thrombotic complications. *Blood Coagulation & Fibrinolysis* 2002;13:187–192. [PubMed: 11943931]
133. Rajkumar SV, Gertz MA, Witzig TE. Life-threatening toxic epidermal necrolysis with thalidomide therapy for myeloma. *N Engl J Med* 2000;343:972–973. [PubMed: 11012329]
134. Badros AZ, Siegel E, Bodenner D, et al. Hypothyroidism in patients with multiple myeloma following treatment with thalidomide. *American Journal of Medicine* 2002;112:412–413. [PubMed: 11904117]
135. Barlogie B, Desikan R, Eddlemon P, et al. Extended survival in advanced and refractory multiple myeloma after single-agent thalidomide: identification of prognostic factors in a phase 2 study of 169 patients. *Blood* 2001;98:492–494. [PubMed: 11435324]
136. Dimopoulos MA, Anagnostopoulos A, Weber D. Treatment of plasma cell dyscrasias with thalidomide and its derivatives. *J Clin Oncol* 2003;21:4444–4454. [PubMed: 14645435]
137. Adams J. Development of the proteasome inhibitor PS-341. *Oncologist* 2002;7:9–16. [PubMed: 11854543]
138. Adams J. Proteasome inhibition in cancer: development of PS-341. *Semin Oncol* 2001;28:613–619. [PubMed: 11740819]
139. Adams J. Proteasome inhibitors as new anticancer drugs. *Curr Opin Oncol* 2002;14:628–634. [PubMed: 12409653]
140. Ciechanover A. The ubiquitin-proteasome proteolytic pathway. *Cell* 1994;79:13–21. [PubMed: 7923371]
141. Adams J, Palombella VJ, Elliott PJ. Proteasome inhibition: a new strategy in cancer treatment. *Invest New Drugs* 2000;18:109–121. [PubMed: 10857991]
142. Adams J. Preclinical and clinical evaluation of proteasome inhibitor PS-341 for the treatment of cancer. *Curr Opin Chem Biol* 2002;6:493–500.
143. Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem* 1998;67:425–479. [PubMed: 9759494]
144. Varshavsky A. The ubiquitin system. *Trend Biochem Sci* 1997;22:383–387. [PubMed: 9357313]
145. Craiu A, Gaczynska M, Akopian T, et al. Lactacystin and clasto-lactacystin beta-lactone modify multiple proteasome beta-subunits and inhibit intracellular protein degradation and major histocompatibility complex class I antigen presentation. *J Biol Chem* 1997;272:13437–13445. [PubMed: 9148969]
146. Orłowski M. The multicatalytic proteinase complex, a major extralysosomal proteolytic system. *Biochemistry* 1990;29:10289–10297. [PubMed: 2175651]
147. Coux O, Tanaka K, Goldberg AL. Structure and functions of the 20S and 26S proteasomes. *Annu Rev Biochem* 1996;65:801–847. [PubMed: 8811196]
148. Almond JB, Cohen GM. The proteasome: a novel target for cancer chemotherapy. *Leukemia* 2002;16:433–443. [PubMed: 11960320]
149. Braun BC, Glickman M, Kraft R, et al. The base of the proteasome regulatory particle exhibits chaperone-like activity. *Nature Cell Biology* 1999;1:221–226.
150. Kloetzel PM. Antigen processing by the proteasome. *Nature Rev Mol Cell Biol* 2001;2:179–187. [PubMed: 11265247]
151. Pickart CM, VanDemark AP. Opening doors into the proteasome. *Nature Struct Biol* 2000;7:999–1001. [PubMed: 11062549]comment

152. Groll M, Bajorek M, Kohler A, et al. A gated channel into the proteasome core particle. *Nature Struct Biol* 2000;7:1062–1067. [PubMed: 11062564]comment
153. Lowe J, Stock D, Jap B, Zwickl P, Baumeister W, Huber R. Crystal structure of the 20S proteasome from the archaeon *T. acidophilum* at 3.4 Å resolution. *Science* 1995;268:533–539. [PubMed: 7725097]comment
154. Adams J, Palombella VJ, Sausville EA, et al. Proteasome inhibitors: a novel class of potent and effective antitumor agents. *Cancer Res* 1999;59:2615–2622. [PubMed: 10363983]
155. Orlowski RZ, Eswara JR, Lafond-Walker A, Grever MR, Orlowski M, Dang CV. Tumor growth inhibition induced in a murine model of human Burkitt's lymphoma by a proteasome inhibitor. *Cancer Res* 1998;58:4342–4348. [PubMed: 9766662]
156. Drexler HC. Activation of the cell death program by inhibition of proteasome function. *Proc Natl Acad Sci USA* 1997;94:855–860. [PubMed: 9023346]
157. Delic J, Masdehors P, Omura S, et al. The proteasome inhibitor lactacystin induces apoptosis and sensitizes chemo- and radioresistant human chronic lymphocytic leukaemia lymphocytes to TNF- α -initiated apoptosis. *Br J Cancer* 1998;77:1103–1107. [PubMed: 9569046]comment
158. An B, Goldfarb RH, Siman R, Dou QP. Novel dipeptidyl proteasome inhibitors overcome Bcl-2 protective function and selectively accumulate the cyclin-dependent kinase inhibitor p27 and induce apoptosis in transformed, but not normal, human fibroblasts. *Cell Death Diff* 1998;5:1062–1075.
159. Traenckner EB, Wilk S, Baeuerle PA. A proteasome inhibitor prevents activation of NF- κ B and stabilizes a newly phosphorylated form of I κ B- α that is still bound to NF- κ B. *EMBO J* 1994;13:5433–5441. [PubMed: 7957109]
160. Mori S, Tanaka K, Omura S, Saito Y. Degradation process of ligand-stimulated platelet-derived growth factor beta-receptor involves ubiquitin-proteasome proteolytic pathway. *J Biol Chem* 1995;270:29447–29452. [PubMed: 7493983]
161. Cusack JC Jr, Liu R, Houston M, et al. Enhanced chemosensitivity to CPT-11 with proteasome inhibitor PS-341: implications for systemic nuclear factor- κ B inhibition. *Cancer Res* 2001;61:3535–3540. [PubMed: 11325813]
162. Russo SM, Tepper JE, Baldwin AS Jr, et al. Enhancement of radiosensitivity by proteasome inhibition: implications for a role of NF- κ B. *Int J Rad Oncol Biol Phys* 2001;50:183–193.
163. Shah SA, Potter MW, McDade TP, et al. 26S proteasome inhibition induces apoptosis and limits growth of human pancreatic cancer. *J Cell Biochem* 2001;82:110–122. [PubMed: 11400168]
164. Teicher BA, Ara G, Herbst R, Palombella VJ, Adams J. The proteasome inhibitor PS-341 in cancer therapy. *Clin Cancer Res* 1999;5:2638–2645. [PubMed: 10499643]
165. Frankel A, Man S, Elliott P, Adams J, Kerbel RS. Lack of multicellular drug resistance observed in human ovarian and prostate carcinoma treated with the proteasome inhibitor PS-341. *Clin Cancer Res* 2000;6:3719–3728. [PubMed: 10999766]
166. Bold RJ, Virudachalam S, McConkey DJ. Chemosensitization of pancreatic cancer by inhibition of the 26S proteasome. *J Surg Res* 2001;100:11–17. [PubMed: 11516199]
167. Hideshima T, Richardson P, Chauhan D, et al. The proteasome inhibitor PS-341 inhibits growth, induces apoptosis, and overcomes drug resistance in human multiple myeloma cells. *Cancer Res* 2001;61:3071–3076. [PubMed: 11306489]
168. Mitsiades N, Mitsiades CS, Poulaki V, et al. Molecular sequelae of proteasome inhibition in human multiple myeloma cells. *Proc Natl Acad Sci USA* 2002;99:14374–14379. [PubMed: 12391322]
169. Mitsiades N, Mitsiades CS, Richardson PG, et al. The proteasome inhibitor PS-341 potentiates sensitivity of multiple myeloma cells to conventional chemotherapeutic agents: therapeutic applications. *Blood* 2003;101:2377–2380. [PubMed: 12424198]
170. LeBlanc R, Catley LP, Hideshima T, et al. Proteasome inhibitor PS-341 inhibits human myeloma cell growth in vivo and prolongs survival in a murine model. *Cancer Res* 2002;62:4996–5000. [PubMed: 12208752]
171. Berenson JR, Jagannath S, Barlogie B, et al. Experience with long-term therapy using the proteasome inhibitor, bortezomib, in advanced multiple myeloma (MM). *Proc Am Soc Clin Oncol* 2003;22:581.
172. Richardson PG, Schlossman RL, Weller E, et al. Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma. *Blood* 2002;100:3063–3067. [PubMed: 12384400]

173. Zangari M, Tricot G, Zeldis J, Eddlemon P, Saghafifar F, Barlogie B. Results of Phase I Study of CC-5013 for the Treatment of Multiple Myeloma (MM) Patients Who Relapse after High Dose Chemotherapy (HDCT). *Blood* 2001;775a.A3226
174. Dimopoulos MA, Spencer A, Attal M, et al. Study of Lenalidomide Plus Dexamethasone Versus Dexamethasone Alone in Relapsed or Refractory Multiple Myeloma (MM): Results of a Phase 3 Study (MM-010). *Blood* 2005;106:6.
175. Weber DM, Chen C, Niesvizky R, et al. Lenalidomide plus high-dose dexamethasone provides improved overall survival compared to high-dose dexamethasone alone for relapsed or refractory multiple myeloma (MM): Results of a North American phase III study (MM-009). *Proc Am Soc Clin Oncol* 2006;24:A7521.
176. Orlowski RZ, Nagler A, Sonneveld P, et al. Randomized Phase III Study of Pegylated Liposomal Doxorubicin Plus Bortezomib Compared With Bortezomib Alone in Relapsed or Refractory Multiple Myeloma: Combination Therapy Improves Time to Progression. *J Clin Oncol* 2007;25:3892–3901. [PubMed: 17679727]
177. Streetly M, Jones RW, Knight R, et al. An Update of the Use and Outcomes of the New Immunomodulatory Agent CC-4047 (Actimid) in Patients with Relapsed / Refractory Myeloma. *Blood* 2003;102:236a.
178. Lacy MQ, Hayman SR, Gertz MA, et al. Pomalidomide (CC4047) Plus Low-Dose Dexamethasone (Pom/dex) Is Highly Effective Therapy in Relapsed Multiple Myeloma. *Blood (ASH Annual Meeting Abstracts)* 2008;112:866.
179. Jagannath S, Vij R, Stewart AK, et al. Initial Results of PX-171-003, An Open-Label, Single-Arm, Phase II Study of Carfilzomib (CFZ) in Patients with Relapsed and Refractory Multiple Myeloma (MM). *Blood (ASH Annual Meeting Abstracts)* 2008;112:864.
180. Vij R, Wang M, Orlowski R, et al. Initial Results of PX-171-004, An Open-Label, Single-Arm, Phase II Study of Carfilzomib (CFZ) in Patients with Relapsed Myeloma (MM). *Blood (ASH Annual Meeting Abstracts)* 2008;112:865.
181. Hideshima T, Chauhan D, Podar K, Schlossman RL, Richardson P, Anderson KC. Novel therapies targeting the myeloma cell and its bone marrow microenvironment. *Semin Oncol* 2001;28:607–612. [PubMed: 11740818]
182. Musto P, Petrucci MT, Brinchen S, et al. A multicenter, randomized clinical trial comparing zoledronic acid versus observation in patients with asymptomatic myeloma. *Cancer* 2008;113:1588–1595. [PubMed: 18683218]
183. Cavo M, Zamagni E, Tosi P, et al. Superiority of thalidomide and dexamethasone over vincristine-doxorubicin-dexamethasone (VAD) as primary therapy in preparation for autologous transplantation for multiple myeloma. *Blood* 2005;106:35–39. [PubMed: 15761019]
184. Fermand J-P, Jaccard A, Macro M, et al. A Randomized Comparison of Dexamethasone + Thalidomide (Dex/Thal) vs Dex + Placebo (Dex/P) in Patients (pts) with Relapsing Multiple Myeloma (MM). *Blood* 2006;108:3563.
185. Rajkumar SV, Blood E, Vesole DH, Fonseca R, Greipp PR. Phase III Clinical Trial of Thalidomide Plus Dexamethasone Compared With Dexamethasone Alone in Newly Diagnosed Multiple Myeloma: A Clinical Trial Coordinated by the Eastern Cooperative Oncology Group. *J Clin Oncol* 2006;24:431–436. [PubMed: 16365178]
186. Weber DM, Gavino M, Delasalle K, Rankin K, Giral S, Alexanian R. Thalidomide alone or with dexamethasone for multiple myeloma. *Blood* 1999;94(Suppl 1):604a.A 2686
187. Cavo M, Zamagni E, Tosi P, et al. First-line therapy with thalidomide and dexamethasone in preparation for autologous stem cell transplantation for multiple myeloma. *Haematologica* 2004;89:826–831. [PubMed: 15257934]
188. Rajkumar SV, Rosiñol L, Hussein M, et al. A Multicenter, Randomized, Double-Blind, Placebo-Controlled Study of Thalidomide plus Dexamethasone Versus Dexamethasone as Initial Therapy for Newly Diagnosed Multiple Myeloma. *J Clin Oncol* 2008;26:2171–2177. [PubMed: 18362366]
189. Lokhorst H, van der Holt B, Zweegman S, et al. Final Analysis of HOVON-50 Randomized Phase III Study on the Effect of Thalidomide Combined with Adriamycin, Dexamethasone (AD) and High Dose Melphalan (HDM) in Patients with Multiple Myeloma (MM). *ASH Annual Meeting Abstracts* 2008;112:157.

190. Lacy M, Gertz M, Dispenzieri A, et al. Lenalidomide Plus Dexamethasone (Rev/Dex) in Newly Diagnosed Myeloma: Response to Therapy, Time to Progression, and Survival. *Blood* 2006;108:798.
191. Rajkumar SV, Jacobus S, Callander N, Fonseca R, Vesole D, Greipp P. A Randomized Phase III Trial of Lenalidomide Plus High-Dose Dexamethasone Versus Lenalidomide Plus Low-Dose Dexamethasone in Newly Diagnosed Multiple Myeloma (E4A03): A Trial Coordinated by the Eastern Cooperative Oncology Group. *Blood* 2006;108:799.
192. Richardson PG, Chanan-Khan A, Schlossman RL, et al. Phase II Trial of Single Agent Bortezomib (VELCADE®) in Patients with Previously Untreated Multiple Myeloma (MM). *Blood* 2004;104:100a. [PubMed: 15026315]A336
193. Jagannath S, Durie BG, Wolf J, et al. Bortezomib therapy alone and in combination with dexamethasone for previously untreated symptomatic multiple myeloma. *Br J Haematol* 2005;129:776–783. [PubMed: 15953004]
194. Harousseau J, Attal M, Leleu X, et al. Bortezomib plus dexamethasone as induction treatment prior to autologous stem cell transplantation in patients with newly diagnosed multiple myeloma: results of an IFM phase II study. *Haematologica* 2006;91:1498–1505. [PubMed: 17043025]
195. Harousseau J-L, Marit G, Caillot D, et al. VELCADE/Dexamethasone (Vel/Dex) Versus VAD as Induction Treatment Prior to Autologous Stem Cell Transplantation (ASCT) in Newly Diagnosed Multiple Myeloma (MM): An Interim Analysis of the IFM 2005-01 Randomized Multicenter Phase III Trial. *Blood* 2006;108:56.
196. Barlogie B, Anaissie E, van Rhee F, et al. Incorporating bortezomib into upfront treatment for multiple myeloma: early results of total therapy 3. *Br J Haematol* 2007;138:176–185. [PubMed: 17593024]
197. Palumbo A, Bringhen S, Caravita T, et al. Oral melphalan and prednisone chemotherapy plus thalidomide compared with melphalan and prednisone alone in elderly patients with multiple myeloma: randomised controlled trial. *Lancet* 2006;367:825–831. [PubMed: 16530576]
198. Facon T, Mary JY, Hulin C, et al. Major Superiority of Melphalan - Prednisone (MP) + Thalidomide (THAL) over MP and Autologous Stem Cell Transplantation in the Treatment of Newly Diagnosed Elderly Patients with Multiple Myeloma. *Blood* 2005;106:230a.A780
199. Facon T, Mary JY, Hulin C, et al. Melphalan and prednisone plus thalidomide versus melphalan and prednisone alone or reduced-intensity autologous stem cell transplantation in elderly patients with multiple myeloma (IFM 99-06): a randomised trial. *Lancet* 2007;370:1209–1218. [PubMed: 17920916]
200. Hulin C, Virion J, Leleu X, et al. Comparison of melphalan-prednisone-thalidomide (MP-T) to melphalan-prednisone (MP) in patients 75 years of age or older with untreated multiple myeloma (MM). Preliminary results of the randomized, double-blind, placebo controlled IFM 01-01 trial. *J Clin Oncol (Meeting Abstracts)* 2007;25:8001.
201. Wijermans P, Schaafsma M, van Norden Y, et al. Melphalan + Prednisone Versus Melphalan + Prednisone + Thalidomide in Induction Therapy for Multiple Myeloma in Elderly Patients: Final Analysis of the Dutch Cooperative Group HOVON 49 Study. *Blood (ASH Annual Meeting Abstracts)* 2008;112:649.
202. Waage A, Gimsing P, Juliusson G, Turesson I, Fayers P. Melphalan-Prednisone-Thalidomide to Newly Diagnosed Patients with Multiple Myeloma: A Placebo Controlled Randomised Phase 3 Trial. *ASH Annual Meeting Abstracts* 2007;110:78.
203. Palumbo A, Bringhen S, Liberati AM, et al. Oral melphalan, prednisone, and thalidomide in elderly patients with multiple myeloma: updated results of a randomized controlled trial. *Blood* 2008;112:3107–3114. [PubMed: 18505783]
204. San Miguel JF, Schlag R, Khuageva NK, et al. Bortezomib plus Melphalan and Prednisone for Initial Treatment of Multiple Myeloma. *N Engl J Med* 2008;359:906–917. [PubMed: 18753647]
205. Palumbo A, Falco P, Corradini P, et al. Melphalan, Prednisone, and Lenalidomide Treatment for Newly Diagnosed Myeloma: A Report From the GIMEMA Italian Multiple Myeloma Network. *J Clin Oncol* 2007;25:4459–4465. [PubMed: 17785703]

206. Attal M, Harousseau JL, Stoppa AM, et al. A prospective, randomized trial of autologous bone marrow transplantation and chemotherapy in multiple myeloma. Intergroupe Francais du Myelome. *N Engl J Med* 1996;335:91–97. [PubMed: 8649495]
207. Child JA, Morgan GJ, Davies FE, et al. High-dose chemotherapy with hematopoietic stem-cell rescue for multiple myeloma. *N Engl J Med* 2003;348:1875–1883. [PubMed: 12736280]
208. Blade J, Vesole DH, Gertz M. Transplantation for multiple myeloma: who, when, how often? *Blood* 2003;102:3469–3477. [PubMed: 12893762]
209. Kumar A, Loughran T, Alsina M, Durie BG, Djulbegovic B. Management of multiple myeloma: a systematic review and critical appraisal of published studies. *Lancet Oncology* 2003;4:293–304. [PubMed: 12732167]
210. Moreau P, Facon T, Attal M, et al. Comparison of 200 mg/m² melphalan and 8 Gy total body irradiation plus 140 mg/m² melphalan as conditioning regimens for peripheral blood stem cell transplantation in patients with newly diagnosed multiple myeloma: final analysis of the Intergroupe Francophone du Myélome 9502 randomized trial. *Blood* 2002;99:731–735. [PubMed: 11806971]
211. Bensinger W, Giralt S, Holmberg L, et al. 166H0-DOTMP and high-dose melphalan before autologous peripheral blood stem cell transplantation in patients with multiple myeloma. *Hematol J* 2003;4(Suppl 1):S215–216.
212. Dispenzieri A, Wiseman GA, Lacy MQ, et al. A Phase II Study of High Dose 153-Samarium EDTMP (153-Sm EDTMP) and Melphalan for Peripheral Stem Cell Transplantation (PBSCT) in Multiple Myeloma (MM). *Blood* 2003;102:982a.
213. Femand JP, Ravaud P, Chevret S, et al. High-dose therapy and autologous peripheral blood stem cell transplantation in multiple myeloma: up-front or rescue treatment? Results of a multicenter sequential randomized clinical trial. *Blood* 1998;92:3131–3136. [PubMed: 9787148]
214. Facon T, Mary JY, Harousseau JL, et al. Front-line or rescue autologous bone marrow transplantation (ABMT) following a first course of high dose melphalan (HDM) in multiple myeloma (MM). Preliminary results of a prospective randomized trial (CIAM) protocol. *Blood* 1996;88(Suppl1): 685a.
215. Barlogie B, Kyle R, Anderson K, et al. Comparable Survival in Multiple Myeloma (MM) with High Dose Therapy (HDT) Employing MEL 140 mg/m² + TBI 12 Gy Autotransplants Versus Standard Dose Therapy with VBMCP and No Benefit from Interferon (IFN) Maintenance: Results of Intergroup Trial S9321. *Blood* 2003;102:42a.
216. Blade J, Sureda A, Ribera JM, et al. High-Dose Therapy Autotransplantation/Intensification Versus Continued Conventional Chemotherapy in Multiple Myeloma Patients Responding to Initial Chemotherapy. Definitive Results from PETHEMA after a Median Follow-Up of 66 Months. *Blood* 2003;102:42a.
217. Rajkumar SV, Fonseca R, Lacy MQ, et al. Autologous stem cell transplantation for relapsed and primary refractory myeloma. *Bone Marrow Transplantation* 1999;23:1267–1272. [PubMed: 10414914]
218. Blade J, Esteve J. Treatment approaches for relapsing and refractory multiple myeloma. *Acta Oncologica* 2000;39:843–847. [PubMed: 11145443][Review] [44 refs]
219. Barlogie B, Jagannath S, Vesole DH, et al. Superiority of tandem autologous transplantation over standard therapy for previously untreated multiple myeloma. *Blood* 1997;89:789–793. [PubMed: 9028309]
220. Barlogie B, Jagannath S, Desikan KR, et al. Total therapy with tandem transplants for newly diagnosed multiple myeloma. *Blood* 1999;93:55–65. [PubMed: 9864146]
221. Attal M, Harousseau JL, Facon T, et al. Double Autologous Transplantation Improves Survival of Multiple Myeloma Patients: Final Analysis of a Prospective Randomized Study of the “Intergroupe Francophone du Myelome” (IFM 94). *Blood* 2002;100:5a.
222. Cavo M, Cellini C, Zamagni E, et al. Superiority of Double over Single Autologous Stem Cell Transplantation as First-Line Therapy for Multiple Myeloma. *Blood* 2004;104:155a.A536
223. Femand JP, Alberti C, Marolleau JP. Single versus tandem high dose therapy (HDT) supported with autologous blood stem cell (ABSC) transplantation using unselected or CD34-enriched ABSC: results of a two by two designed randomized trial in 230 young patients with multiple myeloma (MM). *Hematol J* 2003;4(Suppl 1):S59.

224. Goldschmidt H. Single vs. tandem autologous transplantation in multiple myeloma: the GMMG experience. *Hematol J* 2003;4(Suppl1):S61.
225. Sirohi B, Powles R, Singhal S, et al. High-Dose Melphalan and Second Autografts for Myeloma Relapsing after One Autograft: Results Equivalent to Tandem Autotransplantation. *Blood* 2001;98:402a.A1690
226. Mehta J, Singhal S. Graft-versus-myeloma. *Bone Marrow Transplantation* 1998;22:835–843. [PubMed: 9827810]
227. Gahrton G, Svensson H, Cavo M, et al. Progress in allogeneic bone marrow and peripheral blood stem cell transplantation for multiple myeloma: a comparison between transplants performed 1983–93 and 1994–8 at European Group for Blood and Marrow Transplantation centres. *Br J Haematol* 2001;113:209–216. [PubMed: 11360893]
228. Einsele H, Schafer HJ, Hebart H, et al. Follow-up of patients with progressive multiple myeloma undergoing allografts after reduced-intensity conditioning. *Br J Haematol* 2003;121:411–418. [PubMed: 12716363]
229. Crawley C, Lalancette M, Szydlo R, et al. Outcomes for reduced-intensity allogeneic transplantation for multiple myeloma: an analysis of prognostic factors from the Chronic Leukaemia Working Party of the EBMT. *Blood* 2005;105:4532–4539. [PubMed: 15731182]
230. Maloney DG, Molina AJ, Sahebi F, et al. Allografting with nonmyeloablative conditioning following cytoreductive autografts for the treatment of patients with multiple myeloma. *Blood* 2003;102:3447–3454. [PubMed: 12855572]
231. Garban F, Attal M, Michallet M, et al. Prospective comparison of autologous stem cell transplantation followed by dose-reduced allograft (IFM99-03 trial) with tandem autologous stem cell transplantation (IFM99-04 trial) in high-risk de novo multiple myeloma. *Blood* 2006;107:3474–3480.10.1182/blood-2005-09-3869 [PubMed: 16397129]
232. Bruno B, Rotta M, Patriarca F, et al. A Comparison of Allografting with Autografting for Newly Diagnosed Myeloma. *N Engl J Med* 2007;356:1110–1120.10.1056/NEJMoa065464 [PubMed: 17360989]
233. Rosinol L, Perez-Simon JA, Sureda A, et al. A prospective PETHEMA study of tandem autologous transplantation versus autograft followed by reduced-intensity conditioning allogeneic transplantation in newly diagnosed multiple myeloma. *Blood* 2008;112:3591–3593. [PubMed: 18612103]
234. Mateos M-V, Hernandez J-M, Hernandez M-T, et al. Bortezomib plus melphalan and prednisone in elderly untreated patients with multiple myeloma: results of a multicenter phase 1/2 study. *Blood* 2006;108:2165–2172. [PubMed: 16772605]
235. Jagannath S, Richardson PG, Sonneveld P, et al. Bortezomib appears to overcome the poor prognosis conferred by chromosome 13 deletion in phase 2 and 3 trials. *Leukemia* 2006;21:151–157. [PubMed: 17096017]
236. Sagaster V, Ludwig H, Kaufmann H, et al. Bortezomib in relapsed multiple myeloma: response rates and duration of response are independent of a chromosome 13q-deletion. *Leukemia* 2006;21:164–168. [PubMed: 17096015]
237. Myeloma Trialists' Collaborative G. Interferon as therapy for multiple myeloma: an individual patient data overview of 24 randomized trials and 4012 patients. *Br J Haematol* 2001;113:1020–1034. [PubMed: 11442498]
238. Berenson JR, Crowley JJ, Grogan TM, et al. Maintenance therapy with alternate-day prednisone improves survival in multiple myeloma patients. *Blood* 2002;99:3163–3168. [PubMed: 11964279]
239. Attal M, Harousseau J-L, Leyvraz S, et al. Maintenance therapy with thalidomide improves survival in patients with multiple myeloma. *Blood* 2006;108:3289–3294.10.1182/blood-2006-05-022962 [PubMed: 16873668]
240. Spencer A, Prince M, Roberts AW, Bradstock KF, Prosser IW. First Analysis of the Australasian Leukaemia and Lymphoma Group (ALLG) Trial of Thalidomide and Alternate Day Prednisolone Following Autologous Stem Cell Transplantation (ASCT) for Patients with Multiple Myeloma (ALLG MM6). *ASH Annual Meeting Abstracts* 2006;108:58.

241. Offidani M, Polloni C, Corvatta L, et al. Thalidomide-Dexamethasone vs Interferon- α -Dexamethasone as Maintenance Treatment after ThaDD Induction for Multiple Myeloma (MM): Final Analysis of a Prospective, Randomized Study. ASH Annual Meeting Abstracts 2007;110:532.
242. Abdelkefi A, Ladeb S, Torjman L, et al. Single autologous stem-cell transplantation followed by maintenance therapy with thalidomide is superior to double autologous transplantation in multiple myeloma: results of a multicenter randomized clinical trial. *Blood* 2008;111:1805–1810. [PubMed: 17875806]
243. Maiolino A, Hungria VT, Oliveira-Duarte G, et al. Thalidomide + Dexamethasone as Maintenance after Single Autologous Stem Cell Transplantation Improves Progression-Free Survival (PFS) in Advanced Multiple Myeloma. A Prospective Brazilian Randomized Trial. ASH Annual Meeting Abstracts 2008;112:3703.
244. Morgan GJ, Jackson GH, Davies FE, et al. Maintenance Thalidomide May Improve Progression Free but Not Overall Survival; Results from the Myeloma IX Maintenance Randomisation. *Blood* (ASH Annual Meeting Abstracts) 2008;112:656.
245. Barlogie B, Tricot G, Anaissie E, et al. Thalidomide and hematopoietic-cell transplantation for multiple myeloma. *N Engl J Med* 2006;354:1021–1030. [PubMed: 16525139]see comment
246. Gertz MA, Lacy MQ, Inwards DJ, et al. Early harvest and late transplantation as an effective therapeutic strategy in multiple myeloma. *Bone Marrow Transplantation* 1999;23:221–226. [PubMed: 10084252]
247. Alexanian R, Barlogie B, Dixon D. High-dose glucocorticoid treatment of resistant myeloma. *Annals of Internal Medicine* 1986;105:8–11. [PubMed: 3717812]
248. Gertz MA, Garton JP, Greipp PR, Witzig TE, Kyle RA. A phase II study of high-dose methylprednisolone in refractory or relapsed multiple myeloma. *Leukemia* 1995;9:2115–2118. [PubMed: 8609725]
249. Gucalp R, Theriault R, Gill I, et al. Treatment of cancer-associated hypercalcemia. Double-blind comparison of rapid and slow intravenous infusion regimens of pamidronate disodium and saline alone. *Arch Intern Med* 1994;154:1935–1944. [PubMed: 8074597]
250. Major P, Lortholary A, Hon J, et al. Zoledronic acid is superior to pamidronate in the treatment of hypercalcemia of malignancy: a pooled analysis of two randomized, controlled clinical trials. *J Clin Oncol* 2001;19:558–567. [PubMed: 11208851]
251. Berenson JR, Lichtenstein A, Porter L, et al. Efficacy of pamidronate in reducing skeletal events in patients with advanced multiple myeloma. Myeloma Aredia Study Group. *N Engl J Med* 1996;334:488–493. [PubMed: 8559201]see comments
252. Berenson JR, Rosen LS, Howell A, et al. Zoledronic acid reduces skeletal-related events in patients with osteolytic metastases. *Cancer* 2001;91:1191–1200. [PubMed: 11283917]erratum appears in *Cancer* 2001 May 15;91(10):1956
253. Rosen LS, Gordon D, Kaminski M, et al. Zoledronic acid versus pamidronate in the treatment of skeletal metastases in patients with breast cancer or osteolytic lesions of multiple myeloma: a phase II, double blind, comparative trial. *Cancer J* 2001;7:377–387. [PubMed: 11693896]
254. Berenson JR, Hillner BE, Kyle RA, et al. American Society of Clinical Oncology Clinical Practice Guidelines: The role of bisphosphonates in multiple myeloma. *J Clin Oncol* 2002;20:3719–3736. [PubMed: 12202673]
255. Marx RE. Pamidronate (Aredia) and zoledronate (Zometa) induced avascular necrosis of the jaws: a growing epidemic. *Journal of Oral & Maxillofacial Surgery* 2003;61:1115–1117. [PubMed: 12966493]see comment
256. Kademani D, Koka S, Lacy MQ, Rajkumar SV. Primary surgical therapy for osteonecrosis of the jaw secondary to bisphosphonate therapy. *Mayo Clinic Proceedings* 2006;81:1100–1103. [PubMed: 16901034]
257. Badros A, Weikel D, Salama A, et al. Osteonecrosis of the Jaw in Multiple Myeloma Patients: Clinical Features and Risk Factors. *J Clin Oncol* 2006;24:945–952.10.1200/JCO.2005.04.2465 [PubMed: 16484704]
258. Durie BG, Katz M, Crowley J. Osteonecrosis of the jaw and bisphosphonates. *N Engl J Med* 2005;353:99–102. [PubMed: 16000365]discussion 199-102

259. Ruggiero SL, Mehrotra B, Rosenberg TJ, Engroff SL. Osteonecrosis of the jaws associated with the use of bisphosphonates: a review of 63 cases. *Journal of Oral & Maxillofacial Surgery* 2004;62:527–534. [PubMed: 15122554]
260. Lacy MQ, Dispenzieri A, Gertz MA, et al. Mayo clinic consensus statement for the use of bisphosphonates in multiple myeloma. *Mayo Clinic Proceedings* 2006;81:1047–1053. [PubMed: 16901028]
261. Dimopoulos MA, Kastritis E, Anagnostopoulos A, et al. Osteonecrosis of the jaw in patients with multiple myeloma treated with bisphosphonates: evidence of increased risk after treatment with zoledronic acid. *Haematologica* 2006;91:968–971. [PubMed: 16757414]
262. Melton LJ 3rd, Kyle RA, Achenbach SJ, Oberg AL, Rajkumar SV. Fracture risk with multiple myeloma: a population-based study. *Journal of Bone & Mineral Research* 2005;20:487–493. [PubMed: 15746994]
263. Fourney DR, Schomer DF, Nader R, et al. Percutaneous vertebroplasty and kyphoplasty for painful vertebral body fractures in cancer patients. *Journal of Neurosurgery* 2003;98:21–30. [PubMed: 12546384]
264. Yussim E, Schwartz E, Sidi Y, Ehrenfeld M. Acute renal failure precipitated by non-steroidal anti-inflammatory drugs (NSAIDs) in multiple myeloma. *Am J Hematol* 1998;58:142–144. [PubMed: 9625583]
265. Wu MJ, Kumar KS, Kulkarni G, Kaiser H. Multiple myeloma in naproxen-induced acute renal failure. *N Engl J Med* 1987;317:170–171. [PubMed: 3600707]
266. Johnson WJ, Kyle RA, Pineda AA, O'Brien PC, Holley KE. Treatment of renal failure associated with multiple myeloma. Plasmapheresis, hemodialysis, and chemotherapy. *Arch Intern Med* 1990;150:863–869. [PubMed: 2183734]
267. Garton JP, Gertz MA, Witzig TE, et al. Epoetin alfa for the treatment of the anemia of multiple myeloma. A prospective, randomized, placebo-controlled, double-blind trial. *Arch Intern Med* 1995;155:2069–2074. [PubMed: 7575066]
268. Dammacco F, Castoldi G, Rodger S. Efficacy of epoetin alfa in the treatment of anaemia of multiple myeloma. *Br J Haematol* 2001;113:172–179. [PubMed: 11328297]erratum appears in *Br J Haematol* 2001 Sep;114(3):738
269. Oken MM, Pomeroy C, Weisdorf D, Bennett JM. Prophylactic antibiotics for the prevention of early infection in multiple myeloma. *American Journal of Medicine* 1996;100:624–628. [PubMed: 8678082]
270. Gertz MA, Kyle RA. Hyperviscosity syndrome. *Journal of Intensive Care Medicine* 1995;10:128–141. [PubMed: 10155178]
271. Rajkumar SV, Kyle RA. Multiple myeloma: diagnosis and treatment. *Mayo Clinic Proceedings* 2005;80:1371–1382. [PubMed: 16212152]see comment
272. Kyle RA, Therneau TM, Rajkumar SV, et al. Long-term follow-up of IgM monoclonal gammopathy of undetermined significance. *Blood* 2003;102:3759–3764. [PubMed: 12881316]
273. Gobbi PG, Baldini L, Broglia C, et al. Prognostic validation of the international classification of immunoglobulin M gammopathies: a survival advantage for patients with immunoglobulin M monoclonal gammopathy of undetermined significance? *Clin Cancer Res* 2005;11:1786–1790. [PubMed: 15756000]
274. Kyle RA, Therneau TM, Rajkumar SV, et al. Long-term follow-up of IgM monoclonal gammopathy of undetermined significance. *Semin Oncol* 2003;30:169–171. [PubMed: 12720130]
275. Baldini L, Goldaniga M, Guffanti A, et al. Immunoglobulin M monoclonal gammopathies of undetermined significance and indolent Waldenstrom's macroglobulinemia recognize the same determinants of evolution into symptomatic lymphoid disorders: proposal for a common prognostic scoring system. *J Clin Oncol* 2005;23:4662–4668. [PubMed: 16034042]
276. Owen RG, Treon SP, Al-Katib A, et al. Clinicopathological definition of Waldenstrom's macroglobulinemia: consensus panel recommendations from the Second International Workshop on Waldenstrom's Macroglobulinemia. *Semin Oncol* 2003;30:110–115. [PubMed: 12720118]
277. Dimopoulos MA, Mouloupoulos LA, Maniatis A, Alexanian R. Solitary plasmacytoma of bone and asymptomatic multiple myeloma. *Blood* 2000;96:2037–2044. [PubMed: 10979944]

278. Dimopoulos MA, Kiamouris C, Moulopoulos LA. Solitary plasmacytoma of bone and extramedullary plasmacytoma. *Hematol Oncol Clin North Am* 1999;13:1249–1257. [PubMed: 10626148]
279. Dispenzieri A, Kyle RA, Lacy MQ, et al. POEMS syndrome: definitions and long-term outcome. *Blood* 2003;101:2496–2506. [PubMed: 12456500]
280. Browman GP, Bergsagel D, Sicheri D, et al. Randomized trial of interferon maintenance in multiple myeloma: a study of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 1995;13:2354–2360. [PubMed: 7666094]
281. Fassas AB, Spencer T, Sawyer J, et al. Both hypodiploidy and deletion of chromosome 13 independently confer poor prognosis in multiple myeloma. *Br J Haematol* 2002;118:1041–1047. [PubMed: 12199783]
282. Barlogie B, Kyle RA, Anderson KC, et al. Standard chemotherapy compared with high-dose chemoradiotherapy for multiple myeloma: final results of phase III US Intergroup Trial S9321. *J Clin Oncol* 2006;24:929–936. [PubMed: 16432076]
283. Attal M, Harousseau JL, Facon T, et al. Single versus double autologous stem-cell transplantation for multiple myeloma. *N Engl J Med* 2003;349:2495–2502. [PubMed: 14695409]see comment
284. Blade J, Rosinol L, Sureda A, et al. High-dose therapy intensification compared with continued standard chemotherapy in multiple myeloma patients responding to the initial chemotherapy: long-term results from a prospective randomized trial from the Spanish cooperative group PETHEMA. *Blood* 2005;106:3755–3759. [PubMed: 16105975]
285. Rajkumar SV, Jacobus S, Callander N, et al. A Randomized Trial of Lenalidomide Plus High-Dose Dexamethasone (RD) Versus Lenalidomide Plus Low-Dose Dexamethasone (Rd) in Newly Diagnosed Multiple Myeloma (E4A03): A Trial Coordinated by the Eastern Cooperative Oncology Group. *ASH Annual Meeting Abstracts* 2007;110:74.
286. Harousseau JL, Mathiot C, Attal M, et al. VELCADE/Dexamethasone (Vel/D) Versus VAD as Induction Treatment Prior to Autologous Stem Cell Transplantation (ASCT) in Newly Diagnosed Multiple Myeloma (MM): Updated Results of the IFM 2005/01 Trial. *ASH Annual Meeting Abstracts* 2007;110:450.
287. Cavo M, Patriarca F, Tacchetti P, et al. Bortezomib (Velcade(R))-Thalidomide-Dexamethasone (VTD) vs Thalidomide-Dexamethasone (TD) in Preparation for Autologous Stem-Cell (SC) Transplantation (ASCT) in Newly Diagnosed Multiple Myeloma (MM). *ASH Annual Meeting Abstracts* 2007;110:73.
288. Harousseau JL, Attal M. The role of stem cell transplantation in multiple myeloma. *Blood Reviews* 2002;16:245–253. [PubMed: 12350367]
289. Fermand JP, Katsahian S, Divine M, et al. High-dose therapy and autologous blood stem-cell transplantation compared with conventional treatment in myeloma patients aged 55 to 65 years: long-term results of a randomized control trial from the Group Myelome-Autogreffe. *J Clin Oncol* 2005;23:9227–9233. [PubMed: 16275936]
290. Palumbo A, Bringhen S, Petrucci MT, et al. Intermediate-dose melphalan improves survival of myeloma patients aged 50 to 70: results of a randomized controlled trial. *Blood* 2004;104:3052–3057. [PubMed: 15265788]
291. Sonneveld P, van der Holt B, Vellenga E, et al. Intensive Versus Double Intensive Therapy in Untreated Multiple Myeloma: Final Analysis of the HOVON 24 Trial. *ASH Annual Meeting Abstracts* 2005;106:2545.

A.

* Bortezomib containing regimens preferred in renal failure

B. **Figure 1.**

Approach to the treatment of newly diagnosed myeloma in patients who are candidates for transplant (A) and those who are not candidates for transplant (B). Footnote for Figure 1A. Rd, lenalidomide plus low dose dexamethasone; CR, complete response; VGPR, very good partial response; ASCT, autologous stem cell transplantation Footnote for Figure 1B. MPT, melphalan, prednisone, thalidomide; VMP, bortezomib, melphalan, prednisone.

Table 1

Diagnostic Criteria for Plasma Cell Disorders

Disorder	Disease Definition	References
Monoclonal gammopathy of undetermined significance (MGUS)	<p>All 3 criteria must be met:</p> <ul style="list-style-type: none"> Serum monoclonal protein < 3gm/dL Clonal bone marrow plasma cells <10%, and Absence of end-organ damage such as hypercalcemia, renal insufficiency, anemia, and bone lesions (CRAB) that can be attributed to the plasma cell proliferative disorder 	38
Smoldering multiple myeloma (also referred to as asymptomatic multiple myeloma)	<p>Both criteria must be met:</p> <ul style="list-style-type: none"> Serum monoclonal protein (IgG or IgA) ≥ 3gm/dL and/or clonal bone marrow plasma cells $\geq 10\%$, and Absence of end-organ damage such as lytic bone lesions, anemia, hypercalcemia, or renal failure that can be attributed to a plasma cell proliferative disorder 	38
Multiple Myeloma	<p>All 3 criteria must be met except as noted:</p> <ul style="list-style-type: none"> Clonal bone marrow plasma cells $\geq 10\%$ Presence of serum and/or urinary monoclonal protein (except in patients with true non-secretory multiple myeloma), and Evidence of end organ damage that can be attributed to the underlying plasma cell proliferative disorder, specifically <ul style="list-style-type: none"> Hypercalcemia: Serum calcium ≥ 11.5 mg/dL or Renal insufficiency: Serum creatinine > 1.73 mmol/L) Anemia: Normochromic, normocytic with a hemoglobin value of >2 g/dL below the lower limit of normal or a hemoglobin value <10 g/dL Bone lesions: Lytic lesions, severe osteopenia or pathologic fractures 	38-271
Waldenström's macroglobulinemia	<p>Both criteria must be met:</p> <ul style="list-style-type: none"> IgM monoclonal gammopathy (regardless of the size of the M protein), and >10% bone marrow lymphoplasmacytic infiltration (usually intertrabecular) by small lymphocytes that exhibit plasmacytoid or plasma cell differentiation and a typical immunophenotype (eg., surface IgM+, CD5+/-, CD10-, CD19+, CD20+, CD23-) that satisfactorily excludes other lymphoproliferative disorders including chronic lymphocytic leukemia and mantle cell lymphoma. <p>Note: IgM MGUS is defined as</p> <ul style="list-style-type: none"> Serum IgM monoclonal protein <3gm/dL, and bone marrow lymphoplasmacytic infiltration <10%, and No evidence of anemia, constitutional symptoms, hyperviscosity, lymphadenopathy, or hepatosplenomegaly. 	272-276

Disorder	Disease Definition	References
	<p>Smoldering Waldenström's macroglobulinemia (also referred to as indolent or asymptomatic Waldenström's macroglobulinemia) is defined as</p> <ul style="list-style-type: none"> • Serum IgM monoclonal protein ≥ 3 gm/dL and/or bone marrow lymphoplasmacytic infiltration $\geq 10\%$, and • No evidence of end-organ damage such as anemia, constitutional symptoms, hyperviscosity, lymphadenopathy, or hepatosplenomegaly that can be attributed to a lympho-plasma cell proliferative disorder. 	
Solitary Plasmacytoma	<p>All 4 criteria must be met</p> <ul style="list-style-type: none"> • Biopsy proven solitary lesion of bone or soft tissue with evidence of clonal plasma cells • Normal bone marrow with no evidence of clonal plasma cells • Normal skeletal survey and MRI of spine and pelvis (except for the primary solitary lesion) • Absence of end-organ damage such as hypercalcemia, renal insufficiency, anemia, or bone lesions (CRAB) that can be attributed to a lympho-plasma cell proliferative disorder 	277-278
Systemic AL Amyloidosis	<p>All 4 criteria must be met:</p> <ul style="list-style-type: none"> • Presence of an amyloid-related systemic syndrome (such as renal, liver, heart, gastrointestinal tract, or peripheral nerve involvement) • Positive amyloid staining by Congo red in any tissue (e.g., fat aspirate, bone marrow, or organ biopsy) • Evidence that amyloid is light-chain related established by direct examination of the amyloid (immunohistochemical staining, direct sequencing, etc), and • Evidence of a monoclonal plasma cell proliferative disorder (serum or urine M protein, abnormal free light chain ratio, or clonal plasma cells in the bone marrow). <p>Note: Approximately 2-3 percent of patients with AL amyloidosis will not meet the requirement for evidence of a monoclonal plasma cell disorder listed above; the diagnosis of AL amyloidosis must be made with caution in these patients.</p>	39
POEMS Syndrome	<p>All 3 criteria must be met:</p> <ul style="list-style-type: none"> • Presence of a monoclonal plasma cell disorder • Peripheral neuropathy, and • At least one of the following seven features: osteosclerotic bone lesions, Castleman's disease, organomegaly, endocrinopathy (excluding diabetes mellitus or hypothyroidism), edema, typical skin changes, and papilledema. <p>Note: Not every patient meeting the above criteria will have POEMS syndrome; the features should have a temporal relationship to each other and no other attributable cause. The absence of osteosclerotic lesions should make the diagnosis suspect. Elevations in plasma or serum levels of vascular endothelial growth factor and thrombocytosis are common features of the syndrome and are helpful when the diagnosis is difficult.</p>	279

Modified, reproduced with permission from Rajkumar SV, Dispenzieri A, Kyle RA, Mayo Clinic Proc 2006;81:693-703. © Mayo Clinic Proceedings.³⁹

Primary translocations in Monoclonal Gammopathy of Undetermined Significance (MGUS) and Multiple Myeloma

Table 2

Chromosomal translocation	Gene dysregulated by translocation ²⁸⁰
t 11;14	CCND1 (cyclin D1)
t 4;14	FGFR-3 and MMSET
t 6;14	CCND3 (cyclin D3)
t 14;16	C-maf
t 14;20	mafB

Reproduced with permission from Rajkumar SV, Kyle RA². Mayo Clinic Proc 2005;80:1371-82. © Mayo Clinic Proceedings.

Table 3

Staging of Myeloma

Staging System	Key Stage Characteristics	Comments
Durie-Salmon Staging System		Is more correlated with biologic stage and tumor burden of disease than the international staging system. Disadvantage is complexity, and subjective nature or interpretation of bone findings
Stage I	All of the following: <ul style="list-style-type: none"> • 0-1 bone lesion, • hemoglobin >10g/dL, • normal calcium, and • low M protein production 	Low M protein defined as IgG <5gm/dL, IgA <3g/dL, urine M spike >4000 mg/24 hours
Stage II	Not fitting stage I or II	
Stage III	One or more of the following: <ul style="list-style-type: none"> • Advanced bone lesions, • hemoglobin <8.5g/dL, • increased calcium, or • high M protein production 	High M protein defined as IgG >7gm/dL, IgA >5g/dL, urine M spike >12,000 mg/24 hours
Subclassification		
A	<ul style="list-style-type: none"> • Serum creatinine <2.0 mg/dL 	
B	<ul style="list-style-type: none"> • Serum creatinine ≥2.0 mg/dL 	
International Staging System		Simple to use, but most renal failure patients will be stage III regardless of tumor burden. Does not apply to monoclonal gammopathy of undetermined significance or smoldering myeloma.
Stage I	<ul style="list-style-type: none"> • Serum β-2 microglobulin <3.5 mg/L and albumin ≥3.5 g/dL 	
Stage II	<ul style="list-style-type: none"> • Not fitting stage I or II 	

Table 4

Major Prognostic Factors in Myeloma

Independent Prognostic Factors	
1	Performance status
2	International Staging System Stage (includes beta-2 microglobulin and albumin)
3	Durie-Salmon Staging System Stage
4	Deletion 13 or hypodiploidy by karyotyping
5	Translocations t(4;14) or t(14;16)
6	Deletion 17p
7	Serum lactate dehydrogenase (LDH)
8	Plasmablastic morphology
9	Bone marrow plasma cell labeling index (PCLI)
Additional Prognostic Factors	
1	Age
2	C-reactive protein
3	Serum creatinine
4	Platelet count

Table 5
Mayo Risk-Stratification for Myeloma²⁷¹

High-risk characteristic	Percentage of newly diagnosed patients with the abnormality	7,45,281
Conventional Cytogenetics:		
Deletion of chromosome 13 (monosomy)	14%	
Hypodiploidy	9%	
Either hypodiploidy or deletion 13	17%	
Fluorescent in-situ hybridization (FISH):		
t(4;14)	15%	
t(14;16)	5%	
17p-	10%	
Plasma cell labeling index (PCLI) studies: PCLI ≥3%	6%	
Any one of the above high-risk abnormalities	25-30%	

Reproduced with permission from Rajkumar SV, Kyle RA². Mayo Clinic Proc 2005;80:1371-82. © Mayo Clinic Proceedings.

Table 6

Definition of Measurable Disease

One or more of the following:
<ul style="list-style-type: none">• Serum monoclonal (M) protein ≥ 1gm/dL• Urine M protein ≥ 200 mg/24 hours• Involved free light chain (FLC) level ≥ 10mg/dL plus abnormal FLC kappa/lambda ratio (<0.26 or >1.65)[*]

^{*} The FLC assay is not to be used for response assessment in patients who have evidence of measurable disease in the serum or urine by M protein levels.

International Myeloma Working Group Uniform Response criteria for multiple myeloma ⁸¹

Table 7

Response subcategory	Response criteria
Complete response* (CR)	<ul style="list-style-type: none"> Negative immunofixation of serum and urine, and Disappearance of any soft tissue plasmacytomas, and < 5% plasma cells in bone marrow
Stringent complete response (sCR)	<p>CR as defined above plus</p> <ul style="list-style-type: none"> Normal FLC ratio, and Absence of clonal cells in bone marrow by immunohistochemistry or immunofluorescence
Very good partial response (VGPR)*	<ul style="list-style-type: none"> Serum and urine M-component detectable by immunofixation but not on electrophoresis, or ≥ 90% or greater reduction in serum M-component plus urine M-component <100 mg per 24 h
Partial Response (PR)	<ul style="list-style-type: none"> ≥ 50% reduction of serum M-protein and reduction in 24-h urinary M-protein by ≥ 90% or to <200 mg per 24 h If the serum and urine M-protein are unmeasurable a ≥ 50% decrease in the difference between involved and uninvolved FLC levels is required in place of the M-protein criteria If serum and urine M-protein are unmeasurable, and serum free light assay is also unmeasurable, ≥ 50% reduction in bone marrow plasma cells is required in place of M-protein, provided baseline percentage was ≥ 30% In addition to the above criteria, if present at baseline, ≥ 50% reduction in the size of soft tissue plasmacytomas is also required
Stable disease (SD)	<ul style="list-style-type: none"> Not meeting criteria for CR, VGPR, PR or progressive disease
Progressive Disease (PD)*	<p>Increase of 25% from lowest response value in any one or more of the following:</p> <ul style="list-style-type: none"> Serum M-component (absolute increase must be ≥ 0.5 g/dl)[†] and/or Urine M-component (absolute increase must be ≥ 200 mg/24 h) and/or Only in patients without measurable serum and urine M-protein levels: the difference between involved and uninvolved FLC levels (absolute increase must be > 10 mg/dL) Bone marrow plasma cell percentage (absolute % must be ≥ 10%) Definite development of new bone lesions or soft tissue plasmacytomas or definite increase in the size of existing bone lesions or soft tissue plasmacytomas Development of hypercalcemia (corrected serum calcium >11.5 mg/dl) that can be attributed solely to the plasma cell proliferative disorder

All response categories (CR, sCR, VGPR, PR) require two consecutive assessments made at anytime before the institution of any new therapy; complete and PR and SD-categories also require no known evidence of progressive or new bone lesions if radiographic studies were performed. Radiographic studies are not required to satisfy these response requirements. Bone marrow assessments need not be confirmed.

* **Note clarification to IMWG criteria for coding CR and VGPR in patients in whom the only measurable disease is by serum FLC levels:** CR in such patients a normal FLC ratio of 0.26-1.65 in addition to CR criteria listed above. VGPR in such patients is defined as a >90% decrease in the difference between involved and uninvolved free light chain FLC levels.

Adapted with permission from Durie et al. International uniform response criteria for multiple myeloma. *Leukemia* 2006;20:1467-73 ⁸¹

Table 8
New drugs with significant single-agent activity in multiple myeloma

Agent	Usual Starting Dose	Postulated Mechanism of Action	Side-effects
Thalidomide	50-200 mg orally days 1-28 every 4 weeks	Anti-angiogenesis, immunomodulation, and inhibition of tumor necrosis factor alpha. Direct cytotoxicity by inducing free radical mediated DNA damage.	Sedation, fatigue, skin rash, bradycardia, peripheral neuropathy and constipation are the most common side effects. Deep vein thrombosis when combined with dexamethasone or chemotherapy. Less frequent but serious side-effects are seizures, cardiac arrhythmia, and Stevens Johnson syndrome.
Bortezomib	1.3mg/m ² intravenously days 1, 4, 8, 11 every 21 days	Inhibits the ubiquitin-proteasome catalytic pathway in cells by binding directly with the 20S proteasome complex.	The most common toxicities are gastrointestinal, transient cytopenias, fatigue, and peripheral neuropathy.
Lenalidomide*	25 mg orally days 1-21 every 28 days	Analog of thalidomide. Immunomodulation and anti-angiogenesis	The most common adverse effects are grade 3 or 4 thrombocytopenia and neutropenia.

Reproduced with permission from Rajkumar SV, Kyle RA². Mayo Clinic Proc 2005;80:1371-82. © Mayo Clinic Proceedings.

Table 9
Results Of Selected Randomized Trials That Have Had A Major Impact On Myeloma Therapy

Trial	Disease Stage	Treatment comparison	Total Number of patients studied	CR/VGPR (%)	Median EFS (months)	Median OS (months)	Comments
Stem Cell Transplantation							
IFM 90206	Newly Diagnosed	Conv. chemo	100	13	18	44	Established role of ASCT
MRC VII207	Newly Diagnosed	ASCT	100	38	28	57	Confirmed role of ASCT
		Conv. chemo	201	8	20	42	
MAG 90213	Newly Diagnosed	ASCT	200	44	32	54	Demonstrated delayed ASCT as an alternative
		Delayed ASCT	94	21	13	65	
United States Intergroup S9321282	Newly Diagnosed	Early ASCT	91	32	39	64	Demonstrated delayed ASCT as an alternative; dampened enthusiasm for allogeneic SCT
		Delayed ASCT	255	15*	21	64	
		Early ASCT	261	17*	25	58	
		Allogeneic SCT	36	17*	NR	6	
IFM 94283	Newly diagnosed	Single ASCT	199	42	25	48	Established role of tandem ASCT if CR/VGPR not achieved with first ASCT
		Double ASCT	200	50	30	58	
PETHEMA284	Post-induction (responding patients only)	Conv. chemo	83	11*	33	66	Demonstrated that ASCT may have limited value in patients responding well to induction
		ASCT	81	30*	42	61	
Italian232	Newly diagnosed	Tandem ASCT	82	N/A	29	54	Demonstrated efficacy of single ASCT followed by non- myeloablative SCT in selected patients
		Single ASCT followed by non- myeloablative SCT	80	N/A	35	80	

New therapies							
Italian GIMEMA group197	Newly diagnosed	MP	126	12	27	64% at 3yrs	Demonstrated potential value of MPT over the classic MP regimen
		MPT	129	36	54	80% at 3yrs	
IFM 99-06199	Newly diagnosed	MP	196	7	18	33	Changed standard of care in elderly to MPT (after 3 decades of MP)
		MPT	125	47	28	52	
APEX84	Relapsed, refractory (1-3 prior therapies)	Tandem intermediate-dose ASCT	126	43	19	38	Led to full approval of bortezomib in the US
		Bortezomib	333	13	6**	NR	
		Dex	336	2	3**	NR	

Trial	Disease Stage	Treatment comparison	Total Number of patients studied	CR/VGPR (%)	Median EFS (months)	Median OS (months)	Comments
MM-010174	Relapsed, refractory (1-3 prior therapies)	Len/ Dex	351	15*	11**	Not reached	Pivotal trial establishing role of lenalidomide
		Placebo/Dex		3*	5**		
MM-009175	Relapsed, refractory (1-3 prior therapies)	Len/ Dex	341	13*	11*	30	Led to approval of lenalidomide in the US
		Placebo/Dex		1*	5*	20	

Abbreviations: ASCT, autologous hematopoietic stem cell transplantation; Conv. chemo, conventional chemotherapy; CR, complete response; Dex, dexamethasone; ECOG, Eastern Cooperative Oncology Group; EFS, event free survival; GIMEMA, Gruppo Italiano Malattie Ematologiche dell'Adulto; IFM, InterGroupe Francophone du Myélome; Len, Lenalidomide; MAG, Myélome Autogreffe; MM, multiple myeloma; MP, melphalan plus prednisone; MPT, melphalan, prednisone, thalidomide; MRC, Medical Research Council; NR, not reported; OS, overall survival; PETHEMA, Programa para el Estudio y Tratamiento de las Hemopatías Malignas; PFS, progression free survival; SCT, stem cell transplantation; US, United States; VGPR, very good partial response.

* Denotes CR only; VGPR not reported.

** PFS data not available; numbers reflect median time to progression

This Table was originally published in Blood. Kyle RA and Rajkumar SV. Multiple Myeloma. Blood. 2008; 111;2962-2972. © the American Society of Hematology

Dosing Schedule and Response Rates of Commonly used Chemotherapy Regimens in Myeloma

Table 10

Regimen	Suggested Dosing Schedule*	Response rate in newly diagnosed myeloma	CR+VGPR [‡] rate in newly diagnosed myeloma
Transplant Candidates			
Thalidomide- Dexamethasone ^{**} (Thal/Dex) ^{185,188}	Thalidomide 100-200mg oral days 1-28	63%	44%
	Dexamethasone 40mg oral days 1, 8, 15, 22 every 28 days		
	Repeated every 4 weeks × 4 cycles as pre-transplant induction therapy; or continued for up to one year or progression if used as primary therapy		
Lenalidomide-Dexamethasone (Rev/low-dose Dex) ²⁸⁵	Lenalidomide 25mg oral days 1-21 every 28 days	71%	42%
	Dexamethasone 40mg oral days 1, 8, 15, 22 every 28 days		
	Repeated every 4 weeks × 2-4 cycles as pre-transplant induction therapy; or continued till progression if used as primary therapy		
Bortezomib-Dex (Vel/Dex) ²⁸⁶	Bortezomib 1.3mg/m ² intravenous days 1, 4, 8, 11	80%	47%
	Dexamethasone 40mg oral days 1-4, 9-12		
	Reduce dexamethasone to days 1-4 only after first 2 cycles		
Bortezomib-Thalidomide-Dexamethasone (VTD) ^{**287}	Repeated every 3 weeks × 2-4 cycles as pre-transplant induction therapy	93%	60%
	Bortezomib 1.3mg/m ² intravenous days 1, 4, 8, 11		
	Thalidomide 200mg oral days 1-21		
VDT-PACE ^{‡196}	Dexamethasone 20 mg day of and day after bortezomib	Not reported	Not reported
	Repeated every 3 weeks × 3 cycles as pre-transplant induction therapy		
	Dexamethasone 40 mg/day PO daily days 1-4		
	Thalidomide 100 mg/day days 1-28 [†]		
	Cisplatin 10 mg/m ² /day intravenous infusion over 24 hours days 1-4		
	Cyclophosphamide 400 mg/m ² /day intravenous infusion over 24 hours days 1-4		
	Etoposide 40 mg/m ² /day intravenous infusion over 24 hours days 1-4		
	Doxorubicin 10 mg/m ² /day intravenous infusion over 24 hours days 1-4		
	Bortezomib 1 mg/m ² /day intravenous on days 1,4,8,11		

Note:

- 1 For IV infusion, each 24 hour dose of cisplatin is mixed with cyclophosphamide and etoposide in 1000ml of normal saline. Doxorubicin is administered as a separate IV infusion via central line, with each 24 hour dose mixed in 250 ml of normal saline

Regimen	Suggested Dosing Schedule [*]	Response rate in newly diagnosed myeloma	CR+VGPR [†] rate in newly diagnosed myeloma
Non-Transplant Candidates	2 For patients with weight greater than 60 kg, all doses will be based on calculated body weight (actual weight + ideal body weight divided by 2) and not to exceed a BSA of 2m ² .		
	3 Prophylaxis for deep vein thrombosis with low molecular weight heparin is recommended. In addition prophylaxis with levofoxacin, fluconazole, and acyclovir is recommended to prevent infection. Prophylaxis against gastric ulceration is needed. Growth factors (either filgrastim (starting day 5) or pegfilgrastim (one dose on day 5) are needed.		
	Melphalan 9mg/m ² oral days 1-4 Prednisone 60mg/m ² oral days 1-4 Repeated every 6 weeks until plateau	35	8%
Melphalan-Prednisone- Thalidomide (MPT) ^{199†}	An alternative schedule is melphalan 8-10 mg oral days 1-7 with prednisone 60 mg days 1-7 repeated every 6 weeks until plateau		
	Melphalan 0.25mg/kg oral days 1-4 Prednisone 2mg/kg oral days 1-4 Thalidomide 100-200mg oral days 1-28 Repeated every 6 weeks × 12 cycles	76%	47%
	Melphalan 9 mg/m ² oral days 1-4 Prednisone 60 mg/m ² oral days 1 to 4 Bortezomib 1.3mg/m ² intravenous days 1, 4, 8, 11, 22, 25, 29, 32 Repeated every 42 days × 4 cycles followed by maintenance therapy as given below:		
Bortezomib-Melphalan-Prednisone (VMP) ²⁰⁴	Melphalan 9 mg/m ² oral days 1-4 Prednisone 60 mg/m ² oral days 1 to 4 Bortezomib 1.3mg/m ² intravenous days 1, 8, 15, 22 Repeated every 35 days × 5 cycles	71%	41%

^{*} Starting and subsequent doses need to be adjusted for performance status, renal function, blood counts, and other toxicities;

^{**} Dexamethasone dose listed is lower than used in trial.

[†] Thalidomide dose listed is lower than used in trial

[‡]CR, complete response; VGPR, very good partial response; VDT-PACE, velcade, dexamethasone, thalidomide, cisplatin (platinum), Adriamycin, cyclophosphamide, and etoposide.

Table 11
Randomized Trials Comparing Conventional Chemotherapy Versus Single Autologous Stem Cell Transplantation

Trial	No. of patients	CR/VGPR, %		PFS, mo		OS, mo	
		CCT	ASCT	CCT	ASCT	CCT	ASCT
IFM90 ^{206,288}	200	13	38*	18	28*	44	57*
MRC7 ²⁰⁷	401	8	44*	20	32*	42	54*
MAG91 ²⁸⁹	190	4	6	19	25*	48	48
MAG90 ^{**213}	185	57	20	13	39	64	65
PEETHMA ²⁸⁴	164	11	30*	33	42	61	66
S9321 ^{**282}	516	17	15	7 yr 14%	7 yr 17%	7 yr 38%	7 yr 38%
MMSG ^{†,290}	194	6	25*	16	28*	42	58+*
HOVON- 24 ^{‡,291}	303	13	28*	23	24*	50	55

ASCT, autologous stem cell transplant; CCT, conventional chemotherapy; CR/VGPR, complete response or very good partial response; ORR, any response greater than or equal to a partial response; PFS, progression free survival; OS, overall survival;

* Significant

** Trials testing early versus delayed autologous stem cell transplant

[†]Transplant arm is 2 low dose (melphalan 100 mg/m2) autologous stem cell transplants

[‡]Could be considered “double transplant” since both arms received melphalan 70 mg/m2 × 2 without stem cell support as induction therapy

Reproduced with permission from Dispenzieri, A et al. Mayo Clinic Proc 2007;82:323-341. © Mayo Clinic Proceedings.