Published in final edited form as:

Org Lett. 2008 November 20; 10(22): 5135–5138. doi:10.1021/ol8020308.

Design and Synthesis of Stereochemically Defined Novel Spirocyclic P2-Ligands for HIV-1 Protease Inhibitors

Arun K. Ghosh^{*,†}, Bruno D. Chapsal[†], Abigail Baldrige[†], Kazuhiko Ide[‡], Yashiro Koh[‡], and Hiroaki Mitsuya^{‡,§}

[†]Departments of Chemistry and Medicinal Chemistry, Purdue University, West Lafayette, IN 47907

[‡]Department of Hematology and Infectious Diseases, Kumamoto University School of Medicine, Kumamoto 860-8556, Japan

[§]Experimental Retrovirology Section, HIV and AIDS Malignancy Branch, Center for Cancer Research, National Cancer Institute, Bethesda, Maryland 20892

Abstract

The synthesis of a series of stereochemically defined spirocyclic compounds and their use as novel P2-ligands for HIV-1 protease inhibitors are described. The bicyclic core of the ligands was synthesized by an efficient *n*Bu₃SnH-promoted radical cyclization of a 1,6-enyne followed by oxidative cleavage. Structure-based design, synthesis of ligands and biological evaluations of the resulting inhibitors are reported.

The introduction of highly active antiretroviral therapy (HAART) in 1996, in combination with HIV-1 protease inhibitors and reverse transcriptase inhibitors has dramatically changed the management of HIV/AIDS. ¹ The advent of HAART has significantly reduced morbidity and mortality and has improved the quality of life for HIV-infected patients, particularly in developed nations. ² Despite this important breakthrough, current and future management of HIV/AIDS is being challenged by the rapid emergence of multi-drug-resistant HIV-1 strains and drug-related side effect. ³ Consequently, developments of novel and effective treatment regimens are critically important.

In our continuing effort to design a new generation of HIV-1 protease inhibitors (PIs) that combat drug-resistance, we developed a series of exceedingly potent PIs. A number of these nonpeptidyl PIs have shown superb antiviral activity and drug-resistance profiles. In our structure-based design strategies, we introduced the 'backbone binding concept' with the presumption that an inhibitor that makes maximum interactions in the protease active site, particularly hydrogen bonding with the backbone atoms may retain its potency against mutant strains. ⁴ Darunavir (**1**, Figure 1), which has been approved by the FDA for the treatment of patients harboring multi-drug-resistant HIV-1 strains, has emerged from this approach. ^{5, 6} Our detailed X-ray structural analysis of protein-ligand complexes revealed an extensive hydrogen

bonding network with HIV-1 protease backbone atoms, and most notably with the designed bis-THF P2-ligand.⁷

More recently, we reported another novel PI, GRL-06579 (**2**) which features a stereochemically defined bicyclic hexahydrocyclopentylfuran (Cp-THF) P2-ligand in the hydroxyethylsulfonamide isostere core.⁸ The X-ray crystallographic analysis of **2**-bound HIV-1 protease documented extensive hydrogen bonding interactions including the Cp-THF oxygen with the backbone atoms in the S2-subsite.⁸

The favorable drug-resistance profile of this PI containing a Cp-THF ligand logically prompted us to design several structural analogs. We set out to introduce new functionalities on this bicyclic core that could create additional interactions within the enzyme catalytic site. The 3-position of the Cp-THF ligand appeared particularly suitable for this purpose, because of its proximity to the flap region and the S2-subsite of the protease. Based upon our analysis of the X-ray crystal structure of **2**-bound protease, we planned to investigate the effect of a structurally constrained spirocyclic motif at the 3-position of the Cp-THF ring. We speculated that a cyclic ether oxygen or an oxazolidinone carbonyl oxygen may be positioned in this cyclic motif to accept a hydrogen bond from the enzyme active site residues or backbone NH. Such a functionality would fill in the hydrophobic pocket in the S2-subsite as well. Furthermore, this structural feature may improve the pharmacological profile of these inhibitors.^{9,10}

We initially set out to synthesize a series of spirocyclic Cp-THF-derived P2-ligands and their corresponding HIV-protease inhibitors (**3a-e**). A synthetic strategy was devised so that all analogs could be synthesized from a common precursor that gives rapid access to new polycyclic molecular probes. The general synthesis of the bicyclic core of our new P2-ligands was accomplished in enantiomerically pure form as shown in Scheme 1. Optically active monoacetate **4**, was obtained in 95% ee by desymmetrization of the corresponding *meso*-diacetate with acetyl cholinesterase.¹¹ Protection of alcohol **4** as a TBS ether followed by methanolysis of the acetyl group furnished compound **5**. Propargylation of **5** using propargyl bromide in the presence of NaH provided alkyne **6** in excellent yield.

The construction of the bicyclic core was accomplished by an intramolecular radical cyclization of enyne **6** using *n*Bu₃SnH and AIBN in benzene at reflux. This provided vinyl stannane **7** as a mixture of *cis/trans* diastereoisomers (2:1), along with trace amounts of the olefin, which presumably formed during purification on silica gel. The mixture of isomers was directly oxidized with a catalytic amount of OsO₄ in the presence of NaIO₄ and 2,6-lutidine to afford the key-intermediate, ketone **8** in 77% yield.

We first turned our attention to the synthesis of spirocyclic ketal **9** and ether **12**. Molecular modeling of the corresponding inhibitors, suggested that the ligand oxygens could be within hydrogen bonding distance to the NH amide bonds of both Asp30 and Asp29 residues.

Spirocyclic dioxolane ligand **9** was obtained in 86% yield by treatment of ketone **8** with ethylene glycol in benzene with a catalytic amount of *p*-TsOH. Synthesis of ether **12** was achieved in four consecutive steps starting from ketone **8**. Reaction of **8** with homoallyl magnesium bromide furnished compound **10** in 76% yield. Ozonolysis of the terminal alkene and refluxing of the resulting crude aldehyde in benzene/methanol followed by azeotropic distillation of the excess methanol afforded methyl acetal **11** as a mixture (1:0.85) of diastereoisomers. Reduction of this acetal intermediate **11** furnished the desired alcohol **12** by applying a one-pot procedure involving: (1) TMS-protection of the alcohol with hexamethyldisilazane and (2) subsequent reduction of the acetal with triethylsilane.¹²

We have designed spirocyclic oxazolidinone ligands that could potentially exploit polar interactions with the backbone atoms and residues in the HIV-1 protease active site. Their

respective syntheses are highlighted in Scheme 3 and 4. Cyanohydrin **13** was synthesized in 82% yield from ketone **8**. LiAlH_4 -reduction of the cyanide provided the corresponding amine, which exhibited partial TMS-deprotection. Therefore, the crude mixture was directly submitted to the next steps with (1) formation of the methyl carbamate derivative and (2) removal of the silyl ethers with TBAF in THF. The resulting diol **14** was then treated with NaH in THF to give oxazolidinone ligand **15** in 51% yield for 4 steps (from **13**).

Synthesis of oxazolidinone **18** started with vinylstannane **7** (Scheme 4). Proto-destannylation of **7** was carried out with CSA in CH_2Cl_2 . Epoxidation of the resulting olefin with *m*-CPBA gave epoxide **16** as a major diastereomer (93:7 ratio). Opening of the epoxide with *para*-methoxybenzylamine gave amino alcohol **17** in 67% yield. The carbonyl was installed using triphosgene and Et_3N in CH_2Cl_2 . Deprotection of the TBS-group provided the desired oxazolidinone **18**.

The synthesis of polycyclic PIs are shown in Scheme 5. Various synthetic ligands were reacted with 4-nitrophenylchloroformate and pyridine to form the corresponding activated carbonates, **19a-e**. Reaction of the respective active carbonate with known⁸ amine **20** in the presence of diisopropylethylamine afforded PIs **3a-d**. For the synthesis of inhibitor **3e**, amine **20** was reacted with active carbonate **19e** to provide urethane **21**. Removal of the PMB group from **21** by exposure to ceric ammonium nitrate (CAN) afforded inhibitor **3e**.

We examined all inhibitors for their enzymatic potency as well as cellular activity; and the results are displayed in Table 1. As shown, most inhibitors exhibited excellent enzymatic potency. Dioxolane-based analogue **3a** displayed a K_i value of 0.16 nM. Inhibitor **3b** contains the spirocyclic methyl acetal as a mixture (1:0.85 ratio) of diastereomers. These diastereomers were separated by HPLC and the stereochemical identity of each diastereomer was determined by extensive NOESY experiments. Diastereomer **3b**-(*S*) showed an enzymatic K_i of 0.81 nM. The **3b**-(*R*) isomer is slightly more potent (K_i = 0.38 nM). The removal of the methoxy group from **3b** resulted in inhibitor **3c** which has shown loss of enzyme inhibitory activity. Both inhibitors **3b** and **3c** have shown comparable antiviral activity. We have examined stereochemically defined oxazolidinone derivatives as P2-ligands. Inhibitor **3d** displayed a K_i of 0.29 nM. Diastereomeric inhibitor **3e** is slightly more potent than **3d** in both enzyme inhibitory as well as in antiviral assays (IC_{50} = 21 nM in MT-2 cells). The inhibitors in Table 1 in general, are significantly less potent than UIC-PI (TMC-126),¹⁴ the corresponding methoxysulfonamide derivative of darunavir or Cp-THF-containing inhibitor **2**.⁸

In conclusion we have designed and synthesized a series of inhibitors containing stereochemically defined novel spirocyclic P2-ligands. The syntheses of these ligands were carried out from the key intermediate **8** which was efficiently prepared in optically active form by using a radical cyclization as the key step. The spirooxazolidinone-derived inhibitor **3e** is the most potent inhibitor in this series. While these inhibitors contain novel P2-ligands, it appears that the spirocyclic motif at the 3-position of the Cp-THF ring resulted in a significant reduction in potency. Further design and optimization of the ligand binding site interactions are in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support by the National Institutes of Health (GM 53386, AKG) is gratefully acknowledged. This work was also supported in part by the Intramural Research Program of the Center for Cancer Research, National Cancer Institute, National Institutes of Health, and in part by a Grant-in-aid for Scientific Research (Priority Areas) from the Ministry

of Education, Culture, Sports, Science, and Technology of Japan (Monbu Kagakusho) and a Grant for Promotion of AIDS Research from the Ministry of Health, Welfare, and Labor of Japan. We thank Mr. David D. Anderson (Purdue University) for his help with the HPLC and NOESY analysis.

References

1. Sepkowitz KA. N Engl J Med 2001;344:1764. [PubMed: 11396444]
2. Palella FJ, Delaney KM, Moorman AC, Loveless MO, Fuhrer J, Satten GA, Aschman DJ, Homberg SD. N Engl J Med 1998;338:853. [PubMed: 9516219]
3. Boden D, Markowitz M. Antimicrob Agents Chemother 1998;42:2775. [PubMed: 9797203]
4. Ghosh AK, Chapsal BD, Weber IT, Mitsuya H. Acc Chem Res 2008;41:78. [PubMed: 17722874]
5. Ghosh AK, Kincaid JF, Cho W, Walters DE, Krishnan K, Hussain KA, Koo Y, Cho H, Rudall C, Holland L, Buthod J. Bioorg Med Chem Lett 1998;8:687. [PubMed: 9871583]
6. Koh Y, Nakata H, Maeda K, Ogata H, Bilcer G, Devasamudram T, Kincaid JF, Boross P, Wang Y-F, Tie Y, Volarath P, Gaddis L, Harrison RW, Weber IT, Ghosh AK, Mitsuya H. Antimicrob Agents Chemother 2003;47:3123. [PubMed: 14506019]
7. Kovalevsky AY, Liu F, Leshchenko S, Ghosh AK, Louis JM, Harrison RW, Weber IT. J Mol Biol 2006;363:161. [PubMed: 16962136]
8. Ghosh AK, Sridhar PR, Leshchenko S, Hussain AK, Li J, Kovalevsky AY, Walters DE, Wedekind JE, Grum-Tokars V, Das D, Koh Y, Maeda K, Gatanaga H, Weber IT, Mitsuya H. J Med Chem 2006;49:5252. [PubMed: 16913714]
9. Veber DF, Johnson SR, Cheng H-Y, Smith BR, Ward KW, Kopple KD. J Med Chem 2002;45:2615. [PubMed: 12036371]
10. Lipinski CA. J Pharmacol Toxicol Methods 2000;44:235. [PubMed: 11274893]
11. Deardorff DR, Windham CQ, Craney CL. Org Synth 1996;9:487.
12. Yoo SJ, Kim HO, Lim Y, Kim J, Jeong LS. Bioorg Med Chem 2002;10:215. [PubMed: 11738624]
13. Toth MV, Marshall GR. Int J Pep Protein Res 1990;36:544.
14. Ghosh AK, Sridhar P, Kumaragurubaran N, Koh Y, Weber IT, Mitsuya H. ChemMedChem 2006;1:937.

Figure 1.
Structure of HIV Protease Inhibitors

Scheme 1.
Synthesis of bicyclic ketone **8**

Scheme 2.
Synthesis of spirocyclic ketal and ether ligands

Scheme 3.
Synthesis of spirooxazolidinone ligand **15**

Scheme 4.
Synthesis of spirooxazolidinone ligand **18**

Scheme 5.
Synthesis of inhibitors **3a-e**

Table 1

Enzymatic and antiviral activity of PIs

Inhibitor	R	K _i (nM)
-----------	---	---------------------

3a

0.16

Inhibitor	R	K _i (nM)
3b		3b-(S) iso: (X=OMe, Y=H):0.38 3b-(R) iso: (X=H,Y=OMe):0.38
3c		2.22
3d		0.29
3e		0.17

^a K_i determined following protocol as described by Toth and Marshall, mean values of at least 4 determinations.¹³

^b MT-2 cells (2 × 10⁴/mL) were exposed to 100 TCID₅₀ of HIV-1_{LAJ} and cultured in the presence of various concentrations of PIs, and the IC₅₀s were determined by using the MTT assay on day 7 of culture.⁶

^c Tested as a 1:0.85 mixture.