Published in final edited form as:

J Liq Chromatogr Relat Technol. 2009 January 1; 32(2): 273–280. doi:10.1080/10826070802603369.

PREPARATIVE ISOLATION OF ISORHAMNETIN FROM *STIGMA MAYDIS* USING HIGH-SPEED COUNTERCURRENT CHROMATOGRAPHY

Xiaoying Cao^a, Yun Wei^{a,*}, and Yoichiro Ito^b

^a Applied Chemistry Department, Faculty of Sciences, Beijing University of Chemical Technology, 15 Beisanhuan East Road, Chaoyang District, Beijing 100029, China

^b Bioseparation Technology Laboratory, Biochemistry and Biophysics Center, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland, USA

Abstract

Preparative high-speed countercurrent chromatography (HSCCC) has been successfully used for the isolation and purification of isorhamnetin from *Stigma maydis*. This was achieved in two stages: the first separation was performed with a two-phase solvent system composed of n-hexane-ethyl acetate-methanol-water (HEMW) at a volume ratio of 5:5:5:5, yielding isorhamnetin at 65.6 %, which is followed by the second run using a two-phase solvent system composed of HEMW 5:5:6:4, v/v. From 700 mg of the crude extract 11.8 mg of isorhamnetin was obtained at a high purity of 98%. The final identification was performed by MS, ¹H-NMR and ¹³C-NMR spectra.

Keywords

High-speed countercurrent chromatography; Preparative isolation; Isorhamnetin; *Stigma maydis*

1. Introduction

Stigma maydis is the capital of *Zea mays* L., containing saponin, alkaloid, and flavones which can be used in the treatment of diuresis [1]. Flavones and similar structural compounds have been recently shown to inhibit blood glucose and aspergillus activity[2].

However, the separation of these active compounds from natural sources using conventional support-based chromatography can have the problem of irreversible adsorption to the support. High-speed countercurrent chromatography (HSCCC), being a support-free liquid-liquid partition chromatographic technique, eliminates such adsorption problems [3], and has been widely used in preparative separation of natural products [4–6]. Some researches have focused on the separation of flavone from *Stigma maydis* by macroporous resin [7,8]. The present paper describes the successful preparative separation and purification of Isorhamnetin from the crude extract of *Stigma maydis* which is achieved by two-stage high-speed countercurrent chromatography process with two-phase solvent systems composed of n-hexane-ethyl acetate-methanol-water at volume ratios of 5:5:5:5 and 5:5:6:4.

Corresponding author. Prof. Yun WEI, Applied Chemistry Department, Faculty of Sciences, Beijing University of Chemical Technology, 15 Beisanhuan East Road, Chaoyang District, Beijing 100029, P. R. China. Tel & fax: 0086 10 64442928. weiyun@mail.buct.edu.cn.

2. Experimental

2.1 Materials

All organic solvents used for HSCCC were of analytical grade and purchased from Beijing Chemical Factory (Beijing, China). Methanol used for HPLC analysis was of chromatographic grade and purchased from Tianjin Huaxi Special Reagent Factory (Tianjin, China).

Stigma maydis was gifted from Enkang Factory (Hebei province, China).

2.2 Preparation of Sample

About 100g of dried *Stigma maydis* were ground, and 7.5g amount of the dried powder was extracted twice by refluxing for 3 hour with 800ml of 80% ethanol solution, and then extracted with ethyl acetate, concentrated to dryness under reduced pressure and concentrated to dryness under reduced pressure yielding 3 g of a crude sample that contained isorhamnetin.

2.3 Preparation of two-phase solvent system and sample solutions

The solvent systems utilized in the phase system optimization study were prepared by mixing n-hexane-ethyl acetate-methanol-water (5:5:5:5 or 5:5:6:4, v/v), or ethyl acetate-methanol-water (10:1:10, v/v). The mixture were equilibrated in a separating funnel at room temperature, the two phases were separated shortly before use.

The sample solutions were prepared by dissolving the crude extract in the lower phase at the appropriate concentrations.

2.4 The choice of two-phase system

Three two-phase systems including ethyl acetate-methanol-water (10:1:10 v/v), n-hexane-ethyl acetate-methanol-water (5:5:6:4, v/v), and n-hexane-ethyl acetate-methanol-water (5:5:5:5, v/v) have been used.

Each HSCCC peak has been determined by HPLC.

2.5 Two – stage preparative HSCCC

Two-stage preparative HSCCC was performed on the Model GS 10AB HSCCC instrument as follows: the multiplayer coiled column was first entirely filled with the upper phase of the selected two-phase solvent system composed of n-hexane-ethyl acetate-methanol-water with volume ratios of 5:5:5:5. The lower phase was then pumped into the head end of the inlet column at the selected flow-rate, while the apparatus was running at a revolution speed of 800 rpm. After hydrodynamic equilibrium was established, as indicated by a clear mobile phase eluting at the tail outlet, the sample solution at the selected concentration was injected through the sample port. The effluent from the tail end of the column was continuously monitored with a UV detector at 254 nm. Each peak fraction was collected according to the chromatogram.

For the second preparative HSCCC stage, the peak 3 from the first stage were concentrated to dryness under reduced pressure, then injected again onto the HSCCC and using the upper phase of n-hexane-ethyl acetate-methanol-water (5:5:6:4, v/v) as the stationary phase and the upper phase as the mobile phase at a flow rate of 2.0 ml/min.

2.6 HPLC analyses and identification of HSCCC peak fractions

The crude extract of *Stigma maydis* and HSCCC peak fractions were each analyzed by HPLC using a YMC-PACK ODS column (150 × 4.6 mm) at room temperature. The mobile phase, composed of methanol and 0.05% H₃PO₄ aqueous solution (50:50, v/v), was isocratically

eluted at a flow-rate of 1.0 ml/min and the effluent monitored at 254nm and 360nm by a DAD detector.

Identification of the target compound (Isorhamnetin) was based on EI-MS, ^1H -NMR, and ^{13}C -NMR spectra.

3. Results and discussion

The chromatogram was obtained by the solvent system ethyl acetate-methanol-water (10:1:10) is shown in Fig. 1a. An impurity peak was eluted at the solvent front while the target compound was retained in the column. Fig. 1b was obtained using a less polar solvent system composed of n-hexane-ethyl acetate-methanol-water (5:5:6:4, v/v). Although the main peak contained isorhamnetin, HPLC analysis showed a large amount of impurity. The solvent system of n-hexane-ethyl acetate-methanol-water (5:5:5:5, v/v) gave the best result as shown in Fig. 1c

Fig. 2 shows the HPLC analysis of the crude sample, in which the crude extract contained about 2.25 % of isorhamnetin based on the external standard curve. In order to shorten the separation time, a two-stage HSCCC process was developed: quick separation of isorhamnetin from *Stigma maydis* was achieved using a HEMW (5:5:5:5, v/v) solvent system where acceptable rapid fractionation was achieved in two hours yielding 19.7mg of isorhamnetin at 65.6% purity based on the external standard curve. Then, the portion (shaded) of the main peak containing isorhamnetin was subjected to the second HSCCC separation using the HEMW (5:5:6:4, v/v) two-phase solvent system. As shown in Fig. 3, two peaks were resolved and the second peak (shaded) contained 11.8 mg of isorhamnetin at a high purity of 98% based on the external standard curve.

The isorhamnetin separation with HEMW (5:5:5:5, v/v) is better than that with HEMW (5:5:6:4, v/v) at the first preparative HSCCC stage. The reason may be that the crude extract contains a large amount of water-soluble materials which should be removed using a two-phase solvent system containing less methanol. After these polar impurities have been removed, better separation of isorhamnetin from the rest of lipophilic material is achieved by increasing the methanol concentration in the second solvent system.

The structural identification of isorhamnetin was carried out by EI-MS, ^1H -NMR and ^{13}C -NMR Spectra as follows: the EI-MS: m/z 316, 301, 287, 245, 153, 142, 128, 108, 69. It showed the molecular ion at m/z 316, which is in agreement with the molecular formula $\text{C}_{16}\text{H}_{12}\text{O}_7$ of isorhamnetin.

Isorhamnetin: ^1H -NMR (500 MHz, DMSO) δ ppm: 3.84 (3H, s, OCH_3), 6.20 (1H, d, 6-H), 6.48 (1H, d, 8-H), 6.94 (1H, d, 5'-H), 7.69 (1H, m, 6'-H), 7.75 (1H, d, 2'-H), 9.40 (1H, s, OH), 9.73 (1H, s, OH), 10.76 (1H, s, OH), 12.46 (1H, s, OH).

Isorhamnetin: ^{13}C -NMR (500 MHz, DMSO) δ ppm: 148.61 (2-C), 135.62 (3-C), 175.72 (4-C), 156.05 (5-C), 98.11 (6-C), 163.71 (7-C), 93.53 (8-C), 160.32 (9-C), 102.91 (10-C), 121.84 (1'-C), 111.56 (2'-C), 146.43 (3'-C), 147.21 (4'-C), 115.33 (5'-C), 121.64 (6'-C), 55.63 (OCH_3).

Conclusions

The results of our studies demonstrate successful preparative separation of isorhamnetin from *Stigma maydis* using two-step HSCCC operation. A 700 mg of the crude extract was first separated with a two-phase solvent system composed of n-hexane-ethyl acetate-methanol-water at a volume ratio of 5:5:5:5 to improve the purify of isorhamnetin from 2.3 % to 65.7 %. This was followed by the second run using a slightly less polar solvent system of n-hexane-

ethyl acetate-methanol-water at a volume ratio of 5:5:6:4 to remove the rest of the impurities, yielding 11.8 mg of isorhamnetin at a high purity of 98%.

Acknowledgments

Financial support from Beijing Natural Science Foundation (Project 2073028) is gratefully acknowledged.

References

1. Muneer AA, Salman W, Husni T. Journal of Ethnopharmacology 2003;85:257–260. [PubMed: 12639749]
2. Wang YL, Li XG. Special Wild Economic Animal and Plant Research 2004;2:42–45.
3. Ito Y, Crit CRC. Rev Analytical Chemistry 1986;17:65–143.
4. Chen F, Li HB, Ngok R-Wong S, Ji B, Jiang Y. Journal of Chromatography A 2005;1064:183–186. [PubMed: 15739885]
5. Köhler N, Winterhalter P. Journal of Chromatography A 2005;1072:217–222. [PubMed: 15887491]
6. Shi SL, Zhang YP, Huang KL, Liu SQ, Zhao Y. Food Chemistry 2008;108(1):402–406.
7. Ren SC, Ding XL. Journal of Henan University of Technology (Natural Science Edition) 2007;28(4): 34–36.
8. Zhang HE, Xu DP. Journal of Chinese medical materials 2007;30(2):164–166.

Fig. 1.

Chromatogram of the crude extract from *stigma maydis* by HSCCC.

Solvent system a: ethyl acetate-methanol-water (10:1:10, v/v), sample: 700mg dissolved in 10 ml lower phase of solvent system.

Solvent system b: n-hexane-ethyl acetate-methanol-water (5:5:6:4, v/v), sample: 800mg dissolved in 10 ml lower phase of solvent system.

Solvent system c: n-hexane-ethyl acetate-methanol-water (5:5:5:5, v/v), sample: 700mg dissolved in 10 ml lower phase of solvent system. Peak 3: Isorhamnetin at the purity 65.64% Stationary phase: upper organic phase, mobile phase: lower aqueous phase; flow-rate: 2.0 ml/min; revolution speed: 800rpm

HPLC conditions: YMC-PACK ODS column (150×4.6 mm) at column temperature of room temperature. The mobile phase, composed of methanol and water including 0.05% H_3PO_4 (50:50, v/v), was isocratically eluted at a flow-rate of 1.0 ml/min and the effluent monitored at 254nm and 360nm by a DAD detector.

Fig. 2.
HPLC analyses of the crude extract from *Stigma maydis* with the chemical structure of isorhamnetin.
HPLC conditions are the same as those in Fig. 1

Fig. 3.

Chromatogram of the second-stage HSCCC

Sample: a shaded portion of the peak in Fig. 1c obtained by the first HSCCC run; solvent system: n-hexane-ethyl acetate-methanol-water (5:5:6:4 v/v); stationary phase: upper organic phase; mobile phase: lower aqueous phase; flow-rate: 2.0ml/min; revolution speed: 800rpm. HPLC conditions are the same as those in Fig. 1

Peak 2: Isorhamnetin at the purity 98%