Published in final edited form as:

J Org Chem. 2008 October 3; 73(19): 7603–7610. doi:10.1021/jo801273q.

A Palladium-Catalyzed Method for the Synthesis of Carbazoles via Tandem C-H Functionalization and C-N Bond Formation

W. C. Peter Tsang, Rachel H. Munday, Gordon Brasche, Nan Zheng, and Stephen L. Buchwald*

Department of Chemistry, Massachusetts Institute of Technology Cambridge, Massachusetts 02139

Abstract

The development of a new method for the assembly of unsymmetrical carbazoles is reported. The strategy involves the selective intramolecular functionalization of an arene C-H bond and the formation of a new arene C-N bond. The substitution pattern of the carbazole product can be controlled by the design of the biaryl amide substrate, and the method is compatible with a variety of functional groups. The utility of the new protocol was demonstrated by the concise synthesis of three natural products from commercially available materials.

Keywords

Palladium; Catalysis; C-H Bond; C-N bond; Carbazole

Introduction

Carbazole alkaloids manifest a variety of biological activities and, hence, they are attractive synthetic targets.^{1,2} The indigenous application of the stem bark of the curry-leaf tree as folk medicine² led to the discovery of the antibacterial, antifungal, and antiviral properties of this class of natural products.^{1,2,3,4} The anti-inflammatory properties of carbazomycin B^{4,5} and the antitumor properties of ellipticine derivatives⁶ encouraged the development of synthetic methods for the preparation of carbazoles. Carvedilol⁷ and carazolol⁸ were identified for their potential as multiple-action anti-hypertensive drugs. It was later realized that their therapeutic value might derive from the properties of the 4-alkoxycarbazole cores that act as potent antioxidants against radical-induced lipid oxidation. Subsequently, new molecules with a 4-alkoxycarbazole core have been developed for the treatment of obesity and type II diabetes.⁹

In addition to their use in medicinal chemistry, carbazoles have attracted increasing attention from the community of material scientists owing to their potential in photophysical and optoelectronic applications. Polyvinylcarbazole (PVK) has been extensively investigated for its use in photorefractive materials and xerography, while *N*-ethylcarbazole (ECz) was recognized as an effective charge-transporting functional plasticizer for PVK and other doped polymers.¹⁰ In the last decade, carbazole compounds were intensively investigated for the development of optoelectronic applications such as polymeric light-emitting diodes (PLED)^{10a,10c,11} and organic light-emitting devices (OLED).^{12,13,14} In particular, it was found that certain carbazole oligomers led to a family of host materials that were suitable for the

electrophosphorescence of blue lights,^{12,13} a critical component for the fabrication of commercial OLED displays.

In view of these important applications, we set out to develop a metal-catalyzed synthesis of unsymmetrical carbazoles. Prior to our investigations, Du Bois, Che, and their respective co-workers had demonstrated that a metal-catalyzed sequence converting an alkyl C-H bond to a C-N bond can be quite efficient.^{15,16,17,18} On the basis of successful applications of direct arylation of arene C-H bonds in recent years¹⁹ and our long-standing interest in the formation of aromatic C-N bonds,²⁰ we felt that it should be possible to realize an overall substitution reaction where an arene C-H bond is replaced with a C-N bond. In late 2005,²¹ our laboratory reported a new strategy for assembling simple unsymmetrical carbazoles by combining sequential palladium-catalyzed C-H functionalization and C-N bond forming reactions, although the substrate scope was limited to molecules containing functional groups that were compatible with the co-oxidant Cu(OAc)₂. This method was recently adopted by Shi as a part of an elegant sequence for the synthesis of unsymmetrical carbazoles such as 4-deoxycarbazomycin B.²² To address the shortcomings of our earlier work, we decided to expand our research to address the issue of the functional group compatibility, to study the reaction mechanism, and to apply the method to the synthesis of naturally occurring carbazoles. Herein we report our progress toward achieving these goals.

Results and Discussion

Catalyst and Oxidation Systems for the Synthesis of Carbazoles

We initiated our investigation by studying the conversion of 2-acetaminobiphenyl (**1**) to *N*-acetylcarbazole (**2**). Substrate **1** was quantitatively converted to **2** in the presence of a stoichiometric amount of Pd(OAc)₂ in toluene at 120 °C after 24 h. As shown in Table 1 (entry 1), a near quantitative yield of **2** can also be obtained with a combination of 5% Pd(OAc)₂ and one equivalent of Cu(OAc)₂ in an atmosphere of oxygen. No product was observed in the absence of Pd(OAc)₂, suggesting that the role of Cu(OAc)₂ was primarily to mediate the reoxidation of the reduced palladium species. Other commercially available palladium precatalysts (entries 4–7) were less efficient than Pd(OAc)₂ in affecting the transformation. Attempts to use other reoxidants (entries 8–12), which have been reported to be effective in mediating other palladium-catalyzed oxidative coupling reactions,²³ were largely unsuccessful.

Air could be used in the place of oxygen in the synthesis of **2** (Table 2, entry 1), although the reaction was significantly slower compared to that using oxygen (entry 2). We subsequently found that in the presence of oxygen, a catalytic amount of Cu(OAc)₂ was sufficient to effect the synthesis of carbazoles. Three 2-substituted carbazoles (Note: the numbering of the rings changes from the starting biaryl amides to the carbazole products) were obtained in >75% yield using only 20% Cu(OAc)₂ for reactions carried out in toluene as the solvent (entries 3–5). Studies of these three cyclizations also revealed the trend: Cu(OAc)₂ as low as 10% could be used in the cases where the “bottom” aryl ring is neutral or electron-rich; on the other hand, it appears that a minimum of 20% Cu(OAc)₂ is necessary in the cases where the bottom ring bears an electron-withdrawing group. Because of the functional group compatibility, the need to use Cu(OAc)₂ as a co-catalyst is a limitation of the method. In our search for improved reaction conditions, we found that when dimethyl sulfoxide (DMSO) was used as the solvent under an atmosphere of oxygen, no Cu(OAc)₂ was necessary and a comparable yield of **2** was obtained (entry 6). Presumably in this case, DMSO is acting as a ligand to support the direct oxidation of Pd(0) to Pd(II) with oxygen.²⁴ Protocols that employed other polar solvents such as *N,N*-dimethylformamide (DMF) gave poor yields of **2** (entry 7).

Having established that acetamide based biaryl amides were efficiently transformed to the carbazoles in the presence of oxygen with either Cu(OAc)₂ or when carried out in DMSO (Table 3, entries 1 and 2), we turned our attention to substrates containing other amine protecting groups. Sulfonamide-based biaryl amides underwent the cyclization just as efficiently as the corresponding acetamide (entries 3 and 4). In contrast, 2-aminobiphenyl protected as other amides such as benzamide, *tert*-butylcarboxycarbonylamide (BOC-amide), and pentafluoropropionamide provided much lower yields of the desired products (entries 5–10). It appeared that part of the reason for the lower yields in these cases was the cleavage of the amide under the reaction conditions prior to complete conversion to the carbazole. This was clearly observed in the case of the N-Boc substrate when the reaction was attempted in DMSO. In this case, both the 2-aminobiphenyl and free carbazole generated appeared to inhibit the palladium-catalyzed transformation. Since the acetamide group is generally easier to remove than the sulfonamide group, we focused on the formation of *N*-acetylcarbazoles in subsequent studies of the substrate scope. The free carbazoles can be liberated in almost quantitative yield (97%) by treatment either with acid (1:5 H₂SO₄:MeOH) or base (DBU) at 60–80 °C (see Supporting Information).

Synthesis of Substituted Biaryl Amides and 9-Acetylcarbazoles

A variety of substituted biaryl acetamide compounds were prepared in near quantitative yield from commercially available 2'-bromoacetanilide and appropriate boronic acids (Scheme 1, equation 1). The Suzuki-Miyaura coupling reaction was carried out in the presence of a bulky biaryl phosphine, 2-dicyclohexylphosphino-2',6'-dimethoxy-1,1'-biphenyl (SPhos), following a procedure recently developed in our laboratory.²⁵ The reaction can be extended to chloroacetanilide substrates in cases where the substituted bromoacetanilide is inaccessible or costly (Scheme 1, equation 2).

Acetylcarbazoles with various substitution patterns can be obtained in excellent yields via the palladium-catalyzed cyclization reaction (Table 4). In particular, the presence of a substituent (C3') adjacent to the acetamide moiety in the “upper” ring did not affect the efficiency of the cyclization reaction (entries 1 and 2). This observation is significant as it allows us to conveniently assemble carbazoles in excellent yields with substitution at both positions *meta* to the biaryl axis, which is the substitution pattern observed in Mukonine and a number of related natural products.^{1,2} Furthermore, the presence of substituents at positions *ortho* or *para* to the biaryl axis did not affect the formation of the carbazole (entries 6–9). Carbazoles with substitution on both rings were also assembled in good yield (entry 10). The 2-methoxy-6-methylcarbazole core is a synthetically useful building block and has recently been elaborated to form Siamenol,²⁶ a 3,6-disubstituted 2-hydroxycarbazole with anti-HIV activity.²⁷

While all the examples in Table 4 were performed on a 0.2 mmol scale in our initial investigation, both steps in our method can be scaled up with no loss in yield, allowing efficient access to the target carbazoles in sufficient quantity from commercial reagents. For example, the biaryl amide 2-acetamino-2'-methoxybiphenyl (entry 6) could be obtained in 97% yield on a 15 mmol scale from 2'-bromoacetanilide and 2-methoxyphenylboronic acid with 1% Pd(OAc)₂, 2% SPhos, and K₃PO₄ in toluene in 30 min at 90 °C. That product was cyclized in the presence of Pd(OAc)₂, DMSO and oxygen and subsequently hydrolyzed to give 9*H*-4-methoxycarbazole in 86% yield in 2 steps on a 5 mmol scale. This gram-scale experiment demonstrated the practicality of our method to access carbazoles such as 9*H*-4-hydroxycarbazole, a major building block for a variety of therapeutically valuable candidates.^{7,8,9}

The cyclization process of the biaryl amide was compatible with a variety of functional groups. Ketone- and methyl ester-containing substrates can be used in the Cu(OAc)₂ protocol (Table 5). On the other hand, biaryl amides with nitro, cyano, and even thioether groups were more

efficiently cyclized in DMSO (entries 7–9). Unfortunately, attempts to extend this cyclization strategy to substrates with heteroaromatic groups in place of either aryl rings have been unsuccessful.

As shown in Table 4, we have been successful in obtaining 3-substituted carbazoles by derivatizing the “upper” ring (C5') in the biaryl amide substrates (entries 3–5). Alternatively, this class of carbazoles could be obtained by derivatizing the “bottom” ring (C3). Following this approach, 3-substituted carbazoles were formed regioselectively in good to excellent yields as steric factors probably govern the regiochemical nature of the process (Table 6). The cyclization process was generally efficient using Cu(OAc)₂ as the reoxidant with one exception (entries 1 and 4–7). In the case of 3-methoxycarbazole, the cyclization process appeared to be more efficient when Cu(OAc)₂ was replaced with DMSO (entries 2 and 3). When the “bottom” ring is substituted at both C3 and C5, the cyclization process was too slow to be synthetically useful.

Mechanism

We previously proposed a tandem electrophilic palladation and C–N bond formation sequence to account for the intramolecular addition of the amide group to the aryl ring (Scheme 2).^{21, 28} In such a pathway, complexation of Pd(OAc)₂ to the amide moiety of the substrate (**1**) could form a palladium amide (**3**) with concomitant release of one molecule of acetic acid. Palladium amide **3** would then undergo electrophilic palladation to provide a six-membered cyclic palladium amide **4**. Reductive elimination produces the carbazole product (**2**) and a Pd(0) species, which would be reoxidized to Pd(II) in the presence of Cu(OAc)₂ or DMSO and oxygen to complete the catalytic cycle.

However, this proposed catalytic cycle appears to be inconsistent with two experimental observations. First, based on the catalytic cycle, the cyclization of 2'-acetamino-3-methoxybiphenyl was expected to be more facile than those of 2'-acetamino-3-(trifluoromethyl)biphenyl and 2'-acetamino-3-(methoxycarbonyl)biphenyl. However, the experimental data did not support this prediction (Table 6, entries 2, 4, and 5). Second, during our initial optimization studies, attempts to effect the cyclization of biaryl amide **1** at temperatures below 120 °C were met with low yields. For example, only 10% of product **2** was observed at 90 °C after 24 h, and no product was observed at 60 °C. In subsequent studies of substituent effects on the cyclization at 90 °C, we were surprised to see that among all the substrates with substituents at C4 examined, only 2'-acetamino-4-methoxybiphenyl cyclized to the carbazole in satisfactory yield, although this substrate would be expected to destabilize the proposed intermediate due to the inductive effect of the methoxy group in the mechanism suggested in Scheme 2 (Table 7, entry 5). This substituent effect became much less pronounced at 120 °C, although 2'-acetamino-4-methoxybiphenyl remained the most reactive substrate. When the biaryl amide is substituted at C3, regardless of the electronic character of the C3 substituent, the carbazole was formed in about 10% yield over a 24 h period when the reaction was conducted at 90 °C (Table 7, entries 6–8).

Concerned by the discrepancy between our initial model and the data discussed above, we proposed two related catalytic pathways that could better explain the data (Scheme 3). Beginning with the formation of palladium amide **3**, intramolecular cyclization could take place by different mechanisms to form two diastereomeric Pd species (**5a** and **5b**). An insertion into the arene via a Heck-like process would form **5a**, which could undergo a trans β-hydrogen elimination to provide carbazole **2**.²⁹ Alternatively, **3** could add to the arene via a Wacker-like process to provide **5b**, which would then undergo a β-hydrogen elimination to provide carbazole **2**. The reoxidation of the Pd(0) species to Pd(II) in the presence of Cu(OAc)₂ or DMSO and oxygen would complete the catalytic cycle. To further probe these three possible pathways for the cyclization process, we undertook NMR studies of stoichiometric reactions,

measurement of inter- and intramolecular kinetic isotope effects, and the examination of substituent effects on the rate of the reaction. We also note that attempts to modify the reaction conditions by the addition of acids or bases slowed down the cyclization. Despite all our efforts, however, no clear picture of the reaction mechanism emerged from these studies.

Target-oriented Synthesis

To demonstrate the efficiency and applicability of this method, we prepared three natural occurring carbazoles (mukonine,³⁰ mukonidine and glycosinine) as shown in Scheme 4. Mukonidine and glycosinine were synthesized from a common acetylcarbazole intermediate **6a** that was formed in 94% yield from biarylamine **5a** using this method. Biarylamine **5a** was obtained in 98% yield using a Suzuki-Miyaura cross coupling protocol recently developed in this lab.²⁵ Demethylation of **6a** followed by acid hydrolysis afforded mukonidine in 85% yield. Separately, reduction of the methyl ester and cleavage of the amide group of **6a** followed by reoxidation of the resulting benzyl alcohol to an aldehyde provided glycosinine in 74% yield. The synthesis of mukonine followed a sequence similar to those of mukonidine and glycosinine. Biarylamine **5b** was obtained in 95% yield using our protocol for a Suzuki-Miyaura cross coupling between phenylboronic acid and aryl iodide **7** (obtained in 3 steps from commercially available 4-amino-3-methoxybenzoic acid following published procedures³¹). Treatment of **5b** with Pd(OAc)₂, Cu(OAc)₂, and oxygen in toluene followed by acid hydrolysis afford mukonine in 99% yield in two steps. In short, mukonidine and glycosinine were synthesized in 79% and 68% overall yield respectively in 4 steps from commercially available reagents, while mukonine was synthesized in 74% overall yield in 6 steps from commercially available reagents.

Conclusion

The results presented in this paper describe the first applications of intramolecular conversion of an arene C-H bond to a C-N bond in the elaboration of synthetically useful molecules from commercially available and user-friendly reagents. This C-H functionalization strategy allows the assembly of carbazoles with a variety of substitution patterns and functional groups under relatively mild conditions. The utility of this method has been unambiguously demonstrated by the high efficiency displayed in the syntheses of mukonidine, glycosinine, and mukonine. The observation that DMSO could be used in place of Cu(OAc)₂ broadens the substrate scope of this reaction by increasing the number of compatible functional groups. For example, both electron-donating and electron-withdrawing groups are generally tolerated in the synthesis of these carbazoles at 120 °C. Even strong electron-withdrawing groups such as keto, ester, nitro, and cyano groups can all be incorporated into the carbazoles. Additionally, this method proves particularly effective for the synthesis of unsymmetrical carbazoles and is complementary to other syntheses of carbazole alkaloids. This feature differentiates it from known methods for the synthesis of carbazoles.^{19,32,33,34} For example, the introduction of substituents by traditional electrophilic aromatic substitution of 9H-carbazole is normally limited to 3 and 6 positions that are *para* to the NH moiety.^{12d,35} Our method allows efficient installation of substituents at positions *ortho* and *para* to the biaryl axis of the amide substrate and leads to carbazoles that are substituted at 2 and/or 4 positions. While the Pd-catalyzed cyclization of *ortho*-halo diarylamines to carbazoles generally favors 2-substituted carbazoles over 4-substituted ones,^{33,34} our method permits extremely efficient synthesis of 4-substituted carbazoles. Using this method, the carbazole cores substituted from C1 to C8 can be prepared. Although we have undergone a number of studies to probe the mechanism for the cyclization process, no clear picture about its detailed course has emerged from these studies.

Experimental Section

General Procedure for the Synthesis of Carbazoles in the Presence of anhydrous Cu(OAc)₂ (Protocol A)

An oven-dried Schlenk tube (15 mL in volume) was cooled under vacuum and Ar. Biarylamide (0.2 mmol), Pd(OAc)₂ (2.3 mg, 0.01 mmol, 5%), and powdered molecular sieves (40 mg, activated 3 Å) were added under air. The tube was evacuated and refilled with Ar. Anhydrous Cu(OAc)₂ (36.3 mg, 0.2 mmol) was loaded into the Schlenk tube in a nitrogen-filled glovebox. Under a positive Ar pressure, toluene (2 mL) was added via syringe. The reaction mixture was sonicated and degassed under a weak vacuum and refilled with O₂ from the double manifold (this sequence was carried out three times). The sealed Schlenk tube was lowered into an oil bath at 120 °C and stirred for 12–24 h. After cooling to room temperature, distilled water (1 mL), ammonium hydroxide solution (3 mL, 30%), and EtOAc (2 mL) were added to the reaction mixture. The organic phase was separated, and the aqueous phase was further extracted with EtOAc (3 × 2 mL). The combined organic layers were dried over anhydrous Na₂SO₄ and filtered through Celite, and concentrated. The residue was purified by silica gel chromatography using a mixture of hexanes and EtOAc to provide the desired carbazole. A representative example is as follows.

9-Acetyl-2-acetylcarbazole

Following Protocol A (12 h), an off-white powder was obtained after silica gel chromatography (gradient; hexanes:EtOAc, 95:5 to 60:40), yielding 44 mg (87%, mp 110 °C). ¹H NMR (500 MHz, CDCl₃) δ 8.91 (s, 1H), 8.22 (d, *J* = 8.4 Hz, 1H), 8.12–7.98 (m, 3H), 7.57 (t, *J* = 7.9 Hz, 1H), 7.44 (t, *J* = 7.5 Hz, 1H), 2.95 (s, 3H), 2.74 (s, 3H). ¹³C NMR (125 MHz, CDCl₃) δ 198.04, 170.31, 140.08, 138.70, 136.22, 130.46, 128.96, 125.61, 124.24, 124.19, 121.01, 119.84, 116.92, 116.50, 28.07, 27.19. IR (KBr plate, CDCl₃) ν 3004, 1693, 1678, 1418, 1368, 1326, 1304, 1229, 1020, 749. Anal. Calcd. for C₁₆H₁₃NO₂: C, 76.48; H, 5.21. Found: C, 76.30; H, 5.25.

General Procedure for the Synthesis of Carbazoles in Dimethyl Sulfoxide (DMSO) (Protocol C)

An oven-dried Schlenk tube (15 mL in volume) was cooled under vacuum or Ar. Biarylamide (0.2 mmol), Pd(OAc)₂ (2.3 to 4.6 mg, 0.01 to 0.02 mmol, 5 to 10 %), and powdered molecular sieves (40 mg, activated 3 Å) were added under air. The tube was evacuated and refilled with Ar. Under a positive Ar pressure, DMSO (2 mL) was added via syringe. The reaction mixture was sonicated and degassed under a weak vacuum and refilled with O₂ from the double manifold (this sequence was carried out three times). The sealed Schlenk tube was lowered into an oil bath at 120 °C and stirred for 24 h. After cooling to room temperature, distilled water (5 mL) and EtOAc (3 mL) were added to the reaction mixture. The organic phase was separated, and the aqueous phase was further extracted with EtOAc (3 × 3 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered through Celite, and concentrated. The residue was purified by silica gel chromatography using a mixture of hexanes and EtOAc to provide the desired carbazole. A representative example is as follows.

9-Acetyl-2-methylthiocarbazole

Following Protocol C (5% Pd(OAc)₂, 24 h), a white powder was obtained after silica gel chromatography (gradient; hexanes:EtOAc, 95:5 to 90:10), yielding 43.0 mg (84%, mp 79 °C). ¹H NMR (500 MHz, CDCl₃) δ 8.27 (s, 1H), 8.09 (d, *J* = 8.4 Hz, 1H), 7.95 (d, *J* = 7.7 Hz, 1H), 7.89 (d, *J* = 8.2 Hz, 1H), 7.47 (t, *J* = 7.8 Hz, 1H), 7.39 (t, *J* = 7.5 Hz, 1H), 7.32 (dd, *J* = 8.1, 1.5 Hz, 1H), 2.89 (s, 3H), 2.61 (s, 3H). ¹³C NMR (125 MHz, CDCl₃) δ 170.31, 139.57, 138.59, 138.34, 127.20, 126.53, 124.05, 123.97, 122.94, 120.03, 119.90, 116.11, 114.73, 28.02,

16.75. IR (KBr plate, CDCl_3) ν 2919, 1693, 1596, 1435, 1411, 1367, 1325, 1303, 1272, 1213, 1189, 1020, 765, 747. Anal. Calcd. for $\text{C}_{15}\text{H}_{13}\text{NOS}$: C, 70.56; H, 5.13. Found: C, 70.40; H, 5.17.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We thank Johnson & Johnson (Focused Giving Grant), the National Institutes of Health (GM58160), and the National Cancer Institute (Cancer Training Grant CA 09112) for supporting this research. G.B. thanks the Deutscher Akademischer Austauschdienst (DAAD) for support through a postdoctoral fellowship. N.Z. thanks the National Institutes of Health for a postdoctoral fellowship (F32-GM074407).

REFERENCE

1. (a) Knölker H-J. *Top. Curr. Chem* 2005;244:115–148. (b) Knölker H-J, Reddy KR. *Chem. Rev* 2002;102:4303–4427. [PubMed: 12428991]
2. Chakraborty, DP. *The Alkaloids: Chemistry and Pharmacology*. Cordell, GA., editor. Vol. Vol. 44. New York: Academic; 1993. p. 257–364. Chakraborty, DP.; Roy, S. *Progress in the Chemistry of Organic Natural Products*. Herz, W.; Kirby, GW.; Steglich, W.; Tamm, Ch, editors. Vol. Vol. 57. New York: Springer-Verlag; 1991. p. 72–152. Bhattacharyya, P.; Chakraborty, DP. *Progress in the Chemistry of Organic Natural Products*. Herz, W.; Grisebach, H.; Kirby, GW.; Tamm, Ch, editors. Vol. Vol. 52. New York: Springer-Verlag; 1987. p. 159–209. Chakraborty, DP. *Progress in the Chemistry of Organic Natural Products*. Herz, W.; Grisebach, H.; Kirby, GW., editors. Vol. Vol. 34. New York: Springer-Verlag; 1977. p. 299–371.
3. (a) Krah MP, Jäger A, Krause T, Knölker H-J. *Org. Biomol. Chem* 2006;4:3215–3219. [PubMed: 17036106] (b) Meragelman KM, McKee TC, Boyd MR. *J. Nat. Prod* 2000;63:427–428. [PubMed: 10757740] (c) TePaske MR, Gloer JB, Wicklow DT, Dowd PF. *J. Org. Chem* 1989;54:4743–4746. (d) Kondo S, Katayama M, Marumo S. *J. Antibiot* 1986;39:727–730. [PubMed: 3733523] (e) Chakraborty DP, Bose PK. *Experientia* 1965;21:1340.
4. Sakano K-I, Ishimaru K, Nakamura S. *J. Antibiot* 1980;33:683–689. [PubMed: 7410212]
5. (a) Crich D, Rumthao S. *Tetrahedron* 2004;60:1513–1516. (b) Hook DJ, Yacobucci JJ, O'Connor S, Lee M, Kerns E, Krishnan B, Matson J, Hesler G. *J. Antibiot* 1990;43:1347–1348. [PubMed: 2258335]
6. (a) Guthrie RW, Brossi A, Mennona FA, Mullin JG, Kierstead RW, Grunberg E. *J. Med. Chem* 1975;18:755–760. [PubMed: 1151997] (b) Cranwell PA, Saxton JE. *J. Chem. Soc* 1962:3482–3487. (c) Woodward RB, Iacobucci GA, Hochstein FA. *J. Am. Chem. Soc* 1959;81:4434–4435.
7. (a) Cheng H-Y, Randall CS, Holl WW, Constantinides PP, Yue T-L, Feuerstein GZ. *Biochim. Biophys. Acta* 1996;1284:20–28. [PubMed: 8865810] (b) Yue T-L, Cheng H-Y, Lysko PG, McKenna PJ, Feuerstein R, Gu J-L, Lysko KA, Davis LL, Feuerstein G. *J. Pharmacol. Exp. Ther* 1992;263:92–98. [PubMed: 1357162]
8. (a) Dubois EA, van den Bos JC, Doornbos T, van Doremalen PAM, Somsen GA, Vekemans JAJM, Janssen AGM, Batink HD, Boer GJ, Pfaffendorf M, van Royen EA, van Zwieten PA. *J. Med. Chem* 1996;39:3256–3262. [PubMed: 8765508] (b) Costin B, O'Donnell SR, Wanstall JC. *J. Pharm. Pharmacol* 1983;35:590–592. [PubMed: 6138410] (c) Cohen ML, Ruffolo RR Jr, Wiley KS. *J. Pharmacol. Exp. Ther* 1980;215:325–331. [PubMed: 6108357] (d) Innis RB, Corrêa FMA, Snyder SH. *Life Sci* 1979;24:2255–2264. [PubMed: 41147]
9. (a) Czeskis BA, Wheeler WJ. *J. Label. Compd. Radiopharm* 2005;48:407–419. (b) Nakajima Y, Hamashima H, Washizuka K-I, Tomishima Y, Ohtake H, Imamura E, Miura T, Kayakiri H, Kato M. *Bioorg. Med. Chem. Lett* 2005;15:251–254. [PubMed: 15603933] (c) Kumar R, Ramachandran U, Srinivasan K, Ramarao P, Raichur S, Chakrabarti R. *Bioorg. Med. Chem* 2005;13:4279–4290. [PubMed: 15869880] (d) Sum F-W, Wong V, Han S, Largis E, Mulvey R, Tillett J. *Bioorg. Med. Chem. Lett* 2003;13:2191–2194. [PubMed: 12798332]

10. (a) Morin J-F, Leclerc M, Adès D, Siove A. *Macromol. Rapid Commun* 2005;26:761–778. (b) Ostroverkhova O, Moerner WE. *Chem. Rev* 2004;104:3267–3314. [PubMed: 15250742] (c) Grazulevicius JV, Strohrriegl P, Pielichowski J, Pielichowski K. *Prog. Polym. Sci* 2003;28:1297–1353. (d) Zhang Y, Wada T, Sasabe H. *J. Mater. Chem* 1998;8:809–828. (e) Meerholz K, Volodin BL, Sandalphon, Kippelen B, Peyghambarian N. *Nature* 1994;371:497–500.
11. Friend RH, Gymer RW, Holmes AB, Burroughes JH, Marks RN, Taliani C, Bradley DDC, Dos Santos DA, Brédas JL, Lögdlund M, Salaneck WR. *Nature* 1999;397:121–128.
12. Müllen, K.; Scherf, U., editors. *Organic Light Emitting Devices: Synthesis, Properties, and Applications*. Weinheim: Wiley-VCH; 2006. (b) Grigalevicius S. *Synth. Met* 2006;156:1–12. (c) Brunner K, van Dijken A, Börner H, Bastiaansen JJAM, Kiggen NMM, Langeveld BMW. *J. Am. Chem. Soc* 2004;126:6035–6042. [PubMed: 15137768] (d) Thomas KRJ, Lin JT, Tao Y-T, Ko C-W. *J. Am. Chem. Soc* 2001;123:9404–9411. [PubMed: 11562223] (e) Baldo MA, O'Brien DF, You Y, Shoustikov A, Sibley S, Thompson ME, Forrest SR. *Nature* 1998;395:151–154.
13. (a) Tsai M-H, Hong Y-H, Chang C-H, Su H-C, Wu C-C, Matoliukstyte A, Simokaitiene J, Grigalevicius S, Grazulevicius JV, Hsu C-P. *Adv. Mater* 2007;19:862–866. (b) Yeh S-J, Wu M-F, Chen C-T, Song Y-H, Chi Y, Ho M-H, Hsu S-F, Chen CH. *Adv. Mater* 2005;17:285–289. (c) Tokito S, Iijima T, Suzuki Y, Kita H, Tsuzuki T, Sato F. *Appl. Phys. Lett* 2003;83:569–571. (d) Holmes RJ, Forrest SR, Tung Y-J, Kwong RC, Brown JJ, Garon S, Thompson ME. *Appl. Phys. Lett* 2003;82:2422–2424. (e) Adachi C, Kwong RC, Djurovich P, Adamovich V, Baldo MA, Thompson ME, Forrest SR. *Appl. Phys. Lett* 2001;79:2082–2084.
14. (a) Chen C-T. *Chem. Mater* 2004;16:4389–4400. (b) Lamansky S, Djurovich P, Murphy D, Abdel-Razzaq F, Lee H-E, Adachi C, Burrows PE, Forrest SR, Thompson ME. *J. Am. Chem. Soc* 2001;123:4304–4312. [PubMed: 11457197] (c) Adachi C, Baldo MA, Forrest SR, Lamansky S, Thompson ME, Kwong RC. *Appl. Phys. Lett* 2001;78:1622–1624. (d) Baldo MA, Lamansky S, Burrows PE, Thompson ME, Forrest SR. *Appl. Phys. Lett* 1999;75:4–6.
15. For recent reviews related to C-H amination, see: Dick AR, Sanford MS. *Tetrahedron* 2006;62:2439–2463. (b) Davies HML, Long MS. *Angew. Chem. Int. Ed* 2005;44:2–4. (c) Sigman MS, Schultz MJ. *Org. Biomol. Chem* 2004;2:2551–2554. [PubMed: 15351815] (d) Stahl S. *Angew. Chem. Int. Ed* 2004;43:3400–3420. (e) Müller P, Fruit C. *Chem. Rev* 2003;103:2905–2919. [PubMed: 12914485]
16. For recent reviews of other C-H functionalization or activation processes, see: Godula K, Sames D. *Science* 2006;312:67–72. (b) Sigman MS, Jensen DR. *Acc. Chem. Res* 2006;39:221–229. [PubMed: 16548511] (b) Kakiuchi F, Chatani N. *Adv. Synth. Catal* 2003;345:1077–1101. (c) Kakiuchi F, Murai S. *Acc. Chem. Res* 2002;35:826–834. [PubMed: 12379135] (d) Ritleng V, Sirlin C, Pfeffer M. *Chem. Rev* 2002;102:1731–1769. [PubMed: 11996548] (e) Jia C, Kitamura T, Fujiwara Y. *Acc. Chem. Res* 2001;34:633–639. [PubMed: 11513570] (f) Bringmann G, Menche D. *Acc. Chem. Res* 2001;34:615–624. [PubMed: 11513568] (g) Dyker G. *Angew. Chem. Int. Ed* 1999;38:1698–1712. (h) Shilov AE, Shul'pin GB. *Chem. Rev* 1997;97:2879–2932. [PubMed: 11851481] (i) Bringmann G, Walter R, Weirich R. *Angew. Chem. Int. Ed. Engl* 1990;29:977–991.
17. (a) Espino CG, Fiori KW, Kim M, Du Bois J. *J. Am. Chem. Soc* 2004;126:15378–15379. [PubMed: 15563154] (b) Fiori KW, Fleming JJ, Du Bois J. *Angew. Chem. Int. Ed* 2004;43:4349–4352. (c) Fleming JJ, Fiori KW, Du Bois J. *J. Am. Chem. Soc* 2003;125:2028–2029. [PubMed: 12590513] (d) Wehn PM, Du Bois J. *J. Am. Chem. Soc* 2002;124:12950–12951. [PubMed: 12405813] (e) Espino CG, Du Bois J. *Angew. Chem. Int. Ed* 2001;40:598–600. (f) Espino CG, Wehn PM, Chow J, Du Bois J. *J. Am. Chem. Soc* 2001;123:6935–6936.
18. (a) Liang J-L, Yuan S-X, Huang J-S, Che C-M. *J. Org. Chem* 2004;69:3610–3619. [PubMed: 15152988] (b) Liang J-L, Yuan S-X, Huang J-S, Yu W-Y, Che C-M. *Angew. Chem. Int. Ed* 2002;41:3465–3468. (c) Reed AA, White MC. *J. Am. Chem. Soc* 2008;130:3316–3318. [PubMed: 18302379]
19. (a) Kalyani D, Deprez NR, Desai LV, Sanford MS. *J. Am. Chem. Soc* 2005;127:7330–7331. [PubMed: 15898779] (b) Daugulis O, Zaitsev VG. *Angew. Chem. Int. Ed* 2005;44:4046–4048. (c) Kakiuchi F, Matsuura Y, Kan S, Chatani N. *J. Am. Chem. Soc* 2005;127:5936–5945. [PubMed: 15839693] (d) Cuny G, Bois-Choussy M, Zhu J. *J. Am. Chem. Soc* 2004;126:14475–14484. [PubMed: 15521768] (e) Campeau L-C, Parisien M, Leblanc M, Fagnou K. *J. Am. Chem. Soc* 2004;126:9186–9187. [PubMed: 15281800] (f) Huang Q, Fazio A, Dai G, Campo MA, Larock RC. *J. Am. Chem. Soc*

- 2004;126:7460–7461. [PubMed: 15198591] (g) Bedford RB, Coles SJ, Hursthouse MB, Limmert ME. *Angew. Chem., Int. Ed* 2003;42:112–114. (h) Terao Y, Wakui H, Satoh T, Miura M, Nomura M. *J. Am. Chem. Soc* 2001;123:10407–10408. [PubMed: 11604000] (i) Ames DE, Opalko A. *Tetrahedron* 1984;40:1919–1925. (j) Åkermark B, Eberson L, Jonsson E, Pettersson E. *J. Org. Chem* 1975;40:1365–1367.
20. For recent reviews of Pd- and Cu-catalyzed carbon-heteroatom bond formation processes, see: Corbet J-P, Mignani G. *Chem. Rev* 2006;106:2651–2710. [PubMed: 16836296] (b) Wolfe JP, Thomas JS. *Curr. Org. Chem* 2005;9:625–655. Jiang, L.; Buchwald, SL. *Metal-Catalyzed Cross-Coupling Reactions*. Vol. 2nd ed. de Meijere, A.; Diederich, F., editors. Vol. Vol. 2. Weinheim: Wiley-VCH; 2004. p. 699–760. (d) Beletskaya IP, Cheprakov AV. *Coord. Chem. Rev* 2004;248:2337–2364. (e) Ley SV, Thomas AW. *Angew. Chem., Int. Ed* 2003;42:5400–5449. Hartwig, JF. *Handbook of Organopalladium Chemistry for Organic Synthesis*. Negishi, E-I.; de Meijere, A., editors. Vol. Vol. 1. New York: Wiley- Interscience; 2002. p. 1051–1096.
21. Tsang WCP, Zheng N, Buchwald SL. *J. Am. Chem. Soc* 2005;127:14560–14561. [PubMed: 16231894]
22. Li B-J, Tian S-L, Fang Z, Shi Z-J. *Angew. Chem. Int. Ed* 2008;47:1115–1118.
23. (a) Thu HY, Yu W-Y, Che C-M. *J. Am. Chem. Soc* 2006;128:9048–9049. [PubMed: 16834374] (b) Yip K-T, Yang M, Law K-L, Zhu N-Y, Yang D. *J. Am. Chem. Soc* 2006;128:3130–3131. [PubMed: 16522078] (c) Dick AR, Hull KL, Sanford MS. *J. Am. Chem. Soc* 2004;126:2300–2301. [PubMed: 14982422] (d) Boele MDK, van Strijdonck GPF, de Vries AHM, Kamer PCJ, de Vries JG, van Leeuwen PWNM. *J. Am. Chem. Soc* 2002;124:1586–1587. [PubMed: 11853427]
24. (a) Steinhoff BA, Fix SR, Stahl SS. *J. Am. Chem. Soc* 2002;124:766–767. [PubMed: 11817948] (b) Chen MS, White MC. *J. Am. Chem. Soc* 2004;126:1346–1347. [PubMed: 14759185]
25. (a) Barder TE, Walker SD, Martinelli JR, Buchwald SL. *J. Am. Chem. Soc* 2005;127:4685–4696. [PubMed: 15796535] (b) Walker SD, Barder TE, Martinelli JR, Buchwald SL. *Angew. Chem. Int. Ed* 2004;43:1871–1876.
26. Krahle MP, Jäger A, Krause T, Knölker H-J. *Org. Biomol. Chem* 2006;4:3215–3219. [PubMed: 17036106]
27. Meragelman KM, McKee TC, Boyd MR. *J. Nat. Prod* 2000;63:427–428. [PubMed: 10757740]
28. Inamoto K, Saito T, Katsuno M, Sakamoto T, Hiroya K. *Org. Lett* 2007;9:2931–2934. [PubMed: 17595097]
29. For one selected example of intramolecular Heck reaction via a trans β -hydrogen elimination, see: Lautens M, Fang Y-Q. *Org. Lett* 2003;5:3679–3682. [PubMed: 14507203]
30. Liégault B, Lee D, Huestis MP, Stuart DR, Fagnou K. *J. Org. Chem* 2008;73:5022–5028. [PubMed: 18543991]
31. Ezquerro J, Pedregal C, Lamas C. *J. Org. Chem* 1996;61:5804–5812.
32. Campeau L-C, Parisien M, Jean A, Fagnou K. *J. Am. Chem. Soc* 2006;128:581–590. [PubMed: 16402846]
33. Liu Z, Larock RC. *Tetrahedron* 2007;63:347–355. [PubMed: 18180808]
34. Bedford RB, Betham M. *J. Org. Chem* 2006;71:9403–9410. [PubMed: 17137367]
35. (a) Zhu Z, Moore JS. *J. Org. Chem* 2000;65:116–123. [PubMed: 10813904] (b) Zhu Z, Moore JS. *Macromolecules* 2000;33:801–807. (c) Buu-Hoi NP, Hoán N. *J. Org. Chem* 1951;16:1327–1332. (d) Buu-Hoi NP, Royer R. *J. Org. Chem* 1951;16:1198–1205. (e) Tucker SH. *J. Chem. Soc* 1926:546–553.

Carbazomycin B

Carvedilol

Ellipticine

Carazolol

Figure 1.
Therapeutically Valuable Carbazole Alkaloids

Scheme 1.
Synthesis of Biaryl Acetamides by Suzuki-Miyaura Coupling

Scheme 2.
Our Initial Proposed Pathway for Carbazole Synthesis

Scheme 3.
Two Alternative Pathways for Carbazole Synthesis

Scheme 4.
Synthesis of Three Natural Occurring Carbazole

Table 1
Investigation of Palladium Precatalysts and Reoxidants

<div><div> </div><div> </div><div>120 ° C</div><div>5% Pd catalyst reoxidant, gas</div><div>solvent,</div></div>						
Entry	Pd source	Reoxidant /equiv	Solvent	Gas atm	Time (h)	Yield (%) ^a
1	Pd(OAc) ₂	Cu(OAc) ₂ / 1	toluene	O ₂	12	92
2	Pd(OAc) ₂		toluene	O ₂	12	5
3		Cu(OAc) ₂ / 1	toluene	O ₂	12	0
4	Pd(O ₂ CCF ₃) ₂	Cu(OAc) ₂ / 1	toluene	O ₂	24	70
5	PdCl ₂	Cu(OAc) ₂ / 1	toluene	O ₂	24	21
6	PdCl ₂ (CH ₃ CN) ₂	Cu(OAc) ₂ / 1	toluene	O ₂	24	27
7	PdCl ₂ (PPh ₃) ₂	Cu(OAc) ₂ / 1	toluene	O ₂	24	67 ^b
8	Pd(OAc) ₂	PPh ₃ / 0.2	toluene	O ₂	12	5
9	Pd(OAc) ₂	pyridine / 0.2	toluene	O ₂	12	1
10	Pd(OAc) ₂	PhI(OAc) ₂ / 0.2	DCE ^c	air	12	25
11	Pd(OAc) ₂	K ₂ S ₂ O ₈ / 5	DCE ^d	air	12	2
12	Pd(OAc) ₂	benzoquinone / 1	toluene	air	12	4

^a GC yield was determined with respect to a calibrated internal standard.

^b About 5–10% diacetylated amide of 2-aminobiphenyl was observed.

^c Slightly higher yield was obtained in dichloroethane than in methylene chloride.

^d The reaction was performed at 80 °C in dichloroethane; the same low yield was observed with or without bases.

Table 2

Evaluation of Reoxidants in the Synthesis of Carbazoles

Entry	R	Reoxidant /equiv	Solvent	Gas atm	Time (h)	Yield (%) ^a
1	H	Cu(OAc) ₂ / 1	toluene	air	24	94
2	H	Cu(OAc) ₂ / 1	toluene	O ₂	12	92 ^b
3	H	Cu(OAc) ₂ / 0.2	toluene	O ₂	12	90
4	OMe	Cu(OAc) ₂ / 0.2	toluene	O ₂	12	88
5	CF ₃	Cu(OAc) ₂ / 0.2	toluene	O ₂	12	76
6	H		DMSO	O ₂	12	92 ^b
7	H		DMF	O ₂	12	11 ^b

^a Isolated yield after silica gel chromatography.

^b GC yield was determined with respect to a calibrated internal standard.

Table 3

Effect of Nitrogen Protecting Groups

Entry	R	Reoxidant /equiv	Solvent	Yield (%) ^a
1	Ac	Cu(OAc) ₂ / 1	toluene	92
2	Ac		DMSO	92
3	SO ₂ Ph	Cu(OAc) ₂ / 1	toluene	81 ^b
4	SO ₂ Ph		DMSO	99
5	BOC	Cu(OAc) ₂ / 1	toluene	34
6	BOC		DMSO	30
7	COPh	Cu(OAc) ₂ / 1	toluene	<10 ^c
8	COPh		DMSO	<10 ^c
9	COCF ₂ CF ₃	Cu(OAc) ₂ / 1	toluene	0
10	COCF ₂ CF ₃		DMSO	0

^a Isolated yield after silica gel chromatography, average of two runs.^b 10% Pd(OAc)₂ was used.^c Yield estimated from GC analysis.

Table 4
Synthesis of Substituted Carbazoles by Cyclization of Biaryl Amides

Entry

Substrate

1

2

Entry

Substrate

3

4

5

6

7

8

Entry

Substrate

9

Entry

Substrate

10

^a Isolated yield after silica gel chromatography, average of two runs.

Table 5

Investigation of Functional Group Compatibility

Entry	R	Reoxidant/ equiv	Solvent	Yield (%) ^a
1	F	Cu(OAc) ₂ / 1	toluene	87
2	Me	Cu(OAc) ₂ / 1	toluene	98
3	OMe	Cu(OAc) ₂ / 1	toluene	97
4	CF ₃	Cu(OAc) ₂ / 1	toluene	94
5	COMe	Cu(OAc) ₂ / 1	toluene	86
6	CO ₂ Me	Cu(OAc) ₂ / 1	toluene	96
7	SMe		DMSO	82
8	NO ₂		DMSO	72 ^b
9	CN		DMSO	84 ^b

^a Isolated yield after silica gel chromatography, average of two runs.^b 10% Pd(OAc)₂ was used.

Table 6

Regioselective Synthesis of 3-substituted Carbazoles

Entry	R	Reoxidant/ equiv	Solvent	Yield (%) ^a
1	Me	Cu(OAc) ₂ / 1	toluene	82 ^b
2	OMe	Cu(OAc) ₂ / 1	toluene	41 ^c
3	OMe		DMSO	59
4	CF ₃	Cu(OAc) ₂ / 1	toluene	86
5	CO ₂ Me	Cu(OAc) ₂ / 1	toluene	99
6	OTIPS	Cu(OAc) ₂ / 1	toluene	81
7	TBS	Cu(OAc) ₂ / 1	toluene	98

^a Isolated yield after silica gel chromatography, average of two runs. Unless otherwise stated, the regioisomeric ratio is >25:1 (GC); only the major product is shown.

^b Selectivity = 18:1 (GC).

^c Selectivity = 16:1 (GC).

Table 7

Reactivity Studies at 90 °C

Entry	R	Temp (°C)	Yield (%) ^a
1	H	120	92
2	2-CF ₃	90	3
3	H	90	10
4	2-Me	90	37
5	2-OMe	90	73
6	3-CF ₃	90	7
7	3-Me	90	14
8	3-OMe	90	8

^a GC yield was determined relative to an internal standard.