Published in final edited form as:

J Org Chem. 2009 September 18; 74(18): 7180–7182. doi:10.1021/jo901111j.

Silylene-Mediated Ring Contraction of Homoallylic Ethers to Form Allylic Silanes

Laura E. Bourque, Pamela A. Haile, and K. A. Woerpel*

Department of Chemistry, University of California, Irvine, Irvine, California 92697-2025

Abstract

(—)-Isopulegol derivatives undergo a ring contraction under silylene-mediated conditions to provide cyclopentane products. Silylene transfer to other homoallylic ethers did not provide the ring contraction products. Allylic silane products were elaborated to determine the stereochemical course of the ring contraction reaction. A mechanism for the transformation is proposed.

Cyclopentane units are found in many natural and non-natural products, including alkaloids, steroids, prostaglandins, triquinanes, and guaianes.^{1–6} The ring contraction of a six-membered carbocyclic compound is an efficient way to assemble a cyclopentane, because the reorganization of the bonds can occur with high selectivity.^{7–12} This reorganization leads to compounds not easily accessed by other syntheses.¹³ In this Note, we report the ring contraction of six-membered carbocyclic homoallylic ethers to form five-membered carbocyclic allylic silanes by treatment with silylene intermediates.

As part of our investigations into silylene transfer reactions to homoallylic ethers,¹⁴ we examined the reaction of (—)-isopulegol derivative **1** under the optimized reaction conditions. Subjection of benzyl ether **1** to cyclohexene silacyclopropane **2** and AgO₂CCF₃ did not provide the expected silylmethyl allylic silane product. Instead, the five-membered ring allylic silane product **3** was isolated as a single stereoisomer (eq 1).

*kwoerpel@uci.edu.

Supporting Information Available Complete experimental procedures, X-ray data, and product characterization. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(1)

Alteration of the protecting group on the homoallylic ether provided similar ring contraction products. When alkyl ether **4** was treated to the reaction conditions, allylic silane **5** was observed (eq 2). Treatment of silyl ether **6** to the reaction conditions provided allylic silane **7** (eq 2). Both allylic silanes were found to be predominantly one stereoisomer, as indicated by ^1H NMR and ^{13}C NMR spectroscopy.^{15,16}

(2)

A variety of homoallylic ethers were subjected to the reaction conditions, but they did not undergo rearrangement. *epi*-Isopulegol derivative **8** underwent silylene transfer to the alkene¹⁷ to form silacyclopropane **9** as a mixture of diastereomers that was not stable to purification (eq 3). The product did not undergo further transformations upon heating. The same result was observed with both five-membered ring ether **10** and seven-membered ring ether **12** (eq 4). The lack of ring-contraction of the five-membered ring silacyclopropane **11** (eq 4) may be the result of too much ring strain involved in contraction to the four-membered ring.¹⁸ Differences in conformational preferences and flexibility between six- and seven-membered rings may cause the seven-membered ring system of silacyclopropane **13** to align orbitals differently,¹⁹ thus preventing ring contraction.

(3)

(4)

The structure and relative stereochemistry of allylic silane **3** was determined by X-ray crystallography of a derivative. Allylic silane **3** underwent hydroboration, followed by oxidation, to form oxasilacyclopentane **14** as a single isomer (eq 5).²⁰ Oxasilacyclopentane **14** was then subjected to modified Tamao—Fleming oxidation conditions^{21–23} to remove the di-*tert*-butylsilylene moiety, providing diol **15** (eq 6). The stereochemistry and connectivity of diol **15** was established by X-ray crystallography.²⁴

(6)

A stepwise mechanism is proposed to account for the contraction of the original six-membered ring to form the five-membered ring (Scheme 1). Silylene transfer to the double bond of **1** would form silacyclopropane adduct **16** as one diastereomer.²⁵ The oxygen atom can then complex to the Lewis-acidic silicon atom²⁶ to form the conformationally favored trans-fused ring system **17** (Figure 1).^{27,28} The cis-fused ring system (**20**) appears to encounter steric hindrance due to a *t*-Bu group being forced over the cyclohexane ring.²⁹ The ring-contracting step forms silacyclopropylcarbinyl cation **18** with inversion of configuration. Carbocation **18** undergoes silacyclopropylcarbinyl cation rearrangement^{30,31} to form ylide **19**. The nucleophilic pentavalent silicon atom can then reform the terminal alkene, providing allylic silane **3**.^{30,31}

In summary, silylene transfer to (—)-isopulegol derivatives provided allylic silanes by a ring contraction. Similar five- and seven-membered ring systems underwent silylene transfer to the alkene, but did not provide ring contraction products. A mechanism for the transformation was proposed utilizing a silylene ylide intermediate and a silacyclopropylcarbinyl rearrangement.

Experimental Section

Allylic silane **3**

To a solution of homoallylic ether **1** (0.394 g, 1.60 mmol) in 8.0 mL of toluene was added cyclohexene silacyclopropane **2** (0.470 g, 2.09 mmol). AgO₂CCF₃ (0.004 g, 0.02 mmol) was then added. The brown solution was then placed under an Ar atmosphere and allowed to stir for 1 hour at 50 °C, at which point the mixture was concentrated *in vacuo*. Purification by flash chromatography (hexanes) provided 0.402 g (65%) of **3** as a colorless oil: ¹H NMR (500 MHz, CDCl₃) δ 7.28–7.35 (m, 4H), 7.19–7.23 (m, 1H), 4.91–5.01 (m, 2H), 4.74–4.76 (m, 2H), 2.22–2.32 (m, 1H), 2.01–2.07 (m, 2H), 1.80–1.93 (m, 1H), 1.61–1.77 (m, 5H), 1.02–1.29 (m, 20H), 0.89 (d, *J* = 6.6, 3H), 0.71–0.79 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 147.7, 142.1, 128.3, 126.8, 126.2, 113.2, 66.6, 44.7, 43.7, 41.1, 34.3, 32.4, 32.3, 30.1, 29.8, 25.0, 23.8, 23.2, 21.5; IR (thin film) 2951, 2860, 2360, 1112, 820, 726 cm^{−1}; HRMS (ESI) *m/z* calcd for C₂₅H₄₂OSiNa (M + Na)⁺ 409.2903, found 409.2910.

Homoallylic ether **4**

To a 0 °C solution of sodium hydride (0.710 g, 29.6 mmol) in 50 mL of THF was added (—)-isopulegol (2.50 mL, 14.8 mmol) dropwise followed by methyl iodide (1.01 mL, 16.3 mmol). The reaction mixture was warmed to room temperature and stirred for twelve hours. The solution was then cooled to 0 °C and diluted with 20 mL of H₂O. The aqueous layer was extracted twice with 20 mL of EtOAc. The combined organic layers were washed with 20 mL of brine, dried over Na₂SO₄, and concentrated *in vacuo*. Purification by flash chromatography (95:5 hexanes:EtOAc) provided 2.35 g (95%) of **4** as a colorless oil: ¹H NMR (500 MHz, CDCl₃) δ 4.77–4.79 (m, 2H), 3.33 (s, 3H), 3.08–3.13 (m, 1H), 2.15–2.17 (m, 1H), 1.98–2.03 (m, 1H), 1.73 (s, 3H), 1.61–1.67 (m, 3H), 1.33–1.44 (m, 2H), 0.93–0.96 (m, 3H), 0.82–0.89

(m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 148.2, 110.9, 80.7, 56.1, 51.9, 39.2, 34.6, 31.6, 31.3, 22.4, 19.6; IR (thin film) 2923, 2869, 1454, 1371, 1106, 885 cm^{-1} ; HRMS (ESI) m/z calcd for $\text{C}_{11}\text{H}_{20}\text{ONa}$ ($\text{M} + \text{Na}$) $^{+}$ 191.1412, found 191.1419.

Allylic silane 5

To a solution of homoallylic ether **4** (0.017 g, 0.01 mmol) in 0.7 mL of benzene- d_6 was added cyclohexene silacyclopropane **2** (0.029 g, 0.13 mmol). AgO_2CCF_3 (0.001 g, 0.005 mmol) was then added. The brown solution was then heated for 1 hour at 50 $^{\circ}\text{C}$, at which point the mixture was concentrated *in vacuo*. Purification by flash chromatography (hexanes) provided 0.022 g (71%) of **5** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 4.72–4.77 (m, 2H), 3.62 (s, 3H), 2.26–2.29 (m, 1H), 2.09–2.12 (m, 1H), 1.93–2.00 (m, 2H), 1.82 (s, 3H), 1.67–1.75 (m, 2H), 1.14–1.20 (m, 3H), 1.10 (s, 9H), 1.09 (s, 9H), 0.98–1.07 (m, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 147.6, 113.1, 53.1, 44.3, 43.4, 41.3, 34.4, 32.5, 32.2, 30.1, 29.6, 27.1, 24.9, 23.1, 21.7; IR (thin film) 2948, 2859, 1469, 1189, 1116, 819 cm^{-1} ; HRMS (ESI) m/z calcd for $\text{C}_{19}\text{H}_{37}\text{OSi}$ ($\text{M} - \text{H}$) $^{+}$ 309.2614, found 309.2623.

Allylic silane 7

To a solution of homoallylic ether **6** (0.027 g, 0.01 mmol) and cyclohexene silacyclopropane **2** (0.029 g, 0.13 mmol) in 0.7 mL of benzene- d_6 was added AgO_2CCF_3 (0.001 g, 0.005 mmol). The brown solution was then heated for 1 hour at 50 $^{\circ}\text{C}$, at which point the mixture was concentrated *in vacuo*. Purification by flash chromatography (hexanes) provided 0.034 g (82%) of **7** as a colorless oil: ^1H NMR (400 MHz, CDCl_3) δ 4.68–4.80 (m, 2H), 2.26–2.32 (m, 1H), 1.91–1.96 (m, 2H), 1.82 (s, 3H), 1.69–1.72 (m, 2H), 1.08 (s, 18H), 1.02–1.06 (m, 5H), 0.97–0.98 (m, 2H), 0.94 (s, 9H), 0.15 (s, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 123.2, 114.8, 44.7, 40.7, 33.7, 33.0, 31.3, 30.1, 29.6, 29.2, 28.2, 27.5, 27.1, 21.5, –1.3, –1.4; IR (thin film) 2952, 2859, 1469, 1253, 1043, 833 cm^{-1} ; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{50}\text{OSi}_2\text{Na}$ ($\text{M} + \text{Na}$) $^{+}$ 433.3298, found 433.3295.

Oxasilacyclopentane 14

To 0 $^{\circ}\text{C}$ solution of allylic silane **3** (0.440 g, 1.10 mmol) in 3.8 mL of THF was added 9-BBN (6.8 mL of a 0.5 M solution in THF, 3.3 mmol) dropwise. The reaction mixture was allowed to warm to room temperature for 8 hours. After cooling to 0 $^{\circ}\text{C}$, 1.5 mL of NaOH (3 N) and 1.5 mL of 30% H_2O_2 was added. The heterogeneous mixture was warmed to room temperature and stirred for an additional 3 hours. The mixture was then saturated with K_2CO_3 and diluted with 5 mL of CH_2Cl_2 . The layers were separated, and the organic layer was washed with 5 mL of brine. The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. Purification by flash chromatography (95:5 hexanes:EtOAc) provided 0.250 g (75%) of **14** as a colorless oil: ^1H NMR (500 MHz, CDCl_3) δ 3.82–3.86 (m, 1H), 3.70–3.72 (m, 1H), 2.23–2.31 (m, 1H), 2.04–2.13 (m, 1H), 1.86–1.97 (m, 3H), 1.72–1.82 (m, 1H), 1.35–1.37 (m, 1H), 1.16–1.26 (m, 2H), 1.11 (s, 9H), 1.00–1.05 (m, 15H), 0.83–0.90 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 73.9, 45.4, 38.8, 37.0, 36.0, 34.6, 33.7, 33.3, 29.6, 29.0, 22.3, 21.3, 21.1, 17.0; IR (thin film) 2948, 2859, 1377, 1080, 985, 822 cm^{-1} ; HRMS (ESI) m/z calcd for $\text{C}_{18}\text{H}_{37}\text{OSi}$ ($\text{M} + \text{H}$) $^{+}$ 297.2614, found 297.2613. Anal. Calcd for $\text{C}_{18}\text{H}_{36}\text{OSi}$: C, 72.60; H, 12.24. Found: C, 72.84; H, 12.30.

Diol 15

To a solution of KH (0.120 g, 2.90 mmol) in 1.5 mL THF was added 18-crown-6 (0.780 g, 2.90 mmol). The solution was cooled to 0 $^{\circ}\text{C}$, and cumene hydroperoxide (88%, 0.49 mL, 2.9 mmol) was added dropwise. The cooling bath was removed, and oxasilacyclopentane **14** (0.150 g, 0.490 mmol) was added by way of cannula in 0.5 mL of THF. The reaction mixture was heated to 50 $^{\circ}\text{C}$ for 2 hours, and Bu_4NF (1.47 mL of a 1.00 M solution in THF, 1.47 mmol)

was added. After 12 hours, the mixture was cooled room temperature, and the solution was diluted with 2 mL of saturated sodium thiosulfate and 5 mL of CH₂Cl₂. The layers were separated, and the organic layer was washed with 5 mL of brine. The organic layer was dried over Na₂SO₄ and concentrated *in vacuo*. Purification by flash chromatography (70:30 hexanes:EtOAc) provided 0.480 g (57%) of diol **15** as a white solid: mp 97 °C; ¹H NMR (500 MHz, CDCl₃) 3.72–3.75 (m, 1H), 3.64–3.68 (m, 1H), 3.56–3.59 (m, 1H), 1.89–2.33 (m, 4H), 1.69–1.81 (m, 2H), 1.56–1.66 (m, 1H), 1.10–1.29 (m, 3H), 0.84–1.04 (m, 7H); ¹³C NMR (125 MHz, CDCl₃) δ 79.9, 68.1, 44.4, 40.3, 37.7, 34.8, 33.8, 28.1, 21.0, 9.3; IR (thin film) 3307, 2917, 2861, 1454, 1029, 820 cm⁻¹; HRMS (ESI) *m/z* calcd for C₁₀H₂₀O₂Na (M + Na)⁺ 195.1361, found 195.1366. Anal. Calcd for C₁₀H₂₀O₂: C, 69.72; H, 11.70. Found: C, 69.49; H, 11.83.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledement

This research was supported by the National Institute of General Medical Sciences of the National Institutes of Health (GM-54909). L.E.B. thanks the Department of Education (GAANN) for a predoctoral fellowship. K.A.W. thanks Amgen and Lilly for awards to support research. We thank Dr. Phil Dennison (UCI) for assistance with NMR spectroscopy, Dr. Joseph W. Ziller (UCI) for X-ray crystallography, and Dr. John Greaves and Ms. Shirin Sorooshian (UCI) for mass spectrometry.

References

- (1). For examples of total synthesis, see: Corey EJ, Cheng X-M. The Logic of Chemical Synthesis. 1989WileyNew York
- (2). For examples of marine natural products, see: Faulkner DJ. Nat. Prod. Rep 2002;19:1–49.49 [PubMed: 11902436]
- (3). For examples of sesquiterpenoids, see: Fraga BM. Nat. Prod. Rep 2001;18:650–673.673 [PubMed: 11820763]
- (4). For examples of monoterpenoids, see: Grayson DH. Nat. Prod. Rep 2000;17:385–419.419 [PubMed: 11014339]
- (5). For examples of diterpenoids, see: Hanson JR. Nat. Prod. Rep 2002;19:125–132.132 [PubMed: 12013275]
- (6). For examples of triterpenoids, see: Connolly JD, Hill RA. Nat. Prod. Rep 2001;18:560–578.578 [PubMed: 11699886]
- (7). For a review concerning Wagner-Meerwein rearrangements, see: Hanson JR. Trost BM. Wagner-Meerwein Rearrangements 1991;Vol. 3PergamonNew York
- (8). Heathcock CH, Ratcliffe R. J. Am. Chem. Soc 1971;93:1746–1757.
- (9). Mann, J. The Favorskii Rearrangement. In: Trost, BM.; Fleming, I., editors. Comp. Org. Synth. Vol. Vol. 3. Pergamon Press; Oxford: 1991. p. 33–64.
- (10). Furniss, BS.; Hannaford, AJ.; Smith, PWG.; Tatchell, AR. Vogel's Textbook of Practical Organic Chemistry. Vol. 5th ed. Longman; London: 1989.
- (11). Ferraz HMC, Silva LF Jr. Tetrahedron Lett 1997;38:1899–1902.
- (12). Mathew F, Myrboh B. Synth. Commun 1996;26:1097–1101.
- (13). For a review concerning ring contractions, see: Redmore D, Gutsche CD. Hart H, Karabastos GJ. Carbocyclic Ring Contraction Reactions 1971;Vol. 3AcademicNew York
- (14). Cleary PA, Woerpel KA. Org. Lett 2005;7:5531–5533. [PubMed: 16288549]
- (15). The relative configurations of **5** and **7** were assigned based on comparisons of their ¹H NMR and ¹³C NMR spectra to those of allylic silane **3**.
- (16). Allylic silane **7** was difficult to purify, so we have not been able to identify minor components of the reaction mixture

- (17). Driver TG, Woerpel KA. J. Am. Chem. Soc 2004;126:9993–10002. [PubMed: 15303873]
- (18). Wiberg KB. Angew. Chem. Int. Ed. Engl 1986;25:312–322.
- (19). Huffman JW, Engle JE. J. Org. Chem 1959;24:1844–1847.
- (20). Fleming I. J. Chem. Soc., Perkin Trans. I 1992:3363–3369.
- (21). Smitrovich JH, Woerpel KA. J. Org. Chem 1996;61:6044–6046.
- (22). Driver TG, Franz AK, Woerpel KA. J. Am. Chem. Soc 2002;124:6524–6525. [PubMed: 12047159]
- (23)a). For reviews of C—Si bond oxidation, see: Tamao K, Larson GL. Advances in Silicon Chemistry 1996;Vol. 3:1–62.62JAIGreenwich, CTb)Fleming I. Chemtracts—Org. Chem 1996;9:1–64.64c) Jones GR, Landais Y. Tetrahedron 1996;52:7599–7662.7662
- (24). The details of the crystallographic studies are provided as Supporting Information
- (25). A similar substrate gave this stereoselectivity using *t*-Bu₂SiCl₂ and Li⁰ to generate the silylene: Driver TG, Franz AK, Woerpel KA. J. Am. Chem. Soc 2002;124:6524–6525.6525 [PubMed: 12047159]
- (26). Sullivan SA, DePuy CH, Damrauer R. J. Am. Chem. Soc 1981;103:480–481.
- (27). *tert*-Butyldimethylsilyl ethers, whose basicities are comparable to that of analogous dialkyl ethers, should be capable of complexation: Blake JF, Jorgensen WL. J. Org. Chem 1991;56:6052–6059.6059
- (28). Belzner J, Dehnert U, Ihmels H, Hubner M, Muller P, Uson I. Chem. Eur. J 1998;4:852–863.
- (29). Both stereoisomers of silacyclopropane **21** encounter steric hindrance. The conformer drawn places the ring methyl group equatorial to avoid 1,3-diaxial interactions
- (30). Tamao K, Nakajima T, Kumada M. Organometallics 1984;3:1655–1660.
- (31). Robinson LR, Burns GT, Barton TJ. J. Am. Chem. Soc 1985;107:3935–3941.

Scheme 1.
Proposed Mechanism for the Formation of Allylic Silane **3**.

Figure 1.
Proposed Conformations of Ylide **20**.