Published in final edited form as:

Angew Chem Int Ed Engl. 2007 ; 46(34): 6492–6494. doi:10.1002/anie.200701386.

A New Strategy for the Stereoselective Synthesis of Isoxazolidines via Pd-Catalyzed Carboetherification of *N*-Butenyldihydroxylamines.^[**]

Michael B. Hay and John P. Wolfe^[*]

University of Michigan, Department of Chemistry, Ann Arbor, MI 48109-1055 (USA)

Keywords

Aryl Halide; Catalysis; Nitrogen Heterocycle; Olefin Insertion; Palladium

Isoxazolidines are frequently used as intermediates in the synthesis of complex molecules,[1] and are found in several interesting biologically active compounds.[2] In addition, the isoxazolidine N-O bond can be easily cleaved under reducing conditions to afford 1,3-amino alcohols, which are also of synthetic utility.[3] The most commonly employed method for the construction of isoxazolidines involves 1,3-dipolar cycloaddition reactions between nitrones and alkenes,[4] which generates the O1-C5 bond and the C3-C4 bond in one step [Eq. (1)]. Although these transformations are very useful, many intermolecular cycloadditions of unactivated alkenes generate mixtures of regioisomers.[4] Moreover, the major stereoisomers typically result from *endo*-addition on the less hindered face of the alkene, and the selective preparation of stereoisomers resulting from *exo*-addition and/or addition to the more substituted alkene face cannot be achieved in a straightforward manner.[4]

In this communication, we describe a new approach to the construction of substituted isoxazolidines that involves palladium-catalyzed carboetherification reactions of *N*-butenyl hydroxylamine derivatives with aryl bromides [Eq. (2)]. This method represents a new strategy for construction of the isoxazolidine ring, in which the O1-C5 bond and a C5'-Ar bond are formed in one step.[5] These transformations also provide access to isoxazolidine stereoisomers that cannot be generated with currently available methods. The reactions appear to proceed via intramolecular alkene insertion into previously unprecedented palladium alkoxyamine intermediates, which may be of utility in other Pd-catalyzed carbon-heteroatom bond-forming processes.

(1)

[**]This research was supported by the NIH-NIGMS (GM-071650). JPW thanks The Camille and Henry Dreyfus Foundation (New Faculty Award, Teacher Scholar Award) and Research Corporation for an Innovation Award. Additional unrestricted support was provided by Eli Lilly, 3M, and Amgen.

[*]Fax: (+)-1-734-763-2307, E-mail: jpwolfe@umich.edu

(2)

In preliminary experiments, we examined Pd-catalyzed reactions of *N*-butenyl hydroxylamines **1-3** with 4-bromobiphenyl using conditions that were employed in our prior studies on Pd-catalyzed carboetherification reactions of γ -hydroxyalkenes.[6-8] As shown below (Table 1), attempts to cyclize unprotected hydroxylamine substrate **1** or *N*-boc-protected derivative **2** were unsuccessful.[9] However, we were gratified to find that treatment of *N*-benzyl-protected substrate **3** with 4-bromobiphenyl and NaOtBu in the presence of catalytic amounts of Pd (OAc)₂ and DPE-Phos[10] afforded the desired product **4** in 80% isolated yield.

With viable reaction conditions and a suitable nitrogen protecting group identified, we examined Pd-catalyzed carboetherification reactions between several different substituted hydroxylamines and a number of aryl bromides. As shown in Table 2, this method is effective with electron-rich (entry 10), electron-neutral (entries 4, 7, 8, and 12), electron-poor (entries 1, 2, 11, and 13), *o*-substituted (entry 8), and heterocyclic (entries 3, 5, 6, and 9) aryl bromides. In addition to the *N*-benzyl protected derivatives described above, hydroxylamine substrates bearing *N*-methyl or *N*-*tert*-butyl groups also undergo cyclization in good yield.

The carboetherification reactions are also effective with substrates bearing substituents along the tether between the hydroxylamine moiety and the alkene. Transformations of these substrates provide access to disubstituted isoxazolidines with moderate to excellent stereocontrol. Importantly, in many cases these cyclizations provide a means to generate isoxazolidines that could not be prepared using 1,3-dipolar cycloaddition methods. For example, Pd-catalyzed reactions of **10** with 4-bromobiphenyl or 3-bromobenzotrifluoride provide **22** and **23** in 78% and 69% isolated yields, with >20:1 diastereoselectivity and regioselectivity (Table 2, entries 12-13). In contrast, a 1,3-dipolar cycloaddition reaction between a nitron and a 3-arylcyclopentene would be expected to occur on the less hindered face of the alkene to afford a different stereoisomer, and would likely generate mixtures of regioisomers.[4] In addition, reactions of **8** with aryl bromides proceed in 82-85% yield and 10:1 dr to afford the 2*R**,3*aS**-hexahydropyrrolo[1,2*b*]isoxazole isomers **18-19** (entries 8-9). However, dipolar cycloadditions between 3,4-dihydropyrrole-1-oxide and allylbenzene derivatives instead generate stereoisomeric 2*S**,3*aS**-hexahydropyrrolo[1,2*b*]isoxazoles.[11] The hydroxylamine carboetherifications can also be used to prepare *trans*-4,5-disubstituted isoxazolidines **12-14** and *cis*-3,5-disubstituted isoxazolidines **15-17** in good yield with 3-5:1 diastereoselectivity.

A plausible mechanism for the isoxazolidine-forming reactions is shown in Scheme 1. These transformations appear to be mechanistically related to Pd-catalyzed carboetherification reactions of γ -hydroxy alkenes with aryl bromides,[6] and are likely initiated by oxidative addition of the aryl bromide to Pd(0) to afford **24**. The L_nPd(Ar)(Br) complex can then be transformed to intermediate **25** via reaction with the hydroxylamine substrate and NaOtBu. Intramolecular *syn*-oxypalladation[6,12] of the tethered alkene moiety of **25** would generate **26**, which can undergo C-C bond-forming reductive elimination[13] to afford the observed isoxazolidine products. The conversion of **10** to *syn*-addition products **22** and **23** is consistent with this hypothesis. Moreover, this model also accounts for the observed stereochemistry of **12-21**, as the *syn*-oxypalladation likely occurs from an organized cyclic transition state in which nonbonding interactions are minimized by pseudoequatorial orientation of the substrate R¹ and

R² groups. This transition state arrangement would provide *cis*-3,5-disubstituted products (R² = H) and *trans*-4,5-disubstituted compounds (R¹ = H).

Although palladium(aryl)alkoxides have been shown to be important intermediates in a number of catalytic processes,[6,14-15] the analogous complexes derived from hydroxylamines (e.g. **25**) are unknown. The reactions described in this paper represent the first examples of catalytic transformations involving Pd(Ar)(ONRR') species. These previously unknown intermediates will likely find additional applications in other metal-catalyzed carbon-heteroatom bond-forming reactions.[15]

In conclusion, we have developed a new stereoselective method for the construction of substituted isoxazolidines via Pd-catalyzed carboetherification reactions of unsaturated hydroxylamine substrates. In many cases the stereochemical outcome of these transformations is complementary to nitrene cycloadditions, and this method provides a new strategic disconnection that can be used for retrosynthetic analysis of substituted isoxazolidines.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- [1]. For a review on the use of isoxazolidines as intermediates in complex molecule synthesis, see:Frederickson M. Tetrahedron 1997;53:403–425.425
- [2]. a) Ishiyama H, Tsuda M, Endo T, Kobayashi J. Molecules 2005;10:312–316. [PubMed: 18007301] b) Minter AR, Brennan BB, Mapp AK. J. Am. Chem. Soc 2004;126:10504–10505. [PubMed: 15327284] c) Palmer GC, Ordy MJ, Simmons RD, Strand JC, Radov LA, Mullen GB, Kinsolving CR, St. Georgiev V, Mitchell JT, Allen SD. Antimicrob. Agents. Chemother 1989;33:895–905. [PubMed: 2764540]
- [3]. a) Lait SM, Rankic DA, Keay BA. Chem. Rev 2007;107:767–796. [PubMed: 17319729] b) Revuelta J, Cicchi S, Brandi A. Tetrahedron Lett 2004;45:8375–8377. c) LeBel NA, Balasubramanian N. J. Am. Chem. Soc 1989;111:3363–3368. d) Iida H, Kasahara K, Kibayashi C. J. Am. Chem. Soc 1986;108:4647–4648.
- [4]. For reviews on 1,3-dipolar cycloaddition reactions, see:a)Confalone PN, Huie EM. Org. React 1988;36:1–173.173b)Gothelf KV, Jorgensen KA. Chem. Commun 2000:1449–1458.1458c) Kanemasa S. Synlett 2002:1371–1387.1387d)Koumbis AE, Gallos JK. Curr. Org. Chem 2003;7:585–628.628e)Merino P. Padwa A. *Science of Synthesis (Houben-Weyl Methods of Molecular Transformations)* 2004;27:511–580.580ThiemeStuttgartf)Pandey G, Banerjee P, Gadre SR. Chem. Rev 2006;106:4484–4517.4517 [PubMed: 17091927]g)Molteni G. Heterocycles 2006;68:2177–2202.2202
- [5]. A recent report described Pd(0)- and Pd(II)-catalyzed syntheses of 2-vinyl isoxazolidines via cyclizations of hydroxylamines bearing tethered allylic acetate groups. In contrast to the reactions described in this paper, the cyclizations of the allyl acetate derivatives do not lead to C-C bond formation concomitant with ring closure. The cyclizations of the allylic acetate derivatives are also mechanistically distinct, as the Pd(0)-catalyzed transformation proceeds via a π -allylpalladium intermediate which is likely captured via an outer-sphere nucleophilic attack on the allyl ligand, and the Pd(II)-catalyzed transformation proceeds via alkene activation followed by outer-sphere *anti*-oxypalladation. No evidence exists for the intermediacy of L_nPd(ONR₂)(allyl) complexes that bear a Pd-O bond. See:a)Merino P, Tejero T, Mannucci V, Prestat G, Madec D, Poli G. Synlett 2007:944–948.948For additional discussion of Pd-catalyzed allylations of nucleophiles containing an N-O bond, see:b)Miyabe H, Yoshida K, Reddy VK, Matsumura A, Takemoto Y. J. Org. Chem 2005;70:5630–5635.5635 [PubMed: 15989347]
- [6]. a) Wolfe JP, Rossi MA. J. Am. Chem. Soc 2004;126:1620–1621. [PubMed: 14871078] b) Hay MB, Hardin AR, Wolfe JP. J. Org. Chem 2005;70:3099–3107. [PubMed: 15822970] c) Hay MB, Wolfe

JP. J. Am. Chem. Soc 2005;127:16468–16476. [PubMed: 16305233] d) Hay MB, Wolfe JP. Tetrahedron Lett 2006;47:2793–2796.

- [7]. For a review on Pd-catalyzed carboetherification and carboamination reactions of γ -hydroxy- or γ -amino alkenes, see: Wolfe JP. Eur. J. Org. Chem 2007:571–582.582
- [8]. Two recent reports have described a related approach to the synthesis of *cis*-3,5-disubstituted isoxazolidines in which the N-C3 bond is generated via Pd-catalyzed carboamination of *O*-homoallyl hydroxylamines. The scope of these transformations has not been fully explored. See: a) Dongol KG, Tay BY. Tetrahedron Lett 2006;47:927–930.930b) Peng J, Lin W, Yuan S, Chen Y. J. Org. Chem 2007;72:3145–3148.3148 [PubMed: 17367194]
- [9]. The major products observed in these reactions resulted from Heck-arylation of the alkene.
- [10]. DPE-Phos = bis(diphenylphosphinophenyl)ether.
- [11]. Iida H, Watanabe Y, Kibayashi C. J. Chem. Soc., Perkin Trans. 1 1985:261–266.
- [12]. For other recent studies on *syn*-alkene insertions into late transition metal-oxygen bonds, see: a) Bryndza HE, Calabrese JC, Wreford SS. Organometallics 1984;3:1603–1604.1604b) Bennett MA, Jin H, Li S, Rendina LM, Willis AC. J. Am. Chem. Soc 1995;117:8335–8340.8340c) Hamed O, Thompson C, Henry PM. J. Org. Chem 1997;62:7082–7083.7083 [PubMed: 11671803] d) Hayashi T, Yamasaki K, Mimura M, Uozumi Y. J. Am. Chem. Soc 2004;126:3036–3037.3037 [PubMed: 15012118] e) Trend RM, Ramtohl YK, Stoltz BM. J. Am. Chem. Soc 2005;127:17778–17788.17788 [PubMed: 16351107] f) Zhao P, Incarvito CD, Hartwig JF. J. Am. Chem. Soc 2006;128:9642–9643.9643 [PubMed: 16866513]
- [13]. a) Milstein D, Stille JK. J. Am. Chem. Soc 1979;101:4981–4991. b) Hartwig JF. Inorg. Chem 2007;46:1936–1947. [PubMed: 17348724]
- [14]. For examples of well-characterized $L_nPd(Ar)(OR)$ complexes, see: a) Mann G, Hartwig JF. J. Am. Chem. Soc 1996;118:13109–13110.13110b) Widenhoefer RA, Zhong HA, Buchwald SL. J. Am. Chem. Soc 1997;119:6787–6795.6795
- [15]. In addition to carboetherification reactions of γ -hydroxyalkenes, synthetically useful reactions that are believed to proceed via $L_nPd(Ar)(OR)$ complexes include Pd-catalyzed *O*-arylation reactions and oxidation reactions. It is likely that the analogous hydroxylamine complexes [$L_nPd(Ar)(ONRR')$] will find similar applications. For lead references on useful catalytic transformations involving $L_nPd(Ar)(OR)$ intermediates, see: a) Vorogushin AV, Huang X, Buchwald SL. J. Am. Chem. Soc 2005;127:8146–8149.8149 [PubMed: 15926842] b) Kataoka N, Shelby Q, Stambuli JP, Hartwig JF. J. Org. Chem 2002;67:5553–5566.5566 [PubMed: 12153253] c) Guram AS, Bei X, Turner HW. Org. Lett 2003;5:2485–2487.2487 [PubMed: 12841761]

1. .

Table 1Carboetherification of *N*-butenyl hydroxylamines^[a]

Hydroxylamine	R	Isoxazolidine	Yield (%) ^[b]
1	H	-	0 ^[c]
2	Boc	-	0 ^[c]
3	Bn	4	80

^[a] Conditions: 1.0 equiv substrate, 1.2 equiv ArBr, 1.2 equiv NaOtBu, 2 mol % Pd(OAc)₂, 2 mol % DPE-Phos, THF (0.125 M), 65 °C.

^[b] Yields represent average isolated yields for two or more experiments.

^[c] Heck arylation products were observed.

Table 2Pd-catalyzed synthesis of substituted isoxazolidines^[a]

$ \begin{array}{c} \text{R}^1\text{-N(OH)-CH=CH-R}^2 + \text{Ar-Br} \xrightarrow[\text{NaOtBu, THF, 65 } ^\circ\text{C}]{\begin{array}{c} 2 \text{ mol } \% \text{ Pd(OAc)}_2 \\ 2 \text{ mol } \% \text{ DPE-Phos} \end{array}} \text{R}^1\text{-N(O)-CH(Ar)-CH-R}^2 \end{array} $				
Entry	Substrate	Isoxazolidine	dr ^[b]	Yield ^[b]
1			-	85%
2			>20:1 (5:1)	56%
3	5		3:1 (3:1)	70%
4	5		>20:1 (3:1)	57%
5			3:1 (3:1)	77%
6	6		3:1 (3:1)	78%
7			3:1 (3:1)	61%
8			>20:1 (10:1)	82%
9	8		>20:1 (10:1)	85%

Entry	Substrate	Isoxazolidine	dr ^[b]	Yield ^[b]
10			19:1 (9:1)	94%
11	9		>20:1 (9:1)	89%
12 ^[d]			>20:1 (>20:1)	78%
13 ^[d]	10		>20:1 (>20:1)	69%

^[a] Conditions: 1.0 equiv substrate, 1.2 equiv ArBr, 1.2 equiv NaOtBu, 2 mol % Pd(OAc)₂, 2 mol % DPE-Phos, THF (0.125 M), 65 °C.

^[b] dr = diastereomeric ratio of isolated material. Numbers in parentheses are the observed diastereomeric ratios for the crude reaction mixture.

^[c] Yields represent average isolated yields for two or more experiments.

^[d] The reaction was conducted at 110 °C in toluene solvent with Pd₂(dba)₃ (1 mol %) used in place of Pd(OAc)₂.