Published in final edited form as:

J Am Chem Soc. 2009 August 26; 131(33): 11707–11711. doi:10.1021/ja9054959.

Allylic C—H Amination for the Preparation of *syn*-1,3-Amino Alcohol Motifs

Grant T. Rice and M. Christina White*

Roger Adams Laboratory, Department of Chemistry, University of Illinois, Urbana, Illinois 61801

Abstract

A highly selective and general Pd/sulfoxide-catalyzed allylic C—H amination reaction en route to *syn*-1,3-amino alcohol motifs is reported. Key to achieving this reactivity under mild conditions is the use of electron-deficient *N*-nosyl carbamate nucleophiles that are thought to promote functionalization by furnishing higher concentrations of anionic species *in situ*. The reaction is shown to be orthogonal to classical C—C bond forming/reduction sequences as well as nitrene-based C—H amination methods.

Introduction

A diverse range of natural products and pharmaceuticals include *syn*-1,3-amino alcohols as dominant motifs. Important advances have been made in asymmetric C—C and C—N bond forming reactions to generate β -amino ketones and β -hydroxy imines.^{1,2} Complementary methods that furnish 1,3-amino alcohols with minimal oxidation state manipulations would provide strategic advantages in streamlining their syntheses. Although metal nitrene systems for aliphatic C—H aminations provide direct routes for accessing *syn*-1,3-amino alcohols,³ analogous allylic C—H aminations face challenges in chemoselectivity and/or reactivity, particularly with terminal olefins.^{3b,c,4} Such reactions would be highly valuable in synthetic planning due to the latent functionality preserved in the unsaturated moiety (*vide infra*). We have recently discovered Pd(II)/sulfoxide catalyst systems that effect predictably selective allylic C—H esterifications,⁵ alkylations,⁶ and aminations^{7,8}, and demonstrated their strategic use in streamlining complex molecule synthesis.⁹ Notably, our group recently reported an intramolecular allylic C—H amination reaction of homoallylic *N*-tosyl carbamates to generate 5-membered vinyl oxazolidinones *en route* to *syn*-1,2-amino alcohol motifs.⁷ We now report that the use of a more electron-deficient *N*-nosyl carbamate nucleophile enables a mild Pd(II)/sulfoxide catalyzed allylic C—H amination reaction with extraordinary chemoselectivities that furnishes vinyl *syn*-1,3-amino alcohol precursors from terminal olefins.

*R groups may contain a wide variety of functional groups including: carbonyls, internal olefins, benzylic C—H's and etheral C—H's

(1)

white@scs.uiuc.edu.

Supporting Information **Available:** Experimental procedures, full characterization, and additional experiments. This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

Results and Discussion

Design Principles

Mechanistic studies indicated that intramolecular allylic C—H amination to form 5-membered vinyl oxazolidinones proceeds *via* Pd(II)/sulfoxide-mediated heterolytic allylic C—H cleavage to generate a π -allyl Pd(OAc) intermediate **A** (Figure 1A)⁷. Intramolecular functionalization with the acidic *N*-tosyl carbamate pro-nucleophile **B** is effectively promoted by palladium carboxylate counterion acting as a base (Figure 1B). This catalytic source of weak base is regenerated during oxidation of Pd(0) with quinone (Figure 1C). However, attempts to promote allylic C—H amination to form a 6-membered oxazinanone with this nucleophile under identical conditions gave poor yields and conversions even after 72 hours (Table 1, entry 1, 75% recovered starting material).^{10,11} Given the higher kinetic barrier for 6- versus 5-membered ring formation, we reasoned that functionalization was likely the problematic step. Decreasing the electron density of the nitrogen is known to assist electrophilic metal catalysis by preventing nucleophile complexation to the metal.¹² We hypothesized that even in cases like ours where metal/nucleophile interactions are not likely to be problematic,^{7,8} switching to a more *electron-poor* amine might promote catalysis by increasing the equilibrium concentration of the active anionic species **C** *in situ* without prohibitively decreasing its nucleophilicity (Figure 1B).¹³

Reaction Optimization

Consistent with the hypothesis that increasing the pro-nucleophile's acidity will lead to an improvement in reactivity, examination of a series of 4-substituted *N*-arylsulfonyl carbamates revealed a positive correlation between pro-nucleophile acidity and product yields.¹⁴ By simply changing from an *N*-tosyl carbamate to an *N*-(4-chlorophenylsulfonyl) carbamate, a doubling in yield was achieved in one-third the reaction time [Table 1, entry 2, 15% (72 h) $\rightarrow$ 38% (24 h)]. Switching to the more acidic *N*-(4-nitrophenylsulfonyl) carbamate group afforded a dramatic positive impact on the reaction rate (72 h $\rightarrow$ 24 h) and yield (15% $\rightarrow$ 67%) of 6-membered ring formation (entry 3).¹⁵ Interestingly, evaluation of the even more acidic *N*-(2-nitrophenylsulfonyl) carbamate group afforded no further improvement, suggesting that decreasing the electron density of a nitrogen pro-nucleophile to increase anion concentration must be balanced with decreasing its nucleophilicity (entry 4). Additional reaction exploration showed that a switch in the solvent from THF to DCE and the inclusion of additives known to promote Pd(0) oxidation (*e.g.* O₂^{16a} and *p*-nitrobenzoic acid^{16b,c}) furnished the *syn*-oxazinanone in good yields and diastereoselectivities (entry 5).

Interestingly, reactivity and selectivity in this system are not strongly impacted by steric bulk adjacent to the carbamate (Table 1, entries 5-7). This is in stark contrast to the allylic C—H amination system furnishing *syn*-1,2-amino alcohol motifs, where one bulky substituent adjacent to the carbamate (generally a branching element) was deemed important for achieving good diastereoselectivity; a bulky quaternary alkyl substituents, albeit proceeding with excellent diastereoselectivities, suffered from poor reactivity (*vide infra*).⁷

Reaction Scope

Significantly, use of this more electron deficient nitrogen nucleophile proved to be a general solution for the formation of a wide range of 6-membered oxazinanones (Table 2). Substrates derived from 2° alcohols having prochiral allylic C—H bonds show good to excellent levels of diastereoselectivity favoring the *syn*-1,3-isomer (**5**, **7-12**). The observed stereochemical outcome is consistent with functionalization proceeding *via* a chair-like transition state.¹³ No conformational biasing element is required for effecting cyclization as evidenced by efficient generation of oxazinanone **4** (70%) derived from a 1° alcohol precursor.^{3c} Even a substrate

originating from a sterically hindered 3° alcohol generated *syn*-1,3-oxazinanone **6** in good yield.

This method is orthogonal to all other current state-of-the-art methods for generating this synthetically important motif. The complimentary nature of this method to C—C and C—N bond forming/reduction sequences is highlighted by the synthesis of oxazinanones **7** and **8** having reduction sensitive proximal ketone and ester functionalities. In contrast to nitrene-based systems, perfect chemoselectivity is seen for allylic C—H amination over benzylic and ethereal C—H amination (**9**, **10**; (+)-**15** and (+)-**16**, Scheme 1).³ Strikingly, this allylic C—H amination method shows unprecedented chemoselectivity for C—H amination of terminal over internal olefins (**11** and **12**). In all cases, *syn*-oxazinanone products of greater than 20:1 diastereomeric ratio can be obtained in good yields after standard column chromatography (see parenthetical yields in Table 2).

Predictable diastereomeric outcomes are crucial for C—H functionalization methods to find use at late stages of complex molecule synthesis. Gratifyingly, for substrates containing multiple stereogenic centers, the diastereomeric outcome of this reaction is controlled by the stereocenter containing the carbamate (Scheme 1). In substrates containing homoallylic (**17**, **19**) or trishomoallylic (**13**, **14**) stereogenic centers the reaction is consistently *syn*-selective relative to the *N*-nosyl carbamate.

Streamlining Synthesis

We have previously demonstrated the ability of predictably selective C—H amination reactions to streamline the synthesis of nitrogen containing molecules by “skipping” oxygenated intermediates that are burdensome to carry through synthetic sequences.^{7,8} The ability of this allylic C—H amination reaction to efficiently access optically enriched *syn*-1,3-amino alcohols is highlighted in the synthesis of (+)-allosedridine **25**, a member of the sedum alkaloids that have shown promising memory enhancing properties.¹⁷ Starting from commercially available enantioenriched bis-homoallylic alcohol (+)-**21**, the nitrogen is introduced at the correct oxidation state *via* C—H amination in only two steps. Major diastereomer (+)-**23** was easily isolated using standard flash column chromatography in 61% yield with >20:1 diastereomeric purity and with no erosion in enantiomeric excess (>99% *ee*). Notably, the *N*-nosyl group can be easily deprotected using mild PhSH/K₂CO₃ conditions to afford the free oxazinanone in 91% yield.¹⁸ Alkylation furnished (–)-**24** whose terminal olefin moieties could be used to forge the piperidine core *via* Grubbs ring-closing metathesis (RCM).¹⁹ Hydrogenation, followed by basic hydrolysis completed the total synthesis of (+)-allosedridine **25** in six steps and 27% overall yield. By avoiding multiple functional group manipulations and exploiting the terminal olefin functionality, an allylic C—H amination route affords the shortest and highest yielding synthesis of (+)-**25** to date.²⁰

syn-1,2-Amino Alcohol Synthesis

Significantly, the discovery that *N*-nosylcarbamate nucleophiles lead to improved reactivity for intramolecular allylic C—H aminations could be used to increase the efficiency of our previously reported reaction for generating 1,2-amino alcohol motifs. Homoallylic *N*-nosyl carbamates **26b** and **26d** underwent Pd(II)/sulfoxide **1** catalyzed intramolecular allylic C—H amination to furnish *anti*-oxazolidinones **27b** and **27d** in comparable yields, diastereoselectivities, and a three-fold decrease in reaction times (72 h → 24 h) to that reported with the analogous *N*-tosyl carbamate substrates. In contrast to the 1,3-amino alcohol system, however, reactivity with sterically congested substrates such as *t*-butyl **26f** and tertiary alcohol derived carbamate **26g** remained low. These results demonstrate a distinct steric limitation for this chemistry in furnishing 5-membered oxazolidinone rings that is not observed in generating 6-membered oxazinanone rings. Gratifyingly, the allylic C—H amination reaction for

generating 1,2-amino alcohol motifs also proceeds with outstanding chemoselectivity. This is illustrated in the preferential C—H amination of terminal over internal olefins in the doubly homoallylic *N*-nosyl carbamate substrate **26h**. Interestingly, *syn*-oxazolidinone **27h** is obtained in both good yields and diastereoselectivities despite the lack of an adjacent branching element, previously deemed to be crucial for obtaining synthetically useful diastereomeric outcomes with this system (Table 3, entry 5).

Conclusion

We report herein the discovery that an electron-deficient *N*-nosyl carbamate nucleophile furnishes a general Pd(II)-sulfoxide catalyzed allylic C—H amination reaction to generate 6-membered *syn*-oxazinanones under mild reaction conditions. The extraordinary chemoselectivity of this method is underscored by its demonstrated ability to selectively aminate allylic C—H bonds of terminal olefins preferentially to internal olefins. This feature, as well as its orthogonality to nitrene-based C—H aminations and C—C bond forming/reduction sequences, makes it a powerful reaction for 1,3-amino alcohol synthesis. This allylic C—H amination reaction is enabled by electronically tuning the pro-nucleophile in order to increase the equilibrium concentration of active anionic species under conditions that employ catalytic amounts of a weak base. The general strategy of decreasing the p*K*_a of a pro-nucleophile to promote functionalization has led to a significant rate increase for the previously reported allylic C—H amination reaction for 1,2-amino alcohol synthesis and has the potential for applications to other electrophilic metal-catalyzed reactions.

Experimental Procedures

General Experimental Procedures

In a one dram vial was added the corresponding carbamate starting material (1 equiv.), phenyl benzoquinone (2 equiv.), *p*-nitrobenzoic acid (0.10 equiv.), 1,2-bis(phenylsulfinyl)ethane (0.05 equiv), Pd(OAc)₂/1,2-bis(phenylsulfinyl)ethane catalyst **1** (0.10 equiv., Aldrich Chemical Company) and a Teflon stir bar. In a separate flask, O₂ gas was simultaneously bubbled through 1,2-dichloroethane for thirty minutes. The oxygenated 1,2-dichloroethane (0.66 M) was then added to the previous one dram vial, O₂ gas was blown over the vial for 5 seconds, and the vial was sealed with a Teflon lined cap. The reaction vial was then vortexed until the solution appeared homogeneous and stirred in a 45° C oil bath for 24 hours. The solution was allowed to cool to room temperature and then transferred using a minimum amount of dichloromethane to a 250 mL separatory funnel. The solution was diluted with 15 mL of diethyl ether and rinsed 1× 15 mL aqueous sodium bisulfite (sat.), 1× 15 mL water, 1× 15 mL 5% aqueous K₂CO₃/H₂O and 1× 15 mL water. The organic layer was collected and dried over MgSO₄, filtered and concentrated *in vacuo*. The crude reaction mixture was purified using flash column chromatography. In general, the major *syn*-diastereomer can be isolated by flash chromatography (25-35% gradient EtOAc/hexanes) directly from the crude reaction mixture.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by NIH/NIGMS (grant no. GM076153) and kind gifts were received by Eli Lilly, Bristol-Myers Squibb, Pfizer, Amgen, Abbott, Boehringer Ingelheim, Roche, the Sloan Foundation, AstraZeneca, and the Camille Dreyfus Teacher-Scholar Award. We thank Strem Chemicals and the Aldrich Chemical Company for a gift of commercial catalyst **1**, P. Gormisky for confirming **2c**, and A. Hamlin for preliminary studies towards the synthesis of (+)-**25**.

References

1. (a) List B. *J Am Chem Soc* 2000;122:9336. (b) Wenzel AG, Jacobsen EN. *J Am Chem Soc* 2002;124:12964. [PubMed: 12405820] (c) Kochi T, Tang TP, Ellman JA. *J Am Chem Soc* 2003;125:11276. [PubMed: 16220949] (d) Matsunaga S, Yoshida T, Morimoto H, Kumagai N, Shibasaki M. *J Am Chem Soc* 2004;126:8777. [PubMed: 15250731] (e) Josephsohn NS, Snapper ML, Hoveyda AH. *J Am Chem Soc* 2004;126:3734. [PubMed: 15038725] (f) Enders D, Grondal C, Vretou M, Raabe G. *Angew Chem Int Ed* 2005;44:4079.
2. For some recent examples of asymmetric conjugate addition reactions with nitrogen nucleophiles to furnish β -amino ketones see: (a) Myers JK, Jacobsen EN. *J Am Chem Soc* 1999;121:8959. (b) Guerin DJ, Miller SJ. *J Am Chem Soc* 2002;124:2134. [PubMed: 11878965] (c) Sibi MP, Prabakaran N, Ghorpade SG, Jasperse CP. *J Am Chem Soc* 2003;125:11796. [PubMed: 14505383] (d) Yamigawa N, Qin H, Matsunaga S, Shibasaki M. *J Am Chem Soc* 2005;127:13419. [PubMed: 16173776] (e) Chen YK, Yoshida M, MacMillan DWC. *J Am Chem Soc* 2006;128:9328. [PubMed: 16848457]
3. (a) Espino CG, Wehn PM, Chow J, Du Bois J. *J Am Chem Soc* 2001;123:6935. (b) Fiori KW, Fleming JJ, Du Bois J. *Angew Chem Int Ed* 2004;43:4349. A sulfamate product analogous to 4 was generated *via* a chemoselective Rh-nitrene based reaction in 43% yield: (c) Zalatan DN, Du Bois J. *J Am Chem Soc* 2008;130:9220. [PubMed: 18582043] (d) Milczek E, Boudet N, Blakey S. *Angew Chem Int Ed* 2008;47:6825. (e) Liang JL, Yuan SX, Huang JS, Yu WY, Che CM. *Angew Chem Int Ed* 2002;41:3465.
4. Interestingly, high chemoselectivities and reactivities have been reported for intermolecular Rh-nitrene allylic aminations of internal olefins: Liang C, Collet F, Robert-Peillard F, Müller P, Dodd RH, Dauban P. *J Am Chem Soc* 2008;130:343. [PubMed: 18072775]
5. Linear allylic C—H acetoxylation: (a) Chen MS, White MC. *J Am Chem Soc* 2004;126:1346. [PubMed: 14759185] Branched allylic C—H esterification: (b) Chen MS, Prabakaran N, Labenz NA, White MC. *J Am Chem Soc* 2005;127:6970. [PubMed: 15884938] Oxidative C—H macrolactonization: (c) Fraunhofer KJ, Prabakaran N, Sirois LE, White MC. *J Am Chem Soc* 2006;128:9032. [PubMed: 16834366] For a linear allylic acetoxylation reaction using DMA solvent see: (d) Mistudome T, Umetani T, Nosaka N, Mori K, Mizugaki T, Ebitani K, Kaneda K. *Angew Chem Int Ed* 2006;45:481.
6. Linear allylic C—H alkylation: (a) Young AJ, White MC. *J Am Chem Soc* 2008;130:14090. [PubMed: 18831588] For a system that also uses Pd(II)/bis-sulfoxide catalysis see: (b) Lin S, Song CX, Cai GX, Wang WH, Shi ZJ. *J Am Chem Soc* 2008;130:12901. [PubMed: 18778061]
7. Intramolecular allylic C—H amination to furnish 1,2-amino alcohol motifs: Fraunhofer KJ, White MC. *J Am Chem Soc* 2007;129:7274. [PubMed: 17516648]
8. Intermolecular allylic C—H amination to furnish linear (*E*)-allylic amines using *N*-(methoxycarbonyl)-*p*-toluenesulfonamide nucleophile: (a) Reed SA, White MC. *J Am Chem Soc* 2008;130:3316. [PubMed: 18302379] (b) Reed SA, Mazzotti AR, White MC. *J Am Chem Soc*. 2009 manuscript in revision. For a similar system using DMA solvent see: (c) Liu G, Yin G, Wu L. *Angew Chem Int Ed* 2008;47:4733.
9. (a) Fraunhofer KF, Bachovchin DA, White MC. *Org Lett* 2005;7:223. [PubMed: 15646963] (b) Covell DJ, Vermeulen NA, Labenz NA, White MC. *Angew Chem Int Ed* 2006;45:8217.
10. Carbamate and *N*-tosyl carbamate substrates have been observed to exclusively form 5-membered ring oxazolidinone products in nitrene based systems: (a) Espino CG, Du Bois J. *Angew Chem Int Ed* 2001;40:598. (b) Lebel H, Huard K, Lectard S. *J Am Chem Soc* 2005;127:14198. [PubMed: 16218610]
11. The inclusion of catalytic amine base, recently discovered to promote intermolecular allylic C—H aminations with *N*-tosyl carbamate nucleophiles (ref. 8b), gave a minor increase in yield for substrate **2a** (15% $\rightarrow$ 37%), however, the diastereoselectivity was significantly diminished (5.1:1 $\rightarrow$ 1.8:1 *syn:anti*).
12. This effect has explicitly been demonstrated in the oxidative amination of olefins *via* electrophilic Pd(II) catalysis: (a) Hegedus LS, McKearin JM. *J Am Chem Soc* 1982;104:2444. (b) Larock RC, Hightower TR, Hasvold LA, Peterson KP. *J Org Chem* 1996;61:3584. [PubMed: 11667199]
13. Pd(0)-mediated allylic substitution of cyclic carbonates to carbamates proceeds *via* anionic nitrogen nucleophiles: Bando T, Harayama H, Fukazawa Y, Shiro M, Fugami K, Tanaka S, Tamaru Y. *J Org Chem* 1994;59:1465.

14. Relative pK_a values were based upon benzoic acid analogues (a)Tao L, Han J, Tao FM. J Phys Chem A 2008;112:775. [PubMed: 18179190](b)Maran F, Celadon D, Severin MG, Vianello E. J Am Chem Soc 1991;113:9320.
15. Heating the reaction with *N*-tosyl carbamate substrates to 65°C also promoted reactivity; however, the yields were lower and inconsistent results were observed.
16. (a)Stahl SS. Angew Chem Int Ed 2004;43:3400. Carboxylic acids have been shown to increase reactivity in oxidative Heck reactions mediated by Pd(II)/bis-sulfoxide, possibly *via* acid-promoted quinone reoxidation of the metal: (b)Delcamp JH, White MC. J Am Chem Soc 2006;128:15076. [PubMed: 17117844](c)Delcamp JH, Brucks AP, White MC. J Am Chem Soc 2008;130:11270. [PubMed: 18671350]
17. Meth-Cohn O, Yu CY, Lestage P, Lebrun MC, Cagniard DH, Renard P. Eur Pat. 20001050531
18. Fukuyama T, Jow C, Cheung M. Tetrahedron Letters 1995;36:6373.
19. (a)Trnka TM, Grubbs RH. Acc Chem Res 2001;34:18. [PubMed: 11170353]For specific reaction conditions see: (b)Anada M, Tanaka M, Washio T, Yamawaki M, Abe T, Hashimoto S. Org Lett 2007;9:4559. [PubMed: 17894504]
20. For previous asymmetric syntheses see: (a) Takahata H, Kubota M, Ikota N. J Org Chem 1999;64:8594.(b)Passarella D, Barilli A, Belinghieri F, Fassi P, Riva S, Sacchetti A, Silvani A, Danieli B. Tetrahedron: Asymmetry 2005;16:2225.

A. Electrophilic C–H Cleavage **B. Nucleophilic Functionalization** **C. Catalyst and Endogenous Base Regeneration**

Figure 1.
Design Principles for Intramolecular C–H Amination

Scheme 1.

Origin of Diastereoselectivity

^a**1** (10 mol%), 1,2-bis(phenylsulfinyl)ethane (5 mol%), *p*-nitrobenzoic acid (10 mol%), phenyl benzoquinone (2 equiv.), oxygenated DCE (0.66 M), 45°C. 24 h. ^bIsolated yield of major *syn*- product. ^cdr of crude reaction.

• Previous Syntheses: 10 Steps (15.8%, >98% ee)^{20a} & 8 Steps (2.7%, 85% ee)^{20b}

Scheme 2.

Total Synthesis of (+)-allosedridine

(a) NsNCO (1.1 equiv.), THF, r.t. (93%); (b) standard conditions; (c) Isolated yield of major *syn*- product; (d) PhSH (1.2 equiv.), K₂CO₃ (3 equiv.), DMF, 0°C, 4 h (91%); (e) *n*-BuLi (1.0 equiv.), THF, -78°C; but-3-enyl trifluoromethanesulfonate (1.2 equiv.), -78°C to 0°C, 2 h (89%); (f) Grubbs (II) (8 mol%), toluene, 65°C, 3 h; H₂, Pd-C, MeOH, r.t., 2 h (80%); (g) KOH/EtOH (1.7 M), 45°C, 2 h (73%).

Table 1

Allylic C-H Amination Reaction Optimization

Entry	R	Time	Isolated Yield ^b	dr ^c
1	<i>p</i> -Tol	72 h	15%	5.1:1
2	<i>p</i> -ClPh	24 h	38%	3.7:1
3	<i>p</i> -NO ₂ Ph (Ns)	24 h	67%	4.4:1
4	<i>o</i> -NO ₂ Ph	24 h	63%	2.6:1

Entry	R'	Isolated Yield ^b	dr ^c	Isolated Syn ^e
5 ^d	<i>i</i> Propyl	80%	6.0:1	65%
6 ^d	Ethyl	87%	4.3:1	67%
7 ^d	<i>t</i> Butyl	84%	6.3:1	68%

^a BisSO Ligand = 1,2-bis(phenylsulfinyl)ethane.

^b Average of two runs.

^c Determined by GC analysis ($R' = p$ -Tol) or ^1H NMR ($R' = p$ -ClPh, p -NO₂Ph, o -NO₂Ph) of crude reaction mixture.

^d Reaction run using 10 mol% p -nitrobenzoic acid and oxygenated DCE.

^e Isolated yield of major *syn*- product, >20:1 *syn:anti*.

Table 2

Allylic C-H Amination Reaction Scope

 $1^\circ, 2^\circ, 3^\circ$

70% Yield

Carbonyls

76% Yield, 3.4:1 dr
(53%, >20:1 dr)^c

Bonzylic C-H

80% Yield, 4.4:1 dr
(63%, >20:1 dr)^C

Internal Olefins

67% Yield, 4.5:1 dr
(52%, >20:1 dr)^c

^a Additives: *p*-nitrobenzoic acid (10 mol%), BisSO Ligand (5 mol%).

^b Reaction run in oxygenated DCE.

^c Isolated yield of major *syn*-product, >20:1 *syn:anti*.

Table 3

1,2-Amination Rate Increase Using *N*-Nosyl Carbamates

Product	R'	Time	Isolated Yield ^b	dr ^c	
	Tosyl (27a)	72 h	76%	6.0:1	
	Nosyl (27b)	24 h	78%	5.0:1	
	Tosyl (27c)	72 h	86%	1.6:1	
	Nosyl (27d)	24 h	79%	1.7:1	
	Tosyl (27e)	72 h	8%	18:1	
	Nosyl (27f)	72 h	20%	>20:1	
	Nosyl (27g)	72h	<1%	--	

J Am Chem Soc. Author manuscript; available in PMC 2010 August 26.

Product	R'	Time	Isolated Yield ^b	dr ^c
	Nosyl	(27h)	68%	5.4:1

^b Average of two runs.

^c Determined by GC analysis (R' = *p*-Tol) or ¹H NMR analysis (R' = *p*-NO₂Ph) of crude reaction mixture.