Before the catheter is tunnelled, a small incision is made in the distal catheter at its junction with the introducer. A syringe containing the local anaesthetic is attached to the other end of the catheter. This is continuously flushed by the assistant during the whole period when the introducer is being withdrawn from the other end of the tunnel. This ensures that the whole track is effectively infiltrated with local anaesthetic. Once the catheter is in place, its distal position can be trimmed according to the size of the child and the distal position of the catheter tip. Thus, the incised segment is removed. Finally, before insertion, the catheter lumen is flushed with saline.

DISCUSSION

Inadvertent catheter damage and inadequate infiltration are thus avoided.

Hanging manoeuvre to facilitate right hepatectomy with control of the right hepatic vein

SM ROBINSON, A AL-MUKHTAR, SAW WHITE
Department of HPB and Transplant Surgery, Freeman Hospital, Heaton Road, Newcastle upon Tyne, UK

CORRESPONDENCE TO

SM Robinson, Department of HPB and Transplant Surgery, Freeman Hospital, Heaton Road, Newcastle upon Tyne, UK
E: sturobinson@doctors.org.uk

BACKGROUND

The hanging manoeuvre facilitates right hepatectomy without prior liver mobilisation.¹ This is advantageous when tumour invades the diaphragm.² The disadvantage is that control of retro-hepatic venous structures is not prior to parenchymal division. Our modification overcomes this by controlling the right hepatic vein prior to parenchymal division.

TECHNIQUE

The anterior surface of the liver is exposed up to the level of the suprahepatic inferior vena cava and the space between the middle and right hepatic veins dissected. Attention is then turned to the infrahepatic inferior vena cava and a clamp is passed, in a cautious manner, posterior to the caudate lobe cranially towards the space created between the middle and right hepatic veins, keeping to the left of the inferior right hepatic vein. A tape is then grasped with the clamp and passed around the hepatic parenchyma.^{1,2} This process is then repeated placing a vessel loop alongside the tape. A space is dissected immediately to the right of the right hepatic vein directed towards this tunnel. The distal end of the vessel loop is then drawn alongside the lateral border of the right hepatic vein so that it is encircled. Following parenchymal transection, the right hepatic vein is divided in the usual manner (Fig. 1).

Figure 1 Illustration of the technique.

DISCUSSION

This modification of the hanging manoeuvre enables control of the right hepatic vein prior to parenchymal transection, avoiding potentially life threatening haemorrhage from inadequate control of the right hepatic vein.

References

1. Belghiti J, Guevara OA, Noun R, Saldinger PF, Kianmanesh R. Liver hanging maneuver: a safe approach to right hepatectomy without liver mobilization. *J Am Coll Surg* 2001; **193**: 109–11.
2. Ogata S, Belghiti J, Varma D, Sommacale D, Maeda A, Dondero F *et al.* Two hundred liver hanging maneuvers for major hepatectomy: a single-center experience. *Ann Surg* 2007; **245**: 31–5.

Nipple aspiration device

GEORGE ALEX
Department of Breast Surgery, Castle Hill Hospital, Cottingham, UK

CORRESPONDENCE TO

George Alex, Department of Breast Surgery, Castle Hill Hospital, Cottingham HU16 5JQ, UK
E: drgeorgealex@doctors.org.uk

BACKGROUND

Nipple discharge, especially if blood-stained, can be a sign of malignancy. About 5% of patients presenting to a symptomatic breast clinic have nipple discharge. It is commonly assessed by squeezing at the region of the nipple areolar complex to elicit any discharge.

TECHNIQUE

The nipple aspiration device is a device for aspirating breast ductal