Published in final edited form as:

Cancer Prev Res (Phila). 2008 November ; 1(6): 476–484. doi:10.1158/1940-6207.CAPR-08-0084.

Gemini Vitamin D Analogs Inhibit Estrogen Receptor Positive and Estrogen Receptor Negative Mammary Tumorigenesis without Hypercalcemic Toxicity

Hong Jin Lee¹, Shiby Paul¹, Nadi Atalla¹, Paul E. Thomas^{1,2,3}, Jennie Lin^{1,3}, Ill Yang³, Brian Buckley³, Gang Lu¹, Xi Zheng¹, You-Rong Lou¹, Allan H. Conney^{1,2}, Hubert Maehr⁴, Milan Uskokovic⁴, and Nanjoo Suh^{1,2}

¹Department of Chemical Biology, Ernest Mario School of Pharmacy, Rutgers, The State University of New Jersey, Piscataway, New Jersey

²The Cancer Institute of New Jersey, New Brunswick, New Jersey

³Environmental and Occupational Health Sciences Institute, Piscataway, New Jersey

⁴BioXell Inc., Nutley, New Jersey

Abstract

Numerous preclinical, epidemiological and clinical studies have suggested the benefits of vitamin D and its analogs for the prevention and treatment of cancer. However, the hypercalcemic effects have limited the use of $1\alpha,25(\text{OH})_2\text{D}_3$, the hormonally active form of vitamin D. To identify vitamin D analogs with better efficacy and low toxicity, we have tested more than 60 novel Gemini vitamin D analogs with a unique structure of two side chains for growth inhibition of breast cancer cells. Our initial studies found that some Gemini analogs are many-fold more active than $1\alpha,25(\text{OH})_2\text{D}_3$ in growth inhibition assay. In vivo experiments were designed to study the inhibitory effect of selected Gemini vitamin D analogs against mammary carcinogenesis by using (a) an N-methyl-N-nitrosourea-induced estrogen receptor (ER)-positive mammary tumor model and (b) an MCF10DCIS.com xenograft model of ER-negative mammary tumors. Among vitamin D analogs we tested, Gemini 0072 [$1\alpha,25$ -dihydroxy-20S-21(3-trideuteromethyl-3-hydroxy-4,4,4-trideuterobutyl)-23-yne-26,27-hexafluoro-19-nor-cholecalciferol] and Gemini 0097 [$1\alpha,25$ -dihydroxy-20R-21(3-trideuteromethyl-3-hydroxy-4,4,4-trideuterobutyl)-23-yne-26,27-hexafluoro-19-nor-cholecalciferol] administration inhibited by 60% the NMU-induced mammary tumor burden compared to the NMU-treated control group, but these compounds were devoid of hypercalcemia toxicity. In an ER-negative xenograft model, Gemini 0097 significantly suppressed tumor growth without hypercalcemia toxicity. We found that the inhibitory effect of Gemini 0097 was associated with an increased level of cyclin dependent kinase inhibitor p21 and the insulin-like growth factor binding protein 3 in both ER-positive and ER-negative mammary tumors. Our results suggest that Gemini vitamin D analogs may be potent agents for the prevention and treatment of both ER-positive and ER-negative breast cancer without hypercalcemia toxicity.

Keywords

Gemini Vitamin D; estrogen receptor; mammary tumorigenesis; IGFBP-3; Smad

Requests for reprints: Nanjoo Suh, Department of Chemical Biology, Ernest Mario School of Pharmacy, 164 Frelinghuysen Road, Rutgers, The State University of New Jersey, Piscataway, NJ 08854. Phone: 732-445-3400 (X226), Fax: 732-445-0687; e-mail: nsuh@rci.rutgers.edu.

INTRODUCTION

In the United States, breast cancer remains the most frequently diagnosed cancer and the second leading cause of cancer death in women according to Cancer Statistics from the American Cancer Society. Because of the complexity and heterogeneity of mammary carcinogenesis (1), many pharmacological agents have been studied for their effects on the prevention of breast cancer. For example, selective estrogen receptor modulators (SERMs) such as tamoxifen and raloxifene have achieved significant reduction of breast cancer incidence in women at high risk (2,3). However, SERMs are not effective in preventing estrogen receptor (ER)-negative breast cancer, which corresponds to at least one third of the breast cancer cases (4). The vitamin D receptor (VDR), a member of the nuclear receptor superfamily, has been suggested as a target for both ER-positive and ER-negative breast cancer prevention (5,6) because it is present in most breast tumors (4), and *VDR* ablation in mice was reported to enhance carcinogen-induced formation of mammary tumors (7). These results suggest a role of vitamin D signaling in the regulation of mammary tumorigenesis.

The ligand for VDR, 1 α ,25-dihydroxyvitamin D₃ (1 α ,25(OH)₂D₃; the key hormone in calcium/phosphate homeostasis), is a hormonally active metabolite synthesized from vitamin D₃ predominantly through hydroxylation by a 25-hydroxylase in the liver and a 1 α -hydroxylase in the kidney (6,8). It has been reported that several nonrenal tissues including breast, colon and prostate also express 1 α -hydroxylase, which can produce 1 α ,25(OH)₂D₃ from 25-hydroxyvitamin D₃. The formation of 1 α ,25(OH)₂D₃ in these tissues may result in inhibition of cell proliferation by (a) arrest in the G1 phase of the cell cycle, (b) induction of apoptosis, and (c) promotion of cell differentiation (9-12). However, hypercalcemic toxicity has limited the use of 1 α ,25(OH)₂D₃ for cancer prevention, and numerous synthetic vitamin D analogs have been developed for better efficacy and less calcemic effects.

Classic synthetic vitamin D analogs have been tested in different animal models to determine their *in vivo* efficacy. For example, synthetic 1 α -hydroxyvitamin D₅ reduced not only the formation of azoxymethane (AOM)-induced aberrant crypt foci (ACF) in CF-1 mice (13), but also the incidence and multiplicity of N-methyl-N-nitrosourea (NMU)-induced mammary tumors in Sprague-Dawley rats (14,15). Treatment with the vitamin D analog Ro26-9114 also caused a significant decrease in total tumor load over the entire gastrointestinal tract in APC^{min} mice (16). In a xenograft model, treatment with 1,25-dihydroxy-16-ene-23-yne vitamin D₃ attenuated the growth of retinoblastoma tumors formed by the subcutaneous injection of Y79 cells, and apoptosis was increased (17). Furthermore, administration of a combination of the vitamin D analog EB1089 with fractionated doses of radiation was significantly more effective for tumor inhibition and apoptosis promotion than either treatment alone for MCF-7 tumor xenografts in athymic mice (18).

To identify vitamin D analogs with better anticancer efficacy and lower toxicity than 1 α ,25(OH)₂D₃, we tested novel Gemini vitamin D analogs, which have a unique structure of two six-carbon chains with a C-20-normal and a C-20-epi side chain (see Fig. 1). They have considerably less toxicity than 1 α ,25(OH)₂D₃ in animals (19,20), and certain Gemini vitamin D analogs had a greater inhibitory effect on cell proliferation than 1 α ,25(OH)₂D₃ in MCF10 breast epithelial cells (21). In animal studies of mouse colorectal cancer cells (MC-26) implanted in BALB/c mice, administration of BXL-01-0072 (see Fig. 1 for the structure) significantly reduced tumor growth at a 10-fold lower dose than did 1 α ,25(OH)₂D₃ (19). This same deuterated Gemini analog also greatly diminished the invasive spread of the tumor into the surrounding muscle compared to 1 α ,25(OH)₂D₃ (20). We also reported that certain Gemini analogs decreased mammary tumor burden in the NMU-induced rat mammary carcinogenesis model (22).

Recently, breast cancer has been suggested to be classified into at least 5 different subtypes, such as luminal A, luminal B, basal A, basal B, and Her-2/neu (23,24). Because of the heterogeneity of breast cancer, we selected two different animal models of breast cancer, a conventional chemoprevention model representing estrogen-receptor (ER) positive luminal cell type and a basal type ER-independent model of breast cancer (24). In the present study, we tested a classic vitamin D analog, Ro-26-2198, and several Gemini vitamin D analogs (Fig. 1) for their effects on NMU-induced ER-positive mammary carcinogenesis and on the *in vitro* growth of cultured ER-negative MCF10DCIS.com human breast epithelial cells. An ER-negative xenograft model with implanted MCF10DCIS.com cells was also employed to investigate the effects of Gemini 0097 on the growth of ER-negative mammary tumors *in vivo*. Molecular mechanisms of action of Gemini vitamin D analogs that inhibited the growth of both ER-positive and ER-negative tumors were studied.

MATERIALS AND METHODS

Reagents

1 α ,25(OH) $_2$ D $_3$ and all Gemini vitamin D analogs (>95% purity) (Fig. 1) were provided by BioXell, Inc. (Nutley, NJ). 1 α ,25(OH) $_2$ D $_3$ and Gemini analogs were dissolved in DMSO. 1 α ,25(OH) $_2$ D $_3$ and Gemini analogs were diluted in 5% horse serum/DMEM/F12 media for [3 H]-thymidine uptake assay. For *in vivo* animal experiments, compounds were diluted in cremophore/PBS (1:8, v/v).

Cell culture

The human MCF10 breast epithelial cell line, MCF10DCIS.com, was provided by Dr. Fred Miller at the Barbara Ann Karmanos Cancer Institute (Detroit, MI). MCF10DCIS.com cells were maintained in DMEM/F12 medium supplemented with 5% horse serum, 1% penicillin/streptomycin, and 1% HEPES solution at 37°C, 5% CO $_2$. The cells were passed every 3–4 days.

NMU-induced mammary tumorigenesis in rats

Female Sprague-Dawley rats (21 $\pm$ 1 days old) were purchased from Taconic Farms (Germantown, NY). Rats were treated with a single intraperitoneal injection (50 mg/kg body weight) of the carcinogen N-methyl-N-nitrosourea (NMU, Sigma, St. Louis, MO). One week after the NMU injection, rats were treated with vehicle or with 1 α ,25(OH) $_2$ D $_3$ or Gemini vitamin D analogs in vehicle via intraperitoneal injection once a day, 5 days a week. The rat weight and tumor size of each animal were measured weekly. Nine weeks after starting vitamin D treatment, the rats were sacrificed and tumors were weighed at autopsy. Tumors and blood samples were collected for further analysis. Serum was collected after centrifugation of clotted blood samples. All animal studies were performed in accordance with an institutionally approved protocol.

Determination of serum calcium level

The concentration of calcium in serum samples was determined using the calcium reagent set (POINTE Scientific, INC. Canton, MI), and we followed the procedure provided by the manufacturer. Briefly, serum (4 μ l) was mixed with the appropriately diluted reagent (200 μ l) in a 96 well plate. After incubating for 1 min, the plate was read at 550 nm using a spectrophotometer. To adjust for differences in hemolysis among samples, serum blanks (serum in water) were prepared and the absorbance reading was subtracted from the test reading. The calcium concentration was calculated from calcium standards provided by the manufacturer.

Animal experiment in xenograft model

MCF10DCIS.com cells were trypsinized and prepared in Hanks' Balanced Salt Solution (Invitrogen, Carlsbad, CA). After the midline incision around the second teat in severe combined immunodeficiency (SCID) mice, 1×10^6 cells were injected in the mammary fat pad. The incision was closed by wound clips and the clips were removed after 4 days. Vehicle control (0.1 ml) or Gemini vitamin D analog 0097 (0.1 $\mu\text{g/kg}$ body weight in 0.1 ml vehicle) was injected intraperitoneally daily from Day 4 until the termination of the experiment. Body weight and tumor size were measured every week. Six weeks after the injection of MCF10DCIS.com cells, tumors were weighed and collected at autopsy for further analysis. All animal studies were performed in accordance with an institutionally approved protocol.

Western blot analysis

Tumor samples were homogenized in RIPA buffer (10 mM Tris-HCl, 5 mM EDTA, 150 mM NaCl, 1% Triton X-100, 1% sodium deoxycholate, 0.1% sodium dodecyl sulfate, 0.1mM Na_3VO_4 , 1% PMSF, 1% aprotinin and 0.1% leupeptin) using a Dounce homogenizer (Wheaton, Millville, NJ), and the protein extracts were electrophoresed in 4–15% gradient gels (Biorad, Hercules, CA) and transferred to a PVDF membrane (PALL, Ann Arbor, MI). The primary antibodies against VDR (Affinity BioReagents, Golden, CO), PARP, cleaved PARP, cleaved caspase-3, pSmad1/5 (Cell Signaling Technology Inc., Beverly, MA), p21, p27, apolipoprotein A-1 (Santa Cruz Biotechnology, Santa Cruz, CA), IGFBP-3 (R&D Systems, Minneapolis, MN), actin (Sigma, St. Louis, MO), and secondary antibodies (Santa Cruz Biotechnology) were used. The band intensity was quantified using TableLab software (ver. 1.10) and normalized with actin.

Sample preparation and 2-D gel electrophoresis

Rat mammary tumor tissue (1–2 gm) was grossly freed of necrotic areas and homogenized in 2 volumes of 50 mM Tris-HCl pH 7.4 at 4°C, 1.15% KCl, 1mM EDTA and 0.2 mM PMSF using a glass-teflon Potter–Elvehjem tissue homogenizer. Cytosolic fractions were isolated as described previously (25) and prepared for 2-D gels by precipitating proteins for 1 hr on ice with 2 volumes of ice cold methanol: acetonitrile (50:50). Samples (500 μg) were applied by cup loading to buffer hydrated (overnight) 24 cm Immobiline IPG strips pH 4–7 (Amersham Biosciences, Piscataway, NJ). Isoelectric focusing was performed at 25°C on an Ettan IPGphor II and the remainder of the 2-D gel electrophoresis was as described previously (26). Protein stained gels were scanned for data analysis using a Molecular Dynamics laser densitometer. Spot patterns were analyzed with Non-Linear Phoretix 2-D software. Selected protein stained spots were excised and in-gel-digested with trypsin and peptides extracted as described previously (27). Tryptic peptides were resolved on a microcapillary column (50 $\times$ 0.3 mm ID, Magic C18) and introduced into a ThermoFinnigan LCQ DECA XP ion trap mass spectrometer through an Electrospray Ionization (ESI) interface operated in positive ion mode by a ThermoFinnigan Surveyor LC system. Data dependent MS/MS spectra were generated and used for a protein identification database search using the ThermoFinnigan TurboSEQUENT program (Bioworks 3.1) after automatic removal of prominent MS/MS peaks due to digestion enzymes and other experimental background compounds.

Quantitative PCR analysis

These procedures have been previously reported (21). Labeled primers, including GAPDH, CYP24A1, and IGFBP-3, were obtained from Applied Biosystems (Foster City, CA).

Immunohistochemistry

Tumor samples from each group were harvested at autopsy and fixed in 10% formalin for 24 hr. They were sectioned, paraffin embedded, and microtomed into 4 μm thick tissue sections.

The slides were incubated overnight at room temperature with antibody to proliferating cell nuclear antigen (PCNA) (1:1000 diluted, BD Pharmingen, San Diego, CA), or cleaved caspase-3 (1:200 diluted, Cell Signaling Technology Inc., Beverly, MA). The slides were incubated with biotinylated secondary antibody, and then with avidin/biotinylated peroxidase complex for 30 min at room temperature (Vector Labs, Burlingame, CA), and were then incubated with 3'-diaminobenzamine (DAB) substrate. The sections were then counterstained with Modified Harris Hematoxylin. The images were taken randomly at using a Zeiss AxioCam HRc camera fitted to a Zeiss Axioskope 2 Plus microscope.

[³H]Thymidine uptake assay

MCF10DCIS.com cells were incubated with compounds in 5% horse serum/DMEM/F12 media for 3 days. One μ Ci of [³H]thymidine was added to each well 3 h before the harvest. Incorporation of [³H]thymidine into the cells was analyzed with a liquid scintillation spectrometer (Beckman Coulter, Fullerton, CA).

Statistical analysis

Statistical significance for average tumor burden, average body weight, and serum calcium level was evaluated using the Student's *t*-test.

RESULTS

Gemini vitamin D analogs inhibit NMU-induced mammary tumorigenesis in rats

The efficacies of a classic vitamin D analog and 6 different Gemini vitamin D analogs on NMU-induced mammary tumorigenesis were tested in rats. The tumors formed with this model are luminal ER-positive (28). The doses of vitamin D analogs in Table 1 were determined based on the maximum tolerated dose in CD-1 mice and our preliminary data (data not shown). As summarized in Table 1, body weights of animals treated with the vitamin D analogs were not significantly different from those in the control group throughout the study. Among 7 different vitamin D analogs, administration of 0072 and 0097 resulted in approximately a 60% reduction of the average tumor burden per rat. Gemini 0072 and 0097 have modification on both side chains with hexafluoride or hexadeuterium functional groups, which may have contributed to better bioavailability and an increased half-life of these compounds in animals. Gemini 0072 and 0097 had stronger activity than 3586 (44% reduction) and 0084 (29% reduction). Other analogs, 2198, 3582, and 0088 had little or no effect on tumor burden (Table 1). Potential toxicities such as body weight loss, diarrhea or blood in the feces were not observed in Gemini vitamin D treated animals.

Gemini vitamin D analog 0097 inhibits tumor growth without hypercalcemia toxicity

Hypercalcemic toxicity has limited the use of naturally occurring $1\alpha,25(\text{OH})_2\text{D}_3$ as a chemopreventive agent although it has been shown to have a potent anti-proliferating activity in different cell types (6,10-12). Among the vitamin D analogs we tested (Table 1), Gemini 0097 exerted the strongest activity in inhibiting tumor growth (Table 1), and it was selected for a further dose dependent study. In Table 2, we compared the efficacy and toxicity of Gemini 0097 to those of $1\alpha,25(\text{OH})_2\text{D}_3$. Table 2 shows that body weight in $1\alpha,25(\text{OH})_2\text{D}_3$ treated groups was comparable to those in the control group, and tumor growth was inhibited 20% and 34% at doses of 0.1 $\mu\text{g/kg}$ body weight and 0.3 $\mu\text{g/kg}$ body weight, respectively. However, we observed that the average level of calcium in serum was increased to 10.9 mg/dl and 11.7 mg/dl by $1\alpha,25(\text{OH})_2\text{D}_3$ at tumor inhibitory doses (Table 2). The normal range of calcium in serum is 8.8–10.4 mg/dL (19). As shown in Table 2, Gemini 0097 at the doses of 0.1 or 0.3 $\mu\text{g/kg}$ body weight significantly inhibited tumor burden per rat as well as tumor multiplicity

but did not show any changes in body weight or serum calcium level compared to the control group, indicating that Gemini 0097 had a better efficacy/toxicity profile than $1\alpha,25(\text{OH})_2\text{D}_3$.

Gemini vitamin D analogs increase markers of cell cycle arrest and apoptosis in NMU-induced mammary tumors

In Western blot analysis, we showed that administration of Gemini 0072 and 0097 upregulated the VDR and increased the level of the CDK inhibitor p21, a well-known inhibitor of cell proliferation in NMU-induced mammary tumors (Fig. 2A). Another CDK inhibitor, p27, was not affected. In addition, administration of 0072 and 0097 increased the level of cleaved PARP and caspase-3, which are markers of apoptosis. These results suggest that administration of 0072 and 0097 inhibit tumor growth by arresting the cell cycle and inducing apoptosis.

Effect of administration of Gemini 0072 and 0097 on proteins in NMU-induced mammary tumors: 2-D gel electrophoresis studies

Proteins in the cytosol of mammary tumors of rats treated with vehicle or Gemini vitamin D analogs 0072 or 0097 were resolved on 2-D electrophoresis gels in order to determine if the inhibition in tumor growth by the analogs was reflected by changes in the proteome. The protein stained 2-D gels resolved approximately 700 – 900 distinct protein spots after subtracting background using Non-Linear Phoretix 2-D software. Comparison of the data derived from tumors of rats treated with Gemini vitamin D analogs 0072 or 0097 to those from tumors of vehicle treated rats revealed relatively few protein spots that were increased or decreased more than 2.0 fold. Some of the protein spots were identified by liquid chromatography-mass spectrometry/ mass spectrometry (LC-MS/MS) of tryptic in-gel-digests of gel plugs (Fig. 2B). One of these proteins, apolipoprotein A-I, which was decreased by treatment with Gemini vitamin D analogs 0072 and 0097 was further investigated on Western blots (Fig. 3B). Apolipoprotein A-I was down-regulated in tumor bearing rats treated with 0072 and 0097. Several other proteins were also identified using LC/MS/MS (Fig. 2B).

Evaluation of proliferating cell nuclear antigen (PCNA) and caspase-3 expression in NMU-induced mammary tumors

All tumors from the control, 0072 or 0097 treated groups shown in Fig. 2B were evaluated and were determined to be adenocarcinomas, and the tumor type, differentiation/grade and stromal responses among the control group and Gemini 0072 and 0097 treated groups were not different when evaluated after H & E staining (data not shown). However, there was a significant difference in PCNA and caspase-3 expression between the control group and the Gemini 0072 and 0097 treated groups, as shown in Fig. 2B. The PCNA staining in tumor tissues was much stronger in the control group (vehicle) than in the Gemini 0072 or Gemini 0097 treated groups. The percentage ($\pm$ S.E.) of PCNA positive cells in tumors from the control group was $62.6 \pm 8.5\%$, whereas the percentages of PCNA positive cells in tumors from the Gemini 0072 and Gemini 0097 treated groups were $32.5 \pm 5.1\%$ (significantly different from the control group, $p = 0.02$) and $45.3 \pm 6.7\%$ ($p = 0.1$), respectively. In addition, the expression of caspase-3 was evaluated in five to six different sections and representative sections of mammary tumors from the control group or the Gemini vitamin D analogs treated groups are shown (Fig. 2B). The percentage ($\pm$ S.E.) of caspase-3 positive cells in tumors from the control group was 1.8 ± 0.4 , whereas the percentages of caspase-3 positive cells in tumors from the Gemini 0072 and Gemini 0097 treated groups were 13.1 ± 2.7 and 15.2 ± 4.3 , respectively (both significantly different from the control group, $p < 0.01$), indicating that the Gemini vitamin D analogs strongly induced apoptosis in mammary tumors.

Effect of administration of Gemini 0097 on the expression of CYP24A1, IGFBP-3 and p21 in NMU-induced mammary tumors

CYP24A1 encoding the 24-hydroxylase adds a hydroxyl group at the carbon 24 position in 25 (OH)D₃ or 1 α ,25(OH)₂D₃, which inactivates and leads to urinary excretion (8). CYP24A1 is a well-known target gene of vitamin D and it has been shown to be induced by 1 α ,25 (OH)₂D₃ or its analog in different cells (6). In this study, the mRNA level for CYP24A1 determined by quantitative PCR was increased by 0097 in tumor tissue dose dependently (Fig. 2C). The expression of insulin-like growth factor binding protein 3 (IGFBP-3), and CDK inhibitor p21 were enhanced by 0097, indicating that 0097 may regulate the IGF pathway to inhibit ER-positive mammary tumorigenesis in rats (Fig. 2C, D).

Inhibitory effect of Gemini 0097 on MCF10DCIS.com growth inhibition *in vitro* and MCF10DCIS.com xenograft tumor growth *in vivo*

Because NMU-induced mammary tumors in rats are ER-positive (28), we investigated the anti-tumor effect of Gemini vitamin D analogs on the growth of ER-negative MCF10DCIS.com cells. We first tested 60 Gemini vitamin D analogs for their growth inhibitory effects on cell growth of MCF10DCIS.com *in vitro*. As shown in Table 3, the growth inhibitory effects of the Gemini vitamin D analogs in the *in vitro* assay were 5–15 times greater than that of 1 α ,25 (OH)₂D₃ based on IC₅₀ values. In an ER-negative MCF10DCIS.com xenograft model, we selected Gemini 0097 for further *in vivo* testing, and we found that administration of Gemini 0097 inhibited the tumor burden by 50% without a significant change in body weight (Fig. 3A) or serum calcium level (data not shown). Interestingly, the expression levels of IGFBP-3 and p21 protein were upregulated by 0097 in MCF10DCIS.com tumors (Fig. 3B), suggesting that IGFBP-3 and p21 may act as common mediators of Gemini vitamin D analogs in inhibiting the growth of both ER-positive and ER-negative mammary tumors. Furthermore, expression of apolipoprotein A-I protein which was suppressed by 0072 and 0097 in NMU-induced rat tumors (Fig. 2B) was also reduced by 0097 in MCF10DCIS.com xenograft tumors (Fig. 3B). The bone morphogenetic proteins (BMPs) pathway was activated by 0097 treatment, which was measured by enhanced levels of phospho-Smad1/5 in tumors (Fig. 3B). Other Smad proteins (Smad1, 2, 3, and 5) were not changed by 0097 treatment (data not shown).

DISCUSSION

Breast cancer in humans is complex involving different histopathologies, genetic changes and clinical outcomes. These observations indicate that a single animal model may not mimic all features of human breast carcinogenesis (1). In the present study, we evaluated the inhibitory effects of Gemini vitamin D analogs on tumorigenesis in two different models: an ER-positive NMU-induced breast cancer model in rats and an ER-negative MCF10DCIS.com xenograft model in immunodeficient mice. NMU-induced mammary tumors showed similar gene expression profiles as were found in luminal type ER-positive, low to intermediate grade human breast cancer (28). Selective estrogen receptor modulators, such as tamoxifen, raloxifene or arzoxifene, have shown inhibitory effects in the NMU model (29), suggesting that this model may be a good ER-positive animal model of breast cancer for testing cancer preventive drugs.

As a xenograft model of breast cancer, we selected the MCF10DCIS.com cell line, which is known to be a basal-like ER-negative cell type (24). MCF10DCIS.com human breast cancer cells are derived from *Ha-ras* oncogene transfection and further passages in mice (30). In addition to genetic *ras* mutation, MCF10DCIS.com cells significantly overexpressed the breast cancer associated marker, Her-2, compared to the parent cells, MCF10A (unpublished data), and mammary tumors generated from MCF10DCIS.com cells were shown to dynamically interact with stroma during progression from *in situ* to invasive cancer (31). We used this cell line for both *in vitro* and *in vivo* studies. Although xenograft models may not be the best model

for testing preventive effects of Gemini vitamin D analogs, MCF10DCIS.com model represents DCIS progression to invasive ductal carcinoma, which can be useful for testing new compounds for inhibition of breast cancer progression.

In this report, we demonstrated that several Gemini vitamin D analogs had better efficacy for growth inhibition of MCF10DCIS.com cells than $1\alpha,25(\text{OH})_2\text{D}_3$ (Table 3) and that certain Gemini vitamin D analogs significantly suppressed tumor growth in both an ER-positive NMU-induced breast cancer model and in an ER-negative MCF10DCIS.com xenograft model without hypercalcemic toxicity (Table 2, Fig. 3). Hypercalcemia is a well-established toxicity induced by $1\alpha,25(\text{OH})_2\text{D}_3$ and for many classical vitamin D derivatives. We found that $1\alpha,25(\text{OH})_2\text{D}_3$ significantly increased the serum calcium level at doses showing tumor growth inhibition (Table 2). In contrast, Gemini 0097 exerted potent efficacy in preventing mammary tumor growth without causing hypercalcemia (Table 2). These results are in agreement with previous reports using a colon cancer model (19,20). The lack of hypercalcemic effects of Gemini vitamin D analogs at anticancer doses may be because of the flexibility of the ligand binding pocket in the vitamin D receptor (32). Recently, it was reported that structural rearrangement of the ligand binding pocket provides space to accommodate the second side chain of Gemini vitamin D analogs, which leads to expansion of the ligand binding pocket volume (33). This conformational flexibility of the ligand binding pocket to accommodate different ligands may result in different ligand-specific cofactor binding and/or selectivity of transcription of target genes (32,34), thus suggesting that Gemini vitamin D analogs may show different regulation of vitamin D target genes when compared with $1\alpha,25(\text{OH})_2\text{D}_3$.

In our mechanistic study, the expression levels of cyclin dependent kinase (CDK) inhibitor p21 and insulin-like growth factor binding protein-3 (IGFBP-3) in tumor tissues from both models were enhanced by Gemini 0097 treatment (Figs. 2 and 3). Interestingly, however, the markers for apoptosis including cleaved PARP and caspase-3 were regulated by Gemini 0072 and 0097 in ER-positive mammary tumors induced by carcinogen (Fig. 2), but not in ER-negative tumors from the xenograft model (data not shown). It has been shown in several studies that $1\alpha,25(\text{OH})_2\text{D}_3$ and its analogs suppress the mRNA and protein synthesis of ER through the negative vitamin D response element (nVDRE) in the ER promoter, which eventually leads to inhibition of cell proliferation through cell cycle arrest and apoptosis (35-37). The involvement of ER in inducing apoptosis by Gemini vitamin D analogs in NMU-induced mammary tumors needs to be further investigated.

Insulin-like growth factor binding proteins (IGFBPs), a group of six different proteins, sequester free insulin-like growth factors (IGFs) by binding with high affinity, which inhibits the IGF signaling pathway having mitogenic activity (38). The expression of IGFBPs has been shown to be up-regulated by $1\alpha,25(\text{OH})_2\text{D}_3$ and certain of its analogs in different cancer cells including prostate (39,40), colon (41), and breast cancer (21). In breast cancer, several studies suggest that IGFBP-3 acts as a tumor suppressive factor (42,43). Recently, the functional vitamin D response element (VDRE) has been identified in the promoter region of the IGFBPs (44,45), indicating that IGFBPs may be primary target genes of $1\alpha,25(\text{OH})_2\text{D}_3$ and its analogs. In the present study, we demonstrated that Gemini vitamin D analog treatment increased the expression of IGFBP-3 in breast tumors from both the carcinogen-induced model and the xenograft model, suggesting that IGFBP-3 induction may be mediated via direct transcriptional regulation. Interestingly, IGFBPs were shown to be involved in the regulation of the CDK inhibitor, p21 (46,47). IGFBP-3 knockdown by siRNA reversed the induction of p21 and growth inhibition by synthetic androgen R1881 in LNCaP human prostate cancer cells, indicating that IGFBP-3 was upstream of p21 (46). These studies support our hypothesis that the Gemini vitamin D analog 0097 inhibits tumor growth, at least in part, through the regulation of the link between IGFBP-3 and p21.

In several previous studies, vitamin D and its analogs have been involved in the regulation of transforming growth factor- β (TGF- β) signaling, which is extensively reviewed in Ref (6). We have also reported that Gemini vitamin D analogs were shown to activate the bone morphogenetic protein (BMP)/Smad signaling pathway via Ras/PKC α , which mediates the proliferation of the MCF10 breast epithelial cells (48,49). In this report, we demonstrate for the first time that the Gemini vitamin D analog 0097 activates Smad signaling *in vivo* in MCF10DCIS.com transplanted tumors (Fig. 3B), suggesting that the regulation of Smad signaling by vitamin D analogs may contribute to the suppression of mammary tumorigenesis. In addition, we showed that Gemini 0097 suppressed the apolipoprotein A-I protein expression in tumors from both animal models that we tested (Fig. 2B and Fig. 3B). Apolipoprotein A-I is one of the high-density lipoproteins, and $1\alpha,25(\text{OH})_2\text{D}_3$ and other VDR modulators were shown to down-regulate the expression of apolipoprotein A-I in human hepatocytes and intestinal cells (50). Since a low serum level of high-density lipoproteins, such as apolipoprotein A-I, has been associated with breast cancer, it will be important to determine whether inhibition of apolipoprotein A-I by vitamin D analogs has an effect on the regulation of mammary tumorigenesis.

In conclusion, we investigated the *in vivo* effect of certain Gemini vitamin D analogs on mammary tumor growth in a chemically-induced breast cancer model and in a xenograft model. We report here that Gemini vitamin D analogs significantly inhibit both ER-positive and ER-negative mammary tumor growth without increasing serum calcium levels. Mechanistic studies showed that the inhibitory activity was associated with the induction of IGFBP-3 and the CDK inhibitor p21 in both animal models. Taken together, these results suggest that Gemini vitamin D analogs may be promising agents for the prevention and treatment of breast cancer without hypercalcemic toxicity.

Acknowledgments

The authors thank BioXcell Inc. for their supports and Dr. Fred Miller at the Karmanos Cancer Institute for the MCF10 cell lines.

Grant Information: This work was supported in part by NIH R03 CA112642, NIH R01 CA127645, NIEHS P30 ES005022, a CINJ new investigator award and BioXcell Inc.

REFERENCES

1. Vargo-Gogola T, Rosen JM. Modelling breast cancer: one size does not fit all. *Nature reviews* 2007;7:659–72.
2. Cuzick J, Powles T, Veronesi U, et al. Overview of the main outcomes in breast-cancer prevention trials. *Lancet* 2003;361:296–300. [PubMed: 12559863]
3. Vogel VG, Costantino JP, Wickerham DL, et al. Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *Jama* 2006;295:2727–41. [PubMed: 16754727]
4. Shen Q, Brown PH. Novel agents for the prevention of breast cancer: targeting transcription factors and signal transduction pathways. *Journal of mammary gland biology and neoplasia* 2003;8:45–73. [PubMed: 14587863]
5. Welsh J, Wietzke JA, Zinser GM, Byrne B, Smith K, Narvaez CJ. Vitamin D-3 receptor as a target for breast cancer prevention. *J Nutr* 2003;133:2425S–33S. [PubMed: 12840219]
6. Deeb KK, Trump DL, Johnson CS. Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nature reviews* 2007;7:684–700.
7. Zinser GM, Suckow M, Welsh J. Vitamin D receptor (VDR) ablation alters carcinogen-induced tumorigenesis in mammary gland, epidermis and lymphoid tissues. *The Journal of steroid biochemistry and molecular biology* 2005;97:153–64. [PubMed: 16111884]

8. Dusso AS, Brown AJ, Slatopolsky E. Vitamin D. *Am J Physiol Renal Physiol* 2005;289:F8–28. [PubMed: 15951480]
9. Hussain-Hakimjee EA, Peng X, Mehta RR, Mehta RG. Growth inhibition of carcinogen-transformed MCF-12F breast epithelial cells and hormone-sensitive BT-474 breast cancer cells by 1 α -hydroxyvitamin D₅. *Carcinogenesis* 2006;27:551–9. [PubMed: 16195238]
10. Chung I, Wong MK, Flynn G, Yu WD, Johnson CS, Trump DL. Differential antiproliferative effects of calcitriol on tumor-derived and matrigel-derived endothelial cells. *Cancer research* 2006;66:8565–73. [PubMed: 16951169]
11. Narvaez CJ, Byrne BM, Romu S, Valrance M, Welsh J. Induction of apoptosis by 1,25-dihydroxyvitamin D₃ in MCF-7 Vitamin D₃-resistant variant can be sensitized by TPA. *The Journal of steroid biochemistry and molecular biology* 2003;84:199–209. [PubMed: 12711004]
12. Bouillon R, Eelen G, Verlinden L, Mathieu C, Carmeliet G, Verstuyf A. Vitamin D and cancer. *The Journal of steroid biochemistry and molecular biology* 2006;102:156–62. [PubMed: 17113979]
13. Murillo G, Mehta RG. Chemoprevention of chemically-induced mammary and colon carcinogenesis by 1 α -hydroxyvitamin D₅. *The Journal of steroid biochemistry and molecular biology* 2005;97:129–36. [PubMed: 16051482]
14. Mehta R, Hawthorne M, Uselding L, et al. Prevention of N-methyl-N-nitrosourea-induced mammary carcinogenesis in rats by 1 α -hydroxyvitamin D(5). *J Natl Cancer Inst* 2000;92:1836–40. [PubMed: 11078761]
15. Mehta RG, Hussain EA, Mehta RR, Das Gupta TK. Chemoprevention of mammary carcinogenesis by 1 α -hydroxyvitamin D₅, a synthetic analog of Vitamin D. *Mutat Res* 2003;523–524:253–64.
16. Huerta S, Irwin RW, Heber D, et al. 1 α ,25-(OH)₂-D₃ and its synthetic analogue decrease tumor load in the Apc(min) Mouse. *Cancer research* 2002;62:741–6. [PubMed: 11830528]
17. Audo I, Darjatmoko SR, Schlamp CL, et al. Vitamin D analogues increase p53, p21, and apoptosis in a xenograft model of human retinoblastoma. *Invest Ophthalmol Vis Sci* 2003;44:4192–9. [PubMed: 14507860]
18. Sundaram S, Sea A, Feldman S, et al. The combination of a potent vitamin D₃ analog, EB 1089, with ionizing radiation reduces tumor growth and induces apoptosis of MCF-7 breast tumor xenografts in nude mice. *Clin Cancer Res* 2003;9:2350–6. [PubMed: 12796405]
19. Spina C, Tangpricha V, Yao M, et al. Colon cancer and solar ultraviolet B radiation and prevention and treatment of colon cancer in mice with vitamin D and its Gemini analogs. *The Journal of steroid biochemistry and molecular biology* 2005;97:111–20. [PubMed: 16154354]
20. Spina CS, Ton L, Yao M, et al. Selective vitamin D receptor modulators and their effects on colorectal tumor growth. *The Journal of steroid biochemistry and molecular biology* 2007;103:757–62. [PubMed: 17368190]
21. Lee HJ, Liu H, Goodman C, et al. Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer. *Biochemical pharmacology* 2006;72:332–43. [PubMed: 16737686]
22. Maehr H, Uskokovic M, Adorini L, et al. Calcitriol derivatives with two different side chains at C-20 III. An epimeric pair of the gemini family with unprecedented antiproliferative effects on tumor cells and renin mRNA expression inhibition. *The Journal of steroid biochemistry and molecular biology* 2007;103:277–81. [PubMed: 17254779]
23. Perou CM, Sorlie T, Eisen MB, et al. Molecular portraits of human breast tumours. *Nature* 2000;406:747–52. [PubMed: 10963602]
24. Hu M, Yao J, Carroll DK, et al. Regulation of in situ to invasive breast carcinoma transition. *Cancer Cell* 2008;13:394–406. [PubMed: 18455123]
25. Thomas PE, Reik LM, Ryan DE, Levin W. Induction of two immunochemically related rat liver cytochrome P-450 isozymes, cytochromes P-450c and P-450d, by structurally diverse xenobiotics. *The Journal of biological chemistry* 1983;258:4590–8. [PubMed: 6403529]
26. Candiano G, Bruschi M, Musante L, et al. Blue silver: a very sensitive colloidal Coomassie G-250 staining for proteome analysis. *Electrophoresis* 2004;25:1327–33. [PubMed: 15174055]
27. Berkelman, T.; Stenstedt, T. 2-D Electrophoresis using immobilized pH gradients: Principles and Methods. 2nd ed. Amersham Bioscience; Uppasla, Sweden: 2002.

28. Chan MM, Lu X, Merchant FM, Iglehart JD, Miron PL. Gene expression profiling of NMU-induced rat mammary tumors: cross species comparison with human breast cancer. *Carcinogenesis* 2005;26:1343–53. [PubMed: 15845649]
29. Suh N, Glasebrook AL, Palkowitz AD, et al. Arzoxifene, a new selective estrogen receptor modulator for chemoprevention of experimental breast cancer. *Cancer research* 2001;61:8412–5. [PubMed: 11731420]
30. Miller FR, Santner SJ, Tait L, Dawson PJ. MCF10DCIS.com xenograft model of human comedo ductal carcinoma in situ. *J Natl Cancer Inst* 2000;92:1185–6. MCF10DCIS.com [PubMed: 10904098]
31. Tait LR, Pauley RJ, Santner SJ, et al. Dynamic stromal-epithelial interactions during progression of MCF10DCIS.com xenografts. *Int J Cancer* 2007;120:2127–34. MCF10DCIS.com [PubMed: 17266026]
32. Norman AW, Manchand PS, Uskokovic MR, et al. Characterization of a novel analogue of 1alpha, 25(OH)(2)-vitamin D(3) with two side chains: interaction with its nuclear receptor and cellular actions. *J Med Chem* 2000;43:2719–30. [PubMed: 10893309]
33. Ciesielski F, Rochel N, Moras D. Adaptability of the Vitamin D nuclear receptor to the synthetic ligand Gemini: remodelling the LBP with one side chain rotation. *The Journal of steroid biochemistry and molecular biology* 2007;103:235–42. [PubMed: 17218092]
34. Weyts FA, Dhawan P, Zhang X, et al. Novel Gemini analogs of 1alpha,25-dihydroxyvitamin D(3) with enhanced transcriptional activity. *Biochemical pharmacology* 2004;67:1327–36. [PubMed: 15013848]
35. Simboli-Campbell M, Narvaez CJ, van Weelden K, Tenniswood M, Welsh J. Comparative effects of 1,25(OH)2D3 and EB1089 on cell cycle kinetics and apoptosis in MCF-7 breast cancer cells. *Breast cancer research and treatment* 1997;42:31–41. [PubMed: 9116316]
36. Stoica A, Saceda M, Fakhro A, Solomon HB, Fenster BD, Martin MB. Regulation of estrogen receptor-alpha gene expression by 1, 25-dihydroxyvitamin D in MCF-7 cells. *Journal of cellular biochemistry* 1999;75:640–51. [PubMed: 10572247]
37. Swami S, Krishnan AV, Feldman D. 1alpha,25-Dihydroxyvitamin D3 down-regulates estrogen receptor abundance and suppresses estrogen actions in MCF-7 human breast cancer cells. *Clin Cancer Res* 2000;6:3371–9. [PubMed: 10955825]
38. Grimberg A, Cohen P. Role of insulin-like growth factors and their binding proteins in growth control and carcinogenesis. *Journal of cellular physiology* 2000;183:1–9. [PubMed: 10699960]
39. Boyle BJ, Zhao XY, Cohen P, Feldman D. Insulin-like growth factor binding protein-3 mediates 1 alpha,25-dihydroxyvitamin d(3) growth inhibition in the LNCaP prostate cancer cell line through p21/WAF1. *The Journal of urology* 2001;165:1319–24. [PubMed: 11257709]
40. Stewart LV, Weigel NL. Role of insulin-like growth factor binding proteins in 1alpha,25-dihydroxyvitamin D(3)-induced growth inhibition of human prostate cancer cells. *The Prostate* 2005;64:9–19. [PubMed: 15651061]
41. Palmer HG, Sanchez-Carbayo M, Ordonez-Moran P, Larriba MJ, Cordon-Cardo C, Munoz A. Genetic signatures of differentiation induced by 1alpha,25-dihydroxyvitamin D3 in human colon cancer cells. *Cancer research* 2003;63:7799–806. [PubMed: 14633706]
42. Tomii K, Tsukuda K, Toyooka S, et al. Aberrant promoter methylation of insulin-like growth factor binding protein-3 gene in human cancers. *Int J Cancer* 2007;120:566–73. [PubMed: 17096329]
43. Pazaitou-Panayiotou K, Kelesidis T, Kelesidis I, et al. Growth hormone-binding protein is directly and IGFBP-3 is inversely associated with risk of female breast cancer. *European journal of endocrinology / European Federation of Endocrine Societies* 2007;156:187–94. [PubMed: 17287408]
44. Peng L, Malloy PJ, Feldman D. Identification of a functional vitamin D response element in the human insulin-like growth factor binding protein-3 promoter. *Molecular endocrinology (Baltimore, Md)* 2004;18:1109–19.
45. Carlberg C, Dunlop TW, Saramaki A, Sinkkonen L, Matilainen M, Vaisanen S. Controlling the chromatin organization of vitamin D target genes by multiple vitamin D receptor binding sites. *The Journal of steroid biochemistry and molecular biology* 2007;103:338–43. [PubMed: 17234401]

46. Peng L, Wang J, Malloy PJ, Feldman D. The role of insulin-like growth factor binding protein-3 in the growth inhibitory actions of androgens in LNCaP human prostate cancer cells. *Int J Cancer* 2007;122:558–66. [PubMed: 17918155]
47. Terrien X, Bonvin E, Corroyer S, Tabary O, Clement A, Henrion Caude A. Intracellular colocalization and interaction of IGF-binding protein-2 with the cyclin-dependent kinase inhibitor p21CIP1/WAF1 during growth inhibition. *The Biochemical journal* 2005;392:457–65. [PubMed: 16131350]
48. Lee HJ, Wislocki A, Goodman C, et al. A novel vitamin D derivative activates bone morphogenetic protein signaling in MCF10 breast epithelial cells. *Mol Pharmacol* 2006;69:1840–8. [PubMed: 16533909]
49. Lee HJ, Ji Y, Paul S, Maehr H, Uskokovic M, Suh N. Activation of bone morphogenetic protein signaling by a Gemini vitamin D3 analogue is mediated by Ras/protein kinase C alpha. *Cancer research* 2007;67:11840–7. [PubMed: 18089814]
50. Wehmeier KR, Mazza A, Hachem S, et al. Differential regulation of apolipoprotein A-I gene expression by vitamin D receptor modulators. *Biochimica et biophysica acta* 2008;1780:264–73. [PubMed: 18082637]

Figure 1.

Structures of 1 α ,25-dihydroxyvitamin D₃ [active vitamin D₃ metabolite, calcitriol or 1 α ,25 (OH)₂D₃], the classic vitamin D analog, Ro-26-2198 (2198, 1 α ,25-dihydroxy-16,23Z-diene-26,27-hexafluoro-19-nor-cholecalciferol) and Gemini vitamin D analogs Ro-438-3582 [Gemini 3582, 1 α ,25-dihydroxy-20S-21(3-hydroxy-3-methyl-butyl)-23-yne-26,27-hexafluoro-cholecalciferol], Ro-438-3586 [Gemini 3586, 1 α ,25-dihydroxy-20R-21(3-hydroxy-3-methyl-butyl)-23-yne-26,27-hexafluoro-cholecalciferol], BXL-01-0084 [Gemini 0084, 1 α ,25-dihydroxy-20-methyl-20S-(4-hydroxy-4-methyl-pentyl)-23E-ene-26,27-hexafluoro-cholecalciferol], BXL-01-0088 [Gemini 0088, 1 α ,25-dihydroxy-20-methyl-20R-(4-hydroxy-4-methyl-pentyl)-23E-ene-26,27-hexafluoro-cholecalciferol], BXL-01-0072

[Gemini 0072, 1 α ,25-dihydroxy-20S-21(3-trideuteromethyl-3-hydroxy-4,4,4-trideuterobutyl)-23-yne-26,27-hexafluoro-19-nor-cholecalciferol], and BXL-01-0097 [Gemini 0097, 1 α ,25-dihydroxy-20R-21(3-trideuteromethyl-3-hydroxy-4,4,4-trideuterobutyl)-23-yne-26,27-hexafluoro-19-nor-cholecalciferol]. The different configuration of epimerization is indicated at the C₂₀ position as either S or R.

Figure 2. Effect of Gemini vitamin D analogs on cell cycle, apoptosis and vitamin D target genes in NMU-induced tumors

NMU-treated rats were given 0072 and 0097 as described in Tables 1 and 2, and tumor samples were analyzed **A**. Gemini vitamin D analogs 0072 and 0097 (0.3 μg/kg body weight/day) enhance VDR, apoptosis and a cell cycle arrest protein in mammary tumors induced by NMU in rats. **B**. Gemini vitamin D analogs 0072 and 0097 (0.3 μg/kg body weight/day) reduce the protein expression level of Apolipoprotein A-I; **1 and 3**. Krt-19 protein; **2**. Cytokeratin (CK19); **4**. no protein match; **5**. Annexin V complexed with glycerophosphoserine; **6**. β-actin, **7** and **11**. Apolipoprotein A-I; **8**. Thioredoxin; **9**. S100 calcium binding protein A6 (calcylin); **10**. Actin; **12**. Heat shock protein 70 cognate (first row). PCNA and caspase-3 staining of tumors are shown. Positive reactions for PCNA and cleaved caspase-3 are noted by a reddish brown precipitate in the cells (second and third rows). **C**. The Gemini vitamin D analog 0097 increases the mRNA level of CYP24A1 encoding 24-hydroxylase and insulin-like growth factor binding protein-3 (IGFBP-3) in NMU-induced mammary tumors. Two separate analyses were performed and combined (statistical significance: * $p < 0.01$, ** $p < 0.001$). **D**. The Gemini vitamin D analog 0097 increases the protein level of IGFBP-3 and p21 in NMU-induced mammary tumors.

Figure 3. The Gemini vitamin D analog 0097 significantly inhibits tumor growth and regulates the level of IGFBP-3, the CDK inhibitor p21, pSmad1/5, and apolipoprotein A-I in ER-negative MCF10DCIS.com xenograft tumors in SCID mice

MCF10DCIS.com cells (1×10^6 cells) were injected into a mammary fat pad. Four days later, the mice received either vehicle (0.1 ml) or Gemini vitamin D analog 0097 (0.1 μ g/kg body weight in vehicle in 0.1 ml) intraperitoneally daily. Six weeks later, tumors were weighed and collected at autopsy for further analysis. **A.** The average body weight and tumor weight per mouse were shown. The number of mice in vehicle control and 0097-treated group was 7 and 5, respectively (statistical significance: *, $p = 0.04$). **B.** The proteins from tumor samples were extracted with RIPA buffer and the expression level of IGFBP-3, CDK inhibitor p21, apolipoprotein A-I, pSmad1/5 and β -actin was analyzed by Western blotting. The bands intensity was quantified as described in Materials and Methods.

Table 1

Prevention of NMU-induced breast cancer by vitamin D analogs

Treatment [*]	Dose (µg/kg body weight)	No. animals	Body weight at autopsy (grams) (mean ± S.E.)	Tumor burden [†] (grams) (mean ± S.E.)
Control (Vehicle)	-	12	242.7 ± 6.5	9.0 ± 2.8
Ro-26-2198 (2198)	0.03	6	224.3 ± 16.0	8.4 ± 4.1
Gemini 3582 (3582)	0.1	6	248.3 ± 10.6	9.2 ± 2.3
Gemini 3586 (3586)	0.3	6	235.0 ± 2.6	5.0 ± 2.0
Gemini 0084 (0084)	0.03	6	247.6 ± 8.1	6.4 ± 3.0
Gemini 0088 (0088)	0.1	6	255.7 ± 9.9	8.9 ± 2.9
Gemini 0072 (0072)	0.3	6	250.8 ± 9.5	3.6 ± 1.7 ($p=0.06^{\ddagger}$)
Gemini 0097 (0097)	0.3	6	262.7 ± 7.0	3.4 ± 1.7 ($p=0.05^{\ddagger}$)

* All rats (21 days old) received NMU (50 mg /kg body weight, i.p.) 1 week before injecting vitamin D analogs i.p. once a day, 5 days a week for 9 weeks.

[†] Tumor burden; average tumor weight per rat at autopsy.

[‡] p is the value for the comparison of tumor burden in rats treated with vitamin D analogs when compared with control rats treated with vehicle alone.

Table 2
Effect of different doses of 1 α ,25(OH) $_2$ D $_3$ (calcitriol) and Gemini 0097 on NMU-induced breast cancer.

Treatment*	Dose (μ g/kg body weight)	No. animals	Body weight at autopsy (grams) (mean $\pm$ S.E.)	Tumor burden [†] (grams) (mean $\pm$ S.E.)	Tumor multiplicity [#] (mean $\pm$ S.E.)	Serum calcium level (mg/dl) (mean $\pm$ S.E.)
Control (Vehicle)	0	30	241.3 $\pm$ 4.0	8.5 $\pm$ 1.7	2.9 $\pm$ 0.4	9.3 $\pm$ 0.9
Calcitriol (1 α ,25(OH) $_2$ D $_3$)	0.03	15	238.3 $\pm$ 4.2	8.8 $\pm$ 1.8	2.9 $\pm$ 0.5	9.3 $\pm$ 0.8
Calcitriol (1 α ,25(OH) $_2$ D $_3$)	0.1	15	241.9 $\pm$ 6.6	6.8 $\pm$ 1.5	2.3 $\pm$ 0.5	10.9 $\pm$ 1.5**
Calcitriol (1 α ,25(OH) $_2$ D $_3$)	0.3	14	225.7 $\pm$ 9.9	5.6 $\pm$ 1.5	2.6 $\pm$ 0.4	11.7 $\pm$ 1.4**
Gemini 0097 (0097)	0.03	15	251.2 $\pm$ 6.1	9.0 $\pm$ 2.6	2.9 $\pm$ 0.5	9.5 $\pm$ 1.0
Gemini 0097 (0097)	0.1	15	248.5 $\pm$ 6.1	4.2 $\pm$ 1.2 ($p=0.02^{\ddagger}$)	1.8 $\pm$ 0.4 ($p=0.03^{\ddagger}$)	9.6 $\pm$ 1.5
Gemini 0097 (0097)	0.3	15	252.5 $\pm$ 5.4	4.2 $\pm$ 1.4 ($p=0.03^{\ddagger}$)	2.3 $\pm$ 0.4	9.9 $\pm$ 1.1

* All rats (21 days old) received NMU (50 mg /kg body weight, i.p.) 1 week before injecting vitamin D analogs i.p. once a day, 5 days a week for 9 weeks.

[†] Tumor burden; average tumor weight per rat at autopsy.

[#] Tumor multiplicity; average number of tumors per rat at autopsy.

[‡] p is the value for the comparison of tumor burden in rats treated with calcitriol or Gemini 0097 when compared with control rats treated with vehicle alone.

** Hypercalcemic. The normal range of calcium in serum is 8.8–10.4 mg/dL (19).

Table 3

IC₅₀ for the inhibitory effect of 1 α ,25(OH)₂D₃ and Gemini vitamin D analogs on the growth of MCF10DCIS.com breast cancer cells.

Vitamin D and analogs	IC ₅₀ (nM) *
1 α ,25(OH) ₂ D ₃	3.2
Gemini 3582 (3582)	0.2
Gemini 3586 (3586)	0.2
Gemini 0072 (0072)	0.4
Gemini 0097 (0097)	0.6

* MCF10DCIS.com cells were treated with vitamin D analogs (0.01, 0.1, 1, 10 and 100 nM) in 5% horse serum/DMEM/F12 media for 3 days. IC₅₀ values were determined using TableCurve 2D v5.1 software.