Published in final edited form as:

Inorg Chem. 2009 March 2; 48(5): 2300–2308. doi:10.1021/ic802180d.

Substrate binding preferences and pK_a determinations of a nitrile hydratase model complex: Variable solvent coordination to [(bmmp-TASN)Fe]OTf

Martin G. O'Toole[†], Brian Bennett[‡], Mark S. Mashuta[†], and Craig A. Grapperhaus^{†,*}

[†]Department of Chemistry, University of Louisville, Louisville, Kentucky 40292

[‡]National Biomedical EPR Center, Department of Biophysics, Medical College of Wisconsin, 8701 Watertown Plank Road, Milwaukee, Wisconsin 53226-0509.

Abstract

The five-coordinate iron-dithiolate complex (N,N'-4,7-bis-(2'-methyl-2'-mercatopropyl)-1-thia-4,7-diazacyclononane)iron(III), [LFe]⁺, has been isolated as the triflate salt from reaction of the previously reported LFeCl with thallium triflate. Spectroscopic characterization confirms an $S = 1/2$ ground state in non-coordinating solvents with room temperature $\mu_{\text{eff}} = 1.78 \mu_B$ and EPR derived g -values of $g_1 = 2.06$, $g_2 = 2.03$ and $g_3 = 2.02$. [LFe]⁺ binds a variety of coordinating solvents resulting in six-coordinate complexes [LFe-solvent]⁺. In acetonitrile the low-spin [LFe-NCMe]⁺ ($g_1 = 2.27$, $g_2 = 2.18$ and $g_3 = 1.98$) is in equilibrium with [LFe]⁺ with a binding constant of $K_{\text{eq}} = 4.03$ at room temperature. Binding of H₂O, DMF, methanol, DMSO and pyridine to [LFe]⁺ yields high-spin six-coordinate complexes with EPR spectra that display significant strain in the rhombic zero-field splitting term E/D . Addition of one equivalent of triflic acid to the previously reported diiron species (LFe)₂O results in the formation of [(LFe)₂OH]OTf, which has been characterized by x-ray crystallography. The aqueous chemistry of [LFe]⁺ reveals three distinct species as a function of pH: [LFe-OH₂]⁺, [(LFe)₂OH]OTf, and (LFe)₂O. The pK_a values for [LFe-OH₂]⁺ and [(LFe)₂OH]OTf are $5.4 \pm .1$ and $6.52 \pm .05$ respectively.

Keywords

Iron-Thiolate; Nitrile Hydratase; EPR; Inorganic Modeling

Introduction

Small molecule model complexes of the non-heme iron enzyme nitrile hydratase (NHase) have been instrumental in elucidating the geometric and electronic structure of the enzyme active site.^{1–11} NHase, which catalyses the hydrolysis of nitriles to amides, possesses either a low-spin Fe(III) or non-corrinoid Co(III) in an unusual nitrogen/sulfur-rich environment with a primary coordination sphere consisting of two deprotonated amides from the peptide backbone, three cysteine derived sulfur donors (now known to exist in three different oxidation states), and a variable coordination site for interacting with substrates/products, Figure 1.^{12–18} Based on spectroscopic and computational results, several catalytic mechanisms have been proposed.

grapperhaus@louisville.edu.

Supporting Information Available: Expanded experimental protocols, EPR and UV/visible spectra, and cyclic voltammetry of [LFe]OTf, and EPR simulations of [LFe-DMSO]OTf in PDF format and crystallographic data in CIF format (CCDC 708538). This material is available free of charge via the Internet at <http://pubs.acs.org>.

4,8,19–21 Despite intensive efforts, it is still uncertain whether water, nitrile, or both substrates coordinate at the active site during catalysis.

The substrate binding affinities of active site models have not provided consistent results.^{22, 23} A high-spin ($S = 5/2$), five-coordinate iron complex with deprotonated amides and thiolate donors coordinates water, but shows no affinity for nitriles.²² A closely related low-spin ($S = 1/2$), five-coordinate iron complex featuring imine nitrogen donors and thiolates exclusively binds nitriles at low temperature, but undergoes ligand degradation upon exposure to water.²³ To our knowledge, a single model complex that coordinates both nitrile and water has not yet been reported. In this manuscript we report the five-coordinate iron dithiolate, N,N'-Bis-(2'-methyl-2'-mercaptopropyl)-1-thia-4,7-diazacyclononane)iron(III) triflate (**[LFe]OTf**). **[LFe]OTf** binds water, nitriles and amides allowing for the first time a direct comparison of substrate (and product) binding affinities in a single model complex. Additionally, the pK_a of the water and hydroxide bound derivatives of **[LFe]OTf** have been evaluated and offer insight regarding the need for low-spin iron at the enzyme active site.

Experimental Section

Materials and Methods

All reagents were obtained from commercially available sources and used as received unless otherwise noted. All solvents were dried and freshly distilled using standard techniques under a nitrogen atmosphere and degassed using freeze-pump-thaw techniques.^{24,25} All reactions were conducted using standard Schlenk techniques under an argon atmosphere or in an argon-filled glovebox unless otherwise noted.²⁶ The complexes [(bmmp-TASN)FeCl], **LFeCl**, and [(bmmp-TASN)Fe]₂O, (**LFe**)₂O, were prepared as previously reported.² Thallium triflate was prepared from thallium carbonate and triflic acid according to published protocols.²⁷

N,N'-Bis-(2'-methyl-2'-mercaptopropyl)-1-thia-4,7-diazacyclononane iron(III) triflate (**[LFe]OTf·0.5CH₂Cl₂**)

To a suspension of 100 mg (0.24 mmol) **LFeCl** in 100 mL dry acetonitrile was added dropwise via cannula a solution of 86 mg (0.24 mmol) thallium triflate in 15 mL acetonitrile. After stirring for 15 hours, the solvent was removed under vacuum and the product extracted into 60 mL dichloromethane followed by filtration through a fritted tube. Removal of solvent from the filtrate yielded **[LFe]OTf·0.5CH₂Cl₂** as a dark blue solid. Yield: 81 mg (0.15 mmol, 63 %). Electronic absorption (dichloromethane (22 °C)): $\lambda_{\text{max}}(\epsilon)$: 274(7601), 317(6740), 427(2570), 504(1400), 623(1600). IR (KBr pellet), cm^{-1} : 3436 (br), 2949 (m), 2912 (m), 2880 (m), 2843 (m), 1450 (m), 1262 (s), 1135 (m), 1102 (m), 1074 (m), 1029 (s), 804 (m), 636 (m). MS-ESI, m/z calcd. for C₁₄H₂₈N₂S₃Fe, [M+]⁺ 376.08; Found, 376.05. Anal. Calcd. for C_{15.5}H₃₁N₂S₄O₃F₃FeCl(**[LFe]OTf·0.5CH₂Cl₂**): C, 31.48; H, 4.95; N, 4.59. Found: C, 31.17; H, 4.92; N, 4.88.

[(LFe)₂OH]Tf

A solution of (**LFe**)₂O (50 mg, 65 μmol) in 100 mL acetonitrile was cooled to $-15\text{ }^{\circ}\text{C}$ in a dry ice/ethylene glycol bath. A solution of 5.7 μL (9.7 mg, 65 μmol) 98% triflic acid in 10 mL acetonitrile was added dropwise via cannula over a 30 min. period. The solution was stirred for 2 hours as it gradually warmed to room temperature. The solution was filtered through a fritted tube and concentrated to 15 mL. Diethyl ether addition led to precipitation of **[(LFe)₂OH]Tf** as a purple solid. Yield: 45 mg (49 μmol , 75%). X-ray quality crystals were obtained by vapor diffusion of diethyl ether into a methanol solution of **2** at $4\text{ }^{\circ}\text{C}$ under an argon atmosphere. Electronic absorption (acetonitrile): $\lambda_{\text{max}}(\epsilon)$: 311(3600), 433(700), 602(1600). IR (KBr pellet), cm^{-1} : 3490 (br), 2958 (m), 2913 (m), 2880 (m), 2847 (m), 1454 (m), 1254 (s), 1147 (m), 1131 (m), 1074 (m), 1029 (s), 976 (m), 641 (s), 518 (m). Anal. Calcd. for

$C_{29}H_{58}N_4S_8O_7F_6Fe_2[(LFe)_2OH]OTf \cdot HOTf$: C, 36.24; H, 5.88; N, 5.64. Found: C, 36.92; H, 5.77; N, 5.57

Physical Methods

Elemental analyses were obtained from Midwest Microlab (Indianapolis, IN). Infrared spectra were recorded on a Thermo Nicolet Avatar 360 spectrometer at 4 cm^{-1} resolution. 1H NMR spectra were obtained on a Varian Inova500 500 MHz spectrometer. Mass spectrometry (ESI-MS) was performed by the Laboratory for Biological Mass Spectrometry at Texas A&M University. The room temperature magnetic moment of $[LFe]OTf$ was determined by the Evan's method in dichloromethane with use of residual CH_2Cl_2 peak and its shifted counterpart in calculation of μ_{eff} .²⁸ Cyclic voltammetry was performed using a PAR 273 potentiostat with a three electrode cell (glassy carbon working electrode, platinum wire counter electrode, and a Ag/AgCl pseudo reference electrode) at room temperature in an argon filled glove box. All potentials were scaled to a ferrocene/ferrocenium standard using an internal reference. Catalytic trials (see Supporting Information) were monitored by gas chromatography using an HP 5890 series II chromatograph fitted with a flame ionization detector using compressed air as a carrier gas with an RTX-1 column (60 m length, i.d. 32, serial number 469083) from Restek Corporation.

Electronic absorption spectra were recorded with an Agilent 8453 diode array spectrometer with air free 1 cm path length quartz cell or in a custom made dewar with a 1 cm path length quartz sample compartment. The equilibrium constant for acetonitrile binding was determined as described in the Supporting Information based on minimization of the R^2 value of the resulting van't Hoff plot using previously reported methods.^{23,29} The pK_a values for $[LFe-OH_2]^+$ and $[(LFe)_2OH]OTf$ were determined spectrophotometrically in a 70:30 water/acetone mixture due to poor solubility of $(LFe)_2O$ in aqueous solution³⁰ using a Corning pH 440 meter with full details in the Supporting Information.

X-band EPR spectra were obtained either on a Bruker EMX EPR spectrometer (77 K, 6.3 mW, modulation amplitude 5 G) or on a Bruker EleXsys E600 spectrometer (20 K, 2 mW, modulation amplitude 10 G) equipped with an ER4116DM TE₀₁₂/TE₁₀₂ dual-mode cavity and an Oxford Instruments ESR900 helium flow cavity. The $S = 1/2$ spectra were modeled using SIMPOW.³¹ Computer simulations of $S = 5/2$ spectra were carried out using matrix diagonalization with XSophe v.1.1.3,³² assuming a spin Hamiltonian $H = \beta g.B.S + S.D.S$ and including distributions ("strains") in the zero-field splitting parameters, σD and $\sigma E/D$.

Crystallographic Studies

A dark purple trapezoidal prism $0.29 \times 0.22 \times 0.05\text{ mm}^3$ crystal of $[(LFe)_2OH]OTf$ was selected for X-ray data collection on a Bruker SMART APEX CCD diffractometer. Frame data were collected (SMART³³ v5.632) and processed (SAINT³⁴ v6.45a) to determine final unit cell parameters: $a = 15.472(6)\text{ \AA}$, $b = 10.629(4)\text{ \AA}$, $c = 25.894(13)\text{ \AA}$, $\beta = 95.172(5)^\circ$, $V = 4241(3)\text{ \AA}^3$, $Z = 4$ and $\rho_{calcd} = 1.477\text{ Mgm}^{-3}$. The raw hkl data were corrected for absorption (SADABS³⁵ v2.10; transmission min./max. = 0.744/0.948; $\mu = 1.082\text{ mm}^{-1}$) and the structure was solved by Patterson methods (SHELXTL³⁶⁻³⁸ (v 6.14) suite of programs) in the space group Ia. For all 9283 unique reflections ($R(int) = 0.0273$) the final anisotropic full matrix least-squares refinement on F^2 for 468 variables converged at $R1 = 0.0420$ and $wR2 = 0.0994$ with a GOF of 1.08. Crystallographic parameters for $[(LFe)_2OH]OTf$ are displayed in Table 1 with additional experimental details provided in the Supporting Information.

Results and Discussion

Synthesis and Characterization

Previously we reported **LFeCl** (L = bmmp-TASN) as a model complex of nitrile hydratase.² The coordinated chloride was not displaced by water or nitrile, although under basic conditions the μ -oxo complex **(LFe)₂O** was obtained. Herein we report metathesis of the chloride with thallium triflate to yield the five-coordinate complex **[LFe]OTf**, Scheme 1. The infrared spectrum of **[LFe]OTf** displays strong absorptions at 1262 and 1029 cm⁻¹ attributed to the triflate counter ion, but is otherwise similar to that of **LFeCl**. The room temperature magnetic moment of **[LFe]OTf** in dichloromethane (1.78 μ_B) is consistent with the spin-only value for low-spin iron(III). Mass spectral analysis of **[LFe]OTf** from dichloromethane, acetonitrile or aqueous solutions displays a major peak at $m/z = 376.08$ as expected for **[LFe]⁺** with no evidence of triflate or solvent coordination in the gas phase.

As a solid **[LFe]OTf** is vivid blue, however, in solution the color depends on equilibrium solvent binding. In water, three species (**[LFe-OH₂]⁺**, **[(LFe)₂OH]⁺**, and **(LFe)₂O**) were observed as a function of pH. **[(LFe)₂OH]OTf** is also obtained by stoichiometric addition of triflic acid to acetonitrile solutions of **(LFe)₂O** with a corresponding bathochromic shift in the low energy absorption band from 553 to 603 nm, Figure 2. The infrared spectrum of **[(LFe)₂OH]OTf** is marked by bands at 1254 and 1029 cm⁻¹ due to the triflate counter ion. Also of note is the absence of the Fe-O-Fe stretch at 800 cm⁻¹ found in the IR spectrum of the precursor μ -oxo complex.² The **[(LFe)₂OH]OTf** complex is soluble in water, acetonitrile, alcohols, and sparingly soluble in dichloromethane, yielding violet colored solutions in each case.

Structural Characterization of **[(LFe)₂OH]OTf**

The complex **[(LFe)₂OH]OTf** crystallizes in the monoclinic space group Ia with one metal containing cation, **[(LFe)₂OH]⁺**, one triflate anion, and a partial occupancy methanol solvent molecule. The position of the hydrogen atom of the bridging hydroxide was determined from an electron density map and refined isotropically. An ORTEP³⁹ representation of **[(LFe)₂OH]⁺** is presented in Figure 3 with selected bond distances and angles given in Table 2. As with all other complexes of **L**, the TASN ligand backbone constrains the two amines and thioether sulfur to a facial coordination mode.^{1,2,11} For each iron center, the three sulfur donors are meridional with each sulfur cis to the hydroxide bridge. An amine nitrogen occupies the position trans to the bridging hydroxide. The iron-ligand bond distances of **[(LFe)₂OH]⁺** are within the ranges expected for high spin Fe(III).^{2,11,40}

Comparison of the structure of **[(LFe)₂OH]OTf** with previously reported crystallographic data for the related μ -oxo complex, **(LFe)₂O**, reveals important distinctions between the two structures.² The Fe-O-Fe angle for **[(LFe)₂OH]⁺** is 166.71(14)°, while the corresponding angle for **(LFe)₂O** is 180.0°. This decrease in Fe-O-Fe bond angle results from protonation of the bridging oxo ligand.⁴¹⁻⁴³ The steric bulk of the ligands force a severe distortion of the idealized trigonal arrangement about the bridging oxygen atom, with Fe-O-H angles compressed to an average of 97°.

Solution Studies

The complex **[LFe]OTf** is soluble in a wide variety of polar and non-polar solvents yielding vivid green (5-coordinate) or blue (six-coordinate) solutions. A low energy charge transfer band near 600 nm, assigned as a thiolate to iron charge transfer, is observed for all six-coordinate complexes assigned as either **[LFe-solvent]⁺** or **LFe-OTf**, depending on reaction conditions.^{44,45} In contrast, the 5-coordinate complex, **[LFe]⁺** displays a lower energy band at 623 nm and a distinct shoulder at 427 nm.

Vacant Binding Site: Five-coordinate, $S = 1/2$

Dichloromethane solutions of $[\text{LFe}]\text{OTf}$ are green in color at all temperatures above $-40\text{ }^{\circ}\text{C}$ indicative of five-coordinate $[\text{LFe}]^+$. The room temperature UV-visible spectrum displays diagnostic bands at 315, 427, and 623 nm. Upon cooling to temperatures below $-40\text{ }^{\circ}\text{C}$ the color changes from green to blue due to coordination of triflate. As shown in Figure S1, the low energy band shifts from 623 nm to 603 nm and the shoulder at 427 nm decreases in intensity.

The lack of triflate coordination at room temperature is also evident in the cyclic voltammogram. In dichloromethane $[\text{LFe}]\text{OTf}$ displays a pseudo-reversible event at -500 mV (vs Fc/Fc^+) assigned as $\text{Fe}^{\text{III/II}}$ and an irreversible oxidation at 510 mV assigned to the ligand, Figure S2. The metal-based reduction potential is shifted by $+560\text{ mV}$ with respect to LFeCl consistent with the loss of a donor atom. The oxidation is similarly shifted.

The EPR spectrum of $[\text{LFe}]\text{OTf}$ in dichloromethane (77 K) indicates incomplete triflate coordination even at low temperature. The spectrum reveals a sharp signal for a low-spin component and a broad complex high-spin signal, Figure S3, that gains intensity as the temperature is lowered, Figure S4. The sharper signal attributable to $[\text{LFe}]^+$ was simulated (SIMPOW) with g -values of 2.06, 2.03, and 2.02, Figure 4.³¹ This narrow g -value spread is atypical of low-spin iron(III)^{46,47} and more akin to that of metal-coordinated thiyl radicals, which show significantly less delocalization onto aromatic ligands than their phenoxyl counterparts.⁴⁸ Computational investigations by our group and others on low-spin iron(III)-thiolates show significant spin-density on sulfur (even in the absence of aromatic ligands) consistent with Fe(II)-thiyl radical character.^{5,11} The observed $S = 1/2$ ground state is consistent with increased Fe-S covalency as the thiolate compensates for the loss of the sixth donor and could result from coupling of either intermediate- or low-spin Fe(II) with the thiyl radical.^{5,11}

Nitrile Binding: Six-coordinate, $S = 1/2$

The low affinity of $[\text{LFe}]^+$ for triflate ensures an accessible coordination site for solvent (substrate) binding. At $40\text{ }^{\circ}\text{C}$ in acetonitrile solutions of $[\text{LFe}]^+$ are green with absorbance bands at 315, 427, and 603 nm similar to the spectrum observed in dichloromethane and consistent with a five-coordinate complex. However, upon cooling to room temperature and below the solution takes on a blue color as the peak at 427 nm loses intensity and the low energy band shifts towards 596 nm, Figure 5. The onset of these changes occurs at significantly higher temperature than in dichloromethane and arises due to nitrile coordination. The apparent equilibrium constants for nitrile binding of 4.7 at 298 K and 25 at $-40\text{ }^{\circ}\text{C}$ are similar to those reported by Kovacs in a related model complex.²³

The coordination of acetonitrile at room temperature is confirmed by cyclic voltammetry. The CV of $[\text{LFe}]\text{OTf}$ in acetonitrile, Figure S2, displays a reversible event at -622 mV (vs. Fc/Fc^+) assigned to the $\text{Fe}^{\text{III/II}}$ couple of $[\text{LFe-NCMe}]^+$ and a chemically irreversible oxidation at 166 mV . The reduction potential is shifted by -120 mV as compared to that of $[\text{LFe}]^+$ consistent with the addition of a weakly coordinating ligand.

The EPR spectrum of $[\text{LFe}]\text{OTf}$ (acetonitrile glass 77 K) also confirms equilibrium binding of acetonitrile, Figure 6A. In addition to the sharp axial signal of $[\text{LFe}]^+$ ($g_1 = 2.04$, $g_2 = 2.02$, $g_3 = 2.01$; 7% relative intensity), the spectrum also displays a rhombic signal ($g_1 = 2.27$, $g_2 = 2.18$, $g_3 = 1.98$; 93% relative intensity) attributed to $[\text{LFe-NCMe}]^+$. The large anisotropy in the g -values is typical of low-spin, six-coordinate Fe(III) complexes.^{46,47}

Similar results are obtained in benzonitrile, Figure 6B. A rhombic signal ($g_1 = 2.28$, $g_2 = 2.18$, $g_3 = 2.00$, relative intensity 85%) attributed to $[\text{LFe-NCPh}]^+$ is observed in addition to the

axial signal of $[\text{LFe}]^+$. The similarity in g-values for the two nitrile bound derivatives $[\text{LFe-NCMe}]^+$ and $[\text{LFe-NCPh}]^+$ implies that the identity of the nitrile donor does not significantly influence the electronic environment. This is consistent with recent computational studies that suggest the nitrile is not significantly polarized upon coordination to iron in NHase in contrast to expectations for a nitrile bound mechanism.⁴⁹

Water and other donor solvents: Six-coordinate, $S = 5/2$

The donor ability of nitriles to $[\text{LFe}]^+$ is limited and low temperatures are required to facilitate coordination. Other solvents including water and DMF more strongly coordinate $[\text{LFe}]^+$ leading to high-spin derivatives. Solutions of $[\text{LFe}]^+$ in H_2O ($\text{pH} < 6.1$), DMF, DMSO, pyridine, or methanol are dark blue at all temperatures with a charge transfer band near 600 nm and no shoulder at 427 nm. The low energy band blue-shifts with increased σ -donor ability of the sixth ligand, although in H-donor solvents the shift is tempered by interactions between the solvent and sulfur, Table S1.

The best evidence for solvent coordination is obtained from EPR spectroscopy. The complex EPR spectra of $[\text{LFe}]\text{OTf}$ in DMSO, pyridine, methanol, DMF and water display multiple broad turning points between 500–4000 G, Figure 7. The observed spectra are similar to those reported in literature for other iron complexes with thiolate or chloride donors.^{22,50,51} Despite the complexity of the spectra, the vast majority of the signal intensity ($> 99\%$ of the spins) can be simulated as a single monomeric highspin iron complex with trace quantities of a $g_{\text{eff}} \sim 4.3$ signal as described below.

The EPR signals shown in Figure 7 are similar to those reported for iron-catecholate complexes⁵² in that the strain in E (here characterized as a strain in E/D) extended the envelope of E/D such that one of the “preferred” values of E/D was exhibited by some of the molecules. These preferred values give rise to anomalously intense features in the EPR spectrum due to partial or total lack of orientation selection in the powder spectrum.⁵³ In the case of the catecholate species, the strains in E were sufficiently high that the spectra exhibited an intense component at $g_{\text{eff}} \sim 4.3$ due to a subpopulation with $E/D \sim 1/3$, although the mean value of E/D was as low as 0.120.

The strain-dependent appearance of the $g_{\text{eff}} \sim 4.3$ resonance can be appreciated by comparing the calculated spectra of Figure 8A and B, that have identical spin Hamiltonian parameters except for the strain in E/D. In the present case, the signals from $[\text{LFe}]\text{OTf}$ exhibit low field lines that are split by only about 170 G (17 mT), compared to about 500 G (50 mT) for the catecholate complexes. Simulations (Figure 8B – E) indicate that the smaller splitting is due to a lower mean E/D of 0.045. One of the consequences of this lower value for E/D is a change in the relative intensities of the two low field lines; g_x and g_y for the $m_S = 3/2$ Kramers' doublet of $S = 5/2$ approach zero as E/D decreases and the spectrum spans an increasingly wide field envelope, leading to a diminution of the intensity of the higher-field line of the pair (the g_z resonance of $m_S = 3/2$, at 1200 G [120 mT], $g_{\text{eff}} \sim 5.7$). As strains in E/D are introduced (Figure 8F – M), this feature appears to grow in intensity again. However, unlike the lowest field resonance of $m_S = 1/2$ at 1000 G (100 mT; $g_{\text{eff}} \sim 6.6$), the line at 1200 G does not become broadened at successively higher values of $\sigma E/D$. In addition, it shifts slightly to a resonance position corresponding to $g_{\text{eff}} = 6.0$ and assumes a typical g_x - g_y derivative shape. Thus the resonance observed is actually due to strain-dependent access of the preferred E/D value of zero, giving rise to an intense resonance due to the g_x and g_y transitions in $m_S = 1/2$. Only when very high E/D-strain is present is the $g_{\text{eff}} \sim 4.3$ line observed (Figure 8M). At such a high value of $\sigma E/D$, only the preferred resonances ($g_{\text{eff}} \sim 6$, $E/D \sim 0$ and $g_{\text{eff}} \sim 4.3$, $E/D \sim 1/3$) are resolved. As expected from the above assignment, the relative intensities of the $g_{\text{eff}} \sim 6$ and $g_{\text{eff}} \sim 4.3$ resonances change upon raising E/D, with the higher value of E/D favoring $g_{\text{eff}} \sim 4.3$.

From the simulations displayed in Figure 8, the high spin spectra of **[LFe]OTf** in H₂O, pyridine, DMSO, methanol and DMF can be explained as containing contributions from a slightly rhombic ($E/D = 0.045$) $S = 5/2$ Fe(III) species that exhibits strain in E/D , and an additional component that is responsible for the $g_{\text{eff}} = 4.3$ line. The strains in the slightly rhombic species result in a broadened resonance at $g_{\text{eff}} \sim 6.6$, due to the E/D envelope associated with the g_x transition in $m_S = 1/2$, and a resonance at $g_{\text{eff}} = 6.0$ due to the g_x and g_y transitions in a subpopulation with the preferred value of $E/D \sim 0$. As is also evident from the simulations, when E/D is sufficiently small to account for the resonance positions observed, and when $\sigma E/D$ is sufficiently small that the $g_{\text{eff}} \sim 6.6$ resonance is resolved, there is no significant contribution to the spectrum from the $m_S = 3/2$ doublet, in contrast to the spectra of iron-catecholate complexes. The $g_{\text{eff}} = 4.3$ line must, therefore, arise from a separate Fe(III) species most likely due to complex degradation. The spectrum of **[LFe]OTf** in DMSO was simulated as such, Figure S5, and integration of the simulations indicated that the slightly rhombic species that exhibit the $m_S = 1/2$ resonances accounted for 85 % of the spins (note that much of the spectral intensity is at high field and is virtually undetectable in the derivative spectrum), and the $g_{\text{eff}} = 4.3$ line accounts for about 15 % of the spins.

The EPR simulations for **[LFe]OTf** in DMSO reveal the majority of the spin density is attributable to the six-coordinate **[LFe-DMSO]OTf**. That the high-spin EPR signal indeed arises from solvent-bound species is evidenced by the spectra of **[LFe-pyridine]OTf**, in which pervasive super-hyperfine coupling to nitrogen is present in the majority of the signal. Similar spectral characteristics displayed in σ -donor solvents are consistent with the formation of six-coordinate high-spin Fe(III) complexes. The spectrum of **[LFe-HOMe]OTf** also displays the sharp signal of **[LFe]⁺** ($g = 2.04, 2.02$, and 2.01) consistent with weak coordination. The predominance of the high spin complex despite the binding of strong σ -donors (e.g. DMSO, pyridine) suggests that π -affects are instrumental in determining the spin state of iron in these complexes.

Solvent competition studies

The unique ability of **[LFe]⁺** to bind both substrates (water and nitriles) as well as product (amide) provides the opportunity to directly probe binding preferences for iron in a nitrile hydratase model complex. The iron displays a preference for water over nitrile. Addition of nitriles to **[LFe-OH₂]⁺** at a pH of 3.0 shows no spectroscopically detectable changes in the UV-visible or EPR spectra. Addition of nitriles to **[(LFe)₂OH]⁺** at a pH of 6.1 (*vide infra*) also reveals no detectable changes. In contrast, addition of pH = 6.1 buffered H₂O (100 equivalents H₂O per Fe) to acetonitrile solutions of **[LFe]⁺** in which the nitrile concentration exceeds the water concentration by three orders of magnitude results in significant spectroscopic changes. The charge transfer band in the UV-visible spectrum broadens and red shifts to 602 nm and the EPR spectrum is silent consistent with formation of the dinuclear complex **[(LFe)₂OH]⁺** and no detectable quantities of **[LFe-NCMe]⁺**.

Although the above experiments clearly demonstrate the affinity of iron for water over nitriles, this preference extends even when sub-stoichiometric quantities of water are added to acetonitrile solutions of **[LFe]OTf**. Careful addition of one equivalent of triflic acid to the μ -oxo complex **(LFe)₂O** in acetonitrile leads to a red shifting of the charge transfer band from 560 nm to 603 nm consistent with protonation of the oxo bridge yielding **[(LFe)₂OH]⁺**, which is EPR silent (*vide infra*). Careful addition of a second equivalent of triflic acid protonates the bridging hydroxide generating one equivalent of water for every two equivalents of **[LFe]OTf**. Water coordination is confirmed spectroscopically by a shift in the low energy band from 603 to 628 nm, Figure S7. The EPR spectrum of the reaction mixture, Figure S8, reveals the expected iron-containing products **[LFe-OH₂]⁺** and **[LFe-NCMe]⁺** (in equilibrium with **[LFe]⁺**).

The relative binding affinity of **[LFe]OTf** for amides as compared to nitrile and water was determined through additions of small amounts of DMF to solutions of **[LFe]OTf** in acetonitrile and pH = 5 buffered water, respectively. The UV-visible and EPR spectra of **[LFe-OH₂]⁺** in buffered aqueous solution remain unchanged upon addition of 10 equivalents DMF, denoting the preference for water over DMF. Addition of 10 equivalents DMF to acetonitrile solutions of **[LFe]OTf** is best monitored by EPR, since the UV-visible spectra of **[LFe]OTf** in the two solvents is similar. The EPR shows no trace of the low spin signals of **[LFe-NCMe]⁺** or **[LFe]⁺**. In fact, the spectrum is dominated by the high-spin signal previously observed for **[LFe-DMF]⁺**. Overall, the competition studies reveal that **[LFe]OTf** prefers catalytically relevant ligands in the order H₂O > amide > nitrile.

Aqueous Chemistry

Given the strong preference for water coordination to **[LFe]OTf**, its aqueous chemistry was further explored. As outlined in Scheme 2, three discrete derivatives of **[LFe]OTf** are accessible as a function of pH. At pH < 5.0 (dilute triflic acid) **[LFe-OH₂]⁺** is present as a dark blue solution with bands at 315 and 613 nm. In addition to deprotonation events described below, the coordinate thiolates of **[LFe-OH₂]⁺** may be protonated but only at pH values lower than 3.00 at which point complex degradation occurs. Similar remarkable acid stability of an iron-thiolate has been previously reported for a NHase model complex with deprotonated amide donors in place of the amines found in **[LFe]OTf**.¹⁰

Titration of aqueous solutions of **[LFe-OH₂]⁺** with NaOH first yields the μ -OH complex **[(LFe)₂OH]⁺**. Monitoring of the reaction by UV-visible spectroscopy reveals a blue shift of the low energy band from 613 to 603 nm. At pH = 6.10, the charge transfer band at 603 nm reaches its maximum intensity and the solution develops an intense violet color. The violet solution is EPR silent suggesting a strong anti-ferromagnetic coupling of the two high-spin iron(III) centers. No intermediate is detectable by UV-visible spectroscopy suggesting the mononuclear **LFe-OH** rapidly proceeds to **[(LFe)₂OH]⁺** once generated. Continued titration of **[(LFe)₂OH]⁺** leads to a further blue shifting of the charge transfer band from 603 nm to 560 nm. The resulting burgundy colored solution is characteristic of the previously reported μ -oxo complex **(LFe)₂O**. Due to the poor water solubility of **(LFe)₂O** reliable pK_a values could not be obtained. The titration was repeated in buffered 70:30 water/acetone mixtures yielding similar results, with the exception that **(LFe)₂O** did not precipitate, Figure S6. From this data, the pK_a of the water bound complex **[LFe-OH₂]⁺** is estimated as 5.4 ± 0.1, while that of **[(LFe)₂OH]⁺** is 6.52 ± .05.

Trials for catalytic activity

The catalytic competency of **[LFe]OTf** towards the hydrolysis of nitriles was evaluated by addition of benzonitrile to room-temperature buffered aqueous solutions of **[LFe]OTf** at pH 6.10 and 3.03. At pH 6.10, **[(LFe)₂OH]⁺** is the dominant iron species in solution, while **[LFe-OH₂]⁺** is predominate at pH 3.03. The reaction mixtures were analyzed by gas chromatography for the presence of hydrolysis products benzamide and benzoic acid in organic extracts of the reaction mixture. No trace of hydrolysis products were detected under either set of conditions.

Conclusions and Relevance to Nitrile Hydratase

The complex **[LFe]OTf** has for the first time allowed direct competition studies to evaluate the relative binding affinities of water, nitriles, and amides in a nitrile hydratase model. The observed ligand binding affinity of H₂O > amide > nitrile reveals that water coordination is favored over nitrile binding. Additionally, the preference for amides over nitriles further decreases the probability of a nitrile bound mechanism, since product release would also be problematic.

The only previously reported iron model complex of nitrile hydratase that coordinates nitriles, $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Et},\text{Pr}))]^+$, cannot bind water due to ligand degradation in aqueous environments.²³ Our model binds nitriles with a similar affinity as $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Et},\text{Pr}))]^+$, but it also binds water much more tightly. The coordination of water to $[\text{LFe}]\text{OTf}$ is similar to that observed in the Mascharak model complex, $[\text{Fe}^{\text{III}}(\text{PyPS})]^-$. However, in that case no nitrile binding was observed precluding a direct and definitive comparison with the $[\text{Fe}^{\text{III}}(\text{S}_2^{\text{Me}_2}\text{N}_3(\text{Et},\text{Pr}))]^+$ model.²² The Mascharak aquo complex is low-spin in contrast to our high-spin model suggesting that either spin state shows a strong affinity for water over nitriles. Our results bridge the prior studies and lend credence to a water-bound mechanism.

While both $[\text{LFe}]\text{OTf}$ and $[\text{Fe}^{\text{III}}(\text{PyPS})]^-$ bind H_2O , the former is high-spin and the latter is low-spin.²² The pK_a values reveal that the high-spin $[\text{LFe}-\text{OH}_2]^+$ ($5.4 \pm .1$) is almost ten times as acidic the low-spin $[\text{Fe}^{\text{III}}(\text{PyPS})(\text{H}_2\text{O})]^-$ (6.3).²² Recently we reported that oxygen sensitivity of iron-thiolates is likely spin-state dependent and question if similar effects may also influence the pK_a of coordinated water.¹¹ From a simple electrostatic approach, the relatively smaller ionic radius of low-spin iron(III) should lower the pK_a of bound-water more efficiently than high-spin iron(III). However, this approach ignores π -interactions between the metal center and water/hydroxide that have been shown to be important for low-spin, but not high-spin, Fe(III).⁴³ The π -interactions between the lone pair on HO^- and a filled/nearly filled “ t_{2g} ” orbital on low-spin Fe(III) would destabilize hydroxide coordination and raise the pK_a . Additionally, the π -interactions may promote the nucleophilic character of the coordinated HO^- similar to effects in metal-thiolates that yield nucleophilic sulfur.^{54,55} Overall, the low-spin state may modulate the pK_a while maintaining significant nucleophilic character. In contrast, the more acidic highspin derivatives may lack a sufficiently nucleophilic hydroxide and even undergo further deprotonation to μ -oxo species in model complexes. Further studies to confirm these hypotheses are underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by the National Science Foundation CAREER Award (CHE-0238137) with continued support by (CHE-0749965). MSM thanks the Kentucky Research Challenge Trust Fund for upgrade of our X-ray facilities. EPR studies were supported in part by an NIH P41 grant (EB001980) to the National Biomedical EPR Center.

References

1. Grapperhaus CA, Patra AK, Mashuta MS. *Inorg. Chem* 2002;41:1039–1041. [PubMed: 11874334]
2. Grapperhaus CA, Li M, Patra AK, Poturovic S, Kozlowski PM, Zgierski MZ, Mashuta MS. *Inorg. Chem* 2003;42:4382–4388. [PubMed: 12844310]
3. Mascharak PK. *Coord. Chem. Rev* 2002;225:201–214.
4. Kovacs JA. *Chem. Rev* 2004;104:825–848. [PubMed: 14871143]
5. Lugo-Mas P, Dey A, Xu L, Davin SD, Benedict J, Kaminsky W, Hodgson KO, Hedman B, Solomon EI, Kovacs JA. *J. Am. Chem. Soc* 2006;128:11211–11221. [PubMed: 16925440]
6. Lee CM, Hsieh CH, Dutta A, Lee GH, Liaw WF. *J. Am. Chem. Soc* 2003;125:11492–11493. [PubMed: 13129340]
7. Kung I, Schweitzer D, Shearer J, Taylor WD, Jackson HL, Lovell S, Kovacs JA. *J. Am. Chem. Soc* 2000;122:8299–8300.
8. Harrop TC, Mascharak PK. *Acc. Chem. Res* 2004;37:253–260. [PubMed: 15096062]
9. Galardon E, Giorgi M, Artaud I. *Chem. Comm* 2004:286–287. [PubMed: 14740040]

10. Artaud I, Chatel S, Chauvin AS, Bonnet D, Kopf MA, Leduc P. *Coord. Chem. Rev* 1999;192:577–586.
11. O'Toole MG, Kreso M, Kozlowski PM, Mashuta MS, Grapperhaus CA. *J. Bio. Inorg. Chem.* 2008In Press.
12. Asano Y, Tani Y, Yamada H. *Agr. Biol. Chem. Tokyo* 1980;44:2251–2252.
13. Endo I, Nakasako M, Nagashima S, Dohmae N, Tsujimura M, Takio K, Odaka M, Yohda M, Kamiya NJ. *Inorg. Biochem* 1999;74:22–22.
14. Endo I, Odaka M, Yohda M. *Trends Biotechnol* 1999;17:244–249. [PubMed: 10354562]
15. Kobayashi M, Shimizu S. *Nat. Biotechnol* 1998;16:733–736. [PubMed: 9702770]
16. Nelson MJ, Jin HY, Turner IM, Grove G, Scarrow RC, Brennan BA, Que LJ. *Am. Chem. Soc* 1991;113:7072–7073.
17. Yamada H, Kobayashi M. *Biosci Biotech Bioch* 1996;60:1391–1400.
18. Shigehiro S, Nakasako M, Dohmae N, Tsujimura M, Tokoi K, Odaka M, Yohda M, Kamiya N, Endo I. *Nat Struct Biol* 1998;5:347–351. [PubMed: 9586994]
19. Hopmann KH, Himo F. *Eur. J. Inorg. Chem* 2008:1406–1412.
20. Hopmann KH, Guo JD, Himo F. *Inorg. Chem* 2007;46:4850–4856. [PubMed: 17497847]
21. Greene SN, Richards NG. *J. Inorg. Chem* 2006;45:17–36.
22. Noveron JC, Olmstead MM, Mascharak PK. *J. Am. Chem. Soc* 2001;123:3247–3259. [PubMed: 11457060]
23. Shearer J, Jackson HL, Schweitzer D, Rittenberg DK, Leavy TM, Kaminsky W, Scarrow RC, Kovacs JA. *J. Am. Chem. Soc* 2002;124:11417–11428. [PubMed: 12236756]
24. Armarego, WLF.; Perrin, DD. *Purification of Laboratory Chemicals*. Vol. 4th ed. Oxford: Butterworth Heinemann; 1996.
25. Gordon AJ, Ford RA. *The Chemist's Companion. A Handbook of Practical Data, Techniques, and References* (1st ed). 1973
26. Shriver, DF.; Drezdon, MA. *The Manipulation of Air-Sensitive Compounds*. Vol. 2nd ed. New York: Wiley; 1986.
27. Woodhouse ME, Lewis FD, Marks TJ. *J. Am. Chem. Soc* 1982;104:5586–5594.
28. Evans DF. *Proc. Chem. Soc* 1959;115:2003–2005.
29. Ellison JJ, Nienstedt A, Shoner SC, Barnhart D, Cowen JA, Kovacs JA. *J. Am. Chem. Soc* 1998;120:5691–5700.
30. Albert, A.; Serjeant, EP. *Ionization Constants of Acids and Bases*. Methuen, New York: Wiley; 1962.
31. Nilges, MJ. Ph.D. Thesis. Urbana: University of Illinois; 1979.
32. Hanson, GR.; Gates, KE.; Noble, CJ.; Mitchell, A.; Benson, S.; Griffin, M.; Burrage, K. In *EPR of Free Radicals in Solids*. In: Shiotani, M.; Lund, A., editors. *Trends in Methods and Applications*. Dordrecht, The Netherlands: Kluwer Press; 2003.
33. SMART. Madison, WI: Bruker Advanced X-ray Solutions, Inc.; 2005.
34. SAINT. Madison, WI: Bruker Advanced X-ray Solutions, Inc.; 2003.
35. SADABS. Germany: Gottingen; 2003.
36. SHELXTL 6.14. Program Library for Structure Solution and Molecular graphics, Bruker Advanced X-Ray Solutions. Masdison, WI: 2000.
37. Sheldrick GM. *Acta Crystallogr., Sect. A: Found. Crystallogr* 2008;A64:112–122.
38. SHELXL-97. Program for the Refinement of Crystal Structures. Gottingen, Germany: University of Gottingen; 1997.
39. Farrugia LJ. *J. Appl. Crystallogr* 1997;30:565.
40. Dey A, Jenney FE, Adams MWW, Johnson MK, Hodgson KO, Hedman B, Solomon EI. *J. Am. Chem. Soc* 2007;129:12418–12431. [PubMed: 17887751]
41. Mosseri S, Mialocq JC, Perly B, Hambright PJ. *Phys. Chem* 1991;95:4659–4663.
42. Evans DR, Mathur RS, Heerwegh K, Reed CA, Xie Z. *Angew. Chem., Int. Ed. Engl* 1997;36:1335–1337.
43. Evans DR, Reed RA. *J. Am. Chem. Soc* 2000;122:4660–4667.

44. Kennepohl P, Neese F, Schweitzer D, Jackson HL, Kovacs JA, Solomon EI. *Inorg. Chem* 2005;44:1826–1836. [PubMed: 15762709]
45. Noveron JC, Herradora R, Olmstead MM, Mascharak PK. *Inorg. Chim. Acta* 1999;285:269–276.
46. Griffith JS. *Proc. R. Soc. London, A* 1956;235:23–26.
47. Taylor CP. *S. Biochim. Biophys. Acta* 1977;491:137–149.
48. Patra AK, Bill E, Bothe E, Chlopek K, Neese F, Weyhermuller T, Stobie K, Ward MD, McCleverty JA, Wieghardt K. *Inorg. Chem* 2006;45:7877–7890. [PubMed: 16961381]
49. Hopmann KH, Guo JD, Himo F. *Inorg. Chem* 2007;46:4850–4856. [PubMed: 17497847]
50. Rodriguez M, Morenster-Badarau, Cesario M, Guilhem J, Keita B, Nadjio L. *Inorg. Chem* 1996;35:7804–7810.
51. Strautmann JBH, George SD, Bothe E, Bill E, Weyhermuller T, Stammel A, Bogge H, Glaser T. *Inorg. Chem* 2008;47:6804–6824. [PubMed: 18582030]
52. Weisser JT, Nilges MJ, Sever MJ, Wilker JJ. *Inorg. Chem* 2006;45:7736–7747. [PubMed: 16961365]
53. Copik AJ, Waterson S, Swierczek SI, Bennett B, Holz RC. *Inorg. Chem* 2005;44:1160–1162. [PubMed: 15732944]
54. Mullins CS, Grapperhaus CA, Kozlowski PM. *J Biol Inorg Chem* 2006;11:617–625. [PubMed: 16724228]
55. Ashby MT, Enemark JH, Lichtenberger DL. *Inorg. Chem* 1988;27:191–197.

Figure 1.
Representation of the active site of nitrile hydratase.

Figure 2.
UV/visible spectrum of the synthesis of $[(\text{LFe})_2\text{OH}]\text{OTf}$ (3 mM) in acetonitrile.

Figure 3. ORTEP view of $[(\text{LFe})_2\text{OH}]^+$ showing 30% probability displacement ellipsoids. Calculated H atoms, solvent and triflate counter ion have been omitted.

Figure 4.

EPR spectrum (77 K) of **[LFe]OTf** in dichloromethane with simulation. Experimental Parameters: Microwave power = 6.3 mW, Modulation Amplitude = 5.35G. Simulation parameters: $g_1 = 2.06$, $g_2 = 2.03$, $g_3 = 2.02$, $W_1 = 21.80$, $W_2 = 39.87$, $W_3 = 21.23$.

Figure 5.
Variable temperature (40 °C to -43 °C) UV/visible spectra of [LFe]OTf (3 mM) in acetonitrile.
Arrows denote changes in the spectra as the temperature is lowered.

Figure 6.

EPR spectra of **[LFe]OTf** in acetonitrile (A) with simulation (B) and **[LFe]OTf** in benzonitrile (C) with simulation (D). Experimental parameters: (A) microwave Power = 1.5 mW, modulation amplitude = 8.09 G; (C) microwave power = 1.99 mW, modulation amplitude = 6.00 G. Simulation parameters: (B) for **[LFe]⁺** $g_1 = 2.04$, $g_2 = 2.01$, $g_3 = 2.02$, $W_1 = 19.40$, $W_2 = 28.15$, $W_3 = 25.80$ and **[LFe-NCMe]⁺** $g_1 = 2.27$, $g_2 = 2.18$, $g_3 = 1.98$, $W_1 = 17.95$, $W_2 = 91.21$, $W_3 = 33.17$; (D) for **[LFe]⁺**: $g_1 = 2.06$, $g_2 = 2.03$, $g_3 = 2.03$, $W_1 = 21.73$, $W_2 = 40.05$, $W_3 = 21.08$ and **[LFe-NCPh]⁺**: $g_1 = 2.28$, $g_2 = 2.18$, $g_3 = 2.00$, $W_1 = 64.78$, $W_2 = 26.50$, $W_3 = 46.33$.

Figure 7. EPR spectra of [LFe]OTf in various solvents. All spectra recorded at 20 K except DMF (77 K).

Figure 8.

Calculated High-Spin Spectra. High-spin spectra were calculated assuming $S = 5/2$, $g_{\text{iso}} = 1.995$, and (A) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.165$, $\sigma E/D = 0.08$; (B) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.165$, $\sigma E/D = 0$; (C) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.125$, $\sigma E/D = 0$; (D) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.085$, $\sigma E/D = 0$; (E) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0$; (F) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.010$; (G) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.015$; (H) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.020$; (I) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.025$; (J) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.035$; (K) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.045$; (L) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.100$; (M) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.045$, $\sigma E/D = 0.320$; (N) $D = 0.45 \text{ cm}^{-1}$, $E/D = 0.130$, $\sigma E/D = 0.320$.

Scheme 1.

Scheme 2.

Table 1
Crystal data and structure refinement for [(LFe)₂OH]OTf.

Identification code	[(LFe) ₂ OH]OTf
Empirical formula	C ₂₉ H ₅₇ F ₃ Fe ₂ N ₄ O ₄ S ₇ · 0.75(CH ₃ OH)
Formula weight	942.94
Temperature	100(2) K
Wavelength	0.71073 Å
Crystal system	Monoclinic
Space group	<i>Ia</i>
Unit cell dimensions	a = 15.472(6) Å b = 10.629(4) Å c = 25.894(13) Å β = 95.172(5)°
Volume	4241(3) Å ³
Z	4
Density (calculated)	1.477 Mg/m ³
Absorption coefficient	1.082 mm ⁻¹
F(000)	1982
Crystal size	0.29 × 0.22 × 0.05 mm ³
Theta range for data collection	2.64 to 28.14°
Crystal color, habit	dark purple trapezoidal prism
Index ranges	-20 ≤ h ≤ 20 -14 ≤ k ≤ 13 -33 ≤ l ≤ 33
Reflections collected	18021
Independent reflections	9283 [R(int) = 0.0273]
Completeness to theta = 28.14°	95.7 %
Absorption correction	SADABS
Max. and min. transmission	0.948 and 0.744
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	9283 / 4 / 468
Goodness-of-fit on F ²	1.08
Final R indices [I > 2σ(I)]	R1 = 0.0388, wR2 = 0.0965
R indices (all data)	R1 = 0.0420, wR2 = 0.0994
Absolute structure parameter	0.030(12)
Largest diff. peak and hole	1.092 and -0.326 e.Å ⁻³

Table 2
 Selected Bond Distances (Å) and Bond Angles (deg) for [(LFe)₂OH]OTf

Fe1-S1	2.3004(12)	Fe1-N1	2.263(3)
Fe1-S2	2.3262(13)	Fe1-N2	2.242(3)
Fe1-S3	2.5385(14)	Fe1-O1	2.002(3)
Fe2-S4	2.3250(12)	Fe2-N3	2.244(3)
Fe2-S5	2.2961(12)	Fe2-N4	2.258(3)
Fe2-S6	2.5565(13)	Fe2-O1	1.998(3)
		O1-H1 ₀	0.841(19)
N1-Fe1-N2	79.10(11)	N1-Fe1-S1	83.56(8)
N2-Fe1-S2	84.11(8)	N1-Fe1-S2	96.11(8)
N2-Fe1-S3	80.71(8)	S1-Fe1-S2	102.22(4)
O1-Fe1-S2	93.91(8)	Fe1-O1-Fe2	166.71(14)
N3-Fe2-N4	78.86(11)	N3-Fe2-S4	84.26(9)
N3-Fe2-S6	81.10(8)	N4-Fe2-S5	83.95(8)
N4-Fe2-S6	80.96(8)	S4-Fe2-S5	100.95(4)
Fe1-O1-H1 ₀	96(3)	Fe2-O1-H1 ₀	98(3)