Published in final edited form as:

Cancer Res. 2009 October 1; 69(19): 7603–7611. doi:10.1158/0008-5472.CAN-09-2066.

Atypical Protein Kinase C α is Required for Bronchioalveolar Stem Cell Expansion and Lung Tumorigenesis

Roderick P. Regala¹, Rebecca K. Davis¹, Alyssa Kunz¹, Andras Khor², Michael Leitges^{3,*}, and Alan P. Fields^{1,*}

¹Department of Cancer Biology, Mayo Clinic Comprehensive Cancer Center, Jacksonville, Florida, United States

²Department of Pathology, Mayo Clinic Comprehensive Cancer Center, Jacksonville, Florida, United States

³Biotechnology Centre of Oslo, University of Oslo, N-0317, Oslo, Norway

Abstract

Protein kinase C α (PKC α) is an oncogene required for maintenance of the transformed phenotype of non-small cell lung cancer (NSCLC) cells. However, the role of PKC α in lung tumor development has not been investigated. To address this question, we established a mouse model in which oncogenic *Kras*^{G12D} is activated by Cre-mediated recombination in the lung with or without simultaneous genetic loss of the mouse PKC α gene, *Prkci*. Genetic loss of *Prkci* dramatically inhibits *Kras*-initiated hyperplasia and subsequent lung tumor formation *in vivo*. This effect correlates with a defect in the ability of *Prkci*-deficient bronchioalveolar stem cells (BASCs) to undergo *Kras*-mediated expansion and morphological transformation *in vitro* and *in vivo*. Furthermore, the small molecule PKC α inhibitor aurothiomalate inhibits *Kras*-mediated BASC expansion and lung tumor growth *in vivo*. Thus, *Prkci* is required for oncogene-induced expansion and transformation of tumor-initiating lung stem cells. Furthermore, aurothiomalate is an effective anti-tumor agent that targets the tumor-initiating stem cell niche *in vivo*. These data have important implications for PKC α as a therapeutic target and for the clinical use of aurothiomalate for lung cancer treatment.

Keywords

lung tumor initiation; aurothiomalate; lung cancer stem cells; *Kras* transformation; transgenic mice

INTRODUCTION

Lung cancer is the leading cause of cancer death in the United States (1). NSCLC is the most prevalent type of lung cancer accounting for ~80% of cases (2). Despite recent treatment advances the five year survival rate of NSCLC patients remains only 15%. Thus, there is a dire need to better understand the key molecular events driving lung tumorigenesis that could lead to more effective means of prevention, diagnosis, prognosis and treatment.

Activating mutations in *Kras* are among the most common molecular changes found in NSCLC tumors (3,4) and several transgenic mouse models have demonstrated the critical role of

*To whom correspondence should be addressed: Alan P. Fields, Ph.D. Department of Cancer Biology Mayo Clinic Comprehensive Cancer Center Griffin Cancer Research Building, Rm. 212 4500 San Pablo Road Jacksonville, Florida 32224 fields.alan@mayo.edu (904) 953-6109 (office) (904) 953-0277 (fax). A.P.F. and M.L. are both communicating authors

oncogenic *Kras* in lung tumor initiation and maintenance. Transgenic mice carrying a conditional oncogenic *Kras*^{G12D} allele activated by Cre-recombinase (*LSL-Kras* mice) develop epithelial hyperplasias, lung adenomas and adenocarcinomas (5,6). A second murine lung cancer model (the *Kras*^{LA} mouse) is based on sporadic *Kras* mutation using a “hit-and-run” gene targeting strategy (7). *Kras*^{LA} mice carry a latent activation oncogenic *Kras*^{G12D} allele activated by spontaneous somatic recombination. *Kras*^{LA} mice invariably develop numerous lung tumors pathologically similar to the human disease (7).

Bronchio-alveolar stem cells (BASCs) are the probable cells of origin of *Kras*-induced lung tumors in these mouse models (8). BASCs function in lung homeostasis and possess characteristics of regional lung stem cells (8). BASCs reside in bronchio-alveolar duct junctions (BADJs), a niche at the boundary between the terminal bronchioles (TBs) and the alveolar space (AS). BASCs express the hematopoietic stem cell marker Sca-1, Clara cell specific protein (CCSP) and surfactant protein C (SPC), but not the hematopoietic and endothelial cell markers CD45 and Pecam (8). BASCs exhibit stem-like self-renewal and the ability to differentiate into Clara cells and alveolar type II (AT2) cells *in vitro* (8). Moreover, BASCs expand in response to *Kras* transformation *in vitro*, and within the BADJ *in vivo*, eventually giving rise to adenomas and carcinomas in the *LSL-Kras* mouse model (8).

We recently demonstrated that PKC ϵ is an oncogene in NSCLC (9,10). PKC ϵ expression is elevated in NSCLC tumors and cell lines, and PKC ϵ is required for the transformed phenotype of NSCLC cells harboring oncogenic *Kras* mutation (9). PKC ϵ activates a novel oncogenic PKC ϵ -Par6-Rac1 signaling axis that is necessary for the maintenance of the transformed phenotype of human NSCLC cells (11). We recently identified aurothiomalate as a potent and selective inhibitor of oncogenic PKC ϵ signaling (12,13). Aurothiomalate inhibits NSCLC cell transformation *in vitro* and tumorigenicity *in vivo* by disrupting the interaction between PKC ϵ and Par6, thereby blocking the PKC ϵ -Par6-Rac1 signaling axis (12,13).

Although PKC ϵ is important in the maintenance of the transformed phenotype of NSCLC cells harboring *Kras* mutation (11,14), the role of PKC ϵ in *Kras*-mediated lung tumor development has not been explored. Here we assess the role of the mouse PKC ϵ gene *Prkci*, in *Kras* mediated lung tumorigenesis using a novel bi-transgenic mouse model harboring conditional *LSL-Kras* and *Prkci* knock out (*Prkci*^{fl/fl}) alleles. *LSL-Kras/Prkci*^{fl/fl} mice exhibit a significant impairment in lung hyperplasias and lung tumors formation, indicating that *Prkci* is necessary for early events in *Kras*-mediated lung tumorigenesis. The effect of *Prkci* deficiency on tumorigenesis correlates with a defect in *Kras*-mediated BASC expansion in the absence of *Prkci*. Furthermore, the PKC ϵ inhibitor aurothiomalate inhibits *Kras*-mediated BASC expansion and transformation *in vitro* and *in vivo*. Our results demonstrate that *Prkci* plays a requisite role in BASC transformation and identify aurothiomalate as a promising therapeutic agent that can target the lung cancer stem cell niche *in vivo*.

MATERIALS AND METHODS

Generation and AdCre treatment of *LSL-Kras/Prkci*^{fl/fl} mice

LSL-Kras mice (B6.129), generated as previously described (5), were mated to mice harboring a floxed *Prkci* allele (*Prkci*^{fl} mice) to generate bi-transgenic *LSL-Kras/Prkci*^{fl/fl} mice. Non-transgenic littermates served as controls in all experiments. Mice (6–8 weeks of age) were administered AdCre (1.5×10^8 IFU/mL) by intratracheal instillation in two 50 μ L aliquots as described previously (15). All animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) of Mayo Clinic.

Histology and immunohistochemistry

Mice were sacrificed, exsanguinated and the lungs perfused with 10% buffered formalin through the right ventricle. The trachea was intubated and instilled with an additional 3 mL of 10% buffered formalin. The lungs were removed intact and fixed overnight in 10% buffered formalin. For histological analysis, lungs were embedded in paraffin, serially sectioned (5 μ m) and stained with hematoxylin and eosin. Immunohistochemical analysis was performed as described (10). Mouse PKC ϵ was detected using a PKC ϵ antibody (Santa Cruz Biotechnology) and visualized using the Envision Plus Dual Labeled Polymer Kit following the manufacturer's instructions (DAKO). In some experiments, the antibody was incubated overnight at 4°C with a 200-fold molar excess of PKC ϵ peptide (Santa Cruz Biotechnology) prior to use in immunohistochemistry to confirm antibody specificity. Slide images were captured and analyzed using the ScanScope scanner and ImageScope software (Aperio Technologies).

BASC isolation and culture

Lung epithelial cells were isolated as described (16). Red blood cells were lysed in RBC lysis buffer (StemCell Technologies) and BASCs isolated using the EasySep® immunomagnetic cell selection procedure (Stem Cell Technologies). Briefly, CD45^{pos} Pecam^{pos} cells were selected out using primary biotinylated antibodies anti-CD45 (BD Pharmingen) and anti-Pecam (BD Pharmingen). CD45^{neg} Pecam^{neg} cells were incubated for 15 minutes at room temperature with FITC-conjugated anti-CD34 (BD Pharmingen) and Sca1-PE labeling reagent (Stem Cell Technologies). Sca1^{pos} CD34^{pos} CD45^{neg} Pecam^{neg} BASCs were selected using the EasySep® FITC and PE selection kits. BASCs were resuspended in BEGM (Lonza, without hydrocortisone) containing 10ng/mL keratinocyte growth factor (PeproTech) and 5% charcoal-stripped fetal bovine serum (FBS) and plated at equal densities on Matrigel-coated (BD Biosciences) tissue culture wells. Cultures were infected with AdCre (7.5×10^7 IFU/mL) the day after plating and medium was changed every 2 days thereafter. Brightfield images of BASC colonies were captured on an Olympus IX71 inverted microscope and colony size was determined using Image-Pro Plus 6.3 software (Media Cybernetics).

Immunofluorescence of BASCs in Matrigel culture was performed as described (17) using the following antibodies: rabbit anti-CCSP (Upstate), goat anti-SP-C (Santa Cruz Biotechnology), rabbit anti-Ki67 (Abcam), Alexa Fluor 488 donkey anti-rabbit IgG, Alexa Fluor 568 donkey anti-goat IgG, and Alexa Fluor 594 goat anti-rabbit IgG (Molecular Probes). Samples were mounted using ProLong Gold anti-fade reagent with DAPI (Molecular Probes) and observed on a Zeiss 510 LSM confocal laser microscope. Images were analyzed with Adobe Photoshop software (Adobe Systems).

BASC colonies were released from Matrigel culture using BD cell recovery solution (BD Biosciences) and total RNA subjected to quantitative PCR as described (18). RNA quantity and integrity were measured using a NanoDrop® ND-1000 spectrophotometer.

Kras and Prkci allele status

Kras allele status in AdCre-treated BASC cultures and adenomas microdissected from *LSL-Kras/Prkci^{fl/fl}* mice was determined by PCR as described (5,19). *Prkci^{fl}* allele status was determined using the following PCR primers: F1 = 5'-CACCAACGGGTTTGCTATCT-3', R1 = 5'-CCAGCAAGACAAAACACCAA-3', R2 = 5'-ATTACAGCAGGGCAAACCTGC-3'.

Treatment of *Kras^{LA2}* mice with aurothiomalate

Three-week-old *Kras^{LA2}* mice were given daily intraperitoneal injections of aurothiomalate (Myochrysine®; Taylor Pharmaceuticals) at 60 mg/kg/day or vehicle control (0.9% sodium

chloride solution, Sigma-Aldrich) for the indicated time period. Tumors were analyzed as described above.

Detection and analysis of BASCs in mouse lung tissues

BASCs were detected in formalin fixed paraffin embedded (FFPE) mouse lung tissues as described previously (20). Lung tissue sections were stained using goat anti-SPC (Santa Cruz Biotechnology) and rabbit anti-CCSP (Upstate) antibodies diluted in 1% BSA (Sigma-Aldrich) at 4°C overnight. The slides were washed thoroughly with PBS and then incubated with Alexa Fluor 488 donkey anti-rabbit IgG and Alexa Fluor 568 donkey anti-goat IgG secondary antibodies (Molecular Probes) diluted in 1% BSA for 1 hour at room temperature. After washing with PBS, the samples were mounted using ProLong Gold antifade reagent with DAPI. Immunofluorescence images were captured on an Olympus BX51 fluorescent microscope using DPController and DPManager software (Olympus), and processed in Adobe Photoshop.

Statistical analysis

The Cochran-Armitage test was used to compare BASC number and distribution *in vivo* using StatsDirect 2.6.1. The Student's *t* test and one-way ANOVA statistical analyses were performed using SigmaStat 3.5. A *P* value of less than 0.05 was considered statistically significant.

RESULTS

Prkci is required for Kras-mediated lung tumorigenesis

To determine whether PKC ϵ is involved in lung tumor formation, we employed the *LSL-Kras* mouse model in which lung tumors are initiated by AdCre-mediated activation of a conditional oncogenic *Kras* allele (5). Since PKC ϵ is overexpressed in primary human NSCLC tumors (10), we assessed whether mouse PKC ϵ protein is also highly expressed in lung tumors from *LSL-Kras* mice. Immunohistochemical analysis revealed light staining for mouse PKC ϵ in normal lung epithelial cells with more intense staining in lung tumor cells from *LSL-Kras* mice (Figure 1A, **right panel**). Thus, mouse PKC ϵ is highly expressed in *Kras*-initiated lung tumors in mice.

We have shown that PKC ϵ is required for transformed growth and invasion of human NSCLC cells *in vitro* and tumorigenicity *in vivo* (9,10). To assess the role of PKC ϵ in oncogenic *Kras*-initiated lung tumorigenesis, we crossed mice harboring a conditional knockout *Prkci* allele (*Prkci^{fl/fl}*) with *LSL-Kras* mice to generate bi-transgenic *LSL-Kras/Prkci^{fl/fl}* mice. In *LSL-Kras/Prkci^{fl/fl}* mice, intratracheal instillation of AdCre leads to simultaneous Cre-mediated activation of the oncogenic *Kras* allele and inactivation of the *Prkci^{fl/fl}* alleles in the same target cells within the lung. Twelve weeks following AdCre administration, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice were assessed for lung hyperplasias and lung tumors (Figure 1B). As expected, *LSL-Kras* mice developed numerous tumors visible on the surface of the lung (Figure 1B, **upper left panel, arrows**). In contrast, *LSL-Kras/Prkci^{fl/fl}* mice exhibited few visible tumors (Figure 1B, **lower left panel**). Pathological examination revealed that *LSL-Kras* mice develop multiple focal adenomas often surrounding bronchioles (Figure 1B, **upper middle and right panels, arrows**) whereas *LSL-Kras/Prkci^{fl/fl}* mice exhibit largely normal lung morphology and fewer lesions (Figure 1B, **lower middle and right panels**). Analysis also revealed a significant reduction in both hyperplasias (Figure 1C) and adenomas (Figure 1D) in *LSL-Kras/Prkci^{fl/fl}* mice when compared to *LSL-Kras* mice. Thus, genetic deletion of *Prkci* inhibits hyperplasia and subsequent tumor formation, indicating a role for *Prkci* in an early step of *Kras*-mediated lung tumorigenesis.

Since *LSL-Kras/Prkci^{fl/fl}* mice formed some lung tumors, albeit in significantly reduced numbers, we determined the status of the *Kras* and *Prkci* alleles in tumors formed in these

mice. Genomic DNA from tumors microdissected from *LSL-Kras/Prkci^{fl/fl}* mice was analyzed for *Kras* and *Prkci* allele status by PCR (Figure 2A and B). As expected, the *LSL-Kras* allele had undergone Cre-mediated recombination (Figure 2A, **top panel**) with concomitant loss of the unrecombined *Kras* allele (Figure 2A, **bottom panel**) in all of the tumors analyzed. In contrast, each tumor contained a recombined (deleted) *Prkci* allele (Figure 2B, **top panel**), and an unrecombined *Prkci* allele (Figure 2B, **bottom panel**), indicating incomplete deletion of the conditional *Prkci^{fl}* alleles. The presence of unrecombined *Prkci* was not due to contamination of the tumor samples by normal lung as indicated by the absence of the unrecombined *LSL-Kras* allele in the same tumor samples (Figure 2A, **bottom panel**). Furthermore, immunohistochemistry confirmed expression of mouse PKC α protein in these tumors (Figure 2C). Our data indicate that lung tumors in *LSL-Kras/Prkci^{fl/fl}* mice arise due to incomplete recombination of the conditional *Prkci^{fl}* alleles. Thus, *Prkci* is critical for *Kras*-induced lung tumorigenesis, and only *Kras*-transformed cells that escape complete inactivation of both *Prkci^{fl}* alleles are able to progress in tumorigenesis.

Prkci is required for Kras-mediated expansion of the bronchio-alveolar stem cell (BASC) niche in vivo

Our data indicate that *Prkci* is required for a very early event(s) in *Kras*-mediated oncogenesis *in vivo*. *Kras* mediated lung tumorigenesis in the *LSL-Kras* mouse model is thought to be initiated by clonal expansion of BASCs, the putative cell of origin of *Kras*-initiated lung tumorigenesis (8). Therefore, we next assessed whether *Prkci*-deficiency affects the expansion of BASCs induced by *Kras in vivo*. BASCs are identified in tissue sections at the BADJ by dual immunofluorescence for SPC and CCSP (Figure 3A). AdCre-treated *LSL-Kras* mice exhibit frequent clusters of SPC-, CCSP-double-positive BASCs at BADJs indicative of BASC expansion (Figure 3A, **upper panels**). In contrast, *LSL-Kras/Prkci^{fl/fl}* mice exhibit significantly fewer BASCs at the BADJs (Figure 3A, **lower panels**). Quantitative analysis of sections from AdCre-treated non-transgenic (Ntg), *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice revealed a significant increase in the percentage of terminal bronchioles (TBs) containing one or more BASCs, and in the average number of BASCs/TB in *LSL-Kras* mice when compared to Ntg mice (Figure 3B). *LSL-Kras/Prkci^{fl/fl}* mice exhibited a significant inhibition in *Kras*-mediated BASC expansion, such that the number and distribution of BASCs in these mice was indistinguishable from that of Ntg mice (Figure 3B). Furthermore, AdCre-treated *Prkci^{fl/fl}* mice exhibited no difference in BASC number or distribution compared to Ntg mice, indicating that *Prkci* deficiency does not overtly affect the BASC niche in the absence of oncogenic *Kras* (Figure 3B). Thus *Prkci* is necessary for *Kras*-mediated BASC expansion *in vivo*, linking the effect of *Prkci*-deficiency on BASC expansion with lung tumorigenesis. Furthermore, the inhibitory effect of *Prkci* deficiency on lung tumorigenesis is not due to a decrease in the number of BASCs in *LSL-Kras/Prkci^{fl/fl}* mice.

Oncogenic Kras induces expression of PKC α and Prkci-dependent proliferative signaling in BASCs in vitro

To further explore the role of *Prkci* in BASC expansion, we isolated and established BASCs from Ntg, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice in primary culture using established procedures (8). Isolated lung epithelial cell populations consisted of a mixture of SPC-single positive AT2 cells (Figure 4A, **left panel, AT**), CCSP-single positive Clara cells (Figure 4A, **left panel, arrows**), and SPC-, CCSP-double positive BASCs (Figure 4A, **left panel, arrowheads**). This BASC population was enriched based on positive staining for the stem cell markers Sca1 and CD34. Consistent with previous reports, >95% of the cells remaining after this selection stained double positive for the BASC markers SPC and CCSP (Figure 4A, **right panel**). Purified BASCs from Ntg, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice were treated with AdCre, and efficient recombination of the *LSL-Kras* and *Prkci^{fl}* alleles was confirmed by PCR (Figure 4B). Consistent with our finding that mouse PKC α protein expression is elevated in

lung tumors from *LSL-Kras* mice, AdCre treatment of BASCs from *LSL-Kras* mice induced a significant increase in PKC α mRNA abundance, whereas BASCs from *LSL-Kras/Prkci^{fl/fl}* mice, as expected, expressed no detectable PKC α mRNA (Figure 4C). We previously identified matrix metalloproteinase 10 (MMP10) as a transcriptional target of the oncogenic PKC α -Par6-Rac1 signaling axis (11), and MMP10 has also been identified as a transcriptional target of oncogenic *Kras* (21,22). Therefore, we assessed MMP10 expression in BASCs from *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice by QPCR. MMP10 mRNA was induced in BASCs from *LSL-Kras* mice but not *LSL-Kras/Prkci^{fl/fl}* mice after AdCre treatment (Figure 4D). Thus oncogenic *Kras* activates oncogenic PKC α signaling in BASCs, and this signaling is inhibited in BASCs from *LSL-Kras/Prkci^{fl/fl}* mice.

Oncogenic *Kras*-mediated BASC expansion and morphological transformation requires *Prkci*

Since BASCs expand and initiate *Kras*-mediated tumors *in vivo* (8,23), we assessed BASCs from *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice for proliferative expansion and morphological transformation after induction of oncogenic *Kras in vitro* (Figure 5). BASCs from Ntg mice treated with AdCre form colonies of organized spherical structures when plated in three dimensional Matrigel culture (Figure 5A, **left panel**). In contrast, BASCs from *LSL-Kras* mice grow as large amorphic, disorganized colonies (Figure 5A, **middle panel**). Interestingly, BASCs from *LSL-Kras/Prkci^{fl/fl}* mice form spherical colonies similar in size and morphology to those from Ntg mice (Figure 5A, **right panel**). Quantitative analysis revealed that BASCs from *LSL-Kras* mice form structures of larger average diameter (Figure 5B) and more amorphic shape (Figure 5C) than BASCs from either Ntg or *LSL-Kras/Prkci^{fl/fl}* mice. Thus the change in colony size and morphology observed in *LSL-Kras* BASCs requires both *Kras* and *Prkci*. Immunofluorescence confocal microscopy revealed that BASCs from Ntg and *LSL-Kras/Prkci^{fl/fl}* mice form hollow spheres consisting of a single layer of epithelial cells with a low proliferative index; in contrast, BASCs from *LSL-Kras* mice form solid masses of highly proliferative cells (Figure 5D, **left panel**). Quantitative analysis demonstrated a significant increase in proliferation of *LSL-Kras* BASCs (% of cells staining positive for Ki67) when compared to Ntg BASCs and that this increase was not seen in *LSL-Kras/Prkci^{fl/fl}* BASCs (Figure 5D, **right panel**). Thus, oncogenic *Kras* induces *Prkci*-dependent proliferative expansion and morphological changes associated with cellular transformation of BASCs *in vitro*.

BASCs exhibit self-renewal and the ability to differentiate into bronchiolar Clara cells and AT2 cells (8). Therefore, we assessed whether the increase in proliferation observed in *LSL-Kras* BASC cultures *in vitro* was due to expansion of undifferentiated BASCs, or alternatively induction of lineage-specific differentiation of BASC into single positive Clara and/or AT2 cells. Confocal immunofluorescence microscopy revealed that BASC colonies from Ntg, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice are made up pre-dominantly of SPC-, CCSP-double positive staining BASCs (Supplemental Figure 1), indicating that *Kras* induces *Prkci*-dependent BASC expansion.

Aurothiomalate blocks *Kras*-mediated, *Prkci*-dependent BASC expansion and morphological transformation *in vitro*

We recently identified aurothiomalate as a highly selective small molecule inhibitor of oncogenic PKC α / λ signaling that exhibits potent anti-tumor activity against human NSCLC cells (12,13). Since *Prkci* is necessary for *Kras*-mediated BASC expansion and morphological transformation *in vivo* and *in vitro*, we assessed whether aurothiomalate inhibits the effects of *Kras* on BASCs. BASCs from *LSL-Kras* mice were transduced with AdCre and plated in three dimensional Matrigel culture in the presence of 20 μ M aurothiomalate or diluent (saline) (Figure 6A, **left and middle panels**). As expected, saline-treated BASCs form large amorphic

colonies; in contrast aurothiomalate-treated cells form small spherical colonies similar to those formed by BASCs from Ntg and *LSL-Kras/Prkci^{fl/fl}* mice (Figure 6A, **right panel**). Analysis revealed that saline-treated BASCs form significantly larger colonies than aurothiomalate-treated BASCs (Figure 6B, **left panel**), that is associated with an increased proliferative index compared to aurothiomalate-treated BASCs (Figure 6B, **right panel**). Thus, aurothiomalate, like genetic loss of *Prkci*, blocks *Kras*-mediated proliferative expansion and morphological transformation of BASCs.

Aurothiomalate inhibits *Kras*-mediated BASC expansion and lung tumorigenesis *in vivo*

We next assessed the effect of aurothiomalate on BASC expansion and tumor formation in *Kras^{LA2}* mice harboring a latent oncogenic *Kras* allele activated by spontaneous recombination *in vivo* (7). Ntg and *Kras^{LA2}* mice were treated with either aurothiomalate (60 mg/kg/day) or diluent (saline) for three weeks at which time mice were sacrificed and assessed for BASC number and distribution as described above. Aurothiomalate has no demonstrable effect on BASC number or distribution in Ntg mice (Figure 6C) consistent with our finding that *Prkci*-deficiency does not overtly affect BASCs in the absence of oncogenic *Kras in vivo*. In contrast, saline-treated *Kras^{LA2}* mice exhibit a significant increase in BASCs/TB that is inhibited by aurothiomalate (Figure 6C). Consistent with the inhibitory effect of aurothiomalate on BASC expansion, aurothiomalate-treated *Kras^{LA2}* mice exhibited a decrease in tumor growth when compared to saline-treated mice after a six week course of aurothiomalate treatment (Figure 6D). Therefore, like genetic loss of *Prkci*, aurothiomalate blocks *Kras*-mediated BASC expansion and inhibits subsequent lung tumor growth *in vivo*.

DISCUSSION

The PKC ϵ oncogene plays a requisite role in the maintenance of the transformed phenotype of NSCLC cells *in vitro* and *in vivo* (10). However, the present study is the first to assess the role of PKC ϵ in lung tumor formation. Our data firmly establish that genetic disruption of *Prkci* has a profound inhibitory effect on oncogenic *Kras*-mediated lung tumor development. Although *Prkci*-deficient mice still develop lung hyperplasias and lung tumors in response to oncogenic *Kras*, albeit at a dramatically reduced incidence, molecular analysis of these tumors demonstrates that these lesions invariably retain an unrecombined *Prkci* allele and express mouse PKC ϵ protein. These data argue that *Prkci* is absolutely required for a very early step in *Kras*-mediated lung tumorigenesis, such that only the occasional *Kras*-transformed cell that escapes complete recombination of both *Prkci^{fl}* alleles (and retains PKC ϵ expression) is competent to progress in tumorigenesis.

The inhibitory effect of *Prkci* deletion on tumorigenesis is reflected in a severe defect in the ability of *Prkci*-deficient BASCs to expand and undergo *Kras*-mediated transformation *in vitro*. Thus *Prkci* is necessary for the first identifiable step in *Kras*-mediated tumorigenesis involving expansion and transformation of the putative lung cancer stem cell in the mouse. Our data provide important new insight into a key pathway involved in BASC expansion in response to an oncogenic stimulus; namely the oncogenic PKC ϵ signaling axis. Our data place *Prkci* among a small cadre of genes known to play a critical role in oncogene-induced BASC expansion, including the polycomb group member Bmi1 (23), the p38 MAPK gene (24), the cyclin-dependent kinase inhibitor p27 (25) and phosphatidylinositol 3-kinase (PI3K) (20).

By virtue of its role in BASC expansion, PKC ϵ is an attractive target for therapeutic intervention in lung cancer. Aurothiomalate, a highly selective inhibitor of oncogenic PKC ϵ / λ signaling (12,13), potently blocks *Kras*-mediated expansion and morphological transformation of BASCs *in vitro* and *in vivo*, and inhibits lung tumor growth *in vivo*. Our data indicate that aurothiomalate suppresses oncogene-mediated tumor induction, at least in part, through inhibition of the expansion of lung cancer stem cells. These data have important implications

for the use of aurothiomalate in cancer treatment. A population of cells within human lung tumors has been identified that exhibits properties of lung cancer stem cells similar to those exhibited by BASCs (26,27). These lung cancer stem cells are capable of self-renewal, serial passage as tumors in immunocompromised mice, and differentiation into highly proliferative tumor cells (26,27). Based on these properties, lung cancer stem cells are thought to be responsible for initiation and long-term maintenance of human lung tumors, much as BASCs are in the mouse lung. Given their central role in tumorigenesis, lung cancer stem cells must be therapeutically targeted to achieve lasting anti-tumor responses. Unfortunately, lung cancer stem cells exhibit intrinsic resistance to commonly used cytotoxic agents (26,27). Our present results demonstrate that aurothiomalate exhibits potent anti-proliferative activity toward the tumor stem cell niche in a relevant pre-clinical lung cancer model. Future studies will be required to assess whether aurothiomalate has similar anti-proliferative effects on human lung cancer stem cells isolated from primary human lung tumors.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

The authors thank Erin Donovan and Suzie Randle for assistance with animal husbandry procedures, and Brandy Edenfield for immunohistochemical analysis of lung tissues. This work was supported by grants from the National Cancer Institute (CA084136-12), the V-Foundation and the American Lung Association/LUNGevity to A.P.F. R.P.R. is recipient of a Ruth A. Kirschstein Post-doctoral Fellowship Award (CA115160) from the National Cancer Institute.

REFERENCES

1. Jemal A, Siegel R, Ward E, et al. Cancer statistics, 2008. *CA Cancer J Clin* 2008;58:71–96. [PubMed: 18287387]
2. Schiller JH. Current standards of care in small-cell and non-small-cell lung cancer. *Oncology* 2001;61 (Suppl 1):3–13. [PubMed: 11598409]
3. Gazdar AF, Minna JD. Molecular detection of early lung cancer. *J Natl Cancer Inst* 1999;91:299–301. [PubMed: 10050857]
4. Salgia R, Skarin AT. Molecular abnormalities in lung cancer. *J Clin Oncol* 1998;16:1207–17. [PubMed: 9508209]
5. Jackson EL, Willis N, Mercer K, et al. Analysis of lung tumor initiation and progression using conditional expression of oncogenic Kras. *Genes Dev* 2001;15:3243–8. [PubMed: 11751630]
6. Meuwissen R, Linn SC, van der Valk M, Mooi WJ, Berns A. Mouse model for lung tumorigenesis through Cre/lox controlled sporadic activation of the Kras oncogene. *Oncogene* 2001;20:6551–8. [PubMed: 11641780]
7. Johnson L, Mercer K, Greenbaum D, et al. Somatic activation of the Kras oncogene causes early onset lung cancer in mice. *Nature* 2001;410:1111–6. [PubMed: 11323676]
8. Kim CF, Jackson EL, Woolfenden AE, et al. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell* 2005;121:823–35. [PubMed: 15960971]
9. Regala RP, Weems C, Jamieson L, Copland JA, Thompson EA, Fields AP. Atypical protein kinase Ciota plays a critical role in human lung cancer cell growth and tumorigenicity. *J Biol Chem* 2005;280:31109–15. [PubMed: 15994303]
10. Regala RP, Weems C, Jamieson L, et al. Atypical protein kinase C iota is an oncogene in human non-small cell lung cancer. *Cancer Res* 2005;65:8905–11. [PubMed: 16204062]
11. Frederick LA, Matthews JA, Jamieson L, et al. Matrix metalloproteinase-10 is a critical effector of protein kinase Ciota-Par6alpha-mediated lung cancer. *Oncogene* 2008;27:4841–53. [PubMed: 18427549]

12. Erdogan E, Lamark T, Stallings-Mann M, et al. Aurothiomalate inhibits transformed growth by targeting the PB1 domain of atypical protein kinase Ciota. *J Biol Chem* 2006;281:28450–9. [PubMed: 16861740]
13. Stallings-Mann M, Jamieson L, Regala RP, Weems C, Murray NR, Fields AP. A novel small-molecule inhibitor of protein kinase Ciota blocks transformed growth of non- small-cell lung cancer cells. *Cancer Res* 2006;66:1767–74. [PubMed: 16452237]
14. Banan A, Fields JZ, Farhadi A, Talmage DA, Zhang L, Keshavarzian A. The beta 1 isoform of protein kinase C mediates the protective effects of epidermal growth factor on the dynamic assembly of F-actin cytoskeleton and normalization of calcium homeostasis in human colonic cells. *J Pharmacol Exp Ther* 2002;301:852–66. [PubMed: 12023512]
15. Fasbender ALJ, Walters RW, Moninger TO, Zabner J, Welsh MJ. Incorporation of adenovirus in calcium phosphate precipitates enhances gene transfer to airway epithelia in vitro and in vivo. *Journal of Clinical Investigation* 1998;102:184–93. [PubMed: 9649572]
16. Rice WR, Konkright JJ, Na CL, Ikegami M, Shannon JM, Weaver TE. Maintenance of the mouse type II cell phenotype in vitro. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L256–64. [PubMed: 12114186]
17. Debnath J, Muthuswamy SK, Brugge JS. Morphogenesis and oncogenesis of MCF-10A mammary epithelial acini grown in three-dimensional basement membrane cultures. *Methods* 2003;30:256–68. [PubMed: 12798140]
18. Regala RP, Thompson EA, Fields AP. Atypical protein kinase C iota expression and aurothiomalate sensitivity in human lung cancer cells. *Cancer Res* 2008;68:5888–95. [PubMed: 18632643]
19. Kissil JL, Walmsley MJ, Hanlon L, et al. Requirement for Rac1 in a Kras induced lung cancer in the mouse. *Cancer Res* 2007;67:8089–94. [PubMed: 17804720]
20. Yang Y, Iwanaga K, Raso MG, et al. Phosphatidylinositol 3-kinase mediates bronchioalveolar stem cell expansion in mouse models of oncogenic Kras-induced lung cancer. *PLoS ONE* 2008;3:e2220. [PubMed: 18493606]
21. Tchernitsa OI, Sers C, Zuber J, et al. Transcriptional basis of KRAS oncogene-mediated cellular transformation in ovarian epithelial cells. *Oncogene* 2004;23:4536–55. [PubMed: 15064704]
22. Zuber J, Tchernitsa OI, Hinemann B, et al. A genome-wide survey of RAS transformation targets. *Nat Genet* 2000;24:144–52. [PubMed: 10655059]
23. Dovey JS, Zacharek SJ, Kim CF, Lees JA. Bmi1 is critical for lung tumorigenesis and bronchioalveolar stem cell expansion. *Proc Natl Acad Sci U S A* 2008;105:11857–62. [PubMed: 18697930]
24. Ventura JJ, Tenbaum S, Perdiguer E, et al. p38alpha MAP kinase is essential in lung stem and progenitor cell proliferation and differentiation. *Nat Genet* 2007;39:750–8. [PubMed: 17468755]
25. Besson A, Hwang HC, Cicero S, et al. Discovery of an oncogenic activity in p27Kip1 that causes stem cell expansion and a multiple tumor phenotype. *Genes Dev* 2007;21:1731–46. [PubMed: 17626791]
26. Eramo A, Lotti F, Sette G, et al. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ* 2008;15:504–14. [PubMed: 18049477]
27. Chen YC, Hsu HS, Chen YW, et al. Oct-4 expression maintained cancer stem- like properties in lung cancer-derived CD133-positive cells. *PLoS ONE* 2008;3:e2637. [PubMed: 18612434]

Figure 1. *Prkci* is necessary for *Kras*-mediated lung tumorigenesis

LSL-Kras and *LSL-Kras/Prkci^{fl/fl}* mice were infected with AdCre, harvested and processed as described in **Materials and Methods**. (A) Staining of a *LSL-Kras* lung tumor for mouse PKC ι protein. (B) Histopathology of *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice. **upper left:** multiple lesions are evident on the surface of *LSL-Kras* mouse lungs (**arrows**). **upper middle:** *LSL-Kras* mice develop numerous focal adenomas (**arrows**) **upper right:** *LSL-Kras* adenomas often surround bronchioles. **lower left:** *LSL-Kras/Prkci^{fl/fl}* mice exhibit a smooth lung surface devoid of visible lesions. **lower middle:** *LSL-Kras/Prkci^{fl/fl}* lungs exhibit fewer and smaller lesions than *LSL-Kras* mice. **lower right:** *LSL-Kras/Prkci^{fl/fl}* lung exhibit largely normal lung morphology. (C) *LSL-Kras/Prkci^{fl/fl}* mice exhibit reduced lung hyperplasias. Mean $\pm$ SE (n=6); * $P < 0.05$. (D) *LSL-Kras/Prkci^{fl/fl}* mice exhibit reduced lung tumor formation. Mean $\pm$ SE (n=12); * $P = 0.007$.

Figure 2. Lung tumors in *LSL-Kras/Prkci^{fl/fl}* mice arise due to incomplete recombination of the *Prkci^{fl}* alleles

Genomic DNA from microdissected *LSL-Kras/Prkci^{fl/fl}* tumors was analyzed for *Kras* (A) and *Prkci* (B) allele status. Each tumor expresses the recombined (A, **top**) but not the unrecombined *Kras* allele (A, **bottom**). Normal lung epithelium (N) and tumor (T) DNA from a *Kras* mouse serve as controls for *Kras* allele recombination. All *LSL-Kras/Prkci^{fl/fl}* tumors analyzed contained a recombined *Prkci* allele (B, **top panel**) as well as an unrecombined *Prkci* allele (B, **bottom panel**), indicating incomplete deletion of the conditional *Prkci^{fl}* alleles. (R) recombined and (U) unrecombined *Prkci* allele controls. (C) *LSL-Kras/Prkci^{fl/fl}* tumors express mouse PKC_λ protein. Adjacent sections were stained with H&E (**left**), PKC_λ antibody (**middle**), or with PKC_λ antibody in the presence of PKC_λ peptide to verify specificity of staining (**right**).

Figure 3. *Prkci* is necessary for *Kras*-mediated expansion of BASCs *in vivo*

LSL-Kras and *LSL-Kras/Prkci^{fl/fl}* mice were harvested 12 weeks after intratracheal instillation of AdCre. Lung tissue sections were subjected to immunofluorescence as described in **Materials and Methods**. (A) Expansion of CCSP (green) and SPC (red) double positive BASCs (outlined) is evident within the terminal bronchioles (TB) of *LSL-Kras* mice (upper panels). Clara cells within the bronchioles stain positive for CCSP but negative for SPC. AS = alveolar space. BASC expansion is not observed in *LSL-Kras/Prkci^{fl/fl}* mice (lower panels). (B) Quantitative analysis of terminal bronchioles for BASCs in lungs from Ntg, *Prkci^{fl/fl}*, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice. Results are expressed as % terminal bronchioles $\pm$ SE. Statistical analysis revealed a significant increase in BASC number and BASC/TB in *LSL-Kras* mice compared to Ntg, *Prkci^{fl/fl}* or *LSL-Kras/Prkci^{fl/fl}* mice ($p < 0.0001$). No statistically significant difference in BASC number or distribution was observed between Ntg, *Prkci^{fl/fl}* and *LSL-Kras/Prkci^{fl/fl}* mice.

Figure 4. *Kras* induces PKC expression and activates oncogenic *Prkci*-mediated signaling in BASCs *in vitro*

(A) Immunofluorescence staining of isolated lung epithelial cells (*left*) and purified BASCs (*right*) from *LSL-Kras* mice for CCSP (*green*), SPC (*red*), and DAPI (*blue*). SPC-positive AT2 cells (AT), CCSP-positive Clara cells (arrows) and SPC-, CCSP-double positive BASCs (arrowheads). (B) *LSL-Kras* and *Prkci*^{fl/fl} allele status in BASCs isolated from Ntg, *LSL-Kras* and *LSL-Kras/Prkci*^{fl/fl} mice in the presence and absence of AdCre. The conditional *Kras* and *Prkci* alleles are efficiently recombined in the presence of AdCre. (C) *Kras* induces PKC γ mRNA expression in BASCs from *LSL-Kras* but not *LSL-Kras/Prkci*^{fl/fl} mice. (D) *Kras* induces MMP10 mRNA in BASCs from *LSL-Kras* but not *LSL-Kras/Prkci*^{fl/fl} mice. Data in C and D represent the mean $\pm$ SE of triplicate determinations. Results are representative of four independent experiments with similar results. * indicates a statistically significant difference from *LSL-Kras* in the absence of Cre, p<0.05; ** indicates a statistically significant difference from *LSL-Kras*, p<0.05.

Figure 5. *Kras* induces *Prkci*-dependent BASC expansion and morphological transformation *in vitro*

BASCs from Ntg, *LSL-Kras* and *LSL-Kras/Prkci^{fl/fl}* mice were treated with AdCre and plated in three dimensional Matrigel culture. (A) Photomicrographs of cultures after four days. BASC colonies were analyzed for diameter (B) and morphology (C). Data in (B) represent the mean $\pm$ SE and are presented as % Ntg control. Data in (C) represent the % of colonies with spherical (filled bar) or amorphic (open bar) morphology. *indicates a statistically significant difference from Ntg; ** indicates a statistically significant difference from *LSL-Kras*; $p < 0.0001$. There was no statistically significant difference in colony diameter or morphology between Ntg and *LSL-Kras/Prkci^{fl/fl}* colonies. (D) Confocal microscopy of BASC colonies stained for DAPI and Ki67 (left). Quantitative analysis of Ki67 staining (right). Data represent the mean $\pm$ SE expressed as the % of Ki67 positive cells; *indicates a statistically significant difference from Ntg; ** indicates a statistically significant difference from *LSL-Kras*; $p < 0.05$.

Figure 6. Aurothiomalate inhibits *Kras*-mediated BASC expansion and morphological transformation *in vitro* and tumor growth *in vivo*

(A) Morphology of BASCs from *LSL-Kras* mice in the presence of 20 μ M aurothiomalate (middle) or saline (left). Analysis of BASC colony morphology in the presence and absence of aurothiomalate (right). * indicates a statistically significant difference from BASCs cultured in the absence of aurothiomalate; $p < 0.001$. (B) Colony diameter (left) and Ki67 staining (right) of BASCs in the presence and absence of aurothiomalate. Data represent the mean $\pm$ SE; * indicates a statistically significant difference from BASCs in the absence of aurothiomalate; $p = 0.01$. (C) Aurothiomalate blocks *Kras*-mediated BASC expansion *in vivo*. Ntg and *Kras*^{LA2} mice were treated with aurothiomalate or saline for 3 weeks and lungs were

analyzed for BASC number and distribution as described in Figure 3. Statistical analysis revealed a significant increase in BASC number and in the mean number of BASC/TB in saline-treated *Kras^{LA2}* mice when compared to aurothiomalate-treated *Kras^{LA2}* mice, Ntg mice treated with aurothiomalate, or Ntg mice treated with saline ($p < 0.0001$). No statistically significant differences in BASC number or distribution were observed between saline-treated Ntg, aurothiomalate-treated Ntg and aurothiomalate-treated *Kras^{LA2}* mice. **(D)** Aurothiomalate inhibits lung tumor growth in *Kras^{LA2}* mice. Data represent the mean $\pm$ SE fold-change in average tumor size after six weeks in the presence of aurothiomalate or saline. *indicates a statistically significant difference from saline-treated *Kras^{LA2}* mice; $p = 0.003$.