Published in final edited form as:

Trends Immunol. 2009 June ; 30(6): 263–270. doi:10.1016/j.it.2009.04.001.

A_{2B} adenosine receptors in immunity and inflammation

György Haskó^{1,2}, Balázs Csóka¹, Zoltán H. Németh^{1,3}, E. Sylvester Vizi², and Pál Pacher⁴

¹ Department of Surgery, University of Medicine and Dentistry of New Jersey-New Jersey Medical School, Newark, NJ 07103, USA

² Department of Pharmacology, Institute of Experimental Medicine, Hungarian Academy of Sciences, H-1083 Budapest, Szigony u. 43, Hungary

³ Department of Surgery, Morristown Memorial Hospital, Morristown, NJ 07960, USA

⁴ Section on Oxidative Stress Tissue Injury, Laboratory of Physiological Studies, National Institute on Alcohol Abuse and Alcoholism, 12420 Parklawn Dr., MSC-8115, Bethesda, MD 20892-8115, USA

Abstract

A_{2B} adenosine receptors are increasingly recognized as important orchestrators of inflammation. A_{2B} receptor activation promotes the inflammatory response of mast cells, epithelial cells, smooth muscle cells and fibroblasts, thereby contributing to the pathophysiology of asthma and colitis. A_{2B} receptor stimulation limits endothelial cell inflammatory responses and permeability and suppresses macrophage activation thereby preventing tissue injury after episodes of hypoxia and ischemia. A_{2B} receptor stimulation also promotes the production of angiogenic cytokines by endothelial cells, mast cells and dendritic cells, aiding granuloma tissue formation and inflammatory resolution, but can also contribute to tumor growth. A_{2B} receptors are, thus, potentially important pharmacological targets in treating immune system dysfunction and inflammation.

Introduction

Adenosine is a primordial regulator of multiple physiological processes and its modulatory actions have been demonstrated in organisms as diverse as *Drosophila* [1], the common mussel *Mytilus edulis* [2], the protozoan *Leishmania mexicana* [3] and mammals such as *Mus musculus* [4,5] and humans [4,5]. Adenosine is an extracellular purine nucleoside signaling molecule, which governs cell and tissue function both in health and disease. Adenosine is formed after the degradation of its precursor, adenosine 5' triphosphate (ATP), a process which can take place both extra- and intracellularly. ATP, a predominantly intracellular molecule, is released from the cell after stressful and injurious events, and is degraded to adenosine via a cascade of ectonucleotidases, including CD39 (nucleoside triphosphate diphosphorylase [NTPDase]) and CD73 (5'-ectonucleotidase [Ecto5'NTase]) [6]. Intracellular ATP metabolized to adenosine is exported from the cell through nucleoside transporters [7]. Cells of the immune system including neutrophils, mast cells, endothelial cells, regulatory T cells and platelets have been appreciated as the most prodigious sources of extracellular adenosine [8]. In addition to serving as a source for adenosine release, immune cells are also among the most widely studied cell types targeted by the regulatory influences of adenosine [9]. Adenosine imparts its immune regulatory actions by binding to and activating four G protein-coupled cell surface receptors, A₁, A_{2A}, A_{2B} and A₃ [7,9]. Although A₁, A_{2A} and A₃ receptors are high-affinity receptors that are activated by physiological extracellular adenosine

concentrations in the submicromolar range, A_{2B} receptors are activated by micromolar levels of adenosine, concentrations that are achieved in tissues that experience ischemia, trauma, inflammation or other types of stressful insults. This low affinity of A_{2B} receptors for the endogenous ligand adenosine and other selective agonists had, until a few years ago, impeded exploration of the role of this receptor subtype in regulating immune function. The recent availability of sophisticated pharmacological and molecular tools has however dramatically invigorated research on the role of A_{2B} receptors in regulating immunity. Here, we review recent advances in understanding the role of A_{2B} receptors in modulating inflammation and immunity, which might provide a basis for the utilization of these receptors as pharmacological targets in treating immune system dysfunction and inflammation.

Distribution of A_{2B} receptors

A new mouse model incorporating targeted deletion of A_{2B} receptors and replacement of exon 1 with a reporter gene has allowed *in vivo* determination of which cell types and tissues express A_{2B} receptors in the mouse [10]. A_{2B} receptors have been found in all tested organs, including spleen, lung, colon and kidney, and in all of these organs the primary site of expression was the vasculature. Smooth muscle cells, endothelial cells and macrophages exhibit a high level of expression [10], and A_{2B} receptors have also been detected on colonic epithelial cells [11]. Earlier studies employing isolated primary cells or cell lines, however, clearly indicated that other cell types, such as mast cells [12,13], lymphocytes [14,15] dendritic cells [16,17] and neutrophils [15] all express A_{2B} receptors.

Regulation of A_{2B} receptor expression

The signaling capacity of A_{2B} receptors depends on several factors, which include A_{2B} receptor density on the cell surface and A_{2B} receptor coupling to downstream intracellular signaling pathways. A_{2B} receptor expression on the cell surface is a highly regulated and dynamic process, which is influenced by metabolic, inflammatory and hormonal clues from the environment and by adenosine itself. Hypoxia is an important stimulus for the upregulation of A_{2B} receptor expression and it has been shown to induce A_{2B} receptor transcription via the oxygen-regulated transcription factor HIF-1 α (hypoxia inducible factor) in endothelial cells [18]. In addition, hypoxia has been documented to increase A_{2B} receptor expression on dendritic cells [19], smooth muscle cells [20] and fibroblasts [21]. Factors present in an inflammatory environment, such as the bacterial product lipopolysaccharide (LPS) [22], the pro-inflammatory cytokines TNF- α [23–25], IL-1 β [23], IFN- γ [23,26], free radicals [25] and the endogenous agonist adenosine [27] have all been shown to increase the expression of A_{2B} receptors. This occurs by varying mechanisms; for example, TNF- α [24,25] and IFN- γ [26] upregulate receptor expression by increasing its transcription, whereas adenosine induces A_{2B} receptor expression on the cell membrane by rapidly recruiting it from its intracellular depot in intestinal epithelial cells via a mechanism involving the docking proteins vesicle-associated membrane protein (VAMP)-2, soluble N-ethylmaleimide-sensitive factor attachment protein (SNAP-23) and syntaxin-3 [28]. Once in the membrane, the A_{2B} receptor associates with the Na⁺/H⁺ exchanger-3 (NHE-3) kinase regulatory protein E3KARP and ezrin, which not only tether the receptor to the membrane but also serve to stabilize it in a signaling complex, facilitating activation of downstream signaling targets, most importantly protein kinase A (PKA) [27].

Signaling pathways of A_{2B} receptors

A_{2B} receptors couple to a variety of intracellular signaling pathways, the best studied of which are Gs-mediated signaling to PKA, resulting in increased cAMP levels and Gq-mediated activation of phospholipase C (PLC) leading to increased protein kinase C (PKC) activation

and elevations in intracellular Ca^{2+} levels [29]. These two pathways are often activated simultaneously, which enables additional levels of fine-tuning of cellular processes. In addition to these two most important signaling pathways, $\text{A}_{2\text{B}}$ receptor has been shown to activate mitogen activated protein (MAP) kinase signaling, in addition to membrane ion channels, the mechanisms of which are poorly understood and sometimes involve the $\beta\gamma$ subunit of G proteins [29]. Because the intracellular pathways used by adenosine receptors are largely cell type specific, we detail each particular pathway when discussing specific immune cell types.

The pharmacology of $\text{A}_{2\text{B}}$ receptors

The pharmacological characterization of $\text{A}_{2\text{B}}$ receptors has lagged behind that of A_1 , $\text{A}_{2\text{A}}$ and A_3 receptors because until recently, potent and selective $\text{A}_{2\text{B}}$ receptor ligands had not been available. NECA (5'-N-ethylcarboxamido-adenosine) is the most widely used $\text{A}_{2\text{B}}$ receptor agonist; however, it is neither selective nor potent. Nevertheless, in those rare cellular systems in which $\text{A}_{2\text{B}}$ receptors are the only adenosine receptors expressed, a role for $\text{A}_{2\text{B}}$ receptors in regulating a particular response can be deduced based on the efficacy of NECA and inefficacy of selective A_1 , $\text{A}_{2\text{A}}$ and A_3 receptor agonists. When $\text{A}_{2\text{B}}$ receptors are co-expressed with other adenosine receptors, $\text{A}_{2\text{B}}$ receptor-mediated responses can be distinguished based on the efficacy of NECA in a manner antagonizable with selective $\text{A}_{2\text{B}}$ receptor antagonists. MRS 1754 (N-(4-cyanophenyl)-2-[4-(2,3,6,7-tetrahydro-2,6-dioxo-1,3-dipropyl-1H-purin-8-yl)phenoxy]-acetamide) and CVT-6883 (3-ethyl-1-propyl-8-[1-(3-trifluoromethylbenzyl)-1H-pyrazol-4-yl]-3,7-dihydropurine-2,6-dione) are recently developed, selective and potent $\text{A}_{2\text{B}}$ receptor antagonists that are very useful in characterizing $\text{A}_{2\text{B}}$ receptors. Enprofylline, IPDX (3-isobutyl-8-pyrrolidinoxanthine) and alloxazine are relatively selective albeit not particularly potent $\text{A}_{2\text{B}}$ receptor antagonists. Theophylline also blocks $\text{A}_{2\text{B}}$ receptors; however, it is not selective and can antagonize all four types of adenosine receptor. Molecular approaches such as the use of $\text{A}_{2\text{B}}$ receptor knockout (KO) mice or siRNA-mediated knock-down of $\text{A}_{2\text{B}}$ receptors in conjunction with pharmacological studies, are able to paint a more accurate picture of the role of $\text{A}_{2\text{B}}$ receptors in physiological processes.

Regulation of cell function by $\text{A}_{2\text{B}}$ receptors

Mast cells

Mast cell activation has long been recognized as a crucial factor in the pathophysiology of asthma and other allergic diseases [30]. Mast cells are also increasingly appreciated as a crucial cell type in initiating and maintaining inflammatory and immune responses, owing to their ability to rapidly release biogenic amines, proinflammatory lipid molecules and cytokines [31]. Adenosine receptors on mast cells have become of considerable interest as therapeutic targets for asthma, based on the observation that inhaled adenosine or its precursor AMP provokes bronchoconstriction in individuals suffering from asthma, furthermore this bronchoconstrictive response can be prevented using mast cell membrane stabilizers or histamine receptor antagonists [32]. The demonstration that enprofylline and theophylline, two clinically used anti-asthmatic agents, block adenosine-induced IL-8 production by the human mast cell line HMC-1 via targeting of $\text{A}_{2\text{B}}$ receptors [33] has aroused interest in studying the role of $\text{A}_{2\text{B}}$ receptors on mast cell function (Figure 1). $\text{A}_{2\text{B}}$ receptor activation evokes the release of this proinflammatory chemokine by a mechanism that involves the MAP kinase p38 [34], PKC [34] and the third intracellular loop of the receptor and not the $\text{G}\alpha$ or $\text{G}\beta$ proteins [35].

Mast cells are among the earliest producers of IL-4 during a Th2 type immune response; however, until recently the stimuli eliciting IL-4 release by mast cells were unknown. It has now been demonstrated [36] that $\text{A}_{2\text{B}}$ receptor stimulation using NECA induces IL-4 production by HMC-1 cells, indicating that adenosine might provide an early signal resulting

in commitment to a Th2 type immune response. The IL-4 response of mast cells stimulated via A_{2B} receptors requires nuclear factor of activated T cells (NFAT)-mediated transcriptional activation of the IL-4 gene via PLC β and elevations in intracellular Ca²⁺ levels [37] (Figure 1). Although Gq proteins seem indispensable for the stimulatory effect of adenosine on NFAT transcriptional activity, simultaneous activation of Gs proteins potentiates the IL-4 response to adenosine by leading to increased intracellular cAMP concentrations and up-regulated NFAT protein levels (Figure 1) [36,37]. In addition, co-culturing B lymphocytes with A_{2B} receptor-stimulated mast cells triggers the former's IgE production, an effect proposed to be secondary to increased secretion of both IL-4 and another Th2 cytokine (IL-13) by the mast cells [36]. The proinflammatory effect of A_{2B} receptor activation that had first been observed in HMC-1 cells was recently also confirmed using murine bone-marrow derived mast cells [13]. Mast cells isolated from wild type (WT) but not A_{2B} receptor KO mice released IL-13 and VEGF (vascular endothelial growth factor) in response to extracellular adenosine. These studies also demonstrated that A_{2B} triggering does not directly affect antigen-induced degranulation of mast cells because adenosine was able to potentiate antigen-induced histamine release by both WT and KO cells. Taken together, A_{2B} receptor activation increases proinflammatory cytokine production by mast cells (Figure 1), which suggests that this mechanism contributes to the increased inflammatory response that is seen in asthmatic patients exposed to inhaled adenosine.

Endothelial cells

Both innate and adaptive immune responses are dependent on the migration of leukocytes across endothelial cells [38]. Endothelial cells orchestrate leukocyte transmigration by producing inflammatory chemokines and cytokines, and upregulating adhesion molecules on their surface. In the resolution phase of inflammation, endothelial cells proliferate to form new blood vessels, a phase governed by angiogenic molecules (Figure 2).

Endothelial cells actively metabolize adenosine and have a large capacity to both take up and release adenosine. A_{2B} receptors are expressed on the surface of endothelial cells [39] and regulate every aspect of endothelial inflammatory processes. *In vivo* studies utilizing A_{2B} KO mice have shown that A_{2B} receptors inhibit the expression of the adhesion molecules ICAM-1 and E-selectin, which results in decreased leukocyte rolling and adhesion [10]. A_{2B} receptor KO mice exposed to hypoxia exhibit increased neutrophil infiltration into tissues revealing an inhibitory role for A_{2B} receptors in neutrophil transmigration *in vivo* [15]. Similarly, pharmacologic studies indicate that neutrophil A_{2B} receptors contribute to the decreased adhesion and transmigration of these cells to endothelial cells [40].

Leukocyte transendothelial migration has the potential to disturb vascular barrier function, resulting in fluid loss and edema. Adenosine-mediated activation of A_{2B} receptors decreases endothelial paracellular permeability during leukocyte transmigration thereby resealing the endothelium and preventing edema formation and fluid loss [15,41] (Figure 2). This protective effect of adenosine is potentiated in a hypoxic environment, which is due to both the increased availability of extracellular adenosine and upregulation of A_{2B} receptors [18], processes that are coordinated by the oxygen-sensitive transcription factor HIF-1 α [42,43]. In addition to the promoter of the A_{2B} receptor, HIF-1 α transactivates the promoter of CD73 [44] leading to further generation of extracellular adenosine. Moreover, HIF-1 α represses the promoter of the equilibrative nucleoside transporter (ENT), which causes increased adenosine availability by limiting the cellular uptake of adenosine [45], and together creates a positive feedback loop of A_{2B} signaling.

Once the acute inflammation has subsided, in-growth of vascular tissue containing capillary loops ensues [46]. The development of this granulation tissue is driven predominantly by the angiogenic factors produced, in a large part, by the endothelial cells themselves. A_{2B} receptors

have a key role in promoting this angiogenic response because activation of A_{2B} receptors by adenosine increases endothelial cell proliferation, chemotaxis and capillary tube formation [47,48]. In this context, A_{2B} receptor activation has been shown to stimulate the release of VEGF, basic fibroblast growth factor and insulin-like growth factor-I by human microvascular endothelial cells, which primarily express A_{2B} receptors [39]. A_{2B} receptors signal through Gq proteins to induce transcriptional VEGF upregulation; however, this process does not require HIF-1 α . Resting human umbilical vein endothelial cells, however, do not respond to NECA with increased VEGF release because they do not express A_{2B} receptors [20]. Exposure of human umbilical vein endothelial cells to hypoxia increases their A_{2B} receptor expression, which upon stimulation promotes the release of VEGF by the cells [20].

Taken together, endothelial A_{2B} receptors seem to prevent endothelial cell-mediated inflammatory events and increase endothelial barrier function at the early stages of tissue injury, and at later phases, A_{2B} receptor activation promotes angiogenesis (Figure 2).

Antigen-presenting cells: macrophages and dendritic cells

Adenosine receptors have been recognized as important regulators of macrophage function; however, A_{2B} receptors have been overshadowed by A_{2A} receptors as the primary adenosine receptors shaping macrophage function [49,50]. In this context, the inhibitory effect of adenosine on TNF- α , IL-12 and MIP-1 α release by Toll-like receptor (TLR)-activated macrophages is mediated primarily by A_{2A} receptors [51,52]. Because the A_{2B} receptor antagonist MRS 1754 reverses the adenosine-induced suppression of TNF- α production only in the absence and not in the presence of functional A_{2A} receptors, it was [53] proposed that A_{2B} receptors become operational only when their effect is not masked by A_{2A} receptors. In addition, stimulation of A_{2B} but not A_{2A} receptors augments TLR-induced IL-10 production by RAW264.7 cells, a macrophage-like cell line that predominantly expresses A_{2B} receptors [54]. In spite of the most important role of A_{2A} receptors in regulating TLR-induced cytokine production, a recent study utilizing macrophages isolated from A_{2B} KO mice showed that adenosine elicits IL-6 production from resting macrophages through A_{2B} receptors [55].

In contrast to cytokine production, A_{2B} receptors down-regulate IFN- γ -induced MHC class II expression and inducible nitric oxide synthase production (iNOS) [26]. IFN- γ upregulates A_{2B} receptor expression on macrophages resulting in an increased responsiveness of macrophages to the stimulatory effects of NECA, but not other agonists, as measured by intracellular cAMP [26]. This increased cAMP in turn down-regulates both MHC class II and iNOS expression at both the mRNA and protein levels. Similarly, A_{2B} receptor activation inhibits monocyte colony stimulating factor (M-CSF)-induced macrophage proliferation by upregulating the cyclin-dependent kinase inhibitor p27^{Kip-1} [56]. This occurs in a PKA-dependent manner and results in the growth arrest of cells at the G₁ phase of the cell cycle [56].

Recent studies employing A_{2B} KO mice and selective A_{2B} receptor agonists and antagonists have defined a novel role for A_{2B} receptors in regulating dendritic cell function. Human monocytes differentiated towards dendritic cells using IL-4 and GM-CSF in the presence of adenosine develop into a cell type which resembles dendritic cells morphologically but is different phenotypically [16]. These ‘adenosine-differentiated’ cells represent an intermediate phase between monocytes and dendritic cells and are impaired in their ability to induce T-cell proliferation and IFN- γ production [16]. However, these intermediate cells are highly angiogenic and secrete VEGF and IL-8, and are also anti-inflammatory because of their robust production of IL-10 and TGF- β . Furthermore, because these cells also highly express the tolerance-inducing enzyme indoleamine 2,3-dioxygenase (IDO), they can render T cells anergic. Finally, A_{2B} KO mice demonstrated that adenosine triggering of this receptor inhibits IL-12 p70 production by TLR-activated bone-marrow-derived dendritic cells [17]. In

conclusion, A_{2B} receptor activation reshapes the function of antigen-presenting cells in a way that produces an anti-inflammatory and tolerance-inducing phenotype.

Intestinal epithelial cells

Intestinal epithelial cells, which were once considered a simple physical barrier to the luminal environment, are now known to be integral to both the discrimination of pathogenic and commensal bacteria and the subsequent regulation of immune responses in the intestinal microenvironment [57]. A_{2B} receptors have long been known to regulate Cl⁻ secretion from intestinal epithelial cells into the lumen, a key process in the development of diarrhea. Several recent studies have highlighted a novel role of A_{2B} receptors in regulating the immune response of intestinal epithelial cells (Figure 3). A_{2B} receptors are the dominant adenosine receptors that are expressed on intestinal epithelial cells, and they are found both on the apical and basolateral surface of these cells [58]. Stimulation of A_{2B} receptors on either surface triggers the release of IL-6 into the apical compartment [59]. A_{2B} receptor stimulation of epithelial cells increases intracellular cAMP levels, which in turn leads to IL-6 transcription via activation of the ATF and CREB and C/EBP β (NF-IL6) transcription factor systems [59]. The physiological relevance of this response is that it provides an amplification mechanism for intestinal inflammation because neutrophils transmigrating through the epithelial cell layer release adenosine, which in turn induces the production of the neutrophil-activating IL-6 [59]. This amplification loop is further enhanced by a rapid increase in the surface expression of A_{2B} receptors after stimulation of the cells with adenosine, which is made possible by prompt recruitment of preformed A_{2B} receptors from intracellular stores [27].

Fibronectins are multifunctional glycoproteins found primarily in the extracellular matrix and can bind a wide spectrum of molecules including heparin, DNA, collagen and the surface of bacteria [60]. Stimulation of A_{2B} receptors on intestinal epithelial cells induces the transcription and subsequent secretion of fibronectin from these cells into the apical compartment [60]. This secreted fibronectin has been shown to aid the adherence and invasion of *Salmonella typhimurium* into intestinal epithelial cells [60]; however, the clinical relevance of this phenomenon remains to be defined.

In summary, A_{2B} receptors possess proinflammatory effects in intestinal epithelial cells (Figure 3), which is consistent with the observation that A_{2B} receptor activation contributes to the pathophysiology of inflammatory bowel disease (IBD) (see later).

Role of A_{2B} receptors in regulating inflammation and immunity in health and disease

Asthma and chronic obstructive pulmonary disease

There is an increasing body of evidence documenting the proinflammatory effects of A_{2B} receptor activation in both rodent and human asthma and chronic obstructive pulmonary disease (COPD), which indicates that A_{2B} receptor antagonists might be of therapeutic value in preventing disease progression in asthma and COPD. This is borne out by recent *in vivo* evidence [61] showing that selective antagonism of adenosine A_{2B} receptors leads to inhibition of airway inflammation and airway reactivity induced by allergen or AMP in a murine asthma model. Although mast cells mediate most of the proinflammatory effects of adenosine in the lung [32], proinflammatory effects of A_{2B} stimulation have also been observed with human bronchial smooth-muscle cells [62], human bronchial epithelial cells [63] and human lung fibroblasts [21], which produce increased levels of IL-6 [21,62] and IL-19 [63] in response to A_{2B} receptor activation.

Adenosine deaminase (ADA)-deficient mice spontaneously develop severe pulmonary inflammation, airway remodeling, fibrosis and enlargement of airspaces in association with increases in adenosine levels in lung tissue [64]. Thus, increases in endogenous adenosine levels can, at least in part, promote the pathophysiological features observed in chronic lung diseases. Treatment of ADA-deficient mice with a selective A_{2B} antagonist was able to markedly limit pulmonary inflammation, fibrosis and airspace enlargement [65]. These findings indicate that A_{2B} receptor signaling is an important contributor to lung injury in environments where endogenous adenosine levels are increased. Collectively, the results of pre-clinical models together with evidence that enprofylline and theophylline are A_{2B} receptor antagonists, indicate that selective antagonism of A_{2B} receptors is an appealing therapeutic approach for managing patients with asthma and COPD.

Vascular injury, hypoxia, preconditioning and ischemia

By generating A_{2B} KO mice, Ravid and coworkers [10,66] showed that A_{2B} receptors are important in moderating vascular injury to various insults, such as systemic endotoxin challenge and guidewire-induced endothelial denudation of the femoral artery, a model of human restenosis after angioplasty. A_{2B} receptor KO mice exhibited increased expression of vascular adhesion molecules, augmented leukocyte rolling and adhesion and elevated levels of pro-inflammatory cytokines [10,66]. Aortic smooth muscle cells derived from A_{2B} receptor KO mice displayed greater proliferation in comparison with controls, which contributed to the increased deleterious neointima formation in the injured femoral artery [66]. Studies enlisting bone marrow chimeric mice showed that it was the loss of A_{2B} receptors on bone-marrow-derived cells that was responsible for the increased inflammation seen in A_{2B} receptor KO mice [10,66].

Extracellular adenosine has long been implicated in the adaptation to hypoxia [67,68] and recent studies have provided evidence that A_{2B} receptors have a key role in dampening hypoxia-induced vascular leak *in vivo*. A_{2B} receptor KO mice exposed to ambient hypoxia had increased vascular permeability in the majority of organs tested, and studies with bone marrow chimeric mice indicated that it was the lack of parenchymal (vascular) and not hematopoietic A_{2B} receptors that caused the increased vascular leak [69]. A_{2B} receptor KO mice suffering from ventilator-induced lung injury demonstrate increased pulmonary vascular leak and edema, which is due to lack of A_{2B} receptors on pulmonary and not hematopoietic cells [69].

Ischemic or hypoxic preconditioning is defined as a rapid, adaptive response to brief, repeated periods of ischemia or hypoxia to render the tissue resistant to the deleterious effects of subsequent severe and long-lasting ischemia or hypoxia, respectively. It is widely accepted that adenosine has a key role in mediating preconditioning, and recent results incriminate A_{2B} receptors as important components of the preconditioning response. Ischemic preconditioning fails to protect the heart [70] and kidney [71] in A_{2B} receptor KO mice, and this lack of preconditioning is associated with increased NF-κB activation and inflammation [70–72].

Inflammatory bowel disease

Crohn's disease and ulcerative colitis are chronic, relapsing inflammatory bowel diseases (IBDs) that are characterized by a dysregulation of the inflammatory functions of intestinal epithelial cells, in addition to poorly controlled activation of both the innate and adaptive immune systems leading to damage of the intestinal mucosa. Although the etiology of IBD remains to be defined, recent studies indicate that the disease results from a dysregulated immune response to luminal antigens and gut microflora in a genetically susceptible host [73]. Consistent with the proinflammatory role of A_{2B} receptor activation in driving intestinal epithelial cell inflammatory responses, A_{2B} receptor blockade [74] or KO [11] suppresses

intestinal inflammation and attenuates the course of disease in murine colitis. The protection caused by A_{2B} receptor inactivation is correlated with decreased production of IL-6 and keratinocyte-derived chemokine, in addition to limited neutrophil accumulation in gut tissue [11,74]. Because A_{2B} receptors are upregulated in gut tissue during both human and murine colitis [24], they are components of a vicious circle that propagates inflammation in the gut.

Cancer

Inflammation in the tumor microenvironment aids in the proliferation and survival of malignant cells, promotes angiogenesis and metastasis and subverts adaptive immune responses. Concentrations of extracellular adenosine are increased in tumor microenvironments [75] because both inflammation and hypoxia, conditions which are present in most tumors, can increase the accumulation of adenosine [76–79]. Recent studies have revealed that A_{2B} receptors can propagate the growth of Lewis lung carcinoma in mice by reprogramming host dendritic cells to produce copious amounts of VEGF, thereby supporting angiogenesis in the tumors [16,80]. It is, however, conceivable that A_{2B} receptors directly affect the growth of other types of tumors because several tumors have been shown to overexpress A_{2B} receptors [81,82]. Further studies will be required to determine whether A_{2B} receptors have tumor growth-regulating roles in other models of cancer development.

Conclusions and perspectives

Research into the role of A_{2B} receptors in regulating immunity and inflammation has gained momentum with the recent availability of KO mouse models and selective receptor ligands. Based on the new information acquired through this research it has become clear that A_{2B} receptors represent potential targets for manipulation in the treatment of immune-mediated and inflammatory diseases. A_{2B} receptors can be advantageous targets for pharmacological intervention because their expression is normally low and is selectively increased in tissues undergoing inflammation. Local increases in A_{2B} receptors in inflamed tissue might thus present an opportunity for selectively targeting tissues with ongoing inflammation. An additional advantage in manipulating the adenosine-A_{2B} receptor system lies in the fact that extracellular adenosine levels are elevated in inflammatory foci and boosting or decreasing these high levels of adenosine can alter signaling through A_{2B} receptors in the local environment without the risk of affecting non-involved tissues.

Before the potential of targeting A_{2B} receptors for selective pharmacological intervention can be realized there are several issues that need to be clarified. One of the issues that could complicate the translation of results from animal models to humans is adenosine-independent signaling by A_{2B} receptors in murine systems. This is illustrated by the observations that mast cells isolated from A_{2B} receptor KO mice exhibit increased degranulation in response to antigen [12,13] and that macrophages obtained from A_{2B} KO mice produce increased levels of TNF- α in the absence of extracellular adenosine [55]. In addition, resting A_{2B} KO animals display increased expression of adhesion molecules and cytokines [10] and enhanced vascular permeability [15], which demonstrates that A_{2B} receptor KO animals have a widely inflammatory phenotype. One potential explanation is that A_{2B} receptors are constitutively active, similar to other G-protein-coupled receptors, such as β_2 -adrenoceptors or δ opioid receptors [83]. Because inverse agonists can block constitutive signaling, use of antagonists that are also inverse agonists can have off-target effects in tissues that are not inflamed and express constitutively active receptors. Another complicating factor is that non-traditional ligands can also activate A_{2B} receptors. For example, the protein netrin is such a novel ligand because it was recently discovered that it can activate A_{2B} receptors and have anti-inflammatory effects [84].

Enprofylline and theophylline, two agents with A_{2B} receptor blocking properties, have been successfully utilized to treat patients suffering from asthma. It is hoped that a better understanding of the role of A_{2B} receptors in regulating immunity will pave the way for additional use of selective modulators of the adenosine-A_{2B} receptor system in the fight to improve human health.

Acknowledgments

This work was supported by the National Institutes of Health Grant R01 GM66189 and the Intramural Research Program of NIH, National Institute on Alcohol Abuse and Alcoholism, as well as Hungarian Research Fund OTKA (K73110) and Marie Curie Grant PIRG02-GA-2007-224767.

References

1. Dolezelova E, et al. A Drosophila adenosine receptor activates cAMP and calcium signaling. *Insect Biochem Mol Biol* 2007;37:318–329. [PubMed: 17368195]
2. Barraco RA, Stefano GB. Pharmacological evidence for the modulation of monoamine release by adenosine in the invertebrate nervous system. *J Neurochem* 1990;54:2002–2006. [PubMed: 2338553]
3. Hansen BD, et al. Evidence for a membrane adenosine receptor in *Leishmania mexicana mexicana* (WR 227). *Adv Exp Med Biol* 1986;195:547–551. [PubMed: 3020925]
4. Jacobson KA, Gao ZG. Adenosine receptors as therapeutic targets. *Nat Rev Drug Discov* 2006;5:247–264. [PubMed: 16518376]
5. Fredholm BB, et al. International Union of Pharmacology. XXV Nomenclature and classification of adenosine receptors. *Pharmacol Rev* 2001;53:527–552. [PubMed: 11734617]
6. Yegutkin GG. Nucleotide- and nucleoside-converting ectoenzymes: important modulators of purinergic signalling cascade. *Biochim Biophys Acta* 2008;1783:673–694. [PubMed: 18302942]
7. Haskó G, et al. Immunomodulatory and neuroprotective effects of inosine. *Trends Pharmacol Sci* 2004;25:152–157. [PubMed: 15019271]
8. Eltzschig HK, et al. Neutrophils as sources of extracellular nucleotides: functional consequences at the vascular interface. *Trends Cardiovasc Med* 2008;18:103–107. [PubMed: 18436149]
9. Haskó G, et al. Adenosine receptors: therapeutic aspects for inflammatory and immune diseases. *Nat Rev Drug Discov* 2008;7:759–770. [PubMed: 18758473]
10. Yang D, et al. The A_{2B} adenosine receptor protects against inflammation and excessive vascular adhesion. *J Clin Invest* 2006;116:1913–1923. [PubMed: 16823489]
11. Kolachala VL, et al. A_{2B} adenosine receptor gene deletion attenuates murine colitis. *Gastroenterology* 2008;135:861–870. [PubMed: 18601927]
12. Hua X, et al. Enhanced mast cell activation in mice deficient in the A_{2B} adenosine receptor. *J Exp Med* 2007;204:117–128. [PubMed: 17200408]
13. Ryzhov S, et al. Effect of A_{2B} adenosine receptor gene ablation on proinflammatory adenosine signaling in mast cells. *J Immunol* 2008;180:7212–7220. [PubMed: 18490720]
14. Mirabet M, et al. Expression of A_{2B} adenosine receptors in human lymphocytes: their role in T cell activation. *J Cell Sci* 1999;112:491–502. [PubMed: 9914161]
15. Eckle T, et al. A_{2B} adenosine receptor dampens hypoxia-induced vascular leak. *Blood* 2008;111:2024–2035. [PubMed: 18056839]
16. Novitskiy SV, et al. Adenosine receptors in regulation of dendritic cell differentiation and function. *Blood* 2008;112:1822–1831. [PubMed: 18559975]
17. Ben Addi A, et al. Modulation of murine dendritic cell function by adenine nucleotides and adenosine: involvement of the A_{2B} receptor. *Eur J Immunol* 2008;38:1610–1620. [PubMed: 18465770]
18. Eltzschig HK, et al. Coordinated adenine nucleotide phosphohydrolysis and nucleoside signaling in posthypoxic endothelium: role of ectonucleotidases and adenosine A_{2B} receptors. *J Exp Med* 2003;198:783–796. [PubMed: 12939345]
19. Zhao P, et al. Hypoxia suppresses the production of MMP-9 by human monocyte-derived dendritic cells and requires activation of adenosine receptor A_{2B} via cAMP/PKA signaling pathway. *Mol Immunol* 2008;45:2187–2195. [PubMed: 18215420]

20. Feoktistov I, et al. Hypoxia modulates adenosine receptors in human endothelial and smooth muscle cells toward an A2B angiogenic phenotype. *Hypertension* 2004;44:649–654. [PubMed: 15452028]
21. Zhong H, et al. Synergy between A2B adenosine receptors and hypoxia in activating human lung fibroblasts. *Am J Respir Cell Mol Biol* 2005;32:2–8. [PubMed: 15472138]
22. Nemeth ZH, et al. cDNA microarray analysis reveals a nuclear factor-kappaB-independent regulation of macrophage function by adenosine. *J Pharmacol Exp Ther* 2003;306:1042–1049. [PubMed: 12766259]
23. Nguyen DK, et al. Th1 cytokines regulate adenosine receptors and their downstream signaling elements in human microvascular endothelial cells. *J Immunol* 2003;171:3991–3998. [PubMed: 14530318]
24. Kolachala V, et al. TNF-alpha upregulates adenosine 2b (A2b) receptor expression and signaling in intestinal epithelial cells: a basis for A2bR overexpression in colitis. *Cell Mol Life Sci* 2005;62:2647–2657. [PubMed: 16322943]
25. St Hilaire C, et al. TNF-alpha upregulates the A2B adenosine receptor gene: the role of NAD(P)H oxidase 4. *Biochem Biophys Res Commun* 2008;375:292–296. [PubMed: 18647598]
26. Xaus J, et al. IFN-gamma up-regulates the A2B adenosine receptor expression in macrophages: a mechanism of macrophage deactivation. *J Immunol* 1999;162:3607–3614. [PubMed: 10092821]
27. Sitaraman SV, et al. The adenosine 2b receptor is recruited to the plasma membrane and associates with E3KARP and Ezrin upon agonist stimulation. *J Biol Chem* 2002;277:33188–33195. [PubMed: 12080047]
28. Wang L, et al. Agonist-induced polarized trafficking and surface expression of the adenosine 2b receptor in intestinal epithelial cells: role of SNARE proteins. *Am J Physiol Gastrointest Liver Physiol* 2004;287:G1100–G1107. [PubMed: 15256361]
29. Feoktistov I, Biaggioni I. Adenosine A2B receptors. *Pharmacol Rev* 1997;49:381–402. [PubMed: 9443164]
30. Bradding P, et al. The role of the mast cell in the pathophysiology of asthma. *J Allergy Clin Immunol* 2006;117:1277–1284. [PubMed: 16750987]
31. Metz M, Maurer M. Mast cells—key effector cells in immune responses. *Trends Immunol* 2007;28:234–241. [PubMed: 17400512]
32. Polosa R, Holgate ST. Adenosine receptors as promising therapeutic targets for drug development in chronic airway inflammation. *Curr Drug Targets* 2006;7:699–706. [PubMed: 16787172]
33. Feoktistov I, Biaggioni I. Adenosine A2b receptors evoke interleukin-8 secretion in human mast cells. An enprofylline-sensitive mechanism with implications for asthma. *J Clin Invest* 1995;96:1979–1986. [PubMed: 7560091]
34. Feoktistov I, et al. Role of p38 mitogen-activated protein kinase and extracellular signal-regulated protein kinase kinase in adenosine A2B receptor-mediated interleukin-8 production in human mast cells. *Mol Pharmacol* 1999;55:726–734. [PubMed: 10101031]
35. Ryzhov SR, et al. Differential role of the carboxy-terminus of the A(2B) adenosine receptor in stimulation of adenylate cyclase, phospholipase Cbeta, and interleukin-8. *Purinergic Signal*. 2009(in press)
36. Ryzhov S, et al. Adenosine-activated mast cells induce IgE synthesis by B lymphocytes: an A2B-mediated process involving Th2 cytokines IL-4 and IL-13 with implications for asthma. *J Immunol* 2004;172:7726–7733. [PubMed: 15187156]
37. Ryzhov S, et al. Cross-talk between G(s)- and G(q)-coupled pathways in regulation of interleukin-4 by A(2B) adenosine receptors in human mast cells. *Mol Pharmacol* 2006;70:727–735. [PubMed: 16707627]
38. Muller WA. Leukocyte-endothelial-cell interactions in leukocyte transmigration and the inflammatory response. *Trends Immunol* 2003;24:327–334. [PubMed: 12810109]
39. Feoktistov I, et al. Differential expression of adenosine receptors in human endothelial cells: role of A2B receptors in angiogenic factor regulation. *Circ Res* 2002;90:531–538. [PubMed: 11909816]
40. Eltzschig HK, et al. Endogenous adenosine produced during hypoxia attenuates neutrophil accumulation: coordination by extracellular nucleotide metabolism. *Blood* 2004;104:3986–3992. [PubMed: 15319286]

41. Lennon PF, et al. Neutrophil-derived 5'-adenosine monophosphate promotes endothelial barrier function via CD73-mediated conversion to adenosine and endothelial A2B receptor activation. *J Exp Med* 1998;188:1433–1443. [PubMed: 9782120]
42. Kong T, et al. HIF-dependent induction of adenosine A2B receptor in hypoxia. *FASEB J* 2006;20:2242–2250. [PubMed: 17077301]
43. Morote-Garcia JC, et al. HIF-1-dependent repression of adenosine kinase attenuates hypoxia-induced vascular leak. *Blood* 2008;111:5571–5580. [PubMed: 18309031]
44. Synnestvedt K, et al. Ecto-5'-nucleotidase (CD73) regulation by hypoxia-inducible factor-1 mediates permeability changes in intestinal epithelia. *J Clin Invest* 2002;110:993–1002. [PubMed: 12370277]
45. Eltzschig HK, et al. HIF-1-dependent repression of equilibrative nucleoside transporter (ENT) in hypoxia. *J Exp Med* 2005;202:1493–1505. [PubMed: 16330813]
46. Gilroy DW, et al. Inflammatory resolution: new opportunities for drug discovery. *Nat Rev Drug Discov* 2004;3:401–416. [PubMed: 15136788]
47. Grant MB, et al. Proliferation, migration, and ERK activation in human retinal endothelial cells through A(2B) adenosine receptor stimulation. *Invest Ophthalmol Vis Sci* 2001;42:2068–2073. [PubMed: 11481274]
48. Adair TH. Growth regulation of the vascular system: an emerging role for adenosine. *Am J Physiol Regul Integr Comp Physiol* 2005;289:R283–R296. [PubMed: 16014444]
49. Haskó G, et al. Shaping of monocyte and macrophage function by adenosine receptors. *Pharmacol Ther* 2007;113:264–275. [PubMed: 17056121]
50. Csoka B, et al. A2A adenosine receptors and C/EBPbeta are crucially required for IL-10 production by macrophages exposed to *Escherichia coli*. *Blood* 2007;110:2685–2695. [PubMed: 17525287]
51. Haskó G, et al. Adenosine inhibits IL-12 and TNF- α production via adenosine A2a receptor-dependent and independent mechanisms. *FASEB J* 2000;14:2065–2074. [PubMed: 11023991]
52. Szabo C, et al. Suppression of macrophage inflammatory protein (MIP)-1 α production and collagen-induced arthritis by adenosine receptor agonists. *Br J Pharmacol* 1998;125:379–387. [PubMed: 9786512]
53. Kreckler LM, et al. Adenosine inhibits tumor necrosis factor- α release from mouse peritoneal macrophages via A2A and A2B but not the A3 adenosine receptor. *J Pharmacol Exp Ther* 2006;317:172–180. [PubMed: 16339914]
54. Nemeth ZH, et al. Adenosine augments IL-10 production by macrophages through an A2B receptor-mediated posttranscriptional mechanism. *J Immunol* 2005;175:8260–8270. [PubMed: 16339566]
55. Ryzhov S, et al. Effect of A2B adenosine receptor gene ablation on adenosine-dependent regulation of proinflammatory cytokines. *J Pharmacol Exp Ther* 2008;324:694–700. [PubMed: 17965229]
56. Xaus J, et al. Adenosine inhibits macrophage colony-stimulating factor-dependent proliferation of macrophages through the induction of p27kip-1 expression. *J Immunol* 1999;163:4140–4149. [PubMed: 10510349]
57. Artis D. Epithelial-cell recognition of commensal bacteria and maintenance of immune homeostasis in the gut. *Nat Rev Immunol* 2008;8:411–420. [PubMed: 18469830]
58. Kolachala VL, et al. Purinergic receptors in gastrointestinal inflammation. *Am J Physiol Gastrointest Liver Physiol* 2008;294:G401–G410. [PubMed: 18063703]
59. Sitaraman SV, et al. Neutrophil-epithelial crosstalk at the intestinal luminal surface mediated by reciprocal secretion of adenosine and IL-6. *J Clin Invest* 2001;107:861–869. [PubMed: 11285305]
60. Walia B, et al. Polarized fibronectin secretion induced by adenosine regulates bacterial-epithelial interaction in human intestinal epithelial cells. *Biochem J* 2004;382:589–596. [PubMed: 15193149]
61. Mustafa SJ, et al. Effect of a specific and selective A(2B) adenosine receptor antagonist on adenosine agonist AMP and allergen-induced airway responsiveness and cellular influx in a mouse model of asthma. *J Pharmacol Exp Ther* 2007;320:1246–1251. [PubMed: 17159162]
62. Zhong H, et al. A(2B) adenosine receptors increase cytokine release by bronchial smooth muscle cells. *Am J Respir Cell Mol Biol* 2004;30:118–125. [PubMed: 12855406]
63. Zhong H, et al. A2B adenosine receptors induce IL-19 from bronchial epithelial cells, resulting in TNF- α increase. *Am J Respir Cell Mol Biol* 2006;35:587–592. [PubMed: 16778150]

64. Blackburn MR, et al. Metabolic consequences of adenosine deaminase deficiency in mice are associated with defects in alveogenesis, pulmonary inflammation, and airway obstruction. *J Exp Med* 2000;192:159–170. [PubMed: 10899903]
65. Sun CX, et al. Role of A2B adenosine receptor signaling in adenosine-dependent pulmonary inflammation and injury. *J Clin Invest* 2006;116:2173–2182. [PubMed: 16841096]
66. Yang D, et al. The A2b adenosine receptor protects against vascular injury. *Proc Natl Acad Sci U S A* 2008;105:792–796. [PubMed: 18184815]
67. Lukashev D, et al. Targeting hypoxia-A(2A) adenosine receptor-mediated mechanisms of tissue protection. *Drug Discov Today* 2004;9:403–409. [PubMed: 15081957]
68. Sitkovsky M, Lukashev D. Regulation of immune cells by local-tissue oxygen tension: HIF1 alpha and adenosine receptors. *Nat Rev Immunol* 2005;5:712–721. [PubMed: 16110315]
69. Eckle T, et al. A2B adenosine receptor signaling attenuates acute lung injury by enhancing alveolar fluid clearance in mice. *J Clin Invest* 2008;118:3301–3315. [PubMed: 18787641]
70. Eckle T, et al. Cardioprotection by ecto-5'-nucleotidase (CD73) and A2B adenosine receptors. *Circulation* 2007;115:1581–1590. [PubMed: 17353435]
71. Grenz A, et al. The reno-vascular A2B adenosine receptor protects the kidney from ischemia. *PLoS Med* 2008;5:e137. [PubMed: 18578565]
72. Khoury J, et al. Antiinflammatory adaptation to hypoxia through adenosine-mediated cullin-1 deneddylation. *J Clin Invest* 2007;117:703–711. [PubMed: 17318263]
73. Bamias G, Cominelli F. Immunopathogenesis of inflammatory bowel disease: current concepts. *Curr Opin Gastroenterol* 2007;23:365–369. [PubMed: 17545770]
74. Kolachala V, et al. Blockade of adenosine A(2B) receptors ameliorates murine colitis. *Br J Pharmacol* 2008;155:127–137. [PubMed: 18536750]
75. Blay J, et al. The extracellular fluid of solid carcinomas contains immunosuppressive concentrations of adenosine. *Cancer Res* 1997;57:2602–2605. [PubMed: 9205063]
76. Lukashev D, et al. From “Hellstrom Paradox” to anti-adenosinergic cancer immunotherapy. *Purinergic Signal* 2007;3:129–134. [PubMed: 18404426]
77. Lukashev D, et al. Hypoxia-dependent anti-inflammatory pathways in protection of cancerous tissues. *Cancer Metastasis Rev* 2007;26:273–279. [PubMed: 17404693]
78. Sitkovsky MV. T regulatory cells: hypoxia-adenosinergic suppression and re-direction of the immune response. *Trends Immunol* 2009;30:102–108. [PubMed: 19201652]
79. Ohta A, et al. A2A adenosine receptor protects tumors from antitumor T cells. *Proc Natl Acad Sci U S A* 2006;103:13132–13137. [PubMed: 16916931]
80. Ryzhov S, et al. Host A(2B) adenosine receptors promote carcinoma growth. *Neoplasia* 2008;10:987–995. [PubMed: 18714400]
81. Panjehpour M, et al. Human breast cancer cell line MDA-MB-231 expresses endogenous A2B adenosine receptors mediating a Ca²⁺ signal. *Br J Pharmacol* 2005;145:211–218. [PubMed: 15753948]
82. Gessi S, et al. Adenosine receptors in colon carcinoma tissues and colon tumoral cell lines: focus on the A(3) adenosine subtype. *J Cell Physiol* 2007;211:826–836. [PubMed: 17348028]
83. Vizi ES. Role of high-affinity receptors and membrane transporters in nonsynaptic communication and drug action in the central nervous system. *Pharmacol Rev* 2000;52:63–89. [PubMed: 10699155]
84. Rosenberger P, et al. Hypoxia-inducible factor-dependent induction of netrin-1 dampens inflammation caused by hypoxia. *Nat Immunol* 2009;10:195–202. [PubMed: 19122655]

Figure 1.

A_{2B} receptor activation on mast cells triggers the release of IL-4 and IL-8. Coupling of A_{2B} receptors to G_q proteins activates PLCβ leading to elevations in intracellular Ca²⁺ levels and activation of the transcription factor NFAT, which in turn transactivates the IL-4 promoter and results in the release of IL-4. Coupling of A_{2B} receptors to G_s proteins and subsequent accumulation of cAMP upregulates NFAT protein levels, amplifying the G_q-mediated IL-4 response. G_q and G_s protein-independent activation of p38 through the third intracellular loop of the A_{2B} receptor leads to increased IL-8 transcription and release. Abbreviations: A_{2B}AR, A_{2B} adenosine receptor; cAMP, cyclic adenosine monophosphate; NFAT, nuclear factor of activated T cells; PLCβ, phospholipase C β.

Figure 2.

Regulation of endothelial cell function by A_{2B} receptors. During acute inflammation, A_{2B} receptors suppress the expression of adhesion molecules on endothelial cells (EC), which leads to decreased leukocyte rolling, adhesion and transmigration. A_{2B} receptors also prevent plasma leakage from the intravascular compartment into the tissue. During inflammatory resolution, A_{2B} receptors facilitate endothelial cell proliferation and migration and promote the formation of blood vessels, effects that are mediated by the release of VEGF (vascular endothelial growth factor).

Figure 3.

A_{2B} receptors trigger the release of IL-6 by intestinal epithelial cells. Transmigrating neutrophils release ATP into the luminal compartment, where it is sequentially degraded to adenosine by CD39 and CD73 which are present on the brush border (apical side) of intestinal epithelial cells. Adenosine (ADO) released from transmigrated neutrophils activates luminal A_{2B} receptors, which results in intracellular cAMP accumulation, and subsequently, increased transcription of IL-6 via activation of the transcription factors CREB and C/EBPβ.