Published in final edited form as:

Exp Biol Med (Maywood). 2008 August ; 233(8): 997–1012. doi:10.3181/0802-RM-70.

Expanded CD34⁺ Human Umbilical Cord Blood Cells Generate Multiple Lymphohematopoietic Lineages in NOD-*scid* IL2 γ ^{null} Mice

Lisa J. Giassi^{*}, Todd Pearson^{*}, Leonard D. Shultz^{†,1}, Joseph Laning[‡], Kristin Biber[‡], Morey Kraus[‡], Bruce A. Woda[§], Madelyn R. Schmidt^{||}, Robert T. Woodland^{||}, Aldo A. Rossini^{*}, and Dale L. Greiner^{*}

^{*}Division of Diabetes, Department of Medicine, University of Massachusetts Medical School, Worcester, Massachusetts 01605

[†]The Jackson Laboratory, Bar Harbor, Maine 04609

[‡]Viacell, Inc., Cambridge, Massachusetts 02142

[§]Department of Pathology, University of Massachusetts Medical School, Worcester, Massachusetts 01655

^{||}Department of Molecular Genetics and Microbiology University of Massachusetts Medical School, Worcester, Massachusetts 01655

Abstract

Umbilical cord blood (UCB) is increasingly being used for human hematopoietic stem cell (HSC) transplantation in children but often requires pooling multiple cords to obtain sufficient numbers for transplantation in adults. To overcome this limitation, we have used an *ex vivo* two-week culture system to expand the number of hematopoietic CD34⁺ cells in cord blood. To assess the *in vivo* function of these expanded CD34⁺ cells, cultured human UCB containing 1×10^6 CD34⁺ cells were transplanted into conditioned NOD-*scid* IL2 γ ^{null} mice. The expanded CD34⁺ cells displayed short- and long-term repopulating cell activity. The cultured human cells differentiated into myeloid, B-lymphoid, and erythroid lineages, but not T lymphocytes. Administration of human recombinant TNF α to recipient mice immediately prior to transplantation promoted human thymocyte and T-cell development. These T cells proliferated vigorously in response to TCR cross-linking by anti-CD3 antibody. Engrafted TNF α -treated mice generated antibodies in response to T-dependent and T-independent immunization which was enhanced when mice were co-treated with the B cell cytokine BLyS. *Ex vivo* expanded CD34⁺ human UCB cells have the capacity to generate multiple hematopoietic lineages and a functional human immune system upon transplantation into TNF α -treated NOD-*scid* IL2 γ ^{null} mice.

Keywords

NOD-*scid* IL2 γ ^{null}; mice; SCID; umbilical cord blood; TNF α ; *ex vivo* expansion; human; immune response

Introduction

Allogeneic hematopoietic stem cell (HSC) transplantation following myeloablative therapy is curative for many malignant, genetic, and autoimmune disorders (1). Sources of stem cells for clinical transplantation include bone marrow, cytokine-mobilized peripheral blood, and umbilical cord blood (UCB) (2–5). Human UCB is increasingly used as a clinical source of HSC due to the advantages including immediate availability, high proliferative capacity, and the low incidence of severe GVHD in recipients (4,6–13). However, successful human HSC transplantation is dependent on the absolute number of CD34⁺ cells transplanted (14–19). In the case of UCB, this often requires the pooling of multiple cords to obtain the optimal CD34⁺ cell dose (17,18,20). Concerns related to the use of pooled cords include the possibility of graft-versus-graft as well as the unpredictable reconstitution of individual cords (17,19, 20).

Many laboratories are culturing UCB to increase the CD34⁺ cell yield from a single cord (21–25). Long-term cultures have been successful in expanding CD34⁺ cell numbers, but maintaining HSC activity has been difficult (26–30). Investigators rely on immunodeficient mice, particularly NOD-*scid* mice, to define the *in vivo* function of the cultured HSC. However, transplantation of cultured human HSC into NOD-*scid* mice has yielded only low engraftment, limited multilineage hematopoietic development, and few if any HSC-derived T lymphocytes (31–33).

Recently, new immunodeficient murine mouse models that lack a functional common IL-2-receptor gamma chain (*IL2r γ ^{null}*) have been found to be superior in their support of human HSC engraftment (34). These animals lack mature lymphocytes and NK cells, express other impairments in innate immunity, and are long-lived, permitting long-term studies of engraftment with human HSCs (34).

In the present study, we used the NOD-*scid* *IL2r γ ^{null}* mouse model to investigate the *in vivo* potential of human UCB stem cells following expansion of CD34⁺ cells in a two-week culture system. Cultured UCB engrafted and developed into myeloid, erythroid, and B-lymphoid lineages, and when the recipients were pre-treated with TNF α , functional T lymphoid cells were generated. Engrafted mice produced antibody in response to immunization with T-dependent and T-independent antigens, which was enhanced by administration of the human B Lymphocyte Stimulator (BLyS) cytokine (also termed BAFF or TNFSF13B).

Materials and Methods

Mice

NOD.CB17-*Prkdc^{scid}* (abbreviated as NOD-*scid*) and NOD.Cg-*Prkdc^{scid}* *IL2r γ ^{tmWjl}/Sz* (abbreviated as NOD-*scid* *IL2r γ ^{null}*) mice have been described (35,36). Male and female 6- to 12-week-old mice of each strain were used in these studies. They were specific pathogen free, housed in microisolator cages, and given autoclaved food and on alternate weeks given autoclaved acidified water or sulfamethoxazole-trimethoprim-medicated water (Goldline Laboratories, Ft. Lauderdale, FL). All animal use was in accordance with the guidelines of the Animal Care and Use Committees of the University of Massachusetts Medical School and The Jackson Laboratory.

Antibodies

Anti-human α/β TCR was obtained from Becton Dickinson (San Jose, CA). Anti-human BDCA2 was obtained from Miltenyi (Auburn, CA). All other antibodies and isotype controls were obtained from BD Pharmingen (San Diego, CA). Antibodies were conjugated with FITC, PE, PerCP, APC, Alexa 405, Pacific Blue, or Alexa700.

Cell Preparation

Single-cell suspensions of bone marrow, spleen, and thymus were prepared from engrafted mice. Whole blood was collected in PBS containing EDTA. RBC in marrow and spleen were removed by lysis with a hypotonic solution. GlyA-stained bone marrow cells were not subjected to red blood cell lysis. Cells counts were performed using a Coulter Counter (Beckman, Miami, FL).

Flow Cytometry

Single cell suspensions or whole blood were preincubated with rat anti-mouse FcRIIb (2.4G2) to block Fc binding and then incubated with appropriate dilutions of specific antibodies. In all wells, PerCP-conjugated anti-mouse CD45 and APC-conjugated anti-human CD45 were included. Labeled cells were fixed in 1% paraformaldehyde for analysis. Whole blood was stained according to manufacturer's instructions (BD Biosciences, San Jose, CA). At least 50,000 events were acquired on either BD Biosciences LSRII or FACSCalibur instruments (BD Biosciences). All cells that were murine CD45⁻ and human CD45⁺ were analyzed for the presence of multilineage differentiation. Bone marrow cells that were negative for murine CD45 were also analyzed for the presence of human RBC and their progenitors. Data acquisition and analysis was performed with BD FACSDiva (BD Pharmingen) and Flow Jo (Tree Star, Inc., Ashland, OR) software.

Ex Vivo Culture of CD34⁺ Human UCB Stem Cells

Cultured UCB stem cells were prepared by Viacell, Inc. (Cambridge, MA). Briefly, red blood cells were removed using a double-density Percoll Gradient (1.05/ 1.077 g/ml). The recovered UCB cells were lineage-depleted using Stem Cell Technologies (Vancouver, British Columbia, Canada) SCT 9 antibody cocktail (CD2, CD3, CD14, CD16, CD19, CD24, CD56, CD66b, GlyA). The lineage-depleted cells were cultured for seven days in Stemspan (Stem Cell Technologies) supplemented with defined lipid cocktail (Gibco, Carlsbad, CA), 50 (μg/mL gentamycin (Abbott Laboratories, Chicago, IL), 100 ng/mL SCF (Amgen, Thousand Oaks, CA), 100 ng/mL Flt3-L (Amgen), and 100 ng/mL Tpo (R&D Systems, Minneapolis, MN). At day seven of culture, non-adherent cells were recovered, lineage depletion was repeated, and cells were cultured for an additional seven days. After two weeks in culture, the percentage and number of cell subsets was determined by flow cytometry.

Engraftment of Mice with Cultured CD34⁺ Human UCB Stem Cells

Adult NOD-*scid* IL2 γ ^{null} and NOD-*scid* mice were irradiated with 2.4 and 3.25 Gy, respectively, from a ¹³⁷Cs source (Gammacell 40, Atomic Energy of Canada, Ottawa, Canada) using previously optimized irradiation doses (36,37). Mice were injected intravenously with cultured cells containing 1×10^6 CD34⁺ cells.

TNF α Treatment

Recombinant human TNFα (R&D Systems) was reconstituted in PBS (10.0 μg/ml) containing 0.1% BSA. A single injection of 50 μl of TNFα (0.5 μg/recipient) was given i.p. immediately after irradiation and 4 hours before injection of human UCB cells.

Immunization

NOD-*scid* IL2 γ ^{null} mice engrafted with cultured CD34⁺ cells 12 weeks earlier were immunized s.c. with 20 μl/mouse containing 1 μg each of 23 pneumococcal serotypes (Pneumovax 23; Merck, White-house Station, NJ). Serum was recovered 14 days after primary immunization. Mice were then injected i.p. with BLyS (10 (μg) or PBS for 4 days, and reboosted with Pneumovax 28 days after primary immunization. BLyS was made from

recombinant material from a cell line provided by Dr. Randolph Noelle (Dartmouth Medical, Lebanon, NH). Serum was recovered 14 days after boosting for analysis. The same mice were then immunized s.c. 8 days later with Tetanus Toxoid Adsorbed USP (0.15 µl; Aventis Pasteur Inc., Swiftwater, PA), and serum was obtained 14 days later. Three mice that had previously been given BLyS (see above) were again treated with BLyS for 4 days and reboosted with Tetanus Toxoid. Serum was recovered 14 days later for analysis.

Human Ig ELISA

Levels of human IgG and IgM were measured in mouse sera using an ELISA assay. Briefly, ELISA plates were coated with the F(ab')₂ fragment of anti-human IgG or IgM (Jackson ImmunoResearch, West Grove, PA) overnight and blocked with 0.5% horse serum and 0.02% azide in PBS. Triplicate serial dilutions of sera were added, followed by biotinylated mouse anti-human IgL kappa or lambda (Ebioscience, San Diego, CA). Streptavidin alkaline phosphatase (Southern Biotech) was added, and the reaction visualized by addition of 1 mg/ml *p*-nitrophenyl phosphate (Sigma-Aldrich) in 1 M diethanolamine, 0.5 mM MgCl₂. Color development was terminated after 30 min with 3M NaOH, and plates were read at 405 nm using an Emax ELISA plate reader (Molecular Devices, Sunnyvale, CA). The levels of human IgG and IgM in the sera were calculated using a standard curve (Jackson ImmunoResearch). Limits of detection were 3 ng/ml for IgM and 2 ng/ml for IgG.

Streptococcus Pneumonia ELISA

Antibodies specific for *Streptococcus pneumoniae* capsular polysaccharide serotypes 4 or 14 were measured in mouse sera as described (38). The levels of antibody in the sera were calculated using a standard (US reference human pneumococcal antiserum; FDA, Rockville, MD).

Tetanus ELISA

Quantification of anti-tetanus antibody used a commercial ELISA kit (SCIMEDX Corporation, Denville, NJ). The ELISA was performed according to the manufacturer's instructions, and developed for visualization using sheep anti-human horseradish peroxidase conjugate. The optical density at 450/620 was measured on a MDS Analytical Technologies Emax Precision Microplate Reader (Toronto, Canada). All tetanus antibody titers were calculated based on standards included with the ELISA kit and are reported as milli-International Units per milliliter (mIU/ml).

Proliferative Response Following Stimulation with Anti-CD3 Antibody

Splenocytes from TNFα-treated NOD-*scid* IL2r^{null} mice engrafted with cultured UCB were plated in anti-CD3 mAb coated wells of a flat bottom 96-well plate (Costar, Cambridge, MA) in triplicate at a concentration of 2.5×10^5 cells/well in 200 µl RPMI supplemented with 10% FBS. PBMC from a normal donor was used as a positive control. Cells were stimulated with plate bound anti-CD3 mAb and soluble anti-CD28/CD49d FastImmune reagent (1 µg/ml, BD Pharmingen, San Diego, CA) for 64 hrs at 37°C in an atmosphere of 5% CO₂, and 1 (µCi of [³H]thymidine (Perkin Elmer, Waltham, MA) was added during the last 16 hrs of culture. [³H]thymidine incorporation was measured using a 1450 Microbeta Trilux Instrument (Wallac, Gaithersburg, MD). The number of CD3⁺ cells in each sample was determined and used to calculate the counts/CD3⁺ T cell. Data are presented as mean cpm/ 1×10^5 human CD3⁺ cells $\pm$ 1 SEM.

Histology

Tissues were fixed in 10% neutral buffered formalin, embedded in paraffin, and sectioned at 5 µm. Antigen retrieval prior to immunohistochemical staining was performed by incubating

the sections in 0.1 mol/l citrate buffer (pH 6.0) in an 800-W microwave oven for 15 min. The slides were stained on a Dako (Carpinteria, CA) automated immunostainer using the EnVision (Dako) staining procedure. The sections were incubated with the EnVision plus Dual Link reagent (a polymer conjugated with goat anti-rabbit Ig or goat anti-mouse Ig and horseradish peroxidase) for 30 min. The sections were reacted with diaminobenzidine and hydrogen peroxide for visualization and counterstained with hematoxylin. Negative controls induced species matched non-binding antibodies. A qualified pathologist (BAW) who was unaware of the treatment of the tissue donor evaluated all tissues.

Statistics

Results are expressed as mean $\pm$ one standard error. Values of specified groups were compared with the unpaired Student's *t* test (GraphPad Software, Version 4.0, San Diego, CA). Statistical significance was assumed for *P* values of <0.05 .

Results

Ex Vivo Expansion of CD34⁺ Human UCB

Phenotype of Cells Recovered from a Two-Week Culture of Human UCB—Human UCB, cord blood cells were cultured for two weeks in the presence of a cytokine cocktail, and the number and phenotype of the nonadherent cells was determined. The average expansion of human CD45⁺ cells and CD34⁺ cells was ~68-fold and ~37-fold, respectively (Table 1). The UCB population put into culture was devoid of mature lymphoid and myeloid populations and contained $32.3 \pm 3.9\%$ CD34⁺ cells (Table 2). Although the percentage of CD34⁺ cells in the population recovered from culture ($20.1 \pm 1.9\%$) was slightly lower than in the input population, the overall expansion led to an increase in the total number of CD34⁺ cells recovered.

Phenotypically, many two-week cultured CD34⁺ cells co-expressed CD33 and CD13, suggesting they were early stem/progenitor cells in the granulocyte and monocyte lineages (Table 2). Less than 1% of the cells recovered from culture were CD3⁺ or CD19⁺, less than 2% of the cells were GlyA⁺, and less than 5% of recovered cells were CD14⁺. Approximately 1% of the total cells were CD34⁺CD7⁺, suggesting a commitment to a T or NK cell lineage. Megakaryocytic lineage progenitors (CD34⁺CD41⁺CD61⁺) comprised ~3% of the recovered cells (Table 2).

Engraftment of Cultured Human UCB in NOD-*scid* and NOD-*scid* IL2 γ ^{null} mice—The NOD-*scid* mouse has been the “gold standard” for *in vivo* analyses of human HSC for over 10 years (36,37,39–43). We and others have recently described new immunodeficient mouse models based on the targeted mutation in the IL-2 receptor common γ chain (34). Therefore, we first compared the ability of cultured UCB to engraft in NOD-*scid* and NOD-*scid* IL2 γ ^{null} mice. These stocks of mice were injected with cultured human UCB cells containing 1×10^6 CD34⁺ cells and engraftment in the bone marrow was quantified twelve weeks later.

Low levels of human CD45⁺ cell engraftment were observed in the bone marrow of NOD-*scid* mice ($0.2 \pm 0.1\%$, $n = 8$). In contrast, robust levels of human CD45⁺ cells were generated in NOD-*scid* IL2 γ ^{null} mice ($24.3 \pm 6.6\%$, $n = 5$, $P < 0.001$). Based on these data, all subsequent experiments were performed using NOD-*scid* IL2 γ ^{null} mice as recipients.

Short-and Long-Term Stem Cell Repopulating Potential of Cultured Human UCB

Two major human stem/progenitor cell populations have been described using immunodeficient mice engrafted with human HSC: one is termed short-term repopulating cell (STRC), the other long-term repopulating cell (LTRC) (44,45).

Short-Term Repopulating Cell Engraftment—Cultured human UCB containing 1×10^6 CD34⁺ cells were injected into conditioned NOD-*scid* IL2 γ^{null} mice, and human cell engraftment in the bone marrow of the recipients was determined 3 weeks later.

Human CD45⁺ cells comprised ~2% of total bone marrow cells (Table 3). To determine their lineage, we analyzed the subsets of human CD45⁺ cells present in the marrow, specifically analyzing for the presence of cell populations that are known to comprise the short-term repopulation cell population in *scid* mice (44,45). Of the CD45⁺ cells, CD34⁺ stem/progenitor cells were generated, including many committed to the myeloid (CD33⁺ and/or CD38⁺) lineage. CD45⁺ cells expressing phenotypic characteristics of early granulocytes, monocytes, and macrophages (CD33⁺CD64⁺) were also observed. In the lymphoid lineage, CD20⁺ B cells were generated, but no detectable CD3⁺ T cells were observed (Table 3). Representative histograms are shown in Figure 1. These data demonstrate that cultured UCB display human STRC capacity when transplanted into NOD-*scid* IL2 γ^{null} mice.

Long-Term Repopulating Cell Engraftment—We next analyzed the subsets of human cells known to comprise the long-term repopulation cell population in *scid* mice (36). We determined if cultured UCB could generate long-term engraftment, multiple lineages of immature and mature cells, and hematopoietic stem cells in NOD-*scid* IL2 γ^{null} mice. Bone marrow and spleen were recovered 12 weeks after injection of cultured UCB and analyzed for the presence of human cells.

Bone Marrow—Human CD45⁺ cells ($23.3 \pm 4.4\%$) expressing phenotypes associated with multilineage hematopoietic cell populations were readily detected in the bone marrow (Table 4). Myeloid progenitors expressing CD33 as well as more mature CD64⁺CD33⁺ myeloid cells were detected. CD14⁺ monocytes, CD41a⁺ megakaryocytes, and BDCA2⁺ dendritic cells were also observed (Table 4). In the lymphoid lineage, CD10⁺CD20⁺ B cells were detected and some B cells coexpressed both CD20 and cell surface IgM. Human CD56⁺ NK cells were also detected (Table 4). These data document that the cultured cells could generate both lymphoid and myeloid lineage populations.

Spleen—Human CD45⁺ cells were also detected in the spleen twelve weeks after injection ($16.2 \pm 4.1\%$). These CD45⁺ cells were comprised of multiple hematopoietic lineages. CD20⁺IgM⁺ B cells and CD33⁺ myeloid progenitor cells were observed. However, no human CD3⁺ T cells were detected (Table 4). Bone marrow recovered from a cohort of these mice was adoptively transferred into secondary NOD-*scid* IL2 γ^{null} mice. Twelve weeks later, human CD45⁺ cells were detected in 6 of 8 of the secondary recipients, suggesting the presence of HSC in the primary hosts. These data demonstrate that cultured UCB display human LTRC activity but fail to generate mature T cells when transplanted into NOD-*scid* IL2 γ^{null} mice.

TNF α Promotes Human CD3⁺ T-Cell Development in NOD-*scid* IL2 γ^{null} Mice Engrafted with Cultured Human UCB

The inability of cultured UCB to generate CD3⁺ T cells in NOD-*scid* IL2 γ^{null} mice suggests that 1) T-cell progenitors fail to survive in culture, 2) an inhibitory component(s) restricting T-cell development is present, or 3) a factor required for the generation of human CD3⁺ T cells in the host is absent. One critical factor may be TNF α , as it has been reported that TNF α

promotes the *in vivo* generation of T cells in NOD-*scid* mice engrafted with human HSC (46).

Human CD45⁺ Cell Engraftment in TNF α -Treated NOD-*scid* IL2 γ ^{null} Mice

Transplanted with Cultured Human UCB—We next determined whether TNF α would enhance the ability of cultured UCB to generate T cells. NOD-*scid* IL2 γ ^{null} mice pretreated with TNF α generated levels of human CD45⁺ cells in the blood ($3.1 \pm 0.8\%$; $7.1 \pm 2.4 \times 10^4$ /ml, $n = 32$) equivalent to mice not given TNF α ($4.4 \pm 1.1\%$; $10.0 \pm 0.2 \times 10^4$ /ml, $n = 27$, $P = \text{N.S.}$, Fig. 2). Levels of human CD45⁺ cells in the bone marrow ($23.3 \pm 4.4\%$; $6.5 \pm 1.3 \times 10^6$, $n = 26$) of TNF α treated mice were also equivalent to that in non-TNF α -treated mice ($25.6 \pm 4.7\%$; $4.8 \pm 1.0 \times 10^6$, $n = 19$, $P = \text{N.S.}$).

The percent of CD45⁺ cells in the spleen of mice pretreated with TNF α ($38.6 \pm 6.2\%$, $n = 11$) was significantly greater than that in mice not treated with TNF α ($16.2 \pm 4.2\%$, $n = 12$, $P < 0.01$). There was also a difference in cell numbers in non-TNF α -treated mice ($2.6 \pm 0.9 \times 10^6$) and TNF α -treated mice ($7.3 \pm 1.8 \times 10^6$, $P < 0.025$). Non-TNF α -treated mice displayed undetectable ($<0.05\%$, $n = 22$) levels of human CD45⁺ cell engraftment in the thymus whereas TNF α treated mice had readily detectable human CD45⁺ cells ($47.9 \pm 10.7\%$ and $8.6 \pm 2.3 \times 10^7$, $P < 0.001$, Fig. 2).

Thymocytopoiesis and Human CD3⁺ T-Cell Development in NOD-*scid* IL2 γ ^{null} Mice Treated with TNF α and Transplanted with Cultured Human UCB

Thymus—In contrast with non-TNF α -treated mice that generated no human thymic engraftment, mice treated with TNF α developed CD3⁺ T cells in the thymus (Fig. 3). The human CD3 thymocytes displayed a normal distribution of immature double-positive CD4⁺CD8⁺ thymocytes and mature single-positive CD4⁺ and CD8⁺ T cells. Histological analysis of thymic sections of TNF α -treated mice supports the flow cytometry data (Fig. 4).

Spleen—In contrast to non-TNF α -treated mice in which only very low levels of splenic CD3⁺ T cells were detected, CD3⁺ T cells were readily detected in the spleen of mice treated with TNF α (Table 4 and Fig. 5, $P < 0.05$). The CD3⁺ T cells were comprised of mostly TCR α / β ⁺ cells ($90.2 \pm 3.7\%$) with low levels of TCR γ / δ ⁺ cells ($2.6 \pm 0.6\%$). The splenic CD3⁺ population consisted of approximately equal percentages of single positive CD4⁺ (42.4 ± 7.3) and CD8⁺ (56.1 ± 6.9) T cells (Fig. 5). Treatment with TNF α did not alter the percent or number of CD20⁺ B cells.

Splenic sections of TNF α -treated mice displayed human CD45⁺ cells predominantly in the white pulp (Fig. 6). Human CD20⁺ B cells as well as human CD5⁺ T cells were also identified. Human B cell maturation was documented by the presence of human CD138⁺ plasma cells, which contain kappa and lambda light chains (Fig. 6).

Function of Human T and B Cells in TNF α -Treated NOD-*scid* IL2 γ ^{null} Mice Engrafted with Cultured Human UCB

To determine whether engrafted B cells were functional, we first quantified the concentrations of human Igs in the serum of mice engrafted 12 weeks earlier with cultured UCB. Levels of human IgM ranged from ~ 50 $\mu\text{g/ml}$ to >250 $\mu\text{g/ml}$ and levels of human IgG ranged from ~ 25 ($\mu\text{g/ml}$) to >250 ($\mu\text{g/ml}$). The level of both IgG and IgM was increased in mice immunized with Pneumovax and treated with BLyS (Fig. 7A). Low but detectable levels of IgM and IgG antibody to serotype 4 and serotype 14 of *Streptococcus pneumoniae* were detected following immunization with Pneumovax, and administration of recombinant human BLyS in mice boosted with Pneumovax significantly increased the concentration of IgM antibody to both serotypes (Fig. 7B).

Splenocytes obtained from TNF α -treated mice were cultured in the presence of human anti-CD3, anti-CD28, and anti-CD49d mAb for 64 hours. When adjusted for the number of human CD3⁺ cells in the cultures, the proliferation of human CD3⁺ cells in the spleen of engrafted TNF α -treated NOD-*scid* IL2 γ ^{null} mice was only slightly lower than that of freshly isolated human T cells (Fig. 8A). Finally, to determine if the functional T and B cells could collaborate to generate a T-dependent antibody response, mice were immunized with tetanus and boosted in the presence of recombinant human BLyS. Anti-tetanus antibody was detected in the serum of 2/7 of the immunized mice and 2/3 of mice boosted in the presence of BLyS (Fig. 8B).

Discussion

We have investigated the functional potential of human cord blood CD34⁺ cells expanded in culture for two weeks to generate *in vivo* a multi-lineage human hematopoietic system and a functional human immune system. Cultured human UCB generated myeloid, erythroid, and lymphoid-lineage cells in NOD-*scid* IL2 γ ^{null} mice. Administration of TNF α facilitated the generation of human T cells. The immune system of these engrafted mice was functional as assessed by the presence of human Ig, and the ability to generate antibodies in response to immunization with either T-independent or T-dependent antigens.

Multiple sources of human HSC are now used for clinical transplantation, including mobilized peripheral blood hematopoietic stem cells, bone marrow, and umbilical cord blood (47,48). However, cord blood is emerging as the HSC source of choice due to the low cellular immunity of the passenger leukocytes, and the ease of identifying HLA-matches due to the observation that cord blood can be transplanted with a greater HLA disparity than can HSC from mobilized stem cells or bone marrow (8,11). One remaining barrier is that clinical transplantation of human UCB cells into adults is limited by the low yield of CD34⁺ cells from a single cord. Currently, adult transplantation of the required CD34⁺ cell dose necessitates the pooling of multiple cords, increasing the risk of GVHD as well as the need for matching to multiple donors [for review, see (49,50)]. Many approaches for *ex vivo* expansion of cord blood cells have been reported (21–25,51–55). In the culture system used in our studies, a substantial proportion of the cells phenotypically appeared to retain multi-lineage potential. To assess their *in vivo* potential, we adoptively transferred the cells into conditioned NOD-*scid* IL2 γ ^{null} mice. The use of this stock of mice was used based on 1) earlier observations showing that human HSC engraft better than in NOD-*scid* mice (36,56) and 2) our initial experiment comparing directly the ability of cultured HSC to engraft in these two strains. Although NOD-*scid* IL2 γ ^{null} mice is the strain we chose for these *in vivo* analyses of human HSC, other strains of immunodeficient mice based on mutations in the IL-2 receptor gamma chain have been reported (34). We recently compared engraftment in BALB/c-*Rag*^{null} IL2 γ ^{null} mice with that of NOD-*scid* IL2 γ ^{null} mice, and have determined that NOD-*scid* IL2 γ ^{null} mice are much better recipients of human HSC (TP, unpublished observations). Moreover, additional modification of NOD-*scid* IL2 γ ^{null} mice is currently underway, including the introduction of transgenes expressing human HLA (57).

Previous studies comparing the *in vivo* engraftment kinetics and developmental potential of cultured UCB cells in immunodeficient mice have reported that cultured cells exhibit delayed engraftment (32,33). In our study, cultured HSC exhibited STRC capacity as evidenced by their ability to generate human myeloid and lymphoid cells within 3 weeks after transplantation. The cultured cells also exhibited LTRC capacity and serial transfer and engraftment of the bone marrow into secondary NOD-*scid* IL2 γ ^{null} recipients suggests the maintenance of true HSC in the cultured population.

Few human T cells are observed following the injection of cultured HSC into NOD-*scid* mice (23,32,33). In those studies, the percentages of human T cells were low, and the possibility of

contaminating human T cells in the injected cell population was not excluded. In one study, cells recovered from the mice required *ex vivo* expansion in the presence of PHA and IL-2 to detect T cells (23). We also observed that few cultured HSC develop into T cells in immunodeficient mice.

We hypothesized that the cultured cells may fail to produce TNF α , which increases T-cell lineage commitment and differentiation of human stem cells in NOD-*scid* mice transplanted with HSC (46). Administration of exogenous TNF α prior to transplantation of cultured HSC augmented human thymus cell engraftment. These thymocytes migrated to the peripheral tissues leading to the generation of mature CD4 and CD8 T cells. How TNF α augments T-cell development is not known, but it has been reported that TNF enhances human T-cell generation in a xenogeneic fetal thymus organ culture, suggesting a direct effect on the engrafted stem cells (58). However, experiments where cultured CD34⁺ cells were treated with TNF α prior to injection into NOD-*scid* IL2 γ ^{null} mice, no increase in T-cell development was observed (LJG and KB, unpublished observations). Alternatively, our data are consistent with the interpretation that TNF could have an indirect effect on the host environment in which the stem cells mature, possibly by activation of innate immunity and the induction of cytokines (59). Multiple cytokines have been shown to enhance human HSC engraftment and differentiation in immunodeficient mice [recently reviewed in (57)].

The human immune system generated by cultured HSC in NOD-*scid* IL2 γ ^{null} mice was functional. Human IgG and IgM were detected in the circulation, and the mice were able to generate both T-independent and T-dependent antibodies in response to immunization with Pneumovax and tetanus, respectively. We note, however, that we only assayed the antibodies generated to 2 of the 23 serotypes of Pneumovax. We chose these two serotypes to analyze because they are two serotypes shared by both the pneumovax23 adult vaccine and the protein conjugated childhood vaccine (38). Also, they elicit different responses in young versus older individuals: the response to serotype 4 is down in aged individuals whereas the response to serotype 14 remains high (60). The ability to generate antibodies following immunization confirms the phenotypic and histological data showing that T cells, B cells, myeloid, and dendritic cells of human origin were generated. Equally important is that the allogeneic immune system that developed from the HSC in the expanded cord blood is tolerant to host, and could lead to the induction of donor-specific tolerance (61,62) or for the treatment of autoimmune disorders (63–66).

In summary, human CD34⁺ UCB cells expanded for two weeks in culture engraft and differentiate into multiple hematopoietic lineages in NOD-*scid* IL2 γ ^{null} mice and generate a functional human immune system. *Ex vivo* expansion of UCB CD34⁺ stem cells may be a promising alternative to pooling multiple cords for adult hematopoietic stem cell transplantation.

Acknowledgments

We thank Linda Paquin, Amy Cuthbert, Jean Leif, and Veronica Zielinski for their technical assistance.

This work was supported by National Institutes of Health Grants HL77642, AI46629, an institutional Diabetes Endocrinology Research Center (DERC) grant DK32520, a Cancer Center Core grant CA34196, the Beta Cell Biology Consortium, and the Juvenile Diabetes Foundation, International. Joseph Laning and Morey Kraus are employees of Viacell, Inc. Kristin Biber is a former employee of Viacell, Inc. Dale Greiner is a consultant and has grant support from Viacell, Inc. Leonard Shultz is a consultant for Viacell, Inc. The contents of this publication are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health.

References

1. Armitage JO. Bone marrow transplantation. N Engl J Med 1994;330:827–838. [PubMed: 8114836]

2. Handgretinger R, Klingebiel T, Lang P, Schumm M, Neu S, Geiselhart A, Bader P, Schlegel PG, Greil J, Stachel D, Herzog RJ, Niethammer D. Megadose transplantation of purified peripheral blood CD34⁺ progenitor cells from HLA-mismatched parental donors in children. *Bone Marrow Transplant* 2001;27:777–783. [PubMed: 11477433]
3. Woodard P, Cunningham JM, Benaim E, Chen X, Hale G, Horwitz E, Houston J, Kasow K, Leung W, Wang W, Yusuf U, Handgretinger R. Effective donor lymphohematopoietic reconstitution after haploidentical CD34⁺-selected hematopoietic stem cell transplantation in children with refractory severe aplastic anemia. *Bone Marrow Transplant* 2004;33:411–418. [PubMed: 14676782]
4. Lu L, Shen R-N, Broxmeyer HE. Stem cells from bone marrow, umbilical cord blood and peripheral blood for clinical application: current status and future application. *Crit Rev Oncol-Hematol* 1996;22:61–78. [PubMed: 8679101]
5. Broxmeyer HE, Douglas GW, Hangoc G, Cooper S, Bard J, English D, Arny M, Thomas L, Boyse EA. Human umbilical cord blood as a potential source of transplantable hematopoietic stem/progenitor cells. *Proc Natl Acad Sci USA* 1989;86:3828–3832. [PubMed: 2566997]
6. Benito AI, Diaz MA, Gonzalez-Vicent M, Sevilla J, Madero L. Hematopoietic stem cell transplantation using umbilical cord blood progenitors: review of current clinical results. *Bone Marrow Transplant* 2004;33:675–690. [PubMed: 14981535]
7. Laughlin MJ, Barker J, Bambach B, Koc ON, Rizzieri DA, Wagner JE, Gerson SL, Lazarus HM, Cairo M, Stevens CE, Rubinstein P, Kurtzberg J. Hematopoietic engraftment and survival in adult recipients of umbilical-cord blood from unrelated donors. *N Engl J Med* 2001;344:1815–1822. [PubMed: 11407342]
8. Bradley MB, Cairo MS. Cord blood immunology and stem cell transplantation. *Hum Immunol* 2005;66:431–446. [PubMed: 15935881]
9. Holyoake TL, Nicolini FE, Eaves CJ. Functional differences between transplantable human hematopoietic stem cells from fetal liver, cord blood, and adult marrow. *Exp Hematol* 1999;27:1418–1427. [PubMed: 10480433]
10. Brunstein CG, Wagner JE. Umbilical cord blood transplantation and banking. *Annu Rev Med* 2006;57:403–417. [PubMed: 16409157]
11. Rubinstein P. Why cord blood? *Hum Immunol* 2006;67:398–404. [PubMed: 16728260]
12. Schoemans H, Theunissen K, Maertens J, Boogaerts M, Verfaillie C, Wagner J. Adult umbilical cord blood transplantation: a comprehensive review. *Bone Marrow Transplant* 2006;38:83–93. [PubMed: 16751788]
13. Majhail NS, Branstein CG, Wagner JE. Double umbilical cord blood transplantation. *Curr Opin Immunol* 2006;18:571–575. [PubMed: 16895752]
14. Gluckman E, Rocha V, Boyer-Chamard A, Locatelli F, Arcese W, Pasquini R, Ortega J, Souillet G, Ferreira E, Laporte JP, Fernandez M, Chastang C. Outcome of cord-blood transplantation from related and unrelated donors. Eurocord Transplant Group and the European Blood and Marrow Transplantation Group. *N Engl J Med* 1997;337:373–381. [PubMed: 9241126]
15. Rubinstein P, Carriet C, Scaradavou A, Kurtzberg J, Adamson J, Migliaccio AR, Berkowitz RL, Cabbad M, Dobrila NL, Taylor PE, Rosenfield RE, Stevens CE. Outcomes among 562 recipients of placental blood transplants from unrelated donors. *N Engl J Med* 1998;339:1565–1569. [PubMed: 9828244]
16. Wagner JE, Barker JN, DeFor TE, Baker KS, Blazar BR, Eide C, Goldman A, Kersey J, Krivit W, MacMillan ML, Orchard PJ, Peters C, Weisdorf DJ, Ramsay NK, Davies SM. Transplantation of unrelated donor umbilical cord blood in 102 patients with malignant and nonmalignant diseases: influence of CD34 cell dose and HLA disparity on treatment-related mortality and survival. *Blood* 2002;100:1611–1618. [PubMed: 12176879]
17. Weinreb S, Delgado JC, Clavijo OP, Yunis EJ, Bayer-Zwirello L, Polansky L, Deluhery L, Cohn G, Yao JT, Stec TC, Higby D, Anderzejewski C. Transplantation of unrelated cord blood cells. *Bone Marrow Transplant* 1998;22:193–196. [PubMed: 9707029]
18. Barker JN, Wagner JE. Umbilical cord blood transplantation: current practice and future innovations. *Crit Rev Oncol Hematol* 2003;48:35–43. [PubMed: 14585482]
19. de Lima M, St John LS, Wieder ED, Lee MS, McMannis J, Karandish S, Giralt S, Beran M, Couriel D, Korbliing M, Bibawi S, Champlin R, Komanduri KV. Double-chimaerism after transplantation of

- two human leucocyte antigen mismatched, unrelated cord blood units. *Br J Haematol* 2002;119:773–776. [PubMed: 12437658]
20. Barker JN, Weisdorf DJ, DeFor TE, Blazar BR, Miller JS, Wagner JE. Rapid and complete donor chimerism in adult recipients of unrelated donor umbilical cord blood transplantation after reduced-intensity conditioning. *Blood* 2003;102:1915–1919. [PubMed: 12738676]
 21. Hofmeister CC, Zhang J, Knight KL, Le P, Stiff PJ. Ex vivo expansion of umbilical cord blood stem cells for transplantation: growing knowledge from the hematopoietic niche. *Bone Marrow Transplant* 2007;39:11–23. [PubMed: 17164824]
 22. Mohamed AA, Ibrahim AM, El Masry MW, Mansour IM, Khroshied MA, Gouda HM, Riad RM. Ex vivo expansion of stem cells: defining optimum conditions using various cytokines. *Lab Hematol* 2006;12:86–93. [PubMed: 16751136]
 23. Kobari L, Giarratana MC, Gluckman JC, Douay L, Rosenzweig M. *Ex vivo* expansion does not alter the capacity of umbilical cord blood CD34⁺ cells to generate functional T lymphocytes and dendritic cells. *Stem Cells* 2006;24:2150–2157. [PubMed: 16946001]
 24. Vanheusden K, Van Coppenolle S, De Smedt M, Plum J, Vandekerckhove B. *In vitro* expanded cells contributing to rapid severe combined immunodeficient repopulation activity are CD34⁺38[−]33⁺90⁺45RA[−]. *Stem Cells* 2007;25:107–114. [PubMed: 16973833]
 25. Ito Y, Hasaouda H, Kitajima T, Kiyono T. *Ex vivo* expansion of human cord blood hematopoietic progenitor cells using glutaraldehyde-fixed human bone marrow stromal cells. *J Biosci Bioeng* 2006;102:467–469. [PubMed: 17189177]
 26. Petzer AL, Zandstra PW, Piret JM, Eaves CJ. Differential cytokine effects on primitive (CD34⁺CD38[−]) human hematopoietic cells: novel responses to Flt3-ligand and thrombopoietin. *J Exp Med* 1996;183:2551–2558. [PubMed: 8676076]
 27. McNiece I, Kubegov D, Kerzic P, Shpall EJ, Gross S. Increased expansion and differentiation of cord blood products using a two-step expansion culture. *Exp Hematol* 2000;28:1181–1186. [PubMed: 11027837]
 28. McNiece I, Jones R, Bearman SI, Cagnoni P, Nieto Y, Franklin W, Ryder J, Steele A, Stoltz J, Russell P, McDermitt J, Hogan C, Murphy J, Shpall EJ. *Ex vivo* expanded peripheral blood progenitor cells provide rapid neutrophil recovery after high-dose chemotherapy in patients with breast cancer. *Blood* 2000;96:3001–3007. [PubMed: 11049977]
 29. Conneally E, Cashman J, Petzer A, Eaves C. Expansion *in vitro* of transplantable human cord blood stem cells demonstrated using a quantitative assay of their lympho- myeloid repopulating activity in nonobese diabetic-*scid/scid* mice. *Proc Natl Acad Sci USA* 1997;94:9836–9841. [PubMed: 9275212]
 30. Ueda T, Tsuji K, Yoshino H, Ebihara Y, Yagasaki H, Hisakawa H, Mitsui T, Manabe A, Tanaka R, Kobayashi K, Ito M, Yasukawa K, Nakahata T. Expansion of human NOD/SCID-repopulating cells by stem cell factor, Flk2/Flt3 ligand, thrombopoietin, IL-6, and soluble IL-6 receptor. *J Clin Invest* 2000;105:1013–1021. [PubMed: 10749580]
 31. Bhatia M, Wang JCY, Kapp U, Bonnet D, Dick JE. Purification of primitive human hematopoietic cells capable of repopulating immune-deficient mice. *Proc Natl Acad Sci USA* 1997;94:5320–5325. [PubMed: 9144235]
 32. Chute JP, Muramoto G, Fung J, Oxford C. Quantitative analysis demonstrates expansion of SCID-repopulating cells and increased engraftment capacity in human cord blood following *ex vivo* culture with human brain endothelial cells. *Stem Cells* 2004;22:202–215. [PubMed: 14990859]
 33. Guenechea G, Segovia JC, Albella B, Lamana M, Ramirez M, Regidor C, Fernandez MN, Bueren JA. Delayed engraftment of nonobese diabetic/severe combined immunodeficient mice transplanted with *ex vivo*-expanded human CD34⁺ cord blood cells. *Blood* 1999;93:1097–1105. [PubMed: 9920860]
 34. Shultz LD, Ishikawa F, Greiner DL. Humanized mice in translational biomedical research. *Nat Rev Immunol* 2007;7:118–130. [PubMed: 17259968]
 35. Shultz LD, Schweitzer PA, Christianson SW, Gott B, Schweitzer IB, Tennent B, McKenna S, Mobraaten L, Rajan TV, Greiner DL, Leiter EH. Multiple defects in innate and adaptive immunologic function in NOD/LtSz-*scid* mice. *J Immunol* 1995;154:180–191. [PubMed: 7995938]
 36. Shultz LD, Lyons BL, Burzenski LM, Gott B, Chen X, Chaleff S, Gillies SD, King M, Mangada J, Greiner DL, Handgretinger R. Human lymphoid and myeloid cell development in NOD/LtSz-

- scidIL2 γ ^{null}* mice engrafted with mobilized human hematopoietic stem cell. *J Immunol* 2005;174:6477–6489. [PubMed: 15879151]
37. Shultz LD, Banuelos S, Lyons B, Samuels R, Burzenski L, Gott B, Lang P, Leif J, Appel M, Rossini A, Greiner DL. NOD/LtSz-Rag1^{null}Prf1^{null} mice: a new model system with increased levels of human peripheral leukocyte and hematopoietic stem cell engraftment. *Transplantation* 2003;76:1036–1042. [PubMed: 14557749]
 38. Wernette CM, Frasc CE, Madore D, Carlone G, Goldblatt D, Plikaytis B, Benjamin W, Quataert SA, Hildreth S, Sikkema DJ, Kayhty H, Jonsdottir I, Nahm MH. Enzyme-linked immunosorbent assay for quantitation of human antibodies to pneumococcal polysaccharides. *Clin Diagn Lab Immunol* 2003;10:514–519. [PubMed: 12853378]
 39. Kollet G, Peled A, Byk T, Ben Hur H, Greiner D, Shultz L, Lapidot T. β 2 microglobulin-deficient (B2m^{null}) NOD/SCID mice are excellent recipients for studying human stem cell function. *Blood* 2000;95:3102–3105. [PubMed: 10807775]
 40. Greiner DL, Shultz LD, Yates J, Appel MC, Perdriest G, Hesselton RM, Schweitzer I, Beamer WG, Shultz KL, Pelsue SC, Leif JH, Rajan TV. Improved engraftment of human spleen cells in NOD/LtSz-*scid/scid* mice as compared with C.B-17-*scid/scid* mice. *Am J Path* 1995;146:888–902. [PubMed: 7717456]
 41. Christianson SW, Greiner DL, Hesselton RM, Leif JH, Wagar EJ, Schweitzer IB, Rajan TV, Gott B, Roopenian DC, Shultz LD. Enhanced human CD4⁺ T cell engraftment in β 2-microglobulin-deficient NOD-*scid* mice. *J Immunol* 1997;158:3578–3586. [PubMed: 9103418]
 42. Christianson SW, Greiner DL, Schweitzer IB, Gott B, Beamer GL, Schweitzer PA, Hesselton RM, Shultz LD. Role of natural killer cells on engraftment of human lymphoid cells and on metastasis of human T-lymphoblastoid leukemia cells in C57BL/6J-*scid* mice and in C57BL/6J-*scidbg* mice. *Cell Immunol* 1996;171:186–199. [PubMed: 8806787]
 43. Lowry PA, Shultz LD, Greiner DL, Hesselton RA, Kittler ELW, Tiarks CY, Rao SS, Reilly J, Leif JH, Ramshaw H, Steward FM, Quesenberry PJ. Improved engraftment of human cord blood stem cells in NOD/LtSz-*scid/scid* mice following irradiation or multiple day injections into unirradiated recipients. *Biol Blood Marrow Transplant* 1996;2:15–23. [PubMed: 9078350]
 44. Glimm H, Eisterer W, Lee K, Cashman J, Holyoake TL, Nicolini F, Shultz LD, von Kalle C, Eaves CJ. Previously undetected human hematopoietic cell populations with short-term repopulating activity selectively engraft NOD/SCID-beta2 microglobulin-null mice. *J Clin Invest* 2001;107:199–206. [PubMed: 11160136]
 45. Shultz LD, Banuelos SJ, Leif J, Appel MC, Cunningham M, Ballen K, Burzenski L, Greiner DL. Regulation of human short-term repopulating cell (STRC) engraftment in NOD/SCID mice by host CD122⁺ cells. *Exp Hematol* 2003;31:551–558. [PubMed: 12829032]
 46. Samira S, Ferrand C, Peled A, Nagler A, Tovbin Y, Ben Hur H, Taylor N, Globerson A, Lapidot T. Tumor necrosis factor promotes human T-cell development in nonobese diabetic/severe combined immunodeficient mice. *Stem Cells* 2004;22:1085–1100. [PubMed: 15536198]
 47. Ringden O, Le Blanc K. Allogeneic hematopoietic stem cell transplantation: state of the art and new perspectives. *APMIS* 2005;113:813–830. [PubMed: 16480452]
 48. Aschan J. Allogeneic haematopoietic stem cell transplantation: current status and future outlook. *Br Med Bull* 2006;77–78:23–36.
 49. Brunstein CG, Baker KS, Wagner JE. Umbilical cord blood transplantation for myeloid malignancies. *Curr Opin Hematol* 2007;14:162–169. [PubMed: 17255794]
 50. Haspel RL, Ballen KK. Double cord blood transplants: filling a niche? *Stem Cell Rev* 2006;2:81–86. [PubMed: 17237545]
 51. Gan OI, Murdoch B, Larochelle A, Dick JE. Differential maintenance of primitive human SCID-repopulating cells, clonogenic progenitors, and long-term culture-initiating cells after incubation on human bone marrow stromal cells. *Blood* 1997;90:641–650. [PubMed: 9226164]
 52. Yamaguchi M, Hirayama F, Kanai M, Sato N, Fukazawa K, Yamashita K, Sawada K, Koike T, Kuwabara M, Ikeda H, Ikebuchi K. Serum-free coculture system for *ex vivo* expansion of human cord blood primitive progenitors and SCID mouse-reconstituting cells using human bone marrow primary stromal cells. *Exp Hematol* 2001;29:174–182. [PubMed: 11166456]

53. Zhang Y, Li C, Jiang X, Zhang S, Wu Y, Liu B, Tang P, Mao N. Human placenta-derived mesenchymal progenitor cells support culture expansion of long-term culture-initiating cells from cord blood CD34⁺ cells. *Exp Hematol* 2004;32:657–664. [PubMed: 15246162]
54. Wang JF, Wang LJ, Wu YF, Xiang Y, Xie CG, Jia BB, Harrington J, McNiece IK. Mesenchymal stem/progenitor cells in human umbilical cord blood as support for *ex vivo* expansion of CD34⁺ hematopoietic stem cells and for chondrogenic differentiation. *Haematologica* 2004;89:837–844. [PubMed: 15257936]
55. Xie CG, Wang JF, Xiang Y, Qiu LY, Jia BB, Wang LJ, Wang GZ, Huang GP. Cocultivation of umbilical cord blood CD34⁺ cells with retro-transduced hMSCs leads to effective amplification of long-term culture-initiating cells. *World J Gastroenterol* 2006;12:393–402. [PubMed: 16489638]
56. Ishikawa F, Yasukawa M, Lyons B, Yoshida S, Miyamoto T, Yoshimoto G, Watanabe T, Akashi K, Shultz LD, Harada M. Development of functional human blood and immune systems in NOD/SCID/IL2 receptor gamma chain null mice. *Blood* 2005;106:1565–1573. [PubMed: 15920010]
57. Pearson T, Shultz LD, Greiner DL. Humanized SCID mice for biomedical research.
58. Smits K, De Smedt M, Naessens E, De Smet G, Stove V, Taghon T, Plum J, Verhasselt B. Tumor necrosis factor promotes T-cell at the expense of B-cell lymphoid development from cultured human CD34⁺ cord blood cells. *Exp Hematol* 2007;35:1272–1278. [PubMed: 17553612]
59. Ware CF. Network communications: lymphotoxins, LIGHT, and TNF. *Annu Rev Immunol* 2005;23:787–819. [PubMed: 15771586]
60. Shriner AK, Smithson SL, Prinz DM, Rabquer B, Khuder S, Goomber R, Westerink MA. Comparison of the human immune response to conjugate and polysaccharide pneumococcal vaccination using a reconstituted SCID mouse model. *Vaccine* 2006;24:7197–7203. [PubMed: 16860443]
61. Scandling JD, Busque S, Dejbakhsh-Jones S, Benike C, Millan MT, Shizuru JA, Hoppe RT, Lowsky R, Engleman EG, Strober S. Tolerance and chimerism after renal and hematopoietic-cell transplantation. *N Engl J Med* 2008;358:362–368. [PubMed: 18216356]
62. Kawai T, Cosimi AB, Spitzer TR, Tolkoff-Rubin N, Suthanthiran M, Saidman SL, Shaffer J, Preffer FI, Ding R, Sharma V, Fishman JA, Dey B, Ko DS, Hertl M, Goes NB, Wong W, Williams WW Jr, Colvin RB, Sykes M, Sachs DH. HLA-mismatched renal transplantation without maintenance immunosuppression. *N Engl J Med* 2008;358:353–361. [PubMed: 18216355]
63. Sorli CH, Greiner DL, Mordes JP, Rossini AA. Stem cell transplantation for treatment of autoimmune diseases. *Graft* 1998;1:71–81.
64. Passweg J, Tyndall A. Autologous stem cell transplantation in autoimmune diseases. *Semin Hematol* 2007;44:278–285. [PubMed: 17961728]
65. Alderuccio F, Siatskas C, Chan J, Field J, Murphy K, Nasa Z, Toh BH. Haematopoietic stem cell gene therapy to treat autoimmune disease. *Curr Stem Cell Res Ther* 2006;1:279–287. [PubMed: 18220873]
66. Marmont AM. Will hematopoietic stem cell transplantation cure human autoimmune diseases? *J Autoimmun* 2008;30:145–150. [PubMed: 18222646]

Figure 1.

Lineage marker expression on engrafted human CD45⁺ cells in NOD-*scid* IL2 γ ^{null} mice. NOD-*scid* IL2 γ ^{null} mice were transplanted with 1×10^6 cultured CD34⁺ human UCB cells. Bone marrow was analyzed by flow cytometry for the presence of human myeloid and lymphoid lineages three (A) and twelve (B) weeks after transplantation and spleen was analyzed twelve weeks (C) after transplantation. Total human cells are shown as the percent positive human CD45⁺ cells. Lineage markers are expressed as percent of human CD45⁺ cells in the bone marrow. CD71 and GlyA populations are expressed as percent of murine CD45⁺ cells. Shown are representative histograms.

Figure 2.

Engraftment of human CD45⁺ cells in non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ ^{null} mice 12 weeks after transplantation of cultured human UCB. Irradiated non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ ^{null} mice were transplanted with 1×10^6 cultured CD34⁺ human UCB cells. (A) Percent and (B) number of human CD45⁺ cells in blood, bone marrow, spleen, and thymus twelve weeks after transplantation. * $P < 0.025$, ** $P < 0.01$, *** $P < 0.001$.

Figure 3.

Phenotype of human CD45⁺ cells in the thymus of non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ ^{null} twelve weeks after transplantation of cultured human UCB. Irradiated non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ ^{null} mice were transplanted with 1×10^6 cultured CD34⁺ human UCB cells. Twelve weeks later, thymi were analyzed by flow cytometry. Representative flow cytometric profiles of human CD45⁺ cell development in (A) non-TNF α -treated and (B) TNF α -treated NOD-*scid* IL2 γ ^{null} mice. Human CD3 percentages are expressed as percent human CD45⁺ cells. Human CD4 and CD8 percentages are expressed as percent human CD3⁺ cells.

Figure 4.

Immunohistochemistry of thymi of TNF α -treated NOD-*scid* IL2 γ^{null} mice twelve weeks after transplantation of cultured human UCB. Irradiated TNF α -treated NOD-*scid* IL2 γ^{null} mice were transplanted with 1×10^6 cultured CD34 $^{+}$ human UCB cells. Twelve weeks later, thymi were analyzed by immunohistochemistry. Thymic sections of TNF α -treated mice contained human CD45 $^{+}$ cells (A), which were localized predominantly in the subcapsular region. Many of these CD45 $^{+}$ cells were CD5 $^{+}$ (B) and expressed either CD4 $^{+}$ (C) and/or CD8 $^{+}$ (D). In addition, CD1a $^{+}$ thymocytes were detected residing in the cortex of the thymus (E). TdT $^{+}$ (F) and CD99 $^{+}$ (G) thymocytes were also present, as were numerous human CD45 $^{+}$ cells that expressed the proliferation antigen Ki-67 (H).

Figure 5.

Human CD45⁺ cell development in the spleen of non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ^{null} mice twelve weeks after transplantation of cultured human UCB. Irradiated non-TNF α -treated and TNF α -treated NOD-*scid* IL2 γ^{null} mice were transplanted with 1×10^6 cultured CD34⁺ human UCB cells. Twelve weeks later, spleen cells were analyzed by flow cytometry. (A) Percentage of human CD3⁺ and CD20⁺ cells in the spleen. Representative flow cytometry profiles of human cell development in (B) non-TNF α -treated and (C) TNF α -treated NOD-*scid* IL2 γ^{null} mice. Lineage markers expressed as percent of human CD45⁺ cells. Human CD4 and CD8 percentages are expressed as percent of human CD3⁺ cells.

Figure 6.

Immunohistochemistry of spleens of TNF α -treated NOD-*scid* IL2 γ^{null} mice twelve weeks after transplantation of cultured human UCB. TNF α -treated NOD-*scid* IL2 γ^{null} mice were transplanted with 1×10^6 cultured CD34 $^{+}$ human UCB cells. Twelve weeks later, the distribution of human lymphocytes in the spleen was determined by immunohistochemistry. Immunoperoxidase staining of human (A) CD45 $^{+}$, (B) CD20 $^{+}$, (C) CD5 $^{+}$; (D, arrows) CD138 $^{+}$, (E, arrows) kappa $^{+}$ and (F, arrows) lambda $^{+}$ cells in spleens of TNF α -treated NOD-*scid* IL2 γ^{null} mice.

Figure 7.

Human immunoglobulin levels and T-independent immune responses in TNF α -treated NOD-*scid* IL2 γ^{null} mice twelve weeks after transplantation of cultured human UCB. TNF α -treated NOD-*scid* IL2 γ^{null} mice were transplanted with 1×10^6 cultured CD34 $^{+}$ human UCB cells. Represented by the dashed line in each panel is the upper limit of the mean of pre-immune sera plus 3 standard deviations. All values above this limit were considered “responders.” The treatment group and the number of responders for each treatment group are shown at the bottom of each panel. Each symbol represents an individual animal. Panel A: Human IgG and IgM antibodies prior to and following Pneumovax immunization in the presence or absence of BLYS treatment. Panel B: Human IgG and IgM antibodies to serotype 4 and 14 following Pneumovax

immunization in the presence or absence of BLyS treatment. Each symbol represents and individual animal. * $P < 0.05$, ** $P < 0.001$.

Figure 8. Human CD3 proliferation and human T-dependent immune responses in TNF α -treated NOD-*scid* IL2 γ^{null} mice twelve weeks after transplantation of cultured human UCB. TNF α -treated NOD-*scid* IL2 γ^{null} mice were transplanted with 1×10^6 cultured CD34 $^{+}$ human UCB cells. Panel A: Proliferation of human CD3 T cells following *ex vivo* stimulation with anti-CD3 twelve weeks after transplantation. Shown are results of triplicate cultures of spleen cells from individual mice displayed as mean $\pm$ SD. Panel B: Human IgG antibodies to tetanus following immunization with tetanus toxoid. Each symbol represents an individual animal. Represented by the dashed line in each panel is the upper limit of the mean of pre-immune sera plus 3 standard deviations. All values above this limit were considered “responders.” The treatment group and the number of responders for each treatment group are shown at the bottom of the panel. Each symbol represents an individual animal.

Table 1**CD34⁺ Cell Expansion in Two-Week Culture^a**

Cell phenotype	Cell input	Cell yield after two week culture	Fold increase in cell number
CD45 cells	$3.8 \pm 0.9 \times 10^6$	$2.6 \pm 1.1 \times 10^8$	68.4-fold
CD34 cells	$7.1 \pm 1.3 \times 10^5$	$2.6 \pm 0.9 \times 10^7$	36.6-fold

^aHuman UCB was cultured as described in Materials and Methods. The number of each cell population after RBC and lineage depletion that was put into culture or recovered after two weeks of culture is shown. Shown are the numbers of CD45⁺ cells (total human leukocytes) and total human CD45⁺CD34⁺ cells (total human CD34⁺ cells). Results represent the mean $\pm$ SEM of at least 30 separate cultures.

Table 2Phenotype of Human UCB Prior to and After a Two-Week Culture^a

Phenotype	Cell population	Input cell population (mean ± SEM)	Cultured cell population (mean ± SEM)
CD34	Hematopoietic progenitor	32.3 ± 3.9	20.1 ± 1.9
CD34/CD33/CD13	Myelomonocytic stem cell	0.1 ± 0.1	12.9 ± 1.2
CD33/CD13	General myeloid precursor	9.0 ± 2.5	53.6 ± 2.7
CD34/CD7	Lymphoid/NK progenitor	0.6 ± 0.4	1.2 ± 0.3
CD41/CD61	Megakaryocyte	13.5 ± 1.8	23.0 ± 1.8
CD34/CD41/CD61	Megakaryocyte progenitor	2.5 ± 1.0	2.7 ± 0.5
CD3	T lymphoid lineage	0.3 ± 0.1	0.8 ± 0.2
CD14	Monocyte	0.3 ± 0.11	4.2 ± 1.0
CD19	B lymphoid lineage	0.9 ± 0.3	0.4 ± 0.1
GlyA	Immature and mature erythrocyte	3.7 ± 2.1	1.6 ± 0.7

^a Human UCB was cultured as described in Materials and Methods. The relative percentage of each cell population after RBC and lineage depletion that was put into culture or recovered after two weeks of culture is shown. Results represent the mean ± SEM of at least 20 separate cultures. All values represent the percent mean ± SEM of human CD45⁺ cells with the exception of GlyA, which were derived from the CD45⁺ population.

Table 3

Lineage Marker Expression on Engrafted Human CD45⁺ Cells in the Bone Marrow of NOD-*scid* IL2 γ ^{null} Mice Three Weeks After Transplantation of Cultured Human UCB^a

Lineage	Phenotype	Percent (mean $\pm$ SEM, <i>n</i> = 11)
Phenotype of CD45 ⁺ cells	CD45	2.3 $\pm$ 0.4
	CD34	17.1 $\pm$ 1.4
	CD33	87.3 $\pm$ 4.4
	CD38	94.1 $\pm$ 1.2
	CD33CD64	23.5 $\pm$ 2.6
	CD20	11.1 $\pm$ 1.7
	CD3	<0.05
	CD71	0.1 $\pm$ 0.1
	GlyA	0.1 $\pm$ 0.1

^a Irradiated NOD-*scid* IL2 γ ^{null} mice were transplanted with cultured human UCB containing 1×10^6 CD34⁺ cells. Bone marrow was analyzed by flow cytometry for the presence of human myeloid and lymphoid lineages three weeks after transplantation. Shown is the percent of human CD45⁺ bone marrow cells and lineage phenotype data are expressed as percent of human CD45⁺ cells.

Lineage Marker Expression on Engrafted Human CD45⁺ Cells in Non-TNF α -Treated and TNF α -Treated NOD-*scid* IL2r γ^{null} Mice Twelve Weeks After Transplantation of Cultured Human UCB^a

Lineage	Phenotype (% of CD45 ⁺ cells)	Bone marrow		Spleen	
		No TNF α (mean $\pm$ SEM, <i>n</i> = 26)	TNF α (mean $\pm$ SEM, <i>n</i> = 19)	No TNF α (mean $\pm$ SEM, <i>n</i> = 12)	TNF α (mean $\pm$ SEM, <i>n</i> = 11)
Stem/progenitor	CD34	10.3 $\pm$ 1.0	9.7 $\pm$ 1.6	N.D.	N.D.
	CD38/CD34	9.9 $\pm$ 1.1	8.7 $\pm$ 1.3	N.D.	N.D.
	CD33	30.0 $\pm$ 3.6	27.5 $\pm$ 3.6	6.0 $\pm$ 1.7	6.4 $\pm$ 2.1
Myeloid	CD33/CD64	12.1 $\pm$ 2.1	7.2 $\pm$ 1.1	<0.05	<0.05
	CD14	5.1 $\pm$ 0.5	6.0 $\pm$ 0.8	<0.05	0.1 $\pm$ 0.1
	CD38	93.0 $\pm$ 2.1	95.1 $\pm$ 1.4	N.D.	N.D.
	CD41a	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	N.D.	N.D.
	B2CA2	3.3 $\pm$ 0.3	3.9 $\pm$ 0.7	N.D.	N.D.
Lymphoid	CD71	0.4 $\pm$ 0.3	2.0 $\pm$ 0.8	N.D.	N.D.
	CD20	26.5 $\pm$ 1.7	21.2 $\pm$ 1.9	81.9 $\pm$ 2.0	70.2 $\pm$ 3.8
	CD20/IgM	15.4 $\pm$ 1.1	11.6 $\pm$ 1.3	66.0 $\pm$ 1.8	49.3 $\pm$ 4.1
	CD20/CD10	14.2 $\pm$ 1.7	11.4 $\pm$ 2.4	N.D.	N.D.
	CD3	<0.05	<0.05	0.1 $\pm$ 0.1	10.2 $\pm$ 6.2
Erythroid	CD56	2.4 $\pm$ 0.5	3.6 $\pm$ 1.0	<0.05	0.1 $\pm$ 0.1
	GlyA	0.5 $\pm$ 0.1	1.7 $\pm$ 0.4	N.D.	N.D.

^a Irradiated non-TNF α -treated and TNF α -treated NOD-*scid* IL2r γ^{null} mice were transplanted with cultured human UCB containing 1×10^6 CD34⁺ cells. Bone marrow and spleen were analyzed by flow cytometry for the presence of human myeloid and lymphoid lineages twelve weeks after transplantation. Data are expressed as percent of human CD45⁺ cells; *n* = 9–12/group; N.D., not determined.