Published in final edited form as:

Curr Prostate Rep. 2008 May 1; 6(2): 67–73. doi:10.1007/s11918-008-0011-5.

Inflammatory Mechanisms Associated with Prostatic Inflammation and Lower Urinary Tract Symptoms

Jennifer L. St. Sauver, PhD and Steven J. Jacobsen, MD, PhD

Abstract

Inflammation is a common finding in histologic prostate specimens obtained from aging men, and accumulating data suggest that inflammation may play an important role in the development of benign prostatic hyperplasia (BPH), and the development and progression of lower urinary tract symptoms (LUTS). Inflammatory processes may contribute to prostatic enlargement directly through stimulation of prostate growth, or, alternatively, through decreasing prostatic apoptosis. Additionally, inflammatory processes may impact other components of the urogenital tract, such as the bladder, and contribute to the LUTS that may be experienced both in the presence and in the absence of prostate enlargement. Current research therefore offers clues about converging inflammatory pathways which may be targeted to improve treatment of BPH and/or LUTS as well as identifying potential targets for prevention of these syndromes.

Introduction

Prostate growth and enlargement and clinically significant LUTS are extremely common in aging men, as prostatic hyperplasia is evident in 50% of men 51 to 60 years of age,[1] while almost half of all men will develop clinically significant LUTS as they age.[2] In 2000, medical visits, treatments, and hospitalizations for BPH alone cost approximately \$1.1 billion, stemming from 87,000 hospitalizations and 4.4 million outpatient visits. Outpatient pharmacotherapy (primarily directed toward ameliorating LUTS) was estimated to cost \$194 million annually between 1996 and 1998.[3] As the population ages, the number of men affected by these conditions will rise dramatically, as over 11 million men are expected to meet American Health Care Policy and Research guidelines for discussing BPH treatment options by 2030.[4] BPH and LUTS therefore have a major impact on public health and medical care expenditures. However, despite the ubiquity of these syndromes among aging men, little is known about the factors that lead to the development or exacerbation of these conditions. Identification of the biologic pathways that contribute to the development and progression of BPH and LUTS would substantially contribute to identifying strategies for improved treatment and prevention that could significantly improve the quality of life of affected men and result in a substantial reduction in health care costs in this population.

Prostate enlargement may result in prostatic compression of the urethra, subsequently leading to LUTS; however, correlations between prostatic enlargement and LUTS severity are relatively modest. Additionally, LUTS may develop in the absence of prostate enlargement, suggesting that multiple factors likely contribute to the development of both prostatic enlargement and LUTS. However, an emerging body of evidence suggests that inflammatory processes may play important roles in the development of both these syndromes. In this review, we will therefore focus on literature examining a role for inflammation in the development of

prostatic enlargement and LUTS, by examining the clinical evidence for such a role, the potential role of inflammation in prostate growth and bladder dysfunction, and how BPH and LUTS therapies might target points in inflammatory pathways.

Clinical Evidence for a Role of Inflammation in Prostate Growth and LUTS

Both chronic and acute inflammation have been frequently noted in prostate biopsy sections and in tissue obtained during prostatic resection for treatment of BPH.[5–8] However, while inflammation is an extremely common observation in prostatic biospecimens, ranging from 43% to 98% in the above studies, it has been more difficult to determine whether the inflammation arose as a normal biologic part of the aging process, or whether inflammation actually contributes to prostatic enlargement, development of LUTS, and/or development of acute urinary retention (AUR).

Clinically, several cross-sectional studies have suggested associations between the presence of inflammatory infiltrates and increased prostate volume. In a retrospective review of tissue specimens obtained from prostatic resection for treatment of BPH, DiSilverio and colleagues found that presence of chronic inflammation increased as prostate volume increased, from 8.9% in prostates 30–39 mls to 61.4% in prostates 80–89 mls.[6] In a prospective, longitudinal study, 2.6% of the men participating in the Medical Therapy of Prostate Symptoms (MTOPS) trial had acute inflammation, while 43% had chronic inflammation, in prostatic biopsy specimens at baseline. Those with acute inflammation also had significantly larger prostate volumes (41.1 mls vs 36.8 mls).[9**]

Clinical studies examining associations between markers of inflammation in development of LUTS have shown much weaker associations than those observed between prostatic inflammation and prostate volume. However, the presence of elevated C-reactive protein, a non-specific marker of inflammation, was associated with a non-significant increased likelihood of having LUTS in men participating in the Third National Health and Nutrition Examination Survey.[10**] Additionally, minor correlations were observed between presence of prostatic inflammation and LUTS in men participating in the Reduction by Dutasteride of Prostate Cancer Events (REDUCE) trial.[11**] Roehrborn, et al. also found that MTOPS participants with acute inflammation in their biopsy specimens were slightly more likely to develop worsening LUTS compared to those without acute inflammation.[9**] These minor correlations between prostatic inflammation and presence of LUTS are perhaps unsurprising given the relatively modest correlations that have been observed between prostate volume and LUTS.[12,13] Finally, several studies have found moderate to strong associations between presence of prostatic inflammation and AUR. In cross-sectional studies, patients referred to urology clinics with AUR were more likely to have evidence of inflammation in prostatic specimens compared to men referred for benign prostatic obstruction.[14*,15*] Intraprostatic inflammation was also present in 70% of men requiring transurethral prostatic resection for AUR compared to 45% of men requiring resection to treat LUTS.[16*] Additionally, Roehrborn, et al., in the same MTOPS participants described above, found that patients with acute inflammation in baseline prostate biopsies were more likely (2.4% vs 0.6%) to develop AUR compared to those without such findings.[9**] Taken together, these studies suggest that inflammatory processes may contribute to the development and/or exacerbation of BPH and LUTS.

Inflammation and Prostate Growth

As described above, several clinical studies have found associations between inflammatory infiltrates in prostate biopsy sections and increased prostate volume.[6,9**] Additionally, in an animal model, administration of an immunostimulator to Wistar rats resulted in prostatic epithelial proliferation.[17] Together with the common finding of inflammatory infiltrates in

prostate biospecimens, it has therefore been hypothesized that inflammation may directly contribute to prostate growth.

The inflammatory infiltrates in prostate tissue have been primarily found to be upregulated macrophages and T lymphocytes,[18,19] Anim, et al. have suggested that initial prostate insults are followed by macrophage responses and subsequent recruitment of T-lymphocytes to the prostate tissue.[20] Other investigators have suggested that BPH might have an autoimmune component, whereby antigenic stimuli may result in the development of a chronic inflammatory response within the prostate that leads to tissue rebuilding and stromal growth in the prostate.[18,21**] Whatever the initial trigger of inflammation, pro-inflammatory cytokines likely play an essential role in attracting and maintaining pro-inflammatory cells in the prostate tissue. Handisurya, et al. found that interleukin (IL) 15 was substantially increased in BPH specimens, and may play an important role in attracting T lymphocytes to prostatic tissue and in stimulating their proliferation.[22] Additionally, macrophage inhibitory cytokine-1 (MIC-1), which inhibits macrophage activity, was downregulated in BPH tissue samples,[23] while MIC-1 levels have been associated with grade of inflammatory changes.[24] Together, these studies suggest that increased attraction of macrophages and T-lymphocytes to prostate tissue, and increased macrophage activity, may be important components in generating and sustaining prostatic inflammation (Figure 1).

The mechanisms by which inflammation may lead to prostatic growth have not been elucidated; however, once activated lymphocytes and macrophages have been attracted to prostatic tissue, they may then secrete a number of pro-inflammatory cytokines that could directly contribute to prostate growth (Figure 1). Steiner and colleagues found that IL-17 mRNA and protein levels were increased in 79% of BPH tissue specimens compared to virtually no increase in prostate specimens obtained from young men.[25] Stimulation of a BPH stromal cell culture with IL-17 also caused a substantial upregulation of the IL-6 and IL-8 cytokines.[25] These results are consistent with studies by other investigators who observed increases in IL-6 and IL-8 in seminal plasma among men with BPH compared to controls,[26*] and with an increased level of IL-6 seen in BPH epithelium compared to the epithelia and stroma of prostates obtained from controls.[27] Interferon gamma (IFN- γ), IL-2, and IL-4 have also been shown to be upregulated in BPH tissue cell lines compared to prostatic stromal cells from controls.[28] These results were supported by Steiner, et al. who also found upregulation of IFN- γ , IL-2 and IL-13 in BPH specimens with inflammatory infiltrates.[29] Unfortunately, most of these studies used control specimens from young (less than 40 years of age), healthy men, so it is unclear whether elevations in these pro-inflammatory cytokines are a normal component of aging, due to an antigenic stimulus, or the result of a chronic inflammatory process.

However, additional studies have indicated that IL-8 in particular is able to induce fibroblast growth factor 2 production in prostate stromal cells.[30], and thereby cause prostate growth. In accordance with this observation, higher IL-8 levels have also been correlated with increased prostate volume.[31] Additionally, Kramer, et al. found that BPH stromal cells were upregulated by IFN- γ . [28] Taken together, these data suggest that pro-inflammatory processes could stimulate the production of specific cytokines that could directly increase the growth and development of prostate stroma (Figure 1).

Pro-inflammatory cytokines are also able to induce additional inflammatory mediators that may contribute to prostate growth. For example, IL-17 can exert a direct influence on cyclooxygenase-2 (COX-2; the enzyme responsible for converting arachidonic acid to prostaglandin) expression through stabilizing COX-2 mRNA and preventing degradation.[32] This stabilization may then result in the presence of higher levels of COX-2 and increased activity of this enzyme. In support of this hypothesis, Wang and colleagues found that expression of COX-2 was highest in BPH tissues that also had moderate to severe T lymphocyte

and macrophage infiltration. COX-2 is a known mediator of cellular growth, and measures of prostatic epithelial cellular proliferation have been found to be highest in areas with high levels of COX-2 expression.[33]

It also appears that COX-2 expression may be induced by inflammation-independent pathways. Lin and colleagues found that the alpha adrenergic pathways may also be involved in up-regulation of COX-2, as stimulation of alpha adrenergic receptors in BPH tissue cultures resulted in an almost 14-fold increase in COX-2 gene expression.[34**] Given these data, COX-2 might represent a convergent mediator of prostate growth in both the inflammatory and alpha-adrenergic pathways.

Decreased apoptosis, and up-regulation of anti-apoptotic factors, has also been observed in BPH tissue specimens. High COX-2 levels have been associated with upregulation of the anti-apoptotic gene bcl-2, and over-expression of bcl-2 in BPH epithelial cells has also been noted. [33,35] Additionally, Gerstenbluth, et al. found intensified bcl-2 staining in ten BPH tissue specimens that had evidence of chronic inflammatory infiltrates.[7] Shariat and colleagues found high levels of survivin, a member of the apoptosis gene family that promotes cellular survival, in BPH stroma compared to normal prostate cells.[36*] Additionally, these investigators found that higher levels of survivin expression were correlated with higher LUTS, decreased maximum urinary flow rates, and increased post-void residual volumes.[36]

Together, these data suggest that inflammatory mediators may contribute to prostatic epithelial and stromal cell growth both directly, through induction of growth via cytokines that stimulate production of prostatic growth factors (Figure 1), and indirectly through decreases in prostate cell death via down-regulation of prostate cell apoptosis.

Potential Anti-Inflammatory Mechanisms of Treatments for BPH

The mechanisms of several common treatments for BPH may exert positive effects at various points in the pathways linked to inflammation in addition to their commonly accepted mechanisms of action (Table 1). For example, finasteride inhibits the conversion of testosterone to di-hydrotestosterone, but has also been shown to increase apoptosis in prostatic tissue,[37] presumably through downregulation of bcl-2.[38] Some α 1-adrenergic receptor inhibitors (doxazosin in particular) have also been shown to cause apoptosis in prostate cells.[39,40*] In the intact human, Roehrborn did not find a reduction in prostate size among men treated for three months with alfuzosin, but did not rule out the possibility that longer-term treatment might impact prostate size, or that apoptosis might not be a feature of alfuzosin.[41*] Use of anti-inflammatory medications in treating BPH has not been widely studied; however, in a group of men residing in Olmsted County, our group found that men who took a non-steroidal anti-inflammatory drug (NSAID) were about two-fold less likely to develop an increased prostate volume compared to those who did not.[42**] Additionally, Minnery and Getzenberg found that ibuprofen could reduce cell viability and induce apoptosis in a BPH cell line.[43*]

Finally, Navarrete and colleagues found that patients taking a common phytotherapy for BPH (Permixon, a sterolic lipid extract of *serenoa repens* or saw palmetto) had fewer inflammatory infiltrates in resected prostate specimens, suggesting that this agent may have anti-inflammatory properties.[44]

Taken together, these data suggest that several drugs may reduce prostate volume by acting at various points in the inflammatory pathway, possibly through direct action on the anti-apoptotic protein bcl-2, indirectly through the COX-2 pathway, or through as yet unidentified mechanisms.

Inflammation and the Bladder

As the development and progression of LUTS can not be completely attributed to prostatic growth, it is likely that LUTS may also occur due to changes in other components of the male urologic system. One obvious component that may contribute to LUTS is the bladder. Bladder outlet obstruction due to prostatic enlargement or to increased smooth muscle tone of the prostate may result in bladder wall thickening and eventual loss of optimal bladder function. However, changes in the bladder may occur apart from prostatic enlargement and may contribute to the LUTS experienced by many aging men. While associations between inflammation and bladder changes have not been examined in as much detail as inflammation and the prostate, a number of studies suggest that inflammation may also play a role in bladder changes and subsequent development of LUTS.

Prostaglandins mediate contractions of both animal and human bladder cells.[45,46] While the methods by which prostaglandins may mediate symptoms of urinary urgency are still unclear, it is clear that cyclooxygenases synthesize prostaglandins from arachadonic acid, suggesting an important role for COX-related pathways in bladder-related pathologies.

Increased prostaglandin production may occur due to mechanical factors rather than infectious, environmental, or autoimmune stimuli, as Park, et al. have found that mechanical stretching of bladder smooth muscle cells, secondary to bladder outlet obstruction in female mice, results in a substantial upregulation of COX-2, and a subsequent increase in PGE2 levels.[47] In addition, bladder smooth muscle cell proliferation was observed following cell stretching, and this response could be inhibited by addition of a COX-2 inhibitor, suggesting that COX-2 may participate in bladder wall thickening via increased bladder cell growth. Therefore, bladder outlet obstruction due to either prostatic enlargement or to increased adrenergic function, could stimulate the COX-2 pathway in the bladder, resulting in increased prostaglandin synthesis and increased LUTS.

Additionally, while bladder muscle cells contract in response to prostaglandins, human detrusor smooth muscle cells also can produce a number of pro-inflammatory cytokines. Upon stimulation with the pro-inflammatory mediators IL-1 beta and tumor necrosis factor, detrusor cells cultured from bladder biopsies were shown to upregulate monocyte chemoattractant protein 1 (MCP-1), IL-6, IL-8, and RANTES.[48,49*] Activated bladder muscle cells may therefore directly contribute to bladder inflammation by recruiting inflammatory cells to the bladder milieu. Further studies are necessary to determine whether bladder inflammation might contribute to LUTS via COX-2 dependent pathways, independent inflammatory pathways, or both.

Effect of Anti-Inflammatory Treatments on Bladder Function

Given the role of COX-2 and prostaglandins in bladder function, a number of investigators have examined the effects of nonsteroidal anti-inflammatory drugs (NSAIDs) on bladder function. Specific NSAIDs have been shown to increase bladder capacity in normal rats, and to reduce micturition frequency in rat models of cystitis.[50] More potent COX-2 inhibitors appear to be more effective at increasing bladder volume capacity in rat models of micturition. [51*] Selective COX-2 inhibitors are also able restore urothelial cell-to-cell interactions that were damaged in rat models of partial bladder outlet obstruction.[52] However, Meini, et al. found that the contractile response of the rat bladder in a model of persistent inflammation was not improved after treatment with COX inhibitors, suggesting that chronic inflammation may change the ability of the bladder to respond to prostaglandin stimuli.[53] If this is the case, men with long-term bladder inflammation may not respond well to NSAID treatments.

Few studies have examined the use of NSAIDs in directly treating LUTS; however, two small trials by Cardozo and colleagues indicated that flurbiprofen and indomethacin could substantially decrease frequency, urgency, and urge incontinence in women with detrusor instability compared to placebo.[54,55] Loxoprofen sodium use once a day improved nocturia symptoms in 74% of patients with BPH,[56] and total urinary symptoms improved in men with BPH who were treated with rofecoxib and finasteride compared to finasteride alone. The positive effects beyond that of finasteride were limited to an initial short-term interval (4 weeks), as, by 24 weeks, both groups showed similar improvements.[57**] Additionally, in support of a positive role for NSAIDs in preventing development of LUTS, we found that men who took a daily NSAID were less likely to develop moderate to severe LUTS.[42]

While these data suggest a promising role for NSAIDs in the treatment (and possibly prevention) of LUTS, NSAID use may not be completely benign in improving LUTS, as the adverse gastrointestinal effects of NSAIDs are well known. Additionally, Verhamme and colleagues have reported that men who were currently taking an NSAID were at increased odds of developing AUR, with the highest risk observed among men who recently began use of the agent.[58**] The authors postulate that in some men, reducing prostaglandin synthesis in the bladder could prevent the bladder from emptying and trigger acute urinary retention. Given the non-specific effects of NSAIDs, further studies are clearly necessary to determine whether this class of drugs might be useful in the treatment and/or prevention of LUTS.

Conclusion

Accumulating evidence suggests that inflammatory processes affecting both the prostate and the bladder may play essential roles in the development and maintenance of prostate growth and LUTS. While the inflammatory triggers and precise mechanisms by which prostate growth and LUTS may develop are still unclear, current research offers clues about converging pathways which may be targeted to improve treatment of these conditions as well as identifying potential targets for prevention of these conditions.

References

1. Berry SJ, Coffey DS, Walsh PC, Ewing LL. The development of human benign prostatic hyperplasia with age. *J Urol* 1984;132(3):474–479. [PubMed: 6206240]
2. Sarma AV, Wei JT, Jacobson DJ, et al. Comparison of lower urinary tract symptom severity and associated bother between community-dwelling black and white men: the Olmsted County Study of Urinary Symptoms and Health Status and the Flint Men's Health Study. *Urology* 2003;61(6):1086–1091. [PubMed: 12809866]
3. Wei JT, Calhoun E, Jacobsen SJ. Urologic diseases in America project: benign prostatic hyperplasia. *J Urol* 2005;173(4):1256–1261. [PubMed: 15758764]
4. Jacobsen SJ, Girman CJ, Guess HA, et al. New diagnostic and treatment guidelines for benign prostatic hyperplasia. Potential impact in the United States. *Arch Intern Med* 1995;155(5):477–481. [PubMed: 7532392]
5. Kohnen PW, Drach GW. Patterns of inflammation in prostatic hyperplasia: a histologic and bacteriologic study. *J Urol* 1979;121(6):755–760. [PubMed: 88527]
6. Di Silverio F, Gentile V, De Matteis A, et al. Distribution of inflammation, pre-malignant lesions, incidental carcinoma in histologically confirmed benign prostatic hyperplasia: a retrospective analysis. *Eur Urol* 2003;43(2):164–175. [PubMed: 12565775]
7. Gerstenbluth RE, Seftel AD, MacLennan GT, et al. Distribution of chronic prostatitis in radical prostatectomy specimens with up-regulation of bcl-2 in areas of inflammation. *J Urol* 2002;167(5):2267–2270. [PubMed: 11956490]
8. Anim JT, Kehinde EO, Prasad A, et al. Relationship between serum prostate specific antigen and the pattern of inflammation in both benign and malignant prostatic disease in Middle Eastern men. *Int Urol Nephrol* 2006;38(1):27–32. [PubMed: 16502049]

- 9**. Roehrborn C, Kaplan SA, Noble WD, et al. The impact of acute or chronic inflammation in baseline biopsy on the risk of clinical progression of BPH: results of the MTOP study (abstract). *J Urol* 2005;173:204. [PubMed: 15592076]This provocative abstract provides important data describing a temporal relationship between presence of prostatic inflammation and development of adverse urologic outcomes
- 10**. Rohrmann S, De Marzo AM, Smit E, et al. Serum C-reactive protein concentration and lower urinary tract symptoms in older men in the Third National Health and Nutrition Examination Survey (NHANES III). *Prostate* 2005;62(1):27–33. [PubMed: 15389816]This is the only currently available study to examine the association between a non-prostatic measure of inflammation and LUTS
- 11*. Nickel, JC.; Roehrborn, CG.; O'Leary, MP., et al. *J Urol*. Vol. 178. 2007. Examination of the relationship between symptoms of prostatitis and histological inflammation: baseline data from the REDUCE chemoprevention trial; p. 896-900.discussion 900–891. This study found small, non-clinically significant associations between presence of prostatic inflammation and symptoms of prostatitis, including presence of urinary symptoms
12. Collins GN, Lee RJ, McKelvie GB, et al. Relationship between prostate specific antigen, prostate volume and age in the benign prostate. *Br J Urol* 1993;71(4):445–450. [PubMed: 7684650]
13. St Sauver JL, Jacobson DJ, Girman CJ, et al. Tracking of longitudinal changes in measures of benign prostatic hyperplasia in a population based cohort. *J Urol* 2006;175(3 Pt 1):1018–1022. [PubMed: 16469608]discussion 1022
- 14*. Tuncel A, Uzun B, Erucar T, et al. Do prostatic infarction, prostatic inflammation and prostate morphology play a role in acute urinary retention? *Eur Urol* 2005;48(2):277–283. [PubMed: 15964132]discussion 283–274. This case-control study found an association between presence of prostatic inflammation and acute urinary retention - independent of prostate volume
- 15*. Kefi A, Koseoglu H, Celebi I, et al. Relation between acute urinary retention, chronic prostatic inflammation and accompanying elevated prostate-specific antigen. *Scand J Urol Nephrol* 2006;40(2):155–160. [PubMed: 16608815]Together with Tuncel *et al*, this case-control study also found an association between presence of prostatic inflammation and acute urinary retention -independent of prostate volume
- 16*. Mishra VC, Allen DJ, Nicolaou C, et al. Does intraprostatic inflammation have a role in the pathogenesis and progression of benign prostatic hyperplasia? *BJU Int* 2007;100(2):327–331. [PubMed: 17617139]These data add to the results of Tuncel *et al* and Kefi *et al* in establishing a role for prostatic inflammation in AUR
17. Kessler OJ, Keisari Y, Servadio C, Abramovici A. Role of chronic inflammation in the promotion of prostatic hyperplasia in rats. *J Urol* 1998;159(3):1049–1053. [PubMed: 9474230]
18. Steiner G, Gessl A, Kramer G, et al. Phenotype and function of peripheral and prostatic lymphocytes in patients with benign prostatic hyperplasia. *J Urol* 1994;151(2):480–484. [PubMed: 7506795]
19. Bostwick DG, de la Roza G, Dundore P, et al. Intraepithelial and stromal lymphocytes in the normal human prostate. *Prostate* 2003;55(3):187–193. [PubMed: 12692784]
20. Anim JT, Udo C, John B. Characterisation of inflammatory cells in benign prostatic hyperplasia. *Acta Histochem* 1998;100(4):439–449. [PubMed: 9842422]
- 21**. Kramer G, Mitteregger D, Marberger M. Is benign prostatic hyperplasia (BPH) an immune inflammatory disease? *Eur Urol* 2007;51(5):1202–1216. [PubMed: 17182170]This interesting review article proposes that BPH meets several characteristic criteria for immune inflammatory diseases
22. Handisurya A, Steiner GE, Stix U, et al. Differential expression of interleukin-15, a pro-inflammatory cytokine and T-cell growth factor, and its receptor in human prostate. *Prostate* 2001;49(4):251–262. [PubMed: 11746271]
23. Kakehi Y, Segawa T, Wu XX, et al. Down-regulation of macrophage inhibitory cytokine-1/prostate derived factor in benign prostatic hyperplasia. *Prostate* 2004;59(4):351–356. [PubMed: 15065082]
24. Taoka R, Tsukuda F, Ishikawa M, et al. Association of prostatic inflammation with down-regulation of macrophage inhibitory cytokine-1 gene in symptomatic benign prostatic hyperplasia. *J Urol* 2004;171(6 Pt 1):2330–2335. [PubMed: 15126815]

25. Steiner GE, Newman ME, Paikl D, et al. Expression and function of pro-inflammatory interleukin IL-17 and IL-17 receptor in normal, benign hyperplastic, and malignant prostate. *Prostate* 2003;56(3):171–182. [PubMed: 12772186]
- 26*. Penna G, Mondaini N, Amuchastegui S, et al. Seminal plasma cytokines and chemokines in prostate inflammation: interleukin 8 as a predictive biomarker in chronic prostatitis/chronic pelvic pain syndrome and benign prostatic hyperplasia. *Eur Urol* 2007;51(2):524–533. [PubMed: 16905241] discussion 533. This manuscript attempts to bridge basic science and clinical data, by both noting an increased level of IL-8 in BPH patients and an association between IL-8 level and urinary symptoms
27. Royuela M, Ricote M, Parsons MS, et al. Immunohistochemical analysis of the IL-6 family of cytokines and their receptors in benign, hyperplastic, and malignant human prostate. *J Pathol* 2004;202(1):41–49. [PubMed: 14694520]
28. Kramer G, Steiner GE, Handisurya A, et al. Increased expression of lymphocyte-derived cytokines in benign hyperplastic prostate tissue, identification of the producing cell types, and effect of differentially expressed cytokines on stromal cell proliferation. *Prostate* 2002;52(1):43–58. [PubMed: 11992619]
29. Steiner GE, Stix U, Handisurya A, et al. Cytokine expression pattern in benign prostatic hyperplasia infiltrating T cells and impact of lymphocytic infiltration on cytokine mRNA profile in prostatic tissue. *Lab Invest* 2003;83(8):1131–1146. [PubMed: 12920242]
30. Giri D, Ittmann M. Interleukin-8 is a paracrine inducer of fibroblast growth factor 2, a stromal and epithelial growth factor in benign prostatic hyperplasia. *Am J Pathol* 2001;159(1):139–147. [PubMed: 11438462]
31. Castro P, Xia C, Gomez L, et al. Interleukin-8 expression is increased in senescent prostatic epithelial cells and promotes the development of benign prostatic hyperplasia. *Prostate* 2004;60(2):153–159. [PubMed: 15162381]
32. Faour WH, Mancini A, He QW, Di Battista JA. T-cell-derived interleukin-17 regulates the level and stability of cyclooxygenase-2 (COX-2) mRNA through restricted activation of the p38 mitogen-activated protein kinase cascade: role of distal sequences in the 3'-untranslated region of COX-2 mRNA. *J Biol Chem* 2003;278(29):26897–26907. [PubMed: 12746433]
33. Wang W, Bergh A, Damber JE. Chronic inflammation in benign prostate hyperplasia is associated with focal upregulation of cyclooxygenase-2, Bcl-2, and cell proliferation in the glandular epithelium. *Prostate* 2004;61(1):60–72. [PubMed: 15287094]
- 34**. Lin, VK.; Wang, SY.; Roehrborn, C. Alpha adrenergic stimulation regulates cox-2 expression in prostate stroma cells. In: Association, AU., editor. American Urologic Association Annual Meeting. Vol. 173. Vol. 390. Lippincott Williams & Wilkins; San Antonio, TX: 2005. This interesting abstract has not been published in manuscript form, but suggests a fascinating association between upregulation of the alpha adrenergic pathway and COX-2
35. Kyprianou N, Tu H, Jacobs SC. Apoptotic versus proliferative activities in human benign prostatic hyperplasia. *Hum Pathol* 1996;27(7):668–675. [PubMed: 8698310]
- 36*. Shariat SF, Ashfaq R, Roehrborn CG, et al. Expression of survivin and apoptotic biomarkers in benign prostatic hyperplasia. *J Urol* 2005;174(5):2046–2050. [PubMed: 16217391] This paper begins to explore the role of apoptotic biomarkers in BPH beyond bcl-2
37. Rittmaster RS, Norman RW, Thomas LN, Rowden G. Evidence for atrophy and apoptosis in the prostates of men given finasteride. *J Clin Endocrinol Metab* 1996;81(2):814–819. [PubMed: 8636309]
38. Huynh H. Induction of apoptosis in rat ventral prostate by finasteride is associated with alteration in MAP kinase pathways and Bcl-2 related family of proteins. *Int J Oncol* 2002;20(6):1297–1303. [PubMed: 12012013]
39. Tahmatzopoulos A, Kyprianou N. Apoptotic impact of alpha1-blockers on prostate cancer growth: a myth or an inviting reality? *Prostate* 2004;59(1):91–100. [PubMed: 14991869]
- 40*. Garrison JB, Kyprianou N. Doxazosin induces apoptosis of benign and malignant prostate cells via a death receptor-mediated pathway. *Cancer Res* 2006;66(1):464–472. [PubMed: 16397262]
- 41*. Roehrborn CG. Three months' treatment with the alpha(1)-blocker alfuzosin does not affect total or transition zone volume of the prostate. *Prostate Cancer Prostatic Dis*. 2005 This study did not find

a clinical association between potential apoptotic effects of alfuzosin and decreased prostate volume; however, three months treatment may not have been long enough to observe any clinical changes in prostate volume

- 42**. St Sauver J, Jacobson D, McGree ME, et al. Protective association between non-steroidal anti-inflammatory drug use and measures of benign prostatic hyperplasia. *American Journal of Epidemiology*. 2006 This paper examines the association between NSAID use and urologic outcomes, finding a protective association between daily use of NSAIDs and prostate volume, LUTS, and AUR, suggesting that NSAIDs may be useful in preventing some of these conditions
- 43*. Minnery CH, Getzenberg RH. Benign prostatic hyperplasia cell line viability and modulation of jm-27 by doxazosin and Ibuprofen. *J Urol* 2005;174(1):375–379. [PubMed: 15947693] These authors provide basic science data to suggest that both doxazosin and ibuprofen act through jm-27 to reduce prostate growth via increased cell death and increased apoptosis
44. Vela Navarrete R, Garcia Cardoso JV, Barat A, et al. BPH and inflammation: pharmacological effects of Permixon on histological and molecular inflammatory markers. Results of a double blind pilot clinical assay. *Eur Urol* 2003;44(5):549–555. [PubMed: 14572753]
45. Maggi CA, Evangelista S, Grimaldi G, et al. Evidence for the involvement of arachidonic acid metabolites in spontaneous and drug-induced contractions of rat urinary bladder. *J Pharmacol Exp Ther* 1984;230(2):500–513. [PubMed: 6431081]
46. Palea S, Toson G, Pietra C, et al. Pharmacological characterization of thromboxane and prostanoid receptors in human isolated urinary bladder. *Br J Pharmacol* 1998;124(5):865–872. [PubMed: 9692770]
47. Park JM, Yang T, Arend LJ, et al. Obstruction stimulates COX-2 expression in bladder smooth muscle cells via increased mechanical stretch. *Am J Physiol* 1999;276(1 Pt 2):F129–136. [PubMed: 9887088]
48. Bouchelouche K, Alvarez S, Andersen L, Nordling J, Horn T, Bouchelouche P. Monocyte chemoattractant protein-1 production by human detrusor smooth muscle cells. *J Urol* 2004;171(1):462–466. [PubMed: 14665956]
- 49*. Bouchelouche K, Alvarez S, Horn T, Nordling J, Bouchelouche P. Human detrusor smooth muscle cells release interleukin-6, interleukin-8, and RANTES in response to proinflammatory cytokines interleukin-1beta and tumor necrosis factor-alpha. *Urology* 2006;67(1):214–219. [PubMed: 16413378] This study provides basic science data indicating that detrusor smooth muscle cells themselves are able to release pro-inflammatory cytokines
50. Takagi-Matsumoto H, Ng B, Tsukimi Y, Tajimi M. Effects of NSAIDs on bladder function in normal and cystitis rats: a comparison study of aspirin, indomethacin, and ketoprofen. *J Pharmacol Sci* 2004;95(4):458–465. [PubMed: 15322343]
- 51*. Angelico P, Guarneri L, Velasco C, et al. Effect of cyclooxygenase inhibitors on the micturition reflex in rats: correlation with inhibition of cyclooxygenase isozymes. *BJU Int* 2006;97(4):837–846. [PubMed: 16536784] This study provides animal model data to suggest that efficacy of COX inhibitors in treating rat micturition conditions is related primarily to action on the COX-2 enzyme
52. Erdogru T, Celik-Ozenci C, Seval Y, et al. The restorative effect of a selective cyclooxygenase-2 inhibitor on urothelial cell-cell interactions after partial bladder outlet obstruction in rats. *BJU Int* 2005;95(4):664–669. [PubMed: 15705100]
53. Meini S, Lecci A, Cucchi P, Catalioto RM, Criscuoli M, Maggi CA. Inflammation modifies the role of cyclooxygenases in the contractile responses of the rat detrusor smooth muscle to kinin agonists. *J Pharmacol Exp Ther* 1998;287(1):137–143. [PubMed: 9765332]
54. Cardozo LD, Stanton SL. A comparison between bromocriptine and indomethacin in the treatment of detrusor instability. *J Urol* 1980;123(3):399–401. [PubMed: 7359644]
55. Cardozo LD, Stanton SL, Robinson H, Hole D. Evaluation of flurbiprofen in detrusor instability. *Br Med J* 1980;280(6210):281–282. [PubMed: 6986948]
56. Araki T, Yokoyama T, Kumon H. Effectiveness of a nonsteroidal anti-inflammatory drug for nocturia on patients with benign prostatic hyperplasia: a prospective non-randomized study of loxoprofen sodium 60 mg once daily before sleeping. *Acta Med Okayama* 2004;58(1):45–49. [PubMed: 15157011]

- 57**. Di Silverio F, Bosman C, Salvatori M, et al. Combination therapy with rofecoxib and finasteride in the treatment of men with lower urinary tract symptoms (LUTS) and benign prostatic hyperplasia (BPH). *Eur Urol* 2005;47(1):72–78. [PubMed: 15582252]discussion 78–79. This small clinical trial provides interesting data to suggest that NSAIDs may be a useful adjunct therapy for treating BPH-related LUTS
- 58**. Verhamme KM, Dieleman JP, Van Wijk MA, et al. Nonsteroidal anti-inflammatory drugs and increased risk of acute urinary retention. *Arch Intern Med* 2005;165(13):1547–1551. [PubMed: 16009872]This large retrospective case-control study provides cautionary data suggesting that, in at least some men, new use of NSAIDs may actually trigger AUR
59. Partin JV, Anglin IE, Kyprianou N. Quinazoline-based alpha 1-adrenoceptor antagonists induce prostate cancer cell apoptosis via TGF-beta signalling and I kappa B alpha induction. *Br J Cancer* 2003;88(10):1615–1621. [PubMed: 12771931]

Figure 1.
Possible Mechanisms By Which Inflammatory Stimuli May Contribute to Prostatic Growth

Table 1
Possible Mechanisms by Which Therapies May Improve BPH and/or LUTS by Interrupting Inflammatory Processes

Type of Drug	Mechanism of Action	Result	References
Finasteride Alpha-Adrenergic Receptor Inhibitors Non-Steroidial Anti- Inflammatory Drugs	Down-regulation of BCL-2	Increased apoptosis of prostate cells	[37,38]
	Decreased function of alpha-adrenergic pathways	Increased apoptosis of prostate cells	[39,40,59]
	Decreased COX-2 activity	Decreased prostate volume	[42]
		Increased apoptosis of prostate cells	[43]
		Increased bladder capacity	[50,51]
Serenoa Repens		Restoration of urothelial cell-cell interactions	[52]
	Unknown mechanisms	Improved LUTS	[54–56]8
	Unknown	Decreased inflammatory infiltrates in the prostate	[44]