Figure 1 The 'neck-line' incision post surgery.

than conventional incisions. The cutaneous innervation of the clavicular region, via subclavicular branches (C4, C5), travels typically in the sagittal plane under the platysma; an incision closer to this plane may reduce postoperative sensory loss. The incision allows adequate exposure of the clavicle for surgical fixation due to tissue mobility, and patients can conceal the scar more effectively with their clothes.

The 'glove cuff' technique for difficult stomas

J HORWOOD, D HAY

Department of General Surgery, Glan Clwyd Hospital, Rhyl, UK

CORRESPONDENCE TO

J Horwood, Department of General Surgery, Glan Clwyd Hospital, Rhyl, Denbighshire LL18 5UJ, UK
E: tipper111@aol.com

In patients with generous abdominal walls and bulky mesenteries, fashioning stomas can be difficult. Struggling to pass the bowel and its mesentery through the stoma can damage the local blood supply. We propose a method which eases this passage. Cut the fingers off a surgical glove, and place the colon in the lumen of the remaining glove cuff (Fig. 1). Pass an instrument through

Figure 1 Colon placed in the glove cuff.

Figure 2 Delivering the colon through the abdominal wall.

the external stomal opening and apply gentle traction to the glove and bowel, delivering the colon through the abdominal wall (Fig. 2). We have found the technique to facilitate passage of the mesentery, reducing trauma to the delicate tissues.

Hand in glove

PADMANABHAN SUBRAMANIAN, MJ METCALFE,
SJ CHADWICK, DJ VAUGHAN

Northwick Park Hospital, The North West London Hospitals NHS Trust, Harrow, Middlesex, UK

CORRESPONDENCE TO

Padmanabhan Subramanian, 46 Aspen Gardens, Hammersmith, London W6 9JE, UK. E: drpad@hotmail.com

Kinking of the radial artery line is a common problem with malpositioning of the hand in the anaesthetised patient during surgery. Line kinking can cause digital ischaemia¹ and altered arterial blood pressure readings. Repositioning the line whilst the patient's hand is under

Figure 1 Hand positioned around inflated latex glove avoids kinking of the radial artery line.

sterile drapes is difficult, disrupts surgery and can prolong the duration of anaesthesia. A simple and cheap solution is to use an inflated latex glove and position the patient's hand with interlocking fingers around it as shown in Figure 1, thus maintaining dorsiflexion of the patient's wrist and so preventing arterial line kinking.

Reference

1. Sferir R, Khoury S, Khoury G, Rustum J, Ghabash M. Ischaemia of the hand after radial artery monitoring. *Cardiovasc Surg* 1996; 4: 456–8.

Using sterile co-polymer overgloves as undergloves prevents hands from 'smelling' after contaminated abdominal surgery

RT PADWICK¹, MJ OSBORNE²

¹Hereford Hospitals NHS Trust, Hereford, UK

²Warwick GI Unit, South Warwickshire Hospitals NHS Trust, Warwick, UK

CORRESPONDENCE TO

Robert Padwick, Specialist Registrar in General Surgery, County Hospital, Union Walk, Hereford HR1 2ER, UK
E: rob.padwick@doctors.net.uk

When performing contaminated abdominal surgery, one of the problems that all surgeons will have experienced is the nasty 'smell' left on their hands after the procedure, even when double-gloving. Surgeons at our institution have found that by using a pair of sterile co-polymer overgloves as undergloves (e.g. Bodyguards®; Medisavers, E: sales@medisavers.co.uk) instead of a pair of latex gloves results in no 'smell' being transmitted to the hands. One of the major problems reported by surgeons when using two pairs of gloves is loss of tactile sensitivity.¹ We have found that these gloves do not significantly affect tactile sensitivity.

Reference

1. Twomey CL. Double gloving: a risk reduction strategy. *Jt Comm J Qual Saf* 2003; 29: 369–78.

A radiolucent retractor for locking screws

ND RILEY, CK SIVAJI

Department of Trauma and Orthopaedic Surgery, Southend University Hospital NHS Trust, Westcliff-on-Sea, Essex, UK

CORRESPONDENCE TO

ND Riley, Flat 6 Roy Square, London E14 8BY, UK
E: ndriley@googlemail.com

During proximal locking of a retrograde femoral intramedullary nail, retraction can be difficult when using traditional radio-opaque retractors. A 50-ml syringe, such as BD Plastipak™ (BD, The Danby Building, Edmund Halley Road, Oxford Science Park, Oxford, UK),

Figure 1 Syringe placed onto saw bone demonstrating retraction technique.

with the catheter tip cut off, either with scissors or a saw, can be placed directly onto bone via the skin incision. This easily retracts the tissues allowing access to the proximal femur with the added benefit of being radiolucent. This allows the surgeon to interpret the fluoroscopy findings accurately and position the proximal locking screw precisely. Furthermore, the surgical assistant is protected from erroneous exposure to radiation whilst screening.

A cost-effective transverse colostomy

E MYERS, D KAVANAGH, D EVOY

Department of Surgery, Saint Vincent's University Hospital, Dublin, Ireland

CORRESPONDENCE TO

Eddie Myers, Department of Surgery, Saint Vincent's University Hospital, Elm Park, Dublin 4, Ireland
E: eddiemyers@eircom.net

A loop transverse colostomy can be performed as an effective method of relieving acute colonic obstruction. In the setting of disseminated intra-abdominal malignancy and resultant abnormal anatomy, the procedure is frequently underestimated. By securing a coin in the right upper quadrant and obtaining a plain film of the abdomen, the location of the proposed colostomy relative to the transverse colon can be estimated pre-operatively. This facilitates less dissection of the peritoneal attachments in order to mobilise the colon to the desired site.

Figure 1 (A) Coin secured in right upper quadrant; (B) pre-operative plain film of the abdomen; (C) loop transverse colostomy in desired site.