Fission Yeast *pk11* Is a Kinesin-related Protein Involved in Mitotic Spindle Function

Alison L. Pidoux, Michel LeDizet, and W. Zacheus Cande*

Department of Molecular and Cell Biology, University of California, Berkeley, California 94720-3200

Submitted December 4, 1995; Accepted July 10, 1996

Monitoring Editor: Mitsuhiro Yanagida

We have used anti-peptide antibodies raised against highly conserved regions of the kinesin motor domain to identify kinesin-related proteins in the fission yeast *Schizosaccharomyces pombe*. Here we report the identification of a new kinesin-related protein, which we have named *pk11*. Sequence homology and domain organization place *pk11* in the Kar3/*ncd* subfamily of kinesin-related proteins. Bacterially expressed *pk11* fusion proteins display microtubule-stimulated ATPase activity, nucleotide-sensitive binding, and bundling of microtubules. Immunofluorescence studies with affinity-purified antibodies indicate that the *pk11* protein localizes to the nucleus and the mitotic spindle. *Pk11* null mutants are viable but have increased sensitivity to microtubule-disrupting drugs. Disruption of *pk11*⁺ suppresses mutations in another kinesin-related protein, *cut7*, which is known to act in the spindle. Overexpression of *pk11* to very high levels causes a similar phenotype to that seen in *cut7* mutants: V-shaped and star-shaped microtubule structures are observed, which we interpret to be spindles with unseparated spindle poles. These observations suggest that *pk11* and *cut7* provide opposing forces in the spindle. We propose that *pk11* functions as a microtubule-dependent motor that is involved in microtubule organization in the mitotic spindle.

INTRODUCTION

Through a series of precise temporally and spatially complex movements, the mitotic spindle ensures the accurate segregation of chromosomes to the two daughter cells. These motility events include the separation of the spindle poles and formation of the bipolar spindle, prometaphase oscillations of chromosomes on kinetochore microtubules (MTs)¹ leading to their congression to the metaphase plate, chromatid separation and movement to the poles (anaphase A), and spindle elongation (anaphase B). At least two mechanisms of force generation are thought to contribute to spindle structure and motility: these are MT polymerization/depolymerization (Koshland *et al.*, 1988; Mitchison, 1988; Coue *et al.*, 1991) and the action of molecular motors of the dynein and kinesin families (McIntosh and Pfarr, 1991; Sawin and Scholey, 1991). Analysis of mutants defective in spindle function and

immunolocalization studies implicate motors of both of these classes in spindle function (Endow and Titus, 1992; Bloom and Endow, 1994). This report is concerned with the role of the kinesin family of motors in mitosis.

Kinesin heavy chain (KHC) was first isolated from squid axoplasm as a plus-end-directed MT motor (Vale *et al.*, 1985) and is thought to be responsible for the transport of vesicles and organelles in a variety of cell types. Since its discovery a decade ago, a large and ever-expanding family of kinesin-related proteins (KRPs) has been identified (Goldstein, 1993). These KRPs share with kinesin a conserved 45-kDa motor domain that has the capacity to bind and hydrolyze ATP, bind MTs, and generate force (Yang *et al.*, 1989). The motor domain may be at the N or C terminus or in the middle of the molecule, and KRPs have been

* Corresponding author: Department of Molecular and Cell Biology, 341 Life Sciences Addition, University of California, Berkeley, CA 94720-3200.

¹ Abbreviations used: DAPI, 4',6-diamidino-2-phenylindole; GST, glutathione S-transferase; KHC, kinesin heavy chain; KRP, kinesin related protein; MT, microtubule; NLS, nuclear localization signal; TBZ, thiabendazole; YES, yeast extract with supplements.

divided into subfamilies on the basis of sequence homology within the motor domain (Goodson *et al.*, 1994). The nonconserved tail regions have been suggested to bind specific cargoes, such as vesicles or chromosomes, and are implicated in dimerization and interaction with accessory subunits (Bloom and Endow, 1994). Although KHC and many of the KRPs that have been tested display plus-end-directed motility (Nislow *et al.*, 1992; Sawin *et al.*, 1992a; Kondo *et al.*, 1994; Noda *et al.*, 1995), several KRPs show minus-end-directed movement, and the majority of these are C-terminal motors (McDonald *et al.*, 1990; Walker *et al.*, 1990; Endow *et al.*, 1994b; Kuriyama *et al.*, 1995).

In some organisms there are very large numbers of KRPs; for example, in *Drosophila* there are reported to be upward of 30 (Endow and Hatsumi, 1991). Kinesins and KRPs are thought to have roles in many cellular processes, including structure and transport of various membrane-bound organelles, nuclear migration, mitosis, and meiosis (reviewed in Bloom and Endow, 1994). It is possible that individual KRPs have very precise or subtle functions or that there is a degree of functional redundancy within the KRP family. One approach toward understanding the relative contributions various KRPs make to spindle motility is to study them in a cell in which the total number of KRPs is limited. The yeasts and other fungi, such as *Aspergillus nidulans*, may offer such a system. In budding yeast, six KRPs have been identified (Kar3p, Cin8p, Kip1p, Kip2p, Smy1p, and RRC805; Meluh and Rose, 1990; Hoyt *et al.*, 1992; Lillie and Brown, 1992; Roof *et al.*, 1992; A. Hoyt, personal communication), and a series of experiments by Hoyt, Saunders, and coworkers has begun to unravel the relative roles and actions of molecular motors in the mitotic spindle (Hoyt *et al.*, 1992; Saunders and Hoyt, 1992; Saunders *et al.*, 1995).

The fission yeast *Schizosaccharomyces pombe* is also an attractive organism for the study of mitosis and the investigation of KRP functions in the spindle. Like budding yeast, *S. pombe* is amenable to genetic analysis, and a large number of mutants affecting the cell cycle, mitosis, and MTs have been identified and studied (Hagan and Hyams, 1988; Yanagida, 1989; Forsburg and Nurse, 1991). The 40–100-kb centromeres of *S. pombe* (Fishel *et al.*, 1988; Chikashige, *et al.*, 1989; Carbon and Clarke, 1990) offer an intermediate in complexity between the simple 125-bp centromeres of *Saccharomyces cerevisiae* and the highly complex heterochromatic centromeres of higher eukaryotes. Serial section EM reconstruction studies of the *S. pombe* spindle indicate that it shares important similarities with the spindles of higher organisms; the chromosomes are attached to the spindle by a bundle (2–4) of kinetochore MTs, and chromatid separation is accompanied by both chromosome-to-pole movement and spindle elongation (Ding *et al.*, 1993). Fluorescence in situ hybridization studies have also demonstrated that

anaphase A occurs in fission yeast (Funabiki *et al.*, 1993). Permeabilized cell models for spindle elongation and for MT nucleation from the spindle pole body have been developed in *S. pombe* (Masuda *et al.*, 1990, 1992). The powerful combination of genetics and in vitro analysis in this organism provides an opportunity to investigate the roles and functional interactions of spindle components, including KRPs. One KRP, cut7, has already been identified in *S. pombe* and is thought to play a role in bipolar spindle formation (Hagan and Yanagida, 1990, 1992).

KRPs have been identified by a variety of approaches, including complementation of mutant phenotypes and more directed approaches based on biochemical purification, PCR, and antibody screening (reviewed in Bloom and Endow, 1994). The KRP motor domain contains several highly conserved regions such as the LAGSE and HIPYR motifs. Anti-peptide antibodies that were raised to these regions (Cole *et al.*, 1992; Sawin *et al.*, 1992b) stain mitotic spindles and kinetochores (Sawin *et al.*, 1992b) and block spindle elongation in vitro (Hogan, *et al.*, 1993). These antibodies have the potential to identify many members of the kinesin family in a particular organism; indeed, they recognize a number of MT-binding proteins in *Xenopus* and sea urchin extracts (Cole *et al.*, 1992; Sawin *et al.*, 1992b) and have been used to isolate KRP clones from mammalian cells and *Xenopus* (Wordeman and Mitchison, 1995; Walczak *et al.*, 1996). We are using the LAGSE and HIPYR antibodies as part of a strategy to identify KRPs in *S. pombe*. Here we report the sequence and analysis of a novel KRP, pkl1, which belongs to the Kar3/ncd subfamily of C-terminal motor proteins. We provide data on its localization, genetics, and biochemical properties that suggest it may function as an MT motor in the mitotic spindle.

MATERIALS AND METHODS

Chemicals, Strains, Media, and S. pombe Techniques

Chemicals were obtained from Sigma Chemical (St. Louis, MO) or Fisher Scientific (Pittsburgh, PA). Benomyl was a gift from DuPont (Boston, MA). *S. pombe* strains used in this study were 972h⁺; PN556 (*h⁺ ade6-M216 ura4-D18 leu1-32*), and PN557 (*h⁻ ade6-M210 ura4-D18 leu-32*) (obtained from Paul Nurse, Imperial Cancer Research Fund, United Kingdom); *h⁺ ade6-M216 leu-32 ura4-D18 his3-D1* (obtained from Kathy Gould, Vanderbilt University, Nashville, TN); *cut7-446 h⁺* (obtained from Mitsuhiro Yanagida, Kyoto University, Japan); *cut7-21, -22, -24* (personal communication, Rachel Bartlett, Paul Nurse, Iain Hagan, Mitsuhiro Yanagida; obtained from Iain Hagan, University of Manchester, United Kingdom). *S. pombe* strains were grown in yeast extract with supplements (YES) or minimal media supplemented as appropriate (Moreno *et al.*, 1991). Standard *S. pombe* techniques were used (Moreno *et al.*, 1991); standard cloning procedures were used (Sambrook *et al.*, 1989).

For comparison of the growth of different strains, log-phase cells were counted with a hemocytometer, and fivefold serial dilutions were made in 96-well plates; the density of cells in the first well was 5×10^6 /ml. Cells were stamped in duplicate onto YES plates (or YES plates containing 12 μ g/ml thiabendazole for the MT-drug

sensitivity experiments), using a 48-pin replicator device (Aladin Enterprises, San Francisco, CA). Plates were incubated at an appropriate temperature for 2–7 d and photographed when significant growth had occurred for at least one strain.

Construction and Screening of Lambda Library

Total RNA was prepared from log-phase 972h[−] cells (Moreno *et al.*, 1991), and poly A⁺ RNA was isolated by using a Fast Track kit (Invitrogen, San Diego, CA), following the manufacturer's instructions. cDNA was synthesized and directionally cloned into the lambda ZAP II vector by using the ZAP-cDNA synthesis kit (Stratagene, La Jolla, CA) and packaged with Gigapack II gold extract (Stratagene), following protocols provided by Stratagene. Primary recombinants (1.25 × 10⁶) were obtained, and the library was amplified once. In all, 80,000 plaques were screened (Sambrook *et al.*, 1989) on Hybond-ECL membrane (Amersham, Arlington Heights, IL) with anti-LAGSE antibody at 2.5 µg/ml (Sawin *et al.*, 1992b) and HRP-conjugated goat anti-rabbit secondary antibody (Jackson ImmunoResearch Laboratories, West Grove, PA) at 0.3 µg/ml and then developed with ECL reagents (Amersham) according to the manufacturer's instructions. After plaque purification of the LAGSE-positive clones, candidates were rescreened with the anti-HIPYR antibody at 5 µg/ml (Sawin *et al.*, 1992b); 55 LAGSE-positive clones were isolated, and six were positive with both antibodies. In vivo excision was performed to obtain positive clones in plasmid form (pBluescript SK[−]).

Sequencing

Six *pk11*⁺ cDNA clones were obtained, the longest of which (pLH-52) was ~3 kb in length and was sequenced. Various subclones, deletion series (Yanisch-Perron *et al.*, 1985), and oligonucleotide primers were made to complete the sequence on both strands with Sequenase 2.0 (United States Biochemical, Cleveland, OH). Sequence analysis was performed with the Sequence Analysis Software Package (Genetics Computer Group, Madison, WI). Sequence comparisons were done with the Bestfit program. The probability of coiled-coil formation was calculated by the method of Lupas *et al.* (1991), using a window size of 28 amino acids.

Disruption of *pk11*⁺

For the partial disruption, pLH-52 was digested with *Nde*I and the overhangs were filled in with Klenow. This vector was ligated with a 1.7-kb *Xho*I fragment bearing the *S. pombe ura4*⁺ gene that had also been blunt-ended. The resulting 3.5-kb fragment was cut out of the vector by using *Sac*I and *Kpn*I. This fragment was used to transform *S. pombe* strains PN556, PN557, and a diploid made by mating these two strains, using the lithium chloride method (Broker, 1987). Transformants were selected on plates lacking uracil. Genomic DNA was prepared from 5 ml of stationary phase cultures (Moreno *et al.*, 1991) and digested with *Eco*RI. After electrophoresis and Southern transfer, DNA was hybridized with a probe made from the 1.4-kb *Hind*III fragment from pLH-52. Diploid disruptants gave two bands with this probe; for haploid disruptants, the wild-type 3.4-kb band was replaced by a 3.9-kb band. This new band also hybridized to a probe made from the *ura4*⁺ gene.

A full-length knockout was also made in the *pk11*⁺ gene. A plasmid, pUR19-gpk11, isolated from an *S. pombe* genomic library (Barbet *et al.*, 1992), was a generous gift of Cindy Troxell and J.R. McIntosh (University of Colorado, Boulder, CO). It contains a 6.9-kb insert bearing the *pk11*⁺ gene. pUR19-gpk11 was mapped (see Figure 4A, bottom panel), digested with *Bam*HI, treated with Klenow, and religated to delete the 1-kb *Bam*HI fragment downstream of *pk11*⁺ and destroy the *Bam*HI site. The resultant plasmid pPHK1 was used as the template in PCR reactions to create the arms for the knockout construct. For the 3' end, an oligonucleotide primer corresponding to the last 19 nucleotides of the *pk11*⁺ coding region, plus a *Bam*HI

site (underlined), was synthesized: 5'-TACTACGGATCCCTATTAACATTATAAACG-3'. This was used in conjunction with the −20 sequencing primer (in the plasmid) to amplify a 2-kb fragment downstream of the *pk11*⁺ gene (see Figure 4A), using Vent DNA polymerase (New England Biolabs, Beverly, MA). This fragment was digested with *Bam*HI and *Bgl*II, and the 1-kb *Bam*HI-*Bgl*II fragment immediately downstream of the *pk11*⁺ gene was cloned into *Bam*HI-digested pBluescript SK[−] (Stratagene) to create pPHK2. For the upstream region, an oligonucleotide primer immediately upstream of the *pk11*⁺ coding region, plus a *Bam*HI site (underlined), was synthesized: 5'-TACTACGGATCCCTATAAGTTAAAAAGAGCG-3'. This was used in combination with the reverse-sequencing primer to amplify a 1.4-kb fragment immediately upstream of *pk11*⁺. This was cut with *Bam*HI and *Pst*I and inserted into *Bam*HI-*Pst*I-cut pPHK2 to create pPHK3. The *his3*⁺ gene was removed from pAF1 as a 2-kb *Bgl*II fragment (Ohi *et al.*, 1996) and cloned into *Bam*HI-cut pPHK3 to create pPHK4. The 4.4-kb insert was removed from the plasmid by using *Sac*I and *Kpn*I; the insert was gel purified and used to transform haploid *S. pombe* strain: *h*⁺ *ade6-M216 leu-32 ura4-D18 his3-D1*. Transformants were selected on plates lacking histidine, and *Sac*I-*Bam*HI-digested genomic DNA was probed with a 1.2-kb *Bam*HI-*Bgl*II fragment from outside the region used for transformation; wild-type DNA gives a 3.6-kb band, whereas the knockout gives a 3-kb band. The status of the knockout was also confirmed with other digests and *his3*⁺ probes and by Western blotting with anti-*pk11* antibodies.

Affinity Purification of Antibodies and Western Blotting

A nonconserved region of *pk11* near the N terminus was chosen for the production of antibodies. The 360-bp *Bgl*II-*Eco*RV fragment was cloned into the *Bam*HI-*Sma*I-digested pGEX-1N (Pharmacia Biotech, Piscataway, NJ; Smith and Johnson, 1988) to allow the expression of a protein fusion with glutathione S-transferase (GST-PKL1-BV). Soluble fusion protein was purified on glutathione agarose beads (Smith and Johnson, 1988). Two New Zealand white rabbits were subcutaneously injected with 0.5 mg of GST-PKL1-BV fusion protein in Freund's complete adjuvant (for the first injection) or incomplete adjuvant at monthly intervals, and test bleeds were taken ~7 d after injection. Bleeds were analyzed by Western blotting, and although there was a reaction against a 96-kDa protein, the antisera recognized many other proteins in *S. pombe* extracts and gave very intense bands at 50 and 70 kDa on Western blots. Affinity purification was therefore necessary. Fusion proteins and bacterial extracts were coupled to Affigel-10 (Bio-Rad Laboratories, Hercules, CA) overnight at 4°C, and affinity purification was performed as described (Harlow and Lane, 1988) with the modifications noted below. Preadsorption on a GST column followed by elution from a GST-PKL1-BV column did not purify anti-PKL1 antibodies away from anti-50/70-kDa antibodies. We reasoned that poor folding of GST-PKL1-BV in *Escherichia coli* caused it to complex with bacterial heat-shock proteins, which then were purified along with GST-PKL1-BV on the glutathione agarose; antibodies raised against these heat-shock proteins in the rabbit then cross-reacted with *S. pombe* heat-shock proteins. Therefore, antiserum was preadsorbed on a column prepared from an extract from bacteria expressing another GST fusion protein and, presumably, heat-shock proteins (a gift from Dr. Margit Foss, Duke University, NC). This depleted serum was passed over the GST-PKL1-BV column, and the specifically bound antibodies were eluted with 100 mM glycine, pH 2.5, or 100 mM triethanolamine, pH 11.5. The peak triethanolamine fraction was used in Western blotting and immunofluorescence experiments described in this paper.

For Western blotting, protein extracts were made by mechanical disruption with glass beads in sample buffer, and proteins were separated on SDS-PAGE gels and transferred to nitrocellulose by semidry blotting. Blots were blocked for 1 h with 5% blotto in PBS plus 0.2% Tween-20 (PBS-T) and incubated with affinity-purified

anti-pkl1 antibodies at a dilution of 1:250 in PBS-T for 1 h at room temperature. Washes were done with PBS-T. HRP-conjugated anti-rabbit secondary antibodies (Jackson ImmunoResearch) were used at 0.3 $\mu\text{g}/\text{ml}$ and detected by enhanced chemiluminescence (ECL) reagents (Amersham) according to the manufacturer's instructions.

Immunofluorescence

Permeabilized cells were prepared as described (Masuda *et al.*, 1990) and stored in liquid nitrogen until use. Cells were spun onto poly-lysine-coated coverslips and fixed with 2.6% paraformaldehyde and 0.2% glutaraldehyde (Electron Microscopy Sciences, Fort Washington, PA). Coverslips were washed three times with PBS, reduced with 2 mg/ml sodium borohydride in 1:1 PBS/methanol, washed, and blocked in PBS, 10 mg/ml bovine serum albumin, and 100 mM lysine-HCl. Coverslips were incubated at 4°C overnight with affinity-purified anti-pkl1 antibody at 1:20 in blocking solution. Coverslips were incubated for 1 h at room temperature with fluorescein-conjugated sheep anti-rabbit secondary antibody (Cappel Research Reagents, Malvern, PA) at 48 $\mu\text{g}/\text{ml}$.

For whole cells, log-phase cultures were fixed with 3% paraformaldehyde for various times ranging from 2 to 30 min and prepared for immunofluorescence as described (Chappell and Warren, 1989). Fixation times of 10–15 min provided the best compromise between good pkl1 staining and preservation of spindle MTs (cytoplasmic MTs were not well preserved). Anti-pkl1 antibody was used at 1:20, and YOL1/34 anti-tubulin antibody (Bioproducts for Science, Indianapolis, IN) was used at 3.5 $\mu\text{g}/\text{ml}$. Fluorescein-conjugated sheep anti-rabbit secondary antibody (Cappel) was used at 48 $\mu\text{g}/\text{ml}$, and rhodamine-conjugated goat anti-rat antibody (Jackson ImmunoResearch) was used at 20 $\mu\text{g}/\text{ml}$. Coverslips were mounted on slides in 90% glycerol containing 1 mg/ml 1,4-diazabicyclo[2,2,2]-octane (Aldrich Chemical, Milwaukee, WI) and 1 $\mu\text{g}/\text{ml}$ 4',6-diamidino-2-phenylindole (DAPI). Slides were observed through a Zeiss Universal Microscope with a Zeiss $\times 100$ Neofluar objective (NA, 1.3; Carl Zeiss, Thornwood, NY) with epifluorescence illumination. Photographs were taken on T-MAX 400 film (Eastman Kodak, Rochester, NY).

Overexpression of pkl1

For mild overexpression, the *pkl1*⁺ cDNA was cloned into the plasmid pSM-2, which contains the SV40 early promoter (Moreno *et al.*, 1991). The *Bam*HI-*Hind*III fragment from pLH-52 containing the 5' end of *pkl1*⁺ was cloned into *Bam*HI-*Hind*III-digested pSM-2. This plasmid was then digested with *Hind*III, and the *Hind*III fragment corresponding to the 3' end of *pkl1*⁺ was cloned into it to create pSM-PKL1. This gives approximately fivefold overexpression, as compared with wild-type. Strains containing pSM-PKL1 were used for Western blotting and immunofluorescence studies.

For greater overexpression, the *pkl1*⁺ cDNA was cloned into the plasmids pREP3X (full-strength *nmt1*⁺ promoter) and pREP81X (weakened *nmt1*⁺ promoter; Basi *et al.*, 1993; Forsburg, 1993). The *nmt1*⁺ promoter is active in the absence of thiamine and repressed in the presence of thiamine. The *pkl1*⁺ cDNA was amplified by PCR from pSM-PKL1 with the following primers: 5'-GGAGCTC-GAGATGCATATGGTAATTGAGAAT-3' and 5'-CCTTGGATC-CGCGGTTAACCCTTTATAATG-3'. This created a 2.5-kb fragment with sites for *Xho*I and *Nsi*I at the 5' end and *Sac*II and *Bam*HI sites at the 3' end (underlined). The *Nsi*I site contains an in-frame ATG; if this ATG is used for initiation of translation, an additional two amino acids, methionine and histidine, would be added at the N terminus of pkl1. The fragment was digested with *Xho*I and *Bam*HI and cloned into pREP3X and pREP81X that had been cut with the same enzymes. The entire insert was verified by sequencing. *S. pombe* strain PN556 was transformed with the resultant plasmids, pREP3X-PKL1 and pREP81X-PKL1, and plated onto selective media containing thiamine. Plasmid transformants were grown to a density of 3–10 $\times 10^6/\text{ml}$ in minimal medium containing 5 $\mu\text{g}/\text{ml}$

thiamine. Cells were resuspended in thiamine-free medium and inoculated into thiamine-free medium at low density ($\sim 2 \times 10^5/\text{ml}$). Several doublings in thiamine-free medium are necessary to lower sufficiently the intracellular thiamine concentration and to induce transcription from the *nmt1*⁺ promoter (Tommasino and Maundrell, 1991). With these conditions, expression began ~ 10 –12 h after the switch to thiamine-free medium. At intervals, samples were taken for cell counting, protein extraction, viability measurements, and immunofluorescence. Cell counting was performed with a hemocytometer. Cell viability was determined by plating cells onto minimal plates supplemented with adenine, uracil, and thiamine (i.e., plasmid selected for, but *nmt1*⁺ promoter repressed). Cells were fixed with paraformaldehyde and glutaraldehyde and processed for immunofluorescence as described (Hagan and Hyams, 1988). The TAT1 anti-tubulin antibody (a kind gift of Professor Keith Gull, University of Manchester, U.K.) was used at 1:15 dilution, and fluorescein-conjugated goat anti-mouse secondary antibody (Sigma) was used at 55 $\mu\text{g}/\text{ml}$. Percentages of different types of spindle structure were determined by counting at least 200 mitotic cells (i.e., those with a spindle or star-shaped structure) for each time point. To determine the degree of overexpression, we made serial (fivefold) dilutions of the pREP3X-PKL1 and pREP3X-PKL1 extracts from the 17-h time point. These dilutions were compared by Western blotting with the anti-pkl1 antibody with pREP81X extract. By inspection of blots, it was determined that pREP81X-PKL1 gives a 10- to 15-fold overexpression, as compared with endogenous pkl1 levels, whereas pREP3X-PKL1 gives a 100-fold overexpression.

MT Pelleting and Bundling

Two fusions were made between GST and the motor domain of pkl1. For GST-PKL1-M, pLH-52 was digested with *Msc*I and *Sma*I and religated. The *Bam*HI-*Eco*RI fragment from the resultant plasmid was then cloned into *Bam*HI-*Eco*RI-digested pGEX-1N; the *pkl1*⁺ fragment used in this construct would encode a polypeptide of 45 kDa. For GST-PKL1-H, the *Hind*III fragment from pLH-52 was cloned into pBluescript SK[−] such that the 3' end of *pkl1*⁺ was closest to the pBluescript *Kpn*I site. The *Bam*HI-*Xho*I fragment from this plasmid was excised and cloned into *Bam*HI-*Xho*I-digested pGEX-20T (this vector was derived from pGEX-2T by modifying the polylinker; it was a gift from Dr. John Armstrong, University of Sussex, U.K.); the *pkl1*⁺ fragment would encode a polypeptide of 52 kDa. GST-PKL1 fusion proteins were purified on glutathione agarose as described (Smith and Johnson, 1988). Tubulin was purified from bovine brain as described previously (Masuda *et al.*, 1990).

For the MT copelleting experiments, GST and fusion proteins were pre-spun at 4°C at 200,000 $\times g$ in a TL100 rotor in a TL100 benchtop ultracentrifuge (Beckman Instruments, Fullerton, CA) to remove aggregates. In a typical 100- μl assay, 7.5 μg of test protein was incubated at room temperature for 20 min in the presence of taxol-stabilized MTs (90 μg of tubulin or buffer) in PEM (100 mM 1,4-piperazinediethanesulfonic acid [PIPES], pH 7, 1 mM ethylene glycol bis(β -aminoethyl)ether-*N,N,N',N'*-tetra-acetic acid [EGTA], and 1 mM MgCl_2) plus 3.3 mM MgCl_2 , 16 μM taxol, and 1 mM guanosine 5'-triphosphate (GTP). In some experiments, ATP or 5'-adenyl-imidodiphosphate (AMP-PNP; Boehringer Mannheim, Indianapolis, IN) was added to 4 mM. Samples were centrifuged at 200,000 $\times g$ at 22°C for 30 min. The supernatant was carefully removed, and the pellet was resuspended in an equal volume of PEM. The supernatant and pellet fractions were analyzed by SDS-PAGE (1/4 of the sample was loaded). For the ATP release experiment, 85 μg of GST-PKL1-M fusion protein was pelleted with MTs in 1.2 ml of buffer at 200,000 $\times g$ (in a TLA100.3 rotor; Beckman), the pellet was carefully rinsed in PEMGT (PEM + 1 mM GTP + 16 μM taxol), and then it was resuspended in PEMGT plus 13 mM MgCl_2 by 10 passes through a 26-gauge needle. The resuspended pellet fraction was divided, and ATP and NaCl were added to different concentrations. After a 20-min incubation at room temperature, the samples were centrifuged and treated as before.

For the MT bundling experiments, rhodamine-labeled MTs were prepared by polymerizing MTs from 9:1 tubulin/rhodamine-tubulin (a kind gift of Dr Tim Mitchison, University of California, San Francisco, CA). The test protein (10–250 ng) was incubated with MTs (200 ng) at 37°C for 20 min in 80 mM PIPES, pH 6.9, 1 mM MgCl₂, and 1 mM EGTA. Samples were then split in two and added to either Mg-ATP (10 mM final concentration) or buffer on a microscope slide and viewed with an epifluorescence microscope. To prevent bleaching and breakage of rhodamine-labeled MTs, an antifade system was present in these experiments (Howard and Hyman, 1993).

ATPase Assay

This spectrophotometric ATPase assay couples ATP hydrolysis to the oxidation of NADH, which is followed by a decrease in absorbance at 340 nm (Catterall and Pederson, 1971). The assay was performed at 37°C, and all components were prewarmed before starting the assay. Two mixes were made: K mix (containing the following assay components [in mM]: 125 KCl, 112.5 imidazole, 12.5 MgCl₂, 5 ATP, 2.5 phospho-enol pyruvate, 1.25 nicotinamide-adenine dinucleotide [NADH], with 7.5 U lactic dehydrogenase and 2.5 U pyruvate kinase) and A mix (containing the following test proteins and prepolymersed MTs or buffer: 400 µg of polymerized tubulin, 0.4 mM GTP, 23 µM taxol, 50 mM PIPES, pH 7, 0.5 mM EGTA, 0.5 mM MgCl₂, 1.25 mM dithiothreitol, and 5–50 µg of test protein). Prewarmed A mix (150 µl) was added to prewarmed K mix (600 µl) in the cuvette, and optical density (OD) readings were taken by a DU-70 spectrophotometer (Beckman) in kinetics mode. The rate was calculated from the slope of absorbance change by using -6.2 OD mM^{-1} .

RESULTS

Isolation of *pk11*⁺ cDNA

We used the anti-peptide antibodies anti-HIPYR and anti-LAGSE, which were raised against highly conserved regions of the motor domain (Sawin *et al.*, 1992b), to screen for novel KRPs in fission yeast. These antibodies recognize several proteins on immunoblots of *S. pombe* extracts in the range that is expected for KRPs (70–140 kDa; see Figure 2B). We used the anti-LAGSE antibody to screen an *S. pombe* expression library and obtained 55 positive clones from the 80,000 plaques screened. Six clones that also cross-reacted with the anti-HIPYR antibody were shown to have overlapping restriction patterns. None of the other clones cross-reacted with the anti-HIPYR antibody, and we have not yet found any additional kinesins in this collection. However, in a similar screen using the anti-HIPYR antibody for the initial screen and anti-LAGSE in a secondary screen, several KRPs have been found: *pk11*⁺, *cut7*⁺, a novel KRP, and some promising candidates (So-ching Brazer, W.Z. Cande, unpublished results).

The longest of the six clones (3 kb) was sequenced and found to contain an open reading frame of 2496 bp. Translation of this open reading frame predicts a protein product of 96 kDa (832 amino acids), which has homology to kinesin. The sequence of *pk11* (*pombe* kinesin-like protein) is presented in Figure 1A, and the locations of the ATP binding region, the

LAGSE, and HIPYR motifs are indicated. The N-terminal half contains two sequences that are similar to the consensus bipartite nuclear localization signal (NLS; Dingwall and Laskey, 1991). The putative motor domain of *pk11* is at the C terminus of the protein, a domain organization it shares with *S. cerevisiae* Kar3p, *A. nidulans* klpA, and *Drosophila* ncd (Goodson *et al.*, 1994). Within the motor domain, *pk11* has the greatest degree of homology to this group of C-terminal motors, sharing 57% identity with klpA (O'Connell *et al.*, 1993) and 54% identity with Kar3p (Figure 1A; Meluh and Rose, 1990), whereas it is only 30–40% identical to KHCs or KRPs belonging to other subfamilies. A conserved motif on the N-terminal side of the motor domain that is specific to this subfamily is also found in *pk11* (Figure 1B; O'Connell *et al.*, 1993). Outside the motor domain there is no significant homology between *pk11* and Kar3-like proteins or any other protein in the GenBank database. Although the N terminus is predicted to be globular, amino acids 196–481 in the central region of the *pk11* protein are predicted to form coiled-coil structures (Figure 1C) that might be involved in dimerization or interaction with other molecules. Hybridization of the *pk11*⁺ cDNA to *S. pombe* P1 and cosmid libraries (Hoheisel *et al.*, 1993) maps *pk11*⁺ to the left arm of chromosome I, close to the centromere but distal to *rad25*⁺ (our unpublished results).

Pk11 Localizes to the Mitotic Spindle

Antibodies were raised to a nonconserved region close to the N terminus of the *pk11* protein (see MATERIALS AND METHODS). Affinity-purified anti-*pk11* antibodies recognize a protein of 96 kDa on Western blots (Figure 2A; WT), in close agreement with the molecular weight predicted from the cDNA sequence. The intensity of this band is increased in extracts from cells overexpressing *pk11* (Figure 2A; OE) and is absent in extracts from a *pk11*-disrupted strain (Figure 2A; ΔP), confirming that this band represents the *pk11* protein.

Pk11 is one of the more minor bands recognized on a Western blot by the anti-LAGSE and anti-HIPYR antibodies. Extracts from wild-type cells contain several proteins in the 70–140-kDa range that cross-react with the anti-LAGSE and anti-HIPYR antibodies (Figure 2B, lanes 1 and 3). When extracts are made from cells overexpressing *pk11* (Figure 2B, lanes 2 and 4), a LAGSE-reactive band of 96 kDa is increased in intensity (Figure 2B, lane 2); for the anti-HIPYR antibody, a band of 96 kDa appears (Figure 2B, lane 4) that comigrates with the *pk11* protein identified with anti-*pk11* antibodies (Figure 2B, lane 5).

Immunofluorescence studies were performed to localize *pk11* in wild-type *S. pombe* cells. The cells used in these experiments were permeabilized with detergent

A

ATAAAAACGCTCTTTTAACTTATAGATGGTAATTGAGAATACAAGGACATTTCCATAAACACAGGGTATAAACACAGGAGATGCTTTAAACACAGATTCTGAGGACTTAAATATATCG 120
M V I E N T K D I S I N T G Y K R Q E D A L N T D S E D L I Y R 32

ACCAAAAAAATTTATTAACAAATCAAGAGGATGCAGTACATGATTGAAATACGAAAAATTCGTATCAAAAAATCACGTTTACAAAGCGACATTAAATGGCAAGAAGAGGACAGCAA 240
P K K I I K T N Q E D A V H D L K Y E N F V S K N H V L Q S D I N G K K R D S N 72

TCGAGACAAGGCCGCTGTAGTCACAGCGCCAATTGCTTCTACTCATGAATCTAATTACGAAGAATCTGTTTCCAAATTTAAAGAGAGTTGGCTGCCTCAATTAAGATCTCATTGAATC 360
R D K A A V V T A P I A S T H E S N Y E E S V S K F K E S W L P Q L K D L I E S 112

TCATAAACCAATATGCAATCAACGTTAGCCTACGAGAGTGATCAAGCTATGCTTCTTCTAATACATTAAAGAGATAAAAGATTAAAAAGCAACAAAAATATTATACAGTTAGA 480
H K T I C E S T L A Y E S D Q A M A S S N T L K R I K D L K S K P K N I I Q L E 152

AAGTTGTCATATGCTATCTAATGGCGAATCGTTTATTATCGAAAGACGAAACCGATGATATGAGTCTGCATCTTGAACCTTAGATCACATCTTCAGGTACAAGAACAGGTTTACGC 600
R L H M L S N G E H R L L S K D E T D D I E S A Y L N L R S H L Q V Q E Q V Y A 192

TGAAAAGGATCATGAATATTCGTTTACAACCTCAATCATACAGAGAGGACGAGAAAAGGCCAAGCAAGATATTTAGAAAACAAAAGAAATCTTCTCTGAATATTCGATATCGAATAT 720
E K D H E Y S L Q L Q S Y R E A A E K A K Q D I L E T K E N L S S E L S I S N I 232

CCAGTTGAAGAGCAAGGAAAGGTTAGAAGCTCGAATGCAAGCTACCAAAAGCTACGTCGAGAACCAAAAGAACTTGCACCTTTATCATGAAAAGAAAACACATTCTCTAGTTTGTAA 840
Q L K E A K E R L E A N A N S Y Q K R E H K E L A L Y H E K T H S K L D A L E 272

TCTAAATGGTGAACGAAATCCTTTGGGATTTGTTGAGAAGCAAGTAAAGTCATATAACATGAATATGCCAATATTTGTGAATCACTTCGGAGAGCTCTAGTGCTGATACAAGGTTTC 960
L N G E R K S F G D F V E N E V K S Y K H E Y A N I C E S L R R A L V L I Q G S 312

TTGTACTGAGAAAAATTTGCGTTTAAAGAAAAAATTTCTGGACTTATTAGAAATGAAACCAAGAAGAAATGATCGAATATCCACATGATTATGAGAATGATTAAACGGTTAAAA 1080
C T E K I L R F K E K I L D L L E M K Q Q E E N D R I S H I D Y E N D L T V K K 352

HindIII
GCTTAAAGGAGAATATCTGAGTTGGAAATGGCAGTAAAGAAATGAATCGGAAAAATCATACTCGGAAAAAGAAATATGAAGAAAAATCTCCTCTGCGTATAGAATTAGAAGCAA 1200
L K R R I S E L E M A V K E Y E S E K S Y S E K E Y E E K I S S L R I E L E D K 392

GTTAGCTGAAATGCATGCTTCGAATAACTGTTAAAGGAAGACATAACACCACCTTACATCTGAAAGAACTTGAAGAACTTTCTAAATATGTTGCAAGTATTCAGATTAAGAAAG 1320
L A E I D L M L K E H K H C H S T S E K L E E L S K Y V A S I Q D K E R 432

MscI
AAATAATGGCAAGCGCATTTGAACTACAAGCTCGAATCCAAGCTGGAACGAGGAGAAATGAAGATATGTATAATAAAGTGTAGCAGAAGAAATATCAGACGAAAGCTGCATAACGA 1440
N N G Q N A L E L Q A R I Q Q L E R R N E D M Y N K L L A E E I I R R K L H N D 472

CATTCAAGAACTAAAGGAAATATTCGAGTTTGTAGGGTTAGACCACTACTCCCAAGTGAAGAAATCCGAATATTCGATTCGTGATGTTTACAATTTCCAGATAAAGACGCACTAGA 1560
I Q E L K G N I R V R L P L P S E E S E Y C I A D V L Q F P D K D A L E 512

GCCACAAAACTTATTCTTAAAGGGCTTAATGTTGAATCATCACTCGGGCATACTTATGATCGTAATTATGAATTTTCCTTTGATCGAGTCTTTGCTCTGAAAGCGATAACTCGTCCGT 1680
P Q K L I L K G P N V E S S L G H T Y D R N Y E F S F D R V F A P E S D N S S V 552

NdeI
TTTTGAGGAAATATCACAGTTAATCAAAAGCGCTATGATGGTTAATGTTAGTATATTTGCATATGGACAACAGGATCGGGTAAGACGTACACAATGTCGTACAGACGGTATGAT 1800
F E E I S Q L I Q S A I D G Y N V S I F A Y G Q T G S G K T Y T M S S Q D G M I 592

AGCGATGTCATTAAGCATATTTAATTTATCTATCAACCTTGAGGGAGAAAGGGTGGGTGATAAAATGAGAGGTCAGTTTGTGAAATTTTACAATGAAACGATTTATGATCTATTAAA 1920
A M S I R K N Y L S T L R E K G V Y K L R G Q F L E I Y N E T I Y D L N L N 632

CAAAGCGAAATGTTGAAAAATCCTAAGCAGATATTCATCATGATGAAAGGAAAGACGAACACAGTGGATAACGTTTCAATAATGATTTTAAATGAAGAGTACTGTTTACAAGAT 2040
K A E M L K N P K H D I H H D E K E R R T T V D N V S I I D F N E E D T V Y K M 672

ClaI
GCTAAATCGAGCAGGAGAAATTCGATTTATAGCAGCGACAAAAGCAATGAACGGTCTTCAAGATCACAACCTGTTTTCATGCTGATATAGATGGTGAATTTCTAGCAACCAACAAAT 2160
L N R A G E N R F I A A T K A N E R S S H T V F M L Y I D G E N S R T K Q I 712

XbaI
ATGTAAAGGTACGCTAAATCTAGTTGATTTGGCAGGTTCTGAGCGTTAAGTTACAGTCAGGAGTGGGGATAGACTACGAGAAACGCAAGCTATTAATAAAGTCTCAGCTGCTCTGG 2280
C K G T L N L V D L A G S E R L S Y S Q A V G D R L R E T Q A I N K S L S C L G 752

CGATGTTATCCATGCATTTGGAATGCCTCAATAGTACCACGAAGGAAAGTCTCACATACCATACCGAAATCAAGCTAACTTATTTGTTAAAAATATTCATTTGGAAGGTTGCCAA 2400
D V I H A L G N A S N S T T K E K S H I P Y R N S K L T Y L L K Y S L G K G A K 792

AACCTGATGTTGTTAATGTGCTCCTTTAAAGTCGCAATTCATGGATACATTAATAGTCTGCGATTTGCAACAAAAGTCAATGATACAAAAGTAGGTTCTATTAAACATTATAACG 2520
T L M F V S P L K S Q F M D T L N S L R F A T K V N D T K V S I K H Y K R 832

HindIII
TTAAGCTTATTTAGAGTAAAAATGTTTACCAAGCATTTAAGATGCTGGGTTTGGTGGAGCAGCTGCTGCTTGTGCTTGAATGATGGCACCTTTAACTTTGCTGAATTTCTTGTGA 2640
.

GATTTTGTGATTTTCGAGGCGTGCACATATAATTTATGAATTAATAAAATGACATACGCAATTCGATTAATACAAATGAAGAAGTACAACCTCGCATTTTTCGAGAGCGATTTTAACTCA 2760
AGGAAATCTCTCTTTTCAAATTTCAACAAACATACATTCTTATTAATGAAATCCAATCTTGTGTTTAAAGTGTGCTTTGCTCTCTCATGACTTTGCTTTACAAGGGTAGGACCGATT 2880
TGCTTATATATATTGAGTCAGGACCCACTTTTCCAATTCCTAATTCGCAATTAACAAAACATTCCTAATATAAAAAAATTTTAAAAA 2969

B

k1pa KLRREETLRRLHNOVELKGNIRVFCVRRLTEN. EGA. .SDAAQTPYD. EGDSKEINIIGPEEKSSFGTVTRKNHNSFDHVPGPSAQNSDVFDEI 496
p1k1 KLLAEELIRRLHNDIQELKGNIRVFCVRRLPES. EESEYCIADVLQPPDKDALEPKLILKGNVSSLGHTYDRNYEFSFDRVPAPESDSSVFEEI 557
kar3 ILIKEETVRLTLHNEQLKGNIRVFCVRRLHALKLENSDLSLNNVEFDNSGVQS.MEVTKIQNTAQVHEPKFDKIPDQQTVDVQFEV 456

k1pa SOLVQSALDGYNVCIFYGQTSKTHM. SSLDGMIPRAVHOIYETATSLLEKGMRYTMEGNFVEVYNNENLNDL.GKAEELDKKLEIRHMQR 590
p1k1 SOLVQSALDGYNVSIFYGQTSKTHM. SSQDMIAHSIKHIFNYLSTLREKGMVYKRGQLEIYNNETIYDL.NKAELKJNPKHDIHDEKE 651
kar3 GOLVQSALDGYNVCIFYGQTSKTHMPLNDGDIIPSTISHFNWINKLTKGMDYKNCFEIYNNENIVLLRSDDNNKEDTSIGLKAEIRHQET 556

k1pa GKTTITDATTVOLESPEMVESLKKRAANRVAATKANERSRSRSHSIFILKLGENDYITGERSBOTLNLVDLAGSERLSHSGATDRLEKTONINRSLSC 690
p1k1 RRTVDNVSIIIDNEEDTVYKLRAGENRFIAATKANERSRSRSHVPLVIDGENSRKQICKGTNLVDLAGSERLSYQAVGDRLEKTONINRSLSC 751
kar3 KTTITNTVTSCKLESSEMEVIELKANKRSTASTASNEHSRSRSHSIFILHLSGSNAKTGAHSYOTLNLVDLAGSERLINVSQVQDRLEKTONINRSLSC 656

k1pa LGDVIALAQG.KKGDIHPYNSKLTLLQPSLGGNSKTLMPVNSPLQHLSETLSLKPAKVVHNTHTGAKQTVRDV* 771
p1k1 LGDVIALGNASNTYKESHIPYNSKLTLLKYSLGKAGKTLMPVNSPLKQPMQDNLNLSRFATKVNNDTKVGSIKHYR* 832
kar3 LGDVIALAQG.DSTRKHIPYNSKLTLLQVSLTGDSKTLMPVNSPSSSHINETLNSLRFASKVNSTRVSRK* 729

C

Figure 1.

in an MT-stabilizing buffer before fixation, a procedure that results in the loss of some soluble cytoplasmic contents (Masuda *et al.*, 1990). Affinity-purified anti-*pk11* antibodies (Figure 3, a–d, top panels) label both short early mitotic spindles (Figure 3, a and b) and longer anaphase spindles (Figure 3, c and d) as the two chromosome masses are separating (DAPI; Figure 3, a–d, bottom panels). Permeabilized *pk11*-disrupted cells showed no evidence of spindle staining with the anti-*pk11* antibodies, indicating that the staining is not due to the presence of a cross-reacting protein in the spindle (our unpublished observations).

We wanted to localize *pk11* in whole cells, partly because we were concerned that the spindle localization in the permeabilized cells could be due to a specific or nonspecific association of *pk11* protein with MTs during the permeabilization procedure. Attempts to immunolocalize *pk11* in whole wild-type cells revealed only diffuse staining throughout the cell. We therefore overexpressed (approximately fivefold) *pk11* from the plasmid pSM-PKL1 and performed immunofluorescence studies with these cells. The most noticeable feature of the staining pattern is coincidence of anti-*pk11* antibody labeling (Figure 3e) with DAPI staining of the DNA (Figure 3f). Differences in the intensity of nuclear staining are probably due to variation in plasmid copy number; there does not seem to be any relationship between staining intensity and position in the cell cycle. As well as the nuclear staining, brighter staining bars were seen in the nuclei of mitotic cells (Figure 3, g–i, top panels); these bars colocalized with anti-tubulin-labeled spindles (Figure 3, g–i, middle panels). The fixation protocol used was a compromise between preserving spindle MTs (long formaldehyde fixation) and preserving *pk11* staining (relatively short fixation with formaldehyde). Short spindles sometimes displayed a bias of staining toward the spindle poles (Figure 3e, arrow), although this pattern was not always observed (Figure 3g). The localization of *pk11* to the mitotic spindle strongly suggests that it may have a role in spindle structure and/or chromosome segregation during mitosis. It is also possible that *pk11* might have a role in meiosis or karyogamy. However, when we used whole cells bearing pSM-PKL1 for anti-*pk11* immunofluorescence, we were unable to detect labeling of meiotic spindles or cytoplasmic MTs.

Figure 2. Western blot analysis of the *pk11* protein. (A) Western blot with affinity-purified anti-*pk11* antibodies. Protein extracts from three strains were electrophoresed on an 8% SDS-polyacrylamide gel, transferred to nitrocellulose, and probed with affinity-purified anti-*pk11* antibodies. WT, wild-type (strain PN556); OE, strain overexpressing *pk11* from a plasmid (wild-type + pSM-PKL1); ΔP , *pk11*-disrupted strain. (B) Anti-LAGSE and anti-HIPYR antibodies recognize several *S. pombe* proteins. Proteins from wild-type (lanes 1, 3, and 5) and wild-type plus pSM-PKL1 (lanes 2 and 4) cells were separated on a 6% SDS-PAGE gel, transferred to nitrocellulose, and probed with various antibodies. Lanes 1 and 2, anti-LAGSE antibody; lanes 3 and 4, anti-HIPYR antibody; lane 5, anti-*pk11* antibody. Markers for molecular mass in kilodaltons are indicated on the right.

Disruption of *pk11*⁺

To determine the effect of removing *pk11* function on cells, we made two disruptions of *pk11*⁺. In the first, a 1255-bp *NdeI* fragment of the cDNA clone was replaced by the *S. pombe* *ura4*⁺ gene, thereby removing 50% of the open reading frame, including part of the motor domain, as diagrammed in Figure 4A, top panel. A genomic clone of *pk11*⁺ (personal communication, C. Troxell and J. R. McIntosh, University of Colorado, Boulder, CO) was used to construct a full-length knockout in which the *pk11*⁺ open reading frame was replaced with the *S. pombe* *his3*⁺ gene (Figure 4A, bottom panel; see MATERIALS AND METHODS). Haploid transformants were analyzed by Southern blotting to determine whether gene replacement had occurred at the *pk11* locus. Strains containing either the partial disruption or the full-length knockout are viable, indicating that *pk11*⁺ is not an essential gene. These strains showed the same phenotypes, and therefore all the data presented are for the full-length disruptant. Analysis of the *pk11*-disrupted strain (*pk11* Δ) by Western blotting showed that the 96-kDa *pk11* band disappeared (see Figure 2A, ΔP). *Pk11* Δ strains showed wild-type growth rates at temperatures from 20 to 36°C, and no gross differences were

Figure 1 (on facing page). Features of the nucleotide and predicted amino acid sequence of *pk11*. (A) Nucleotide sequence of *pk11*⁺ cDNA and predicted amino acid sequence of the *pk11* protein. Relevant restriction sites are indicated. The ATP binding region and the LAGSE and HIPYR regions are underlined. Basic amino acids that may form part of two bipartite nuclear localization signals are marked by asterisks. (B) Alignment of putative motor domains of klpA (amino acids 400–771; O’Connell *et al.*, 1993), *pk11* (amino acids 458–832), and Kar3p (amino acids 365–729; Meluh and Rose, 1990). Identities are indicated by vertical lines; amino acid numbers are indicated. The motif conserved in members of this subfamily (O’Connell *et al.*, 1993) is boxed. (C) The probability of each residue of *pk11* forming part of a coiled-coil structure was determined by using the method of Lupas *et al.* (1991) with a window size of 28 amino acids. These sequence data are available from GenBank/EMBL/DBJ under accession number U63916.

Figure 3. Immunolocalization of *pkl1*. Affinity-purified anti-*pkl1* antibodies were used to immunolocalize *pkl1* in permeabilized wild-type cells (a–d) and in whole cells overexpressing *pkl1* (e–i). (a–d) Wild-type cells were permeabilized in an MT-stabilizing buffer (Masuda *et al.*, 1990), fixed, and labeled for indirect immunofluorescence with anti-*pkl1* (top panels) and DAPI staining (bottom panels). (e–i) Cells containing the pSM-PKL1 plasmid were fixed and then permeabilized and processed for indirect immunofluorescence. (e, and top panels in g–i) Labeling with anti-*pkl1* antibodies. (f, and bottom panels in g–i) DAPI staining of DNA. (g–i, middle panels) Labeling with anti-tubulin antibodies. (e) Shown is a field of cells expressing different levels of *pkl1* in their nuclei; the arrow indicates a cell exhibiting anti-*pkl1* spindle labeling with a poleward bias. Bars, 5 μ m; a–d are different magnification from e–i.

apparent in cell morphology or MT structure (our unpublished observations). Strains disrupted for *pkl1*⁺ were able to mate, form zygotes, and undergo meiosis with either wild-type or *pkl1* Δ strains of the opposite mating type, indicating that, unlike *S. cerevisiae* KAR3, *pkl1*⁺ is not essential for karyogamy (Meluh and Rose, 1990).

Because *pkl1* is predicted to be a microtubule motor protein, we asked whether *pkl1*-disrupted strains have an altered sensitivity to MT-disrupting drugs. The growth of *pkl1* Δ and the parental strain was compared by spotting serial dilutions of log-phase cells on YES plates containing various amounts of the MT-disrupting drug thiabendazole (TBZ). Although *pkl1* Δ and wild-type strains grow well on YES, we found that the growth of *pkl1* Δ colonies was slowed, as compared with the growth of wild-type colonies on YES plus TBZ (Figure 4B) at temperatures between 20 and 25°C. *Pkl1* Δ was also sensitive to another MT-disrupting

drug, benomyl, and in halo assays and viability platings *pkl1* Δ strains showed greater sensitivity to MT-disrupting drugs than wild-type strains (our unpublished results). Although these effects are subtle, it is clear from these experiments that disruption of *pkl1*⁺ causes an increase in sensitivity to MT-disrupting drugs.

Pkl1⁺ Shows a Genetic Interaction with Another KRP-encoding Gene, *cut7*⁺

In both *S. cerevisiae* and *A. nidulans*, genes from the *cut7*/*bimC* group of N-terminal motors have been shown to have a genetic interaction with genes from the Kar3/*ncd* subfamily of C-terminal motors. In *S. cerevisiae*, mutation of both *CIN8* and *KIP1* causes a spindle assembly/stability phenotype at the restrictive temperature, a defect that can be suppressed by deletion of *KAR3* (Saunders and Hoyt, 1992). Deletion

Figure 4. Disruption of *pk11*⁺ causes increased sensitivity to MT-disrupting drugs. (A) Diagram of *pk11*⁺ genomic and cDNA clones. Coding region (5' end at left) is indicated by shading, with the putative motor domain represented by the darker shading. Relevant restriction sites are indicated. For the partial disruption (top panel), a 1255-bp *NdeI* fragment of the cDNA clone was replaced by the *S. pombe ura4*⁺ gene. For the full knockout (bottom panel), the *pk11*⁺ coding region was replaced by the *his3*⁺ gene (see MATERIALS AND METHODS). In this diagram, diagonal stripes represent the arms used for the knockout construct, and heavy stippling represents the probe used in Southern analysis. (B) The growth of wild-type (*h*⁺ *ade6*-M216 *leu1*-32 *ura4*-D18 *his3*-D1) and *pk11*Δ strains was compared on YES plates and on YES plus 12 μg/ml thiabendazole (YES + TBZ). Serial dilutions of the two strains were transferred to the plates and then incubated at 22°C for 3–7 d.

of *KAR3* alone causes a mitotic defect, but strains are viable (Meluh and Rose, 1990). In *A. nidulans*, deletion of *k1pA* suppresses the temperature sensitivity of *bimC4* (Enos and Morris, 1990; O'Connell *et al.*, 1993). Because *pk11* is a sequence homologue of Kar3p and *k1pA*, we sought to determine whether it showed a similar interaction with the *S. pombe* member of the *bimC* family, *cut7* (Hagan and Yanagida, 1990, 1992).

Double mutant strains were made between *pk11*Δ and various alleles of *cut7*. The growth of these double mutant strains was compared with the single mutants and wild-type at various temperatures (Figure 5). Disruption of *pk11*⁺ suppresses the temperature sensitivity of the *cut7* alleles to various degrees: *cut7*-21 and *cut7*-446 are partially suppressed, and *cut7*-22 is almost fully suppressed, whereas *cut7*-24 is not suppressed by deletion of *pk11*⁺. We have examined the pattern of MTs in *cut7*-446 and *cut7*-446*pk11*Δ after 4 h at 28°C (restrictive temperature for the former, semi-permissive for the latter) by anti-tubulin immunofluorescence: almost all mitotic cells in the *cut7*-446 culture contained V- or star-shaped MT structures, and although such structures were also seen in the *cut7*-446*pk11*Δ culture, there was a far higher proportion of normal-looking spindles (our unpublished observations). Therefore, *cut7*⁺ and *pk11*⁺ do show a similar

genetic interaction to those observed in other organisms. These observations in *S. pombe* lend support to the hypothesis that the maintenance of spindle structure is dependent on the existence of balanced opposing forces contributed by plus-end-directed and minus-end-directed microtubule motor proteins (Fuller and Wilson, 1992; Saunders and Hoyt, 1992). This hypothesis received additional support from experiments in which we overexpressed *pk11* in *S. pombe*.

Overexpression of *pk11* to High Levels Causes a Defect in Spindle Bipolarity

Although overexpression of *pk11* to low levels (~5 × endogenous) does not seem to have adverse effects on cells (Figure 3), gross overexpression of the *pk11* protein affected cell growth, viability, and MT organization. *Pk11*⁺ was inserted into the plasmids pREP3X and pREP81X. These multicopy plasmids allow thiamine-repressible expression from the *nmf1*⁺ promoter, which is full strength in pREP3X and weakened in pREP81X. Cells containing pREP3X-PKL1 and pREP81X-PKL1 were grown in the presence of thiamine, and then expression was derepressed by transferring cells to media lacking thiamine. Derepression takes several generations because of thiamine scav-

Figure 5. Disruption of *pk11*⁺ partially suppresses a mutation in *cut7*. Serial dilutions of various strains were transferred to YES plates that then were incubated for 3–7 d at the indicated temperatures (see MATERIALS AND METHODS).

enging, and we began our analysis 10–12 h after the shift to fresh medium. Overexpression of *pk11* slowed the increase in cell density, as shown in the growth curves (Figure 6A, top panel). This was accompanied by a decrease in cell viability (Figure 6A, middle panel). The severity of these effects correlated with the level of *pk11* protein as analyzed by Western blotting (Figure 6A, bottom panel). Although expression from pREP81X-PKL1 was ~10–15 fold higher than in wild-type or in the repressed state, levels of expression from pREP3X-PKL1 were on the order of 100 times higher than that found in wild-type cells.

To determine whether gross overexpression of *pk11* protein was accompanied by changes in MT organization, we prepared cells for immunofluorescence by using an anti-tubulin antibody. In the strains overexpressing *pk11*, we found many cells that were elongated and contained highly condensed chromatin, features typical of mitotically blocked cells. Many cells contained V-shaped or star-shaped MT structures that coincided with the highly condensed chromosomes (Figure 6B). These MT structures were very reminiscent of the aberrant spindles seen in *cut7* mutants at the restrictive temperature (Hagan and Yanagida, 1990), and we interpret them to be spindles that have failed in pole separation. The numbers of these aberrant spindles increased as the level of *pk11* increased, and there was a concomitant decrease in the number of normal-looking spindles in the population (Figure 3C). At 22 h after the switch to thiamine-free medium, there were very few normal-looking spindles in the pREP3X-PKL1 culture. At this time, the pREP81X-PKL1 culture still contained a significant proportion of normal-looking spindles, consistent with the fact that this strain continues to grow in the absence of thiamine, albeit at a reduced rate.

Bacterially Expressed *pk11* Fusion Proteins Exhibit Microtubule-stimulated ATPase Activity

To determine whether *pk11* exhibits the biochemical properties expected of a kinesin-related motor protein, we expressed a series of *pk11* fusion proteins in bacteria and used these for in vitro studies. The C-terminal motor domain plus various portions of the stalk/tail region of *pk11* were fused to GST and expressed in bacteria. We found that fusions containing very large parts of the stalk/tail were not soluble in bacteria, and we therefore used two shorter fusions (Figure 7) for the biochemical studies. We used a spectrophotometric assay (see MATERIALS AND METHODS) to measure the ATPase activity of *pk11* fusions. Although purified GST showed no detectable ATPase activity, GST-PKL1-M and GST-PKL1-H had very low but detectable ATPase activities in this assay. Addition of taxol-stabilized microtubules to the assay enhanced the ATPase activities of GST-PKL1-M and GST-PKL1-H (but not GST) considerably, to 2.1 and 1.7 $\mu\text{mol}/\text{min}/\text{mg}$, respectively. Thus, MTs stimulated the ATPase activity of GST-PKL1-M 120-fold and that of GST-PKL1-H 25-fold.

***Pk11* Fusion Proteins Bind Microtubules**

We performed microtubule copelleting assays to determine whether *pk11* binds directly to MTs. The data presented in Figure 8 are for GST-PKL1-M, although similar results were obtained for GST-PKL1-H. Fusion protein preparations were subjected to a high-speed spin to remove protein aggregates and then mixed with taxol-stabilized MTs and incubated at room temperature before being centrifuged to pellet the MTs. Supernatant and pellet samples were then analyzed by SDS-PAGE (Figure 8A). GST does not pellet in the

Figure 6. Overexpression of *pk11* affects cell growth, viability, and spindle structure. (A) Strains containing the plasmids pREP81X, pREP3X-PKL1, and pREP81X-PKL1 were transferred to thiamine-free medium and grown for several hours to allow derepression of expression. (Top panel) Growth curve of the various strains; (middle panel) viability plots; (bottom panels) Western blots with anti-*pk11* antibodies of extracts taken at time points indicated. The data for the strain containing pREP3X are similar to those for the pREP81X strain. (B) Immunofluorescence of cells bearing pREP81X-PKL1 plasmid. Cells (17-h time point) were fixed and processed for immunofluorescence with the TAT1 anti-tubulin antibody (left panels) and DAPI staining (right panels). A wild-type mitotic cell is shown in the top panels for comparison. Width of panel, 15 μm . (C) Spindle morphology in cultures overexpressing *pk11*. Cells were fixed and processed for anti-tubulin immunofluorescence and DAPI staining at various times after the switch to thiamine-free medium. The column labeled "none" lists data for cells containing pREP3X or pREP81X (taken at the 17-h time point). At least 200 cells judged to be in a mitotic state (i.e., those containing spindles or star-shaped MT structures) were counted for each time point.

absence or presence of MTs (our unpublished observations). In the absence of MTs, all of the GST-PKL1-M fusion protein remains in the supernatant (Figure 8A, lane pair 1). GST-PKL1-M pellets with added MTs either in the absence of nucleotide (apart from the 1 mM GTP present; Figure 8A, lane pair 2) or in the presence of 4 mM AMP-PNP, a nonhydrolyzable analogue of ATP (Figure 8A, lane pair 3). In the presence

of 4 mM ATP, a proportion of the GST-PKL1-M protein remains in the supernatant (Figure 8A, lane pair 4). GST-PKL1-M that had been pelleted with MTs could be released from MTs by the addition of ATP and salt (Figure 8B). Approximately one-third of the GST-PKL1-M is released from the MT pellet with 10 mM ATP (Figure 8B, lane pair 2). The level of protein release can be augmented by addition of 50 mM NaCl

Figure 7. ATPase activity of pkl1 fusion proteins. A diagram of the pkl1 protein is shown in the top left; predicted domains of the protein and amino acid residue numbers are indicated. Diagrams of the fusion proteins are shown below it; amino acid residue numbers are indicated. Both fusion proteins contain part of the region that is predicted to form coiled-coil (stalk). ATPase activities in the absence or presence of MTs and fold stimulation for the fusion proteins GST-PKL1-M and GST-PKL1-H and for GST are indicated.

(Figure 8B, lane pair 4) or 100 mM NaCl (Figure 8B, lane pair 6), whereas these concentrations of NaCl added alone have no effect on release (Figure 8B, lane pairs 3 and 5). These experiments indicate that PKL1 fusion proteins show nucleotide-sensitive MT binding.

To determine whether PKL1 is capable of bundling MTs, we incubated fusion proteins with rhodamine-labeled MTs and observed them with an epifluorescence microscope in the presence or absence of 10 mM ATP. In the absence of ATP, both GST-PKL1-M and GST-PKL1-H caused extensive bundling of MTs (as shown in Figure 9b for GST-PKL1-M), whereas MTs alone, or in those incubated in the presence of GST (Figure 9a), did not show any bundling. MT bundles were disrupted by the addition of ATP (Figure 9c). We tested the possibility that the bundling activity was due simply to dimerization through the GST portion of the fusion protein (Walker *et al.*, 1993) by preparing soluble PKL1-H by thrombin cleavage. These preparations also showed ATP-sensitive bundling of MTs (our unpublished results). Both PKL1-M and PKL1-H fusion proteins include part of the region that is predicted to form coiled-coil, and presumably oligomerization is occurring through this region. The MT pelleting and bundling experiments indicate that bacterially expressed PKL1 fusion proteins bind to MTs in a nucleotide-sensitive manner.

DISCUSSION

Features of the pkl1 Sequence

Using antibodies that recognize conserved regions of the kinesin motor domains, we have identified a novel kinesin-related protein, pkl1. By sequence homology and domain organization, pkl1 clearly belongs to the Kar3/ncd subfamily of C-terminal motors (Goodson *et al.*, 1994). Within the motor domain pkl1 has a high degree of homology to other members of this family (klpA, 57%; Kar3p, 54% identical; ncd 48%), but outside the motor domain there is no significant homology. This lack of homology has become a common theme in the analysis of kinesin-related proteins, and the relationship between sequence homology and con-

servation of function within the kinesin superfamily is far from clear. For instance, *S. cerevisiae* Cin8p/Kip1p, *A. nidulans* bimC, and *S. pombe* cut7 seem to have similar functions in bipolar spindle formation (Enos and Morris, 1990; Hagan and Yanagida, 1990; Hoyt *et al.*, 1992; Roof *et al.*, 1992), and although they fall into the same subfamily on the basis of motor domain homology, they do not have significant homology outside the motor domain. Regions of nonmotor domain homology do exist within the kinesin superfamily. KHCs are well conserved along their length (Yang *et al.*, 1989; Kosik *et al.*, 1990; Wright *et al.*, 1991; Navone *et al.*, 1992); human MKLP1 and hamster CHO1 are very similar for most of their length (Nislow *et al.*, 1992; Kuriyama *et al.*, 1994), and it is likely that they are functional homologues in these evolutionarily close organisms. On the other hand, two members of the "motor-in-the-middle" family, MCAK and KIF2, are conserved along their length (~50%; 73% within the motor domain), yet KIF2 is thought to be involved in vesicle transport (Noda *et al.*, 1995) whereas MCAK localizes to centromeres and is thought to have a mitotic role (Wordeman and Mitchison, 1995). These observations are puzzling because we expect proteins that perform the same function in different organisms to have similar cargoes and accessory subunits and that this would correlate with some sequence conservation in the nonmotor regions. Interestingly, klpA lacks sequence homology to Kar3p outside the motor domain, yet it is able to complement the karyogamy defect of a *kar3Δ* strain to some degree (O'Connell *et al.*, 1993). It will be of interest to determine whether pkl1, which is not required for karyogamy in *S. pombe*, is also able to substitute for Kar3p in this process in *S. cerevisiae*. It is possible that classes of KRPs share tertiary structure of which we are not yet aware. In the meantime, sequence comparisons may provide useful clues about function, but it is important to be aware that this may be an oversimplification.

The closed mitosis of fungi presents an additional protein-targeting problem as compared with higher eukaryotes: getting the components of the spindle into the nucleus when they are required. Pkl1 contains two sequences with similarity to NLSs. Nuclear targeting

Figure 8. Pk11 fusion proteins bind to MTs in a nucleotide-sensitive manner. (A) SDS-PAGE analysis of MT copelleting assay. Incubations were centrifuged and separated into supernatant (S) and pellet (P) fractions and then analyzed on a 12% SDS-PAGE gel with Coomassie blue staining. Lane pair 1, GST-PKL1-M was incubated in buffer alone. Lane pair 2, GST-PKL1-M was incubated in the presence of MTs. Lane pair 3, GST-PKL1-M was incubated in the presence of MTs and 4 mM AMP-PNP. Lane pair 4, GST-PKL1-M was incubated in the presence of MTs and 4 mM ATP. Molecular mass markers in kilodaltons are indicated on the right. (B) ATP and salt-dependent release of pk11 fusion proteins from MTs. An MT pellet containing bound GST-PKL1-M fusion protein was resuspended, split, and incubated in various combinations of ATP and NaCl before being recentrifuged. The supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE and silver staining. Lane pair 1, no added ATP or NaCl; lane pair 2, 10 mM ATP; lane pair 3, 50 mM NaCl; lane pair 4, 10 mM ATP and 50 mM NaCl; lane pair 5, 100 mM NaCl; lane pair 6, 10 mM ATP and 100 mM NaCl. Molecular mass markers in kilodaltons are indicated on the right. The apparent mobility shift seen between the + and - MT lanes was not seen in all similar gels and is probably due to the large quantity of tubulin in the +MT lanes.

sequences are diverse, but all seem to contain basic amino acids. A consensus for bipartite NLSs consists of two clusters of basic amino acids separated by a 10-amino-acid spacer (Dingwall and Laskey, 1991). In each of the putative NLSs identified in pk11, the two basic regions are separated by 14 amino acids; NLSs with such spacing have been shown to function in vivo (Robbins *et al.*, 1991). We found that pk11 localized to the nuclei as well to the spindle in cells over-expressing pk11, suggesting that the NLS-like sequences are functional.

Properties of pk11

Bacterially expressed pk11 displays several properties expected for a kinesin motor protein: MT binding, bundling, and ATPase activity. Pk11 motor domain

fusions display a low basal ATPase activity that is strongly stimulated by the addition of MTs. The ATPase activity of pk11 fusion proteins in the presence of MTs is within the wide range of ATPase activities reported for other members of the kinesin family (Chandra *et al.*, 1993; Bloom and Endow, 1994). A motor domain fragment of bovine KHC shows elevated ATPase activity, as compared with the full-length protein (Kuznetsov *et al.*, 1989); truncations of bacterially expressed ncd have enhanced ATPase activity (Chandra *et al.*, 1993). Such observations suggest that nonmotor regions, or light chains, may modulate motor ATPase activity (Kuznetsov *et al.*, 1989; Hackney *et al.*, 1991). This phenomenon may contribute to the relatively high ATPase activity observed for the pk11 motor domain fusions. Stimulation of ATPase activity by MTs is on the order of 25- to 120-fold for pk11 fusion proteins; values of twofold to >1000-fold have been observed for other kinesins/KRPs (Hackney, 1988; Chandra *et al.*, 1993).

Pk11 fusion proteins exhibit nucleotide-sensitive MT-binding in copelleting assays. GST-PKL1-M binds tightly to MTs in the absence of nucleotide, and relatively high concentrations of ATP in combination with salt are required to release it. Kinesins are thought to be highly processive enzymes that may spend >99% of their time bound to the microtubule (Block *et al.*, 1990). In addition to their ability to bind MTs, PKL1 fusion proteins also bundle MTs in vitro. Bundling activity requires two MT binding sites within the same molecule or complex. Bundling could be due to dimerization of the GST portion of the fusion proteins (Walker *et al.*, 1993), but we still observed bundling with PKL1-H protein that had been cleaved from GST (our unpublished results). Both fusion proteins contain regions of pk11 that are predicted to form coiled-coils, structures implicated in oligomerization. Motor domain fragments of two other KRPs, CHO1 and CHO2, were also observed to cause extensive bundling of MTs in vitro (Kuriyama *et al.*, 1994, 1995). Kar3p and ncd have two MT binding sites per monomer—one in the C-terminal motor domain and the other in a basic, proline-rich region at the N terminus (Meluh and Rose, 1990; Chandra *et al.*, 1993). However, we note that pk11 lacks a similar basic region at its N terminus.

Our attempts to determine the motility properties of pk11 in vitro have thus far been unsuccessful. Although GST-PKL1 fusion proteins bind MTs to glass and bundle MTs in motility assays, we have not observed any MT gliding with the addition of ATP. On the basis of its similarity to the minus-end-directed motors Kar3p, CHO2, and ncd, our assumption is that pk11 is also a minus-end-directed motor (McDonald *et al.*, 1990; Walker *et al.*, 1990; Endow *et al.*, 1994b; Kuriyama *et al.*, 1995).

Figure 9. Pkl1 fusion protein causes bundling of MTs. Rhodamine-labeled MTs were incubated with (a) GST or (b and c) GST-PKL1-M. (c) ATP was added to 10 mM. Bar, 10 μ m.

pkl1 May Function in the Spindle

Our evidence that *pkl1* has a role in the spindle is based on localization data, on the phenotype associated with its overexpression, and on its genetic interaction with *cut7*⁺, which encodes a KRP important in spindle function (Hagan and Yanagida, 1990, 1992). We also have preliminary evidence that moderate overexpression of *pkl1* causes an increase in the rate of mini-chromosome loss (A. Pidoux, W. Z. Cande, L. Clarke and J. Carbon, unpublished results).

Immunofluorescence experiments with both permeabilized cells and whole cells overexpressing *pkl1* indicate that a significant proportion of *pkl1* resides in the mitotic spindle. Our attempts to immunolocalize *pkl1* in whole wild-type cells have not been successful, probably because of a low expression of *pkl1* and high background. Two other C-terminal motors, *ncd* and *CHO2*, have been localized to spindles (Hatsumi and Endow, 1992; Kuriyama *et al.*, 1995). *Pkl1* is the first fungal C-terminal KRP with a specific spindle localization. In contrast, *Kar3- β* -galactosidase fusion proteins have been localized to budding yeast spindle pole bodies and to cytoplasmic MTs after pheromone treatment (Meluh and Rose, 1990), but it has been observed only rarely on spindle MTs (Page *et al.*, 1994). *Cut7* localizes to both mitotic and meiotic spindles, but we have been unable to detect *pkl1* on meiotic spindles by immunofluorescence. Likewise, we have not seen localization to cytoplasmic microtubules. It is not known whether this reflects a true absence of the *pkl1* protein from these structures or is due to technical reasons, such as low abundance or fixation problems.

Pkl1-disrupted cells have an increased sensitivity to the anti-MT drugs benomyl and TBZ. These benzimidazole drugs act by shifting the MT equilibrium toward depolymerization, destabilizing MT structures (Davidse, 1986). Wild-type spindles are presumably able to withstand a certain level of destabilization. Perhaps the increased sensitivity of *pkl1*-disrupted cells to MT-drugs is due to the compound effects of MT destabilization caused by the loss of *pkl1* function and by the drugs. How might loss of a KRP such as *pkl1* destabilize spindle MTs? Motors may contribute to spindle stability through their ability to cross-link MTs into bundles, and we have in vitro evidence that *pkl1* is

capable of bundling MTs. Abolition of a motor function could also disrupt the balance of forces in the spindle (as discussed below). Alternatively, motors could act as the hypothetical coupling molecules that attach MTs to structures such as kinetochores and spindle pole bodies (Desai and Mitchison, 1995; Lombillo *et al.*, 1995a,b); loss of such a motor might disrupt these connections or affect the balance between polymerization and depolymerization at MT ends.

Disruption of *pkl1*⁺ suppresses temperature-sensitive mutations in *cut7*, a KRP with an N-terminal motor domain. It is intriguing that disruption of *pkl1*⁺ shows allele specificity in its suppression of *cut7* and is able to suppress both recessive (*cut7-21*, *-22*) and semi-dominant (*cut7-446*) alleles of *cut7* (Iain Hagan, personal communication), although it does not suppress another recessive allele of *cut7* (*cut7-24*). Strikingly, overexpression of *pkl1* to high levels causes a phenotype similar to that seen in *cut7* mutants at the restrictive temperature. As the level of *pkl1* rises, there is an accumulation of V-shaped and star-shaped MT structures in the cell population. These structures coincide with highly condensed chromatin, and we interpret them to be aberrant spindles that have failed to achieve bipolarity. Such structures could also arise through the collapse of an already formed spindle. Our observations suggest that counteracting forces are created in the spindle by *cut7* (outward pushing) and *pkl1* (inward pulling) and that spindle integrity relies on the maintenance of balance between these forces. Another possibility is that overexpression of *pkl1* causes a *cut7*-like phenotype by preventing binding of *cut7* or other KRPs to MTs.

There is now evidence from three different organisms to support the model proposed by Saunders and Hoyt (1992) that the *Kar3*-like and *bimC*-like classes of motors provide opposing forces in the mitotic spindle and that this balance of forces is required to establish and maintain spindle structure and integrity. Although the actual polarities of the majority of these motors are unknown, the polarity of at least one member from each class has been established: *Xenopus* Eg5 is plus-end-directed (Sawin *et al.*, 1992a) and *Kar3p* is minus-end-directed (Endow *et al.*, 1994b), polarities that are consistent with the model. Where in the spindle do these motors act? *Cut7* is localized at spindle

pole bodies and on spindle MTs and is thought to achieve the reorientation and separation of the spindle poles by moving toward the plus-ends of MTs emanating from the opposite pole (Hagan and Yanagida, 1992). It was proposed that Kar3p would act at kinetochores to produce an inward pulling force (Saunders and Hoyt, 1992), and there is evidence that Kar3p binds to *S. cerevisiae* centromeres (Middleton and Carbon, 1994). A minus-end-directed motor could also produce an inward pulling force if it acted at the spindle poles or in the MT overlap zone. For a minus-end-directed motor to produce an inward "collapsing" force at a position other than the kinetochores, it would need to be attached to one spindle pole or the MTs from that spindle pole and move on MTs emanating from the opposite pole. Our immunolocalization data are not precise enough to answer questions about the sites of force generation, although we do observe a bias of pkl1 staining at the spindle poles in some cells. It is to be hoped that immuno-EM studies will allow a more detailed analysis of pkl1 distribution, although high levels of protein do not necessarily correspond to regions of greatest force generation.

It is becoming increasingly clear that motor proteins do not just move objects about on a static MT lattice but play an active part in spindle organization. We propose that pkl1 is involved in such an organizational function. Pkl1 could also have other functions, such as acting as a governor to regulate the initiation or rate of anaphase B. If pkl1 acted at kinetochores, it could contribute to the polewards movements of chromosomes during prometaphase and/or anaphase A; if it acted at spindle pole bodies, it could help focus MTs at the poles (Endow *et al.*, 1994a). KRPs may turn out to be multifunctional and fulfill several roles in the cell. In addition, they might have redundant or overlapping functions. The nonessential nature of *pkl1*⁺ may be due to functional redundancy with other KRP(s). Another C-terminal class KRP, klp2, has been identified in *S. pombe* (personal communication, Cindy Troxell and J. R. McIntosh, University of Colorado, Boulder, CO). At least one additional KRP has been identified (So-ching Brazier and W.Z.C.). The genetic tractability of fission yeast should facilitate an understanding of the roles of KRPs in the cell and their functional redundancy.

ACKNOWLEDGMENTS

We are very grateful to Cindy Troxell and Dick McIntosh for providing us with the genomic clone of *pkl1*⁺ and for sharing their unpublished work on *klp2*⁺. We thank Ken Sawin, Linda Wordeman, and Tim Mitchison for the anti-LAGSE and anti-HIPYR antibodies. We thank Rachel Bartlett, Kathy Gould, Iain Hagan, Paul Nurse, and Mitsuhiro Yanagida for *S. pombe* strains; Keith Gull for the anti-TAT1 antibody; John Armstrong, Susan Forsburg, and Kathy Gould for plasmids; Margit Foss for fusion protein; Tim Mitchison for rhodamine-tubulin; Doug Kellogg for advice on af-

finity purification; and Ron Vale for advice on the ATPase assay. We also thank past and present members of the Cande lab for helpful discussions and Kathryn Ayscough, Laurie Bost-Usinger, Kevin Hardwick, and Harrison Wein for critical reading of this manuscript. This work was supported by a National Institutes of Health grant GM-23238 to W.Z.C. and a Science and Engineering Research Council (U.K.) postdoctoral research fellowship to A.L.P.

REFERENCES

- Barbet, N., Muriel, W.J., and Carr, A.M. (1992). Versatile shuttle vectors and genomic libraries for use with *Schizosaccharomyces pombe*. *Gene* 114, 59–66.
- Basi, G., Schmid, E., and Maundrell, K. (1993). TATA box mutations in the *Schizosaccharomyces pombe nmt1* promoter affect transcription efficiency but not the transcription start point or thiamine repressibility. *Gene* 123, 131–136.
- Block, S.M., Goldstein, L.S.B., and Schnapp, B.J. (1990). Bead movement by single kinesin molecules studied with optical tweezers. *Nature* 348, 348–352.
- Bloom, G.S., and Endow, S.A. (1994). Motor proteins 1: kinesins. *Protein Profile* 1, 1059–1116.
- Broker, M. (1987). Transformation of intact *Schizosaccharomyces pombe* cells with plasmid DNA. *BioTechniques* 5, 516–518.
- Carbon, J., and Clarke, L. (1990). Centromere structure and function in budding and fission yeasts. *New Biol.* 2, 10–19.
- Catterall, W.A., and Pederson, P.L. (1971). Adenosine triphosphatase from rat liver mitochondria. *J. Biol. Chem.* 246, 4987–4994.
- Chandra, R., Salmon, E.D., Erikson, H.P., Lockhart, A., and Endow, S.A. (1993). Structural and functional domains of the *Drosophila ncd* microtubule motor protein. *J. Biol. Chem.* 268, 9005–9013.
- Chappell, T.G., and Warren, G. (1989). A galactosyltransferase from the fission yeast *Schizosaccharomyces pombe*. *J. Cell Biol.* 109, 2693–2702.
- Chikashige, Y., Kinoshita, N., Nakaseko, Y., Matsumoto, T., Murakami, S., Niwa, O., and Yanagida, M. (1989). Composite motifs and repeat symmetry in *Schizosaccharomyces pombe* centromeres: direct analysis by integration of *NotI* restriction sites. *Cell* 57, 739–752.
- Cole, D.G., Cande, W.Z., Baskin, R.J., Skoufias, D.A., Hogan, C.J., and Scholey, J.M. (1992). Isolation of a sea urchin egg kinesin-related protein using peptide antibodies. *J. Cell Sci.* 101, 291–301.
- Coue, M., Lombillo, V.A., and McIntosh, J.R. (1991). Microtubule depolymerization promotes particle and chromosome movement in vitro. *J. Cell Biol.* 112, 1165–1176.
- Davidse, L.C. (1986). Benzimidazole fungicides: mechanism of action and biological impact. *Annu. Rev. Phytopathol.* 24, 43–65.
- Desai, A., and Mitchison, T.J. (1995). A new role for motor proteins as couplers to depolymerizing microtubules. *J. Cell Biol.* 128, 1–4.
- Ding, R., McDonald, K.L., and McIntosh, J.R. (1993). Three-dimensional reconstruction and analysis of mitotic spindles from the yeast *Schizosaccharomyces pombe*. *J. Cell Biol.* 120, 141–151.
- Dingwall, C., and Laskey, R.A. (1991). Nuclear targeting sequences: a consensus? *Trends Biochem. Sci.* 16, 478–481.
- Endow, S.A., Chandra, R., Komma, D.J., Yamamoto, A.H., and Salmon, E.D. (1994a). Mutants of the *Drosophila ncd* microtubule motor protein cause centrosomal and spindle pole defects in mitosis. *J. Cell Sci.* 107, 859–867.
- Endow, S.A., and Hatsumi, M. (1991). A multimember kinesin gene family in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 88, 4424–4427.

- Endow, S.A., Kang, S.J., Satterwhite, L.L., Rose, M.D., Skeen, V.P., and Salmon, E.D. (1994b). Yeast Kar3 is a minus-end microtubule motor protein that destabilizes microtubules preferentially at the minus-ends. *EMBO J.* 13, 2708–2713.
- Endow, S.A., and Titus, M.A. (1992). Genetic approaches to molecular motors. *Annu. Rev. Cell Biol.* 8, 29–66.
- Enos, A.P., and Morris, N.R. (1990). Mutation of a gene that encodes a kinesin-like protein blocks nuclear division in *A. nidulans*. *Cell* 60, 1019–1028.
- Fishel, B., Amstutz, H., Baum, M., Carbon, J., and Clarke, L. (1988). Structural organization and functional analysis of centromeric DNA in the fission yeast *Schizosaccharomyces pombe*. *Mol. Cell. Biol.* 8, 754–763.
- Forsburg, S.L. (1993). Comparison of *Schizosaccharomyces pombe* expression systems. *Nucleic Acids Res.* 21, 2955–2956.
- Forsburg, S.L., and Nurse, P. (1991). Cell cycle regulation in the yeasts *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*. *Annu. Rev. Cell Biol.* 7, 227–256.
- Fuller, M.T., and Wilson, P.G. (1992). Force and counterforce in the mitotic spindle. *Cell* 71, 547–550.
- Funabiki, H., Hagan, I., Uzawa, S., and Yanagida, M. (1993). Cell cycle-dependent specific positioning and clustering of centromeres and telomeres in fission yeast. *J. Cell Biol.* 121, 961–976.
- Goldstein, L.S.B. (1993). With apologies to Scheherazade: tails of 1001 kinesin motors. *Annu. Rev. Genet.* 27, 319–351.
- Goodson, H.V., Kang, S.J., and Endow, S.A. (1994). Molecular phylogeny of the kinesin family of microtubule motor proteins. *J. Cell Sci.* 107, 1875–1884.
- Hackney, D.D. (1988). Kinesin ATPase: rate limiting ADP release. *Proc. Natl. Acad. Sci. USA* 85, 6314–6318.
- Hackney, D.D., Levitt, J.D., and Wagner, D.D. (1991). Characterization of $\alpha_2\beta_2$ and α_2 forms of kinesin. *Biochem. Biophys. Res. Commun.* 174, 810–815.
- Hagan, I.M., and Hyams, J.S. (1988). The use of cell division cycle mutants to investigate the control of microtubule distribution in the fission yeast *Schizosaccharomyces pombe*. *J. Cell Sci.* 89, 343–358.
- Hagan, I., and Yanagida, M. (1990). Novel potential mitotic motor protein encoded by the fission yeast *cut7⁺* gene. *Nature* 347, 563–566.
- Hagan, I., and Yanagida, M. (1992). Kinesin-related cut7 protein associates with mitotic and meiotic spindles in fission yeast. *Nature* 356, 74–76.
- Harlow, E., and Lane, D. (1988). *Antibodies: A Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Hatsumi, M., and Endow, S.A. (1992). The *Drosophila claret* microtubule motor protein is spindle-associated in meiotic and mitotic cells. *J. Cell Sci.* 103, 1013–1020.
- Hogan, C.J., Wein, H., Wordeman, L., Scholey, J.M., Sawin, K.E., and Cande, W.Z. (1993). Inhibition of anaphase spindle elongation in vitro by a peptide antibody that recognizes kinesin motor domain. *Proc. Natl. Acad. Sci. USA* 90, 6611–6615.
- Hoheisel, J.D., Maier, E., Mott, R., McCarthy, L., Grigoriev, A.V., Schalkwyk, L.C., Nizetic, D., Francis, F., and Lehrach, H. (1993). High-resolution cosmid and P1 maps spanning the 14-Mb genome of the fission yeast *Schizosaccharomyces pombe*. *Cell* 73, 109–120.
- Howard, J., and Hyman, A.A. (1993). Preparation of marked microtubules for the assay of the polarity of microtubule-based motors by fluorescence microscopy. *Motility assays for motor proteins. Methods Cell Biol.* 39, 105–113.
- Hoyt, M.A., He, L., Loo, K.K., and Saunders, W.S. (1992). Two *Saccharomyces cerevisiae* kinesin-related gene products required for mitotic spindle assembly. *J. Cell Biol.* 118, 109–120.
- Kondo, S., Sato-Yoshitake, R., Noda, Y., Aizawa, H., Nakata, T., Matsuura, Y., and Hirokawa, N. (1994). KIF3A is a new microtubule-based anterograde motor in the nerve axon. *J. Cell Biol.* 125, 1095–1107.
- Koshland, D.E., Mitchison, T.J., and Kirshner, M.W. (1988). Polewards chromosome movement driven by microtubule depolymerization in vitro. *Nature* 331, 499–504.
- Kosik, K.S., Orecchio, L.D., Schnapp, B., Inouya, H., and Neve, R.L. (1990). The primary structure and analysis of the squid kinesin heavy chain. *J. Biol. Chem.* 265, 3278–3283.
- Kuriyama, R., Dragas-Granoic, S., Maekawa, T., Vassilev, A., Khodjakov, A., and Kobayashi, H. (1994). Heterogeneity and microtubule interaction of the CHO1 antigen, a mitosis-specific kinesin-like protein: analysis of subdomains expressed in insect sf9 cells. *J. Cell Sci.* 107, 3485–3499.
- Kuriyama, R., Kofron, M., Essner, R., Kato, T., Dragas-Granoic, S., Omoto, C.K., and Khodjakov, A. (1995). Characterization of a minus-end-directed kinesin-like motor protein from cultured mammalian cells. *J. Cell Biol.* 129, 1049–1059.
- Kuznetsov, S.A., Vasiberg, Y.A., Rothwell, S.W., Murphy, D., and Gelfand, V.I. (1989). Isolation of a 45-kDa fragment from the kinesin heavy chain with enhanced ATPase and microtubule-binding activities. *J. Biol. Chem.* 264, 589–595.
- Lillie, S.H., and Brown, S.S. (1992). Suppression of a myosin defect by a kinesin-related gene. *Nature* 356, 358–361.
- Lombillo, V.A., Nislow, C., Yen, T.J., Gelfand, V.I., and McIntosh, J.R. (1995a). Antibodies to the kinesin motor domain and CENP-E inhibit microtubule depolymerization-dependent motion of chromosomes in vitro. *J. Cell Biol.* 128, 107–115.
- Lombillo, V.A., Stewart, R.J., and McIntosh, J.R. (1995b). Minus-end-directed motion of kinesin-coated microspheres driven by microtubule depolymerization. *Nature* 373, 161–164.
- Lupas, A., Van Dyke, M., and Stock, J. (1991). Predicting coiled-coils from protein sequences. *Science* 252, 1162–1164.
- Masuda, H., Hirano, T., Yanagida, M., and Cande, W.Z. (1990). In vitro reactivation of spindle elongation in fission yeast *nuc2* mutant cells. *J. Cell Biol.* 110, 417–425.
- Masuda, H., Sevik, M., and Cande, W.Z. (1992). In vitro microtubule-nucleating activity of spindle pole bodies in fission yeast *Schizosaccharomyces pombe*: cell cycle-dependent activation in *Xenopus* cell-free extracts. *J. Cell Biol.* 117, 1055–1066.
- McDonald, H.B., Stewart, R.J., and Goldstein, L.S.B. (1990). The kinesin-like *ncd* protein of *Drosophila* is a minus-end-directed microtubule motor. *Cell* 63, 1159–1166.
- McIntosh, J.R., and Pfaff, C.M. (1991). Mitotic motors. *J. Cell Biol.* 115, 577–585.
- Meluh, P.B., and Rose, M.D. (1990). *KAR3*, a kinesin-related gene required for yeast nuclear fusion. *Cell* 60, 1029–1042.
- Mitchison, T.J. (1988). Microtubule dynamics and kinetochore function in mitosis. *Annu. Rev. Cell Biol.* 4, 527–550.
- Middleton, K., and Carbon, J. (1994). *KAR3*-encoded kinesin is a minus-end-directed motor that functions with centromere binding proteins (CBF3) on an in vitro yeast kinetochore. *Proc. Natl. Acad. Sci. USA* 91, 7212–7216.
- Moreno, S., Klar, A., and Nurse, P. (1991). Molecular genetic analysis of fission yeast *Schizosaccharomyces pombe*. *Methods Enzymol.* 194, 795–824.

- Navone, F., Niclas, J., Hom-Booher, N., Sparks, L., Bernstein, H.D., McCaffrey, G., and Vale, R.D. (1992). Cloning and expression of a human kinesin heavy chain gene: interactions of the COOH-terminal domain with cytoplasmic microtubules in transfected CV-1 cells. *J. Cell Biol.* 117, 1263–1275.
- Nislow, C., Lombillo, V.A., Kuriyama, R., and McIntosh, J.R. (1992). A plus-end-directed motor enzyme that moves antiparallel microtubules in vitro localizes to the interzone of mitotic spindles. *Nature* 359, 543–547.
- Noda, Y., Sato-Yoshitake, R., Kondo, S., Nangaku, M., and Hirokawa, N. (1995). KIF2 is a new microtubule-based anterograde motor that transports membranous organelles distinct from those carried by kinesin heavy chain or KIF3A-B. *J. Cell Biol.* 129, 157–167.
- O'Connell, M., Meluh, P.B., Rose, M.D., and Morris, N.R. (1993). Suppression of the *bimC4* mitotic spindle defect by deletion of *kfpA*, a gene encoding a KAR3-related kinesin-like protein in *A. nidulans*. *J. Cell Biol.* 120, 153–162.
- Ohi, R., Feoktistova, A., and Gould, K.L. (1996). Construction of vectors and a genomic library for use with his3-deficient strains of *Schizosaccharomyces pombe*. *Gene* (in press).
- Page, B.D., Satterwhite, L.L., Rose, M.D., and Snyder, M. (1994). Localization of the Kar3 kinesin heavy chain-related protein requires the Cik1 interacting protein. *J. Cell Biol.* 124, 507–519.
- Robbins, J., Dilworth, S.M., Laskey, R.A., and Dingwall, C. (1991). Two interdependent basic domains in nucleoplasmin nuclear targeting sequence: identification of a class of bipartite nuclear targeting sequence. *Cell* 64, 615–624.
- Roof, D.M., Meluh, P.B., and Rose, M.D. (1992). Kinesin-related proteins required for assembly of the mitotic spindle. *J. Cell Biol.* 118, 95–108.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Saunders, W.S., and Hoyt, M.A. (1992). Kinesin-related proteins required for structural integrity of the mitotic spindle. *Cell* 70, 451–458.
- Saunders, W.S., Koshland, D., Eshel, D., Gibbons, I.R., and Hoyt, M.A. (1995). *Saccharomyces cerevisiae* kinesin- and dynein-related proteins required for anaphase chromosome segregation. *J. Cell Biol.* 128, 617–624.
- Sawin, K.E., Leguellec, K., Phillippe, M., and Mitchison, T.J. (1992a). Mitotic spindle organization by a plus-end-directed microtubule motor. *Nature* 359, 540–543.
- Sawin, K.E., Mitchison, T.J., and Wordeman, L.G. (1992b). Evidence for kinesin-related proteins in the mitotic apparatus using peptide antibodies. *J. Cell Sci.* 101, 303–313.
- Sawin, K.E., and Scholey, J.M. (1991). Motor proteins in cell division. *Trends Cell Biol.* 1, 122–129.
- Smith, D.B., and Johnson, K.S. (1988). Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* 67, 31–40.
- Tommasino, M., and Maundrell, K. (1991). Uptake of thiamine by *Schizosaccharomyces pombe* and its effect as a transcriptional regulator of thiamine-sensitive genes. *Curr. Genet.* 20, 63–66.
- Vale, R.D., Reese, T.S., and Sheetz, M.P. (1985). Identification of a novel force-generating protein, kinesin, involved in microtubule-based motility. *Cell* 42, 39–50.
- Walczak, C.E., Mitchison, T.J., and Desai, A. (1996). XKCM1: a *Xenopus* kinesin-related protein that regulates microtubule dynamics during mitotic spindle assembly. *Cell* 84, 37–47.
- Walker, J., Crowley, P., Moreman, A.D., and Barrett, J. (1993). Biochemical properties of cloned glutathione S-transferases from *Schistosoma mansoni* and *Schistosoma japonicum*. *Mol. Biochem. Parasitol.* 61, 255–264.
- Walker, R.A., Salmon, E.D., and Endow, S.A. (1990). The *Drosophila claret* segregation protein is a minus-end-directed motor molecule. *Nature* 347, 780–782.
- Wordeman, L., and Mitchison, T.J. (1995). Identification and partial characterization of mitotic centromere-associated kinesin, a kinesin-related protein that associates with centromeres during mitosis. *J. Cell Biol.* 128, 95–105.
- Wright, B.D., Henson, J.H., Wedaman, K.P., Willy, P.J., Morand, J.N., and Scholey, J.M. (1991). Subcellular localization and sequence of sea urchin kinesin heavy chain: evidence for its association with membranes in the mitotic apparatus and the interphase cytoplasm. *J. Cell Biol.* 113, 817–834.
- Yanagida, M. (1989). Gene products required for chromosome separation. *J. Cell Sci. Suppl.* 12, 213–230.
- Yang, J.T., Laymon, R.A., and Goldstein, L.S.B. (1989). A three-domain structure of kinesin heavy chain revealed by DNA sequence and microtubule binding analyses. *Cell* 56, 879–890.
- Yanisch-Perron, C., Vieira, J., and Messing, J. (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* 33, 103–119.