Fast structural dynamics in reduced and oxidized cytochrome c

Weixia Liu, Jon N. Rumbley, S. Walter Englander, and A. Joshua Wand*

Johnson Research Foundation and Department of Biochemistry and Biophysics, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6059

Received 8 November 2008; Revised 31 December 2008; Accepted 7 January 2009

DOI: 10.1002/pro.72

Published online 22 January 2009 proteinscience.org

Abstract: The sub-nanosecond structural dynamics of reduced and oxidized cytochrome c were characterized. Dynamic properties of the protein backbone measured by amide ^{15}N relaxation and side chains measured by the deuterium relaxation of methyl groups change little upon change in the redox state. These results imply that the solvent reorganization energy associated with electron transfer is small, consistent with previous theoretical analyses. The relative rigidity of both redox states also implies that dynamic relief of destructive electron transfer pathway interference is not operational in free cytochrome c.

Keywords: protein dynamics; NMR relaxation; redox state; electron transfer

Introduction

The influence of atomic scale structure on molecular recognition and catalysis continues to be intensively studied, but the role of structural dynamics is most often largely unknown.¹ Protein-mediated electron transfer, integral to many critical biological processes, can be influenced by fast internal protein motions.

Abbreviations: FC, Franck-Condon density of states; O_{NH}^2 and O_{axis}^2 , Lipari-Szabo generalized order parameter for the amide N—H bond vector and the methyl symmetry axis, respectively; T_{DA} , electronic matrix element between the donor (D) and acceptor (A).

Jon N. Rumbley current address is Department of Pharmacy Practice and Pharmaceutical Science, University of Minnesota, Duluth, Minnesota 55812.

Grant sponsor: NIH; Grant numbers: GM35940, DK39806, GM31847; Grant sponsor: Mathers Foundation (SWE).

*Correspondence to: A. Joshua Wand, Department of Biochemistry and Biophysics, 905 Stellar Chance Laboratories, 422 Curie Blvd. Philadelphia, PA 19104-6059. E-mail: wand@mail.med.upenn.edu

Considerable effort has been directed at understanding the principles that govern through-protein electron transport,^{2,3} but the exploration of how the nuclear and electronic structure of protein molecules influence electron tunneling remains a challenging and controversial problem.

Electron transfer between weakly coupled donor and acceptors is usually treated using the non-adiabatic description captured by Fermi's Golden Rule derived from quantum mechanical perturbation theory.⁴

$$k_{\text{ET}} = \frac{2\pi}{\hbar} |T_{\text{DA}}|^2 (\text{FC})$$

where T_{DA} is the electronic matrix element between the donor (D) and acceptor (A) and FC is the Franck-Condon density of states that describes the nuclear control of the energy gap. Studies of the effects of internal protein motions on electron transfer have often focused on the FC factor, which is usually

estimated using static or dynamical views of Marcus theory.^{5,6} In contrast, estimation of the T_{DA} term(s) has largely been based on a static view of protein structure. For example, the Pathways interpretation of electron transfer is a popular approach to the analysis of electron transfer in heterogeneous protein media.^{7,8} In its original form, this treatment assumes that tunneling occurs via a dominant pathway or tube and thereby has two major limitations. It does not accommodate interference among multiple paths and the paths that are revealed arise from a static view of the molecular structure.^{9–11}

The present study was motivated by work that strongly suggests that dynamical effects on the TDA terms can be large.^{10,11} Thus protein dynamics can modulate electron transfer by time-dependent variance of the energy gap or of the TDA terms. It is emphasized that the time scale of the fluctuations of the TDA terms most important for electron transfer is short, nanoseconds. This phenomenon is distinct from so-called gated electron transfer where a relatively slow larger scale conformational or orientational adjustment is required for electron transfer and is rate-limiting.¹²

Most investigations of the influence of sub-ns conformational fluctuations on electron transfer through modulation of the energy gap or TDA terms are based on molecular dynamics simulations. There has been little direct experimental investigation of protein dynamics in electron transfer proteins in this time regime, particularly in multiple redox states. Powerful nuclear magnetic resonance (NMR) relaxation methods now provide experimental access to protein dynamics over a wide range of time scales. Here we use NMR relaxation techniques¹³ to provide a site-resolved view of fast internal motions that are pertinent to the redox kinetics of cytochrome *c*.

Results and Discussion

Cytochrome *c*, the prototypical c-type cytochrome, is a component of the electron transport chain in mitochondria. We employed a H26N,H33N mutant which has superior expression properties but is otherwise largely indistinguishable from the wild-type horse protein.¹⁴ Resonance assignments of the diamagnetic reduced (spin zero) and paramagnetic oxidized (spin [1/2]) states were previously reported.¹⁵ Classical NMR relaxation provides access to the amplitude and time scale of motion through its influence on the spectral density defining observable relaxation parameters such as the spin-lattice relaxation time (T_1).¹³ Here, nitrogen-15 relaxation was used to probe fast motion on the backbone and to characterize molecular tumbling. Deuterium relaxation was used to characterize the motion of methyl-bearing amino acid side chains. Relaxation data were interpreted using the Lipari-Szabo model free formalism.¹⁶ A recent re-evaluation of the model-free treatment reinforces confidence in

its robustness with respect to highly asymmetric side chain motion.¹⁷

The squared generalized order parameters for amide N—H bond vectors (O_{NH}^2) and for methyl group symmetry axes (O_{axis}^2) in reduced and oxidized cytochrome *c* are shown in Figure 1. As for many proteins,²³ the reduced and oxidized cytochrome *c* O_{NH}^2 parameters for residues located in regular secondary structures are generally high. When sites having significant chemical exchange or paramagnetic relaxation contributions to the measured relaxation parameters are excluded, the average O_{NH}^2 parameters for both the reduced ($\langle O_{NH}^2 \rangle \sim 0.927$) and oxidized ($\langle O_{NH}^2 \rangle \sim 0.923$) states point to an unusual rigidity for both forms of the protein. Also the average O_{axis}^2 parameters for the reduced ($\langle O_{axis}^2 \rangle \sim 0.718$) and the oxidized ($\langle O_{axis}^2 \rangle \sim 0.723$) protein are also relatively high compared with most proteins.¹³ Both redox states of the protein are largely devoid of the so-called J-class of methyl sites undergoing sufficiently large excursions to result in rotamer interconversion.¹³ In keeping with previous results,^{24,25} the present results suggest that the presence of large inflexible prosthetic groups suppresses the fast internal motion of proteins.

The O_{NH}^2 parameters of the reduced state are on average slightly higher than their counterparts in the oxidized state ($\langle \Delta O_{NH}^2 \rangle \sim 0.04$; $\langle |\Delta O_{NH}^2| \rangle \sim 0.05$), as they are in yeast iso-1-cytochrome *c*.²⁶ In slight contrast, the O_{axis}^2 parameters of the oxidized state are on average higher than their counterparts in the reduced state ($\langle \Delta O_{axis}^2 \rangle \sim 0.016$; $\langle |\Delta O_{axis}^2| \rangle \sim 0.046$). On average, the oxidized state has slightly smaller amplitude of sub-ns motion than the reduced state. These results are generally in keeping with a vibrational analysis of the cytochrome *c*.^{27,28} They may seem to contrast with the view provided by hydrogen exchange²⁹ and NMR exchange broadening,³⁰ but those probes sample much slower (infrequent) motions. Thus, it seems that the dynamical differences between the two redox states are largely restricted to slower time scales. A similar result has been observed using ¹⁵N-relaxation studies of backbone motion in *Bacillus pasteurii* cytochrome *c*, where the average O_{NH}^2 of the oxidized and reduced states were found to be essentially the same within experimental error. The results presented here also suggest that redox-dependent changes in residual protein entropy, expressed in the motions on the sub-ns time scale,^{1,13,17,36} are small in cytochrome *c*. This contrasts with cyt *b₅* where significant changes in protein dynamics are apparent.³¹

The fact that both redox states of cytochrome *c* are relatively rigid implies that a Pathways-type analysis based on a rigid structural framework is valid. Furthermore, the distributions of dynamics within the two states are closely similar. The near identity of the equilibrium structures of the two redox states^{32–34} suggests that the landscapes near the lowest energy structure of each redox state are also quite similar. These results

Figure 1. The amplitudes of ps-ns dynamics of the main chain N-H (left panels) and of the methyl-bearing side chains (right panels) in reduced and oxidized [H26N,H33N]-horse cytochrome *c* at pH 5.8 and 20°C. Nitrogen-15 T_1 , T_2 and NOE relaxation data were collected magnetic field strengths of 11.7 and 14.1 T essentially as described elsewhere^{18,19}. Deuterium T_1 and $T_{1\rho}$ relaxation in methyl CH_2D isotopomers was measured^{18–20} at magnetic fields of 11.7 and 14.1 Tesla using protein expressed during growth on 50% D_2O . An effective ^{15}N – ^1H bond length of 1.04 Å, a ^{15}N CSA tensor breadth of –170 ppm, and a deuterium quadrupolar coupling constant of 165 kHz were used in the calculations. Rotational anisotropy was determined to be negligible for both redox states. The isotropic form of the spectral density was therefore employed using the determined value for τ_m of 7.2 ns. Squared generalized order parameters and effective correlation times were obtained for individual sites using a grid-search algorithm.²¹ Precision was estimated using Monte Carlo sampling. O^2_{axis} were calculated by dividing the obtained O^2 by 0.111. The g-tensor of oxidized cytochrome *c*²² indicates a negligible paramagnetic contribution to deuterium relaxation and a significant (> 5%) contribution to amide ^{15}N relaxation at only a few sites. Open circles correspond to sites with significant motion on the microsecond-millisecond timescale as detected by a R_1R_2 analysis. Solid triangles correspond to sites having a significant paramagnetic contribution (> 5%) to observed relaxation parameters. These sites are excluded from the redox state comparison (bottom panels). Lines of unit slope are drawn in the bottom panels.

indicate that the reorganization energy associated with electron self-exchange is small, consistent with a previous theoretical analysis.³⁵ Finally, Balabin and Onuchic point out that electron transfer pathways need

not be dynamically activated.¹¹ Insofar as the relevant dynamics occur on the sub-ns time scale, the unusual rigidity of both redox states of cytochrome *c* implies that general dynamic relief of destructive interference

of multiple pathways for electron transfer is not operational in free cytochrome *c*.

Methods

Nitrogen-15 T_1 , T_2 and NOE relaxation data were collected magnetic field strengths of 11.7 and 14.1 T essentially as described elsewhere.^{18,19} Deuterium T_1 and $T_{1\rho}$ relaxation in methyl CH_2D isotopomers was measured^{18–20} at magnetic fields of 11.7 and 14.1 Tesla using protein expressed during growth on 50% D_2O . An effective ^{15}N – ^1H bond length of 1.04 Å, a ^{15}N CSA tensor breadth of -170 ppm, and a deuterium quadrupolar coupling constant of 165 kHz were used in the calculations. Rotational anisotropy was determined to be negligible for both redox states. The isotropic form of the spectral density was therefore employed using the determined value for τ_m of 7.2 ns. Squared generalized order parameters and effective correlation times were obtained for individual sites using a grid-search algorithm.²¹ Precision was estimated using Monte Carlo sampling. O^2_{axis} parameters were calculated by dividing the obtained O^2 by 0.111. The g-tensor of oxidized cytochrome c^{22} indicates a negligible paramagnetic contribution to deuterium relaxation and a significant ($>5\%$) contribution to amide ^{15}N relaxation at only a few sites.

References

- Wand AJ (2001) Dynamic activation of protein function: a view emerging from NMR spectroscopy. *Nat Struct Biol* 8:926–931.
- Gray HB, Winkler JR (2003) Electron tunneling through proteins. *Q Rev Biophys* 36:341–372.
- Page CC, Moser CC, Chen X, Dutton PL (1999) Natural engineering principles of electron tunneling in biological oxidation-reduction. *Nature* 402:47–52.
- Devault D (1980) Quantum-mechanical tunnelling in biological-systems. *Q Rev Biophys* 13:387–564.
- Marcus RA (1956) Theory of oxidation-reduction reactions involving electron transfer. 1. *J Chem Phys* 24:966–978.
- Warshel A, Parson WW (2001) Dynamics of biochemical and biophysical reactions: insight from computer simulations. *Q Rev Biophys* 34:563–679.
- Beratan DN, Onuchic JN, Hopfield JJ (1987) Electron-tunneling through covalent and noncovalent pathways in proteins. *J Chem Phys* 86:4488–4498.
- Beratan DN, Betts JN, Onuchic JN (1991) Protein electron-transfer rates set by the bridging secondary and tertiary structure. *Science* 252:1285–1288.
- Evenson JW, Karplus M (1993) Effective coupling in biological electron-transfer—exponential or complex distance dependence. *Science* 262:1247–1249.
- Balabin IA, Onuchic JN (1998) A new framework for electron-transfer calculations—beyond the pathways-like models. *J Phys Chem B* 102:7497–7505.
- Balabin IA, Onuchic JN (2000) Dynamically controlled protein tunneling paths in photosynthetic reaction centers. *Science* 290:114–117.
- Jeuken LJC (2003) Conformational reorganisation in interfacial protein electron transfer. *Biochem Biophys Acta Bioenerg* 1604:67–76.
- Igumenova TI, Frederick KK, Wand AJ (2006) Characterization of the fast dynamics of protein amino acid side chains using NMR relaxation in solution. *Chem Rev* 106:1672–1699.
- Rumbley JN, Hoang L, Englander SW (2002) Recombinant equine cytochrome *c* in *Escherichia coli*: High-level expression, characterization, and folding and assembly mutants. *Biochemistry* 41:13894–13901.
- Liu W, Rumbley J, Englander SW, Wand AJ (2003) Backbone and side-chain heteronuclear resonance assignments and hyperfine NMR shifts in horse cytochrome *c*. *Prot Sci* 12:2104–2108.
- Lipari G, Szabo A (1982) Model-free approach to the interpretation of nuclear magnetic resonance relaxation in macromolecules. 1. Theory and range of validity. *J Am Chem Soc* 104:4546–4559.
- Frederick KK, Sharp KA, Warischalk N, Wand AJ (2008) Re-evaluation of the model-free analysis of fast internal motion in proteins using NMR relaxation. *J Phys Chem B* 112:12095–12103.
- Farrow NA, Muhandiram R, Singer AU, Pascal SM, Kay CM, Gish G, Shoelson SE, Pawson T, Forman-Kay JD, Kay LE (1994) Backbone dynamics of a free and phosphopeptide-complexed Src homology 2 domain studied by ^{15}N NMR relaxation. *Biochemistry* 33:5984–6003.
- Marlow MS, Wand AJ (2006) Conformational dynamics of calmodulin in complex with the calmodulin-dependent kinase kinase alpha calmodulin-binding domain. *Biochemistry* 45:8732–8741.
- Muhandiram DR, Yamazaki T, Sykes BD, Kay LE (1995) Measurement of H-2 T_1 and $T_1\rho$ relaxation-times in uniformly C-13-labeled and fractionally H-2-labeled proteins in solution. *J Am Chem Soc* 117:11536–11544.
- Dellwo MJ, Wand AJ (1989) Model-independent and model-dependent analysis of the global and internal dynamics of cyclosporine A. *J Am Chem Soc* 111:4571–4578.
- Feng YQ, Roder H, Englander SW (1990) Redox-dependent structure change and hyperfine nuclear magnetic resonance shifts in cytochrome-*c*. *Biochemistry* 29:3494–3504.
- Jarymowycz VA, Stone MJ (2006) Fast time scale dynamics of protein backbones: NMR relaxation methods, applications, and functional consequences. *Chem Rev* 106:1624–1671.
- Flynn PF, Urbauer RJB, Zhang H, Lee AL, Wand AJ (2001) Main chain and side chain dynamics of a heme protein: N-15 and H-2 NMR relaxation studies of R. capsulatus ferrocyclochrome *c* (2). *Biochemistry* 40:6559–6569.
- Liu WX, Flynn PF, Fuentes EJ, Kranz JK, McCormick M, Wand AJ (2001) Main chain and side chain dynamics of oxidized flavodoxin from *Cyanobacterium anabaena*. *Biochemistry* 40:14744–14753.
- Fetrow JS, Baxter SM (1999) Assignment of N-15 chemical shifts and N-15 relaxation measurements for oxidized and reduced iso-1-cytochrome *c*. *Biochemistry* 38:4480–4492.
- Chin JK, Jimenez R, Romesberg FE (2002) Protein dynamics and cytochrome *c*: correlations between ligand vibrations and redox activity. *J Am Chem Soc* 124:1846–1847.
- Sagle LB, Zimmermann J, Matsuda S, Dawson PE, Romesberg FE (2006) Redox-coupled dynamics and folding in cytochrome *c*. *J Am Chem Soc* 128:7909–7915.
- Milne JS, Mayne L, Roder H, Wand AJ, Englander SW (1998) Determinants of protein hydrogen exchange studied in equine cytochrome *c*. *Prot Sci* 7:739–745.

30. Barker PD, Bertini I, Del Conte R, Ferguson SJ, Hajieva P, Tomlinson E, Turano P, Viezzoli MS (2001) A further clue to understanding the mobility of mitochondrial yeast cytochrome *c*—a N-15 T-1 rho investigation of the oxidized and reduced species. *Eur J Biochem* 268: 4468–4476.
31. Dangi B, Blankman JI, Miller CJ, Volkman BF, Guiles RD (1998) Contribution of backbone dynamics to entropy changes occurring on oxidation of cytochrome *b* (5). Can redox linked changes in hydrogen bond networks modulate reduction potentials? *J Phys Chem B* 102: 8201–8208.
32. Takano T, Dickerson RE (1981) Conformation change of cytochrome *c*. 2. Ferricytochrome *c* refinement at 1.8 Å and comparison with the ferrocycytochrome structure. *J Mol Biol* 153:95–115.
33. Berghuis AM, Brayer GD (1992) Oxidation state-dependent conformational changes in cytochrome *c*. *J Mol Biol* 223:959–976.
34. Banci L, Bertini I, Huber JG, Spyroulias GA, Turano P (1999) Solution structure of reduced horse heart cytochrome *c*. *J Biol Inorg Chem* 4:21–31.
35. Muegge I, Qi PX, Wand AJ, Chu ZT, Warshel A (1997) The reorganization energy of cytochrome *c* revisited *J Phys Chem B* 101:825–836.
36. Frederick KK, Marlow MS, Valentine KG, Wand AJ (2007) Conformational entropy in molecular recognition by proteins. *Nature* 448:325–329.