Published in final edited form as:

Physiology (Bethesda). 2009 February ; 24: 58–71. doi:10.1152/physiol.00033.2008.

Maternal Uterine Vascular Remodeling During Pregnancy

George Osol¹ and Maurizio Mandala²

George Osol: George.Osol@uvm.edu; Maurizio Mandala:

¹Department of Obstetrics and Gynecology, University of Vermont College of Medicine, Burlington, Vermont

²University of Calabria, Cosenza, Italy

Abstract

Sufficient uteroplacental blood flow is essential for normal pregnancy outcome and is accomplished by the coordinated growth and remodeling of the entire uterine circulation, as well as the creation of a new fetal vascular organ: the placenta. The process of remodeling involves a number of cellular processes, including hyperplasia and hypertrophy, rearrangement of existing elements, and changes in extracellular matrix. In this review, we provide information on uterine blood flow increases during pregnancy, the influence of placentation type on the distribution of uterine vascular resistance, consideration of the patterns, nature, and extent of maternal uterine vascular remodeling during pregnancy, and what is known about the underlying cellular mechanisms.

The clinical relevance of maternal uterine vascular adaptation during pregnancy is underscored by the fact that its aberrance is associated with several common gestational pathologies, including intrauterine growth restriction, gestational diabetes, and preeclampsia.

In addition to the changes in vessel structure, uterine vascular reactivity is also altered during pregnancy, with the general pattern being one of reduced tone and enhanced vasodilation/blunted vasoconstriction (52,72,111,112,118). Space limitations preclude in-depth consideration of the ionic and enzymatic mechanisms that underlie reactivity, but it is worth noting that, in vivo, uterine vascular resistance (and, therefore, blood flow) is ultimately determined by the combination of vessel size and reactivity.

We also avoid examining the complex angiogenic mechanisms associated with implantation and placentation, other than to consider how hemochorial vs. epitheliochorial placentation influences uterine hemodynamics and vascular remodeling. Although the remodeling of spiral arteries by fetal trophoblast is considered, the main focus of this review is on upstream maternal uterine arteries and veins, since the processes involved in their remodeling have not been reviewed to date. Readers interested in endovascular trophoblast invasion and mechanisms underlying spiral artery remodeling are referred to several reviews on this subject (9,31,71, 101).

Nomenclature Used to Describe Vascular Remodeling

Circumferential remodeling is normally termed inward or outward to denote narrowing vs. widening of the vessel lumen. The term expansive remodeling has also been used and can be substituted for “outward” in describing an increase in circumference. This classification was proposed by Mulvany (86) and is further refined to include consideration of wall mass, which can increase (hypertrophy), decrease (hypotrophy), or remain unchanged (eutrophy). Various patterns of arterial remodeling are shown in FIGURE 1; see legend for additional detail and for consideration of how changes in cross-sectional area, a two-dimensional quantity (e.g., μm^2), relate to changes in wall mass, a three-dimensional quantity (e.g., μm^3).

Uterine Hemodynamics During Pregnancy; Blood Flow Patterns and the Importance of Placentation Type

Uterine vascular anatomy

An overview of comparative uterine vascular anatomy is presented in FIGURE 2, which describes and illustrates the uterine circulation in humans (FIGURE 2A), rodents (FIGURE 2B), and ungulates, such as sheep and pigs. Please see legend for additional detail.

Uterine and placental blood flow during pregnancy

Early (1953–1960) human studies by Assali et al (3,4) and Metcalfe (78) utilizing the diffusion equilibrium principle (most often nitrous oxide, N₂O) or electromagnetic flow probes placed directly on the uterine artery reported that total uteroplacental blood flow (UPBF) increases from a baseline value of 20–50 ml/min to 450–800 ml/min in singleton pregnancies, with values in excess of 1 l/min measured in twin pregnancy. Subsequent measurements of uterine artery blood flow with ¹³³Xe (106), placental metabolic clearance rate techniques (29), and most recently and directly transvaginal Doppler ultrasonography (99) support these early findings. For example, in the latter study, unilateral uterine artery blood flow at *week 36* was calculated to be 353 ml/min, total UPBF would therefore be in excess of 700 ml/min, with some additional increase likely in the remaining few weeks of gestation (the authors' estimate was 921 ml/min).

In experimental animals such as rodents or sheep, relative changes in UPBF are equal to or greater than those of humans, with increases in flow at term ranging from 10- to 100-fold above nonpregnant levels (6,10,15,26,37,38,44,48,56,62,112).

Since blood pressure normally decreases or is unchanged during pregnancy, uterine hemodynamic changes are principally affected by a profound decrease in uterine vascular resistance. This is accomplished by several different but complimentary mechanisms, including circumferential structural enlargement of the entire uterine vascular tree (including the veins), a reduction in vascular tone (vasodilation), and the creation of the placenta. From a systemic standpoint, increases in UPBF are facilitated by the combination of a substantially increased cardiac output and an expanded vascular volume that are characteristic of gestation in every species studied (129).

In humans (4,127) and guinea pigs (6), the increase in UPBF is gradual and fairly linear, whereas in the rat, which has a 22-day gestation, increases in total uterine blood flow are first detectable on or around *day 15*, i.e., the last “trimester” of gestation (26). A similar pattern of augmented flow in the last trimester has been reported in sheep (112). Moreover, as pregnancy advances, there is a progressive increase in the proportion of blood directed to the placenta. This was shown by Dowell and Kauer in the rat (26), where <10% of uterine blood flow was placental on *day 15* of pregnancy; near term (*day 22*), the value increased to 90%. Similar estimates have been made in sheep (112) and primates (65).

Absolute blood flow to the myometrium increases in proportion to uterine mass, whereas relative uterine blood flow (milliliter per minute per 100 g of tissue, excluding the placenta) may fluctuate and decrease somewhat (6,112,137) or remain fairly constant (26,37) during pregnancy.

Influence of placentation on distribution of uterine vascular resistance and blood flow

Mice, rats, guinea pigs, rabbits, and humans share a hemochorial type of placentation, in which intraplacental pressure created by the maternal blood occupying the intervillous space must be kept low enough to avoid compression of the intravillous (human) or intralabyrinthine (rodents) fetal vessels, necessitating the contribution of significant resistance by upstream vessels (83).

To examine this concept directly, Moll and colleagues measured the blood pressure in premyometrial arteries in a number of species (guinea pig, rat, rabbit, sheep) under anesthetized conditions by direct arterial puncture (81,82). A substantial pressure drop occurred in the arcuate and radial arteries of rodents, such that the arterial pressure of maternal blood entering the placenta in rats, rabbits, and guinea pigs was only 8–14 mmHg. Based on this finding, they determined that the majority of uterine vascular resistance was localized in the upstream vessels of the mesometrial arcade. Although comparable measurements have not been carried out in humans, the values are likely similar, since a subsequent study by the same group showed that pressures within the spiral arteries of primates (Rhesus monkeys) were also quite low, ranging from 9 to 15 mmHg (84).

With hemochorial placentation, for a molecule to pass from the maternal to the fetal compartment, only two fetal cell layers (trophoblast and intravillous endothelium) need to be traversed. Conversely, in animals with epitheliochorial placentation such as sheep, both maternal and fetal blood flow through small vessels that are separate from but in close apposition to one another. The site of gaseous and nutrient exchange is therefore between two microvasculatures and, because of the greater resistance inherent in this anatomical arrangement and an increase in the number of cell layers (4 to 6) separating the two circulations, the inflow pressure to an epitheliochorial placenta is significantly higher than that of a hemochorial placenta, e.g., >80 mmHg in the ewe (81). Also, epitheliochorial placentation is less efficient than the hemochorial type (83), which may explain why UPBF in ungulates such as sheep and pigs is relatively higher than that of humans or rodents, with values in excess of 1 l/min reported in studies with sheep (112) and pigs (37). Thus differences in the type of placentation have a major influence on the pattern of afferent (arterial) remodeling and on the pressure head of blood as it enters the placenta.

Pattern, Nature, and Extent of Uterine Vascular Remodeling During Pregnancy

Pattern of circumferential remodeling and changes in vessel cellular properties

During pregnancy, the diameter of the main uterine artery approximately doubles in size in humans (99). This finding is supported by a host of published studies using sheep, pigs, guinea pigs, and rats, where uterine artery diameters also generally increase two- to three-fold (2,45,46,56,75,79,92,99,132,133). This enlargement in arterial caliber occurs most often with little or no thickening of the vascular wall (2,97), with the one apparent exception being the mouse, where media thickness increased significantly during the course of gestation (132,134). With or without wall thickening, the increase in lumen diameter nevertheless results in an increased cross-sectional area; thus the pattern of circumferential remodeling is outward hypertrophic.

Smaller arcuate and radial arteries remodel in a similar pattern, with documented enlargements in luminal caliber ranging from 25 to 220% and with either no change or an increase in wall thickness (20,21,41,74,80,81,97,119,120).

In one study on human myometrial radial arteries from preeclamptic women (94), the pattern of remodeling was reported to be one favoring a smaller lumen and a thicker wall, with no change in cross-sectional area, suggesting rearrangement of existing wall elements around a smaller lumen, i.e., inward eutrophic remodeling.

Since the media occupies the greatest proportion of the wall, luminal enlargement would be accomplished most simply by an increase in vascular smooth muscle cell length (axial hypertrophy), and this does indeed appear to be the case: e.g., unstressed smooth muscle cell length in arcuate vessels from pregnant vs. virgin nonpregnant rats was increased by 20% in rats (20). In guinea pigs (49), smooth muscle cell length increased from 21 to 39 μm (86%),

along with a comparable increase in cellular thickness (from 4.6 to 9.6 μm , or 108%). Morphometric measurements also suggest that elongation of vascular smooth muscle cells occurs in sheep (2). Surprisingly, no human data are available, although hypertrophy of myometrial smooth muscle is well established (e.g., Ref. 11).

In the main uterine artery of the sheep, vascular smooth muscle mass and protein content (including actin and myosin) doubled during pregnancy, with a corresponding (twofold) increase in cell volume (2). These changes in smooth muscle ultrastructure are associated with an increased force-generating ability (stress) per cross-sectional area, suggesting that the concentration of the contractile proteins (myosin and actin) may also increase. As noted by the authors, the magnitude of the increase in stress exceeded the magnitude of the increase in protein content, suggesting the existence of other mechanisms that augment force production, for example, changes in myosin duty cycle or force generation per stroke.

In addition to cellular hypertrophy, there is also strong evidence for hyperplasia within the vascular wall, since increased rates of smooth muscle cell division have been reported in uterine arteries and veins from rats and guinea pigs (20,21,47,56,57,98). Although endothelial hyperplasia has been documented in the rat (20), changes in endothelial hypertrophy (increased cell area or mass) have not yet been detailed and await further study.

Changes in vessel biophysical properties

The biomechanical properties of uterine vessels also change significantly during pregnancy. In an early (1985) study by Griendling et al. (45), the incremental passive elastic modulus, which measures the relationship of stress to strain, increased from 4 to 29 dyn/cm^2 in main uterine arteries from near-term pregnant vs. nonpregnant sheep. Since this is essentially a stress-to-strain ratio, higher values indicate a less compliant material, and the authors interpreted their data as being indicative of increased arterial stiffness (decreased compliance). In the same study, however, arterial distensibility as a function of pressure was clearly increased, and this finding is consistent with other reports of increased distensibility of main uterine arteries from pregnant guinea pigs (56,75). In rats, smaller arcuate and radial arteries are also more distensible than those from their nonpregnant counterparts (97,98), and, in the rabbit, preplacental radial arteries were significantly more distensible than premyometrial vessels (21). In a different study, the distensibility of the main uterine vein was also increased in the rat, and associated with reduced elastin content (98).

Most commonly, changes in compliance are attributed to alterations in extracellular matrix volume, composition, and collagen and elastin fiber orientation. Some species differences may exist in this regard. In main uterine arteries from sheep (45) and pigs (46), collagen content decreased significantly, with no change in elastin; hence, the collagen-to-elastin ratios were decreased. Conversely, Robertson and Manning (109) found a decrease in the elastin content of spiral arteries of pregnant women, although there were no changes in vessels “remote from the placental bed” (presumably upstream uterine arteries, location not denned). In rats, we measured decreased elastin content in the uterine vein (collagen was not measured), and this correlated with significantly increased circumferential distensibility (98).

Spatial and temporal considerations; trophoblast invasion of spiral arteries

The time course and pattern of remodeling may also be specific to vessel size and location and may vary with species. For example, the results of one study in rats suggest that remodeling may begin in the smaller vessels proximal to the sites of placentation or uterine wall and then, as pregnancy progresses, proceeds to the larger, more upstream vessels (20). This observation is based on the mitotic rates of vascular smooth muscle and endothelial cells, which were highest in smaller vessels in mid-pregnancy (*day 16/22*) and in larger upstream vessels later

in gestation (*day 20/22*) and is consistent with the concept of trophoblast invasion of spiral arteries immediately proximal to the placenta being an initiating event in the remodeling process (9,71,101). Corroborative evidence can be found in other papers using guinea pigs as the experimental animal of choice (56,124).

In addition to the influence of gestational age and species on the nature and extent of remodeling, differences in remodeling patterns have been noted in vessels of different size, type, and location. For example, preplacental vessels enlarge more than premyometrial arteries in rabbits (21) and rats (41,47) and may also show significant differences in rates of cell division, distensibility, and reactivity. A photograph of a premyometrial vs. preplacental radial artery in the rat is shown in FIGURE 3 to illustrate the differences in structure.

As can be seen, mesometrial preplacental vessels widen progressively as they approach the placenta and may be several-fold wider at their distal vs. proximal ends. These changes are especially dramatic in the guinea pig, with threefold increases in circumference (or diameter) occurring along the length of a single vessel during pregnancy. Guinea pig preplacental arteries show 50-fold increases in mass and 30-fold increases in nuclear DNA in addition to substantial changes in collagen breakdown (80). Fragmentation of elastic tissues and transformation of cellular wall constituents into a necrotic, homogenous, acellular fibrinoid was also noted (79), as reported in human spiral arteries as well (8,105). Differences in the diameter along the length of the main uterine artery have also been noted, at least in the guinea pig, with a pattern of widening at each end (92). Localized differences in structure may be related to different rates of shear stress, a mechanism likely involved in the expansive remodeling process, as described below in **Shear stress and NO**.

Differences in the structure of spiral arteries immediately proximal to the placenta are normally ascribed to the trophoblast invasion that occurs in association with placentation. In this unique endovascular process, fetal trophoblast cells migrate into the arterial lumen, ablate the endothelium and smooth muscle of the arterial wall, and reorganize the matrix elements (1,7,9,14,22,31,54,71,101). The involvement of other cell types, such as natural killer (NK) cells, a type of lymphocyte, is also likely to play a role in placentation and spiral artery invasion and remodeling (28,115) and has been implicated in the development of preeclampsia (42,116).

This process of endovascular invasion in vessels that completely lose their ability to contract (41) and take on a trumpet-like, splayed shape further decreases resistance and facilitates an increase in flow. Several studies have implicated a reduction in the depth and extent of this process in the genesis of hypertension associated with preeclampsia (9), suggesting that insufficient spiral artery remodeling leads to placental ischemia, triggering the release of placental signals that increase maternal blood pressure to provide more driving force (a higher perfusion pressure) for UPBF.

Axial (longitudinal) remodeling of uterine arteries

In humans, elongation (axial growth) of arcuate arteries must occur as the uterus enlarges circumferentially to accommodate the fetoplacental unit. In women, it is not clear whether this is accomplished by actual longitudinal growth or by progressive straightening of coiled vessels. Tortuosity is characteristic of the human uterine vascular anatomy (32,33), and there are no studies that have attempted to differentiate between vessel straightening due to altered mechanical forces and true growth, which would involve cell division and an increase in wall mass.

In rodents that bear many young (rat and mouse), significant axial growth of the main uterine artery is clearly evident, and the length of the main utero-ovarian artery at term is double or triple that of the nonpregnant state in the rat. The guinea pig, which only gives birth to several

young and has a much longer gestation than the mouse or rat (55 vs. 20 and 22 days, respectively), exhibits comparable elongation of the main uterine artery (92).

Rat and guinea pig mesometrial (arcuate and radial) vessels elongate as well, increasing three- to five-fold in length (80,97). These measurements were made in the unstretched and unstressed state and thus reflect true increases in vessel mass. Taking both circumferential and axial remodeling into account, the actual increase in uterine artery wall mass in the rat is quite substantial, on the order of 300–700%, based on available measurements.

It is interesting to consider the influence of circumferential vs. axial remodeling on uterine hemodynamics, since increases in length would have opposite influences on resistance from those of increased diameter. With an approximate doubling of each parameter, as has been noted in several studies, the effect of widening should predominate, since the relationship between length and resistance is linear (i.e., a doubling of length equals a doubling of resistance), whereas the relationship of diameter (or radius) to resistance is inverse and quadratic (i.e., doubling of diameter theoretically decreases resistance 16-fold), as per Poiseuille's Law. This mathematical formulation relates changes in vessel length, diameter, and blood viscosity to altered flow resistance by: $R = \epsilon L / r^4$, where R is resistance, ϵ is viscosity, L is length, and r is inner radius.

Thus, without considering changes in tone and reactivity, the combined effect of doubling both the length and the diameter of a tube would theoretically reduce resistance by a factor of 8 and, in the absence of changes in viscosity or growth of new vessels, induce a corresponding increase in blood flow. Clearly, expansive circumferential remodeling and decreased distal resistance due to placentation exert a synergistic effect that can result in the many-fold increases in UPBF.

Finally, additional rheological factors, such as a reduction in the viscosity of the blood (95), also occur during pregnancy and contribute to reducing blood flow resistance according to Poiseuille's Law.

Remodeling of uterine veins

The process of expansive remodeling is not limited to arteries, since the veins also enlarge substantially during pregnancy, although the available data are much more limited. The fully distended diameter of the main uterine vein in pregnant vs. nonpregnant rats averaged 1,576 vs. 956 μm , respectively (an increase of 65%), along with a doubling in unstressed length (98). Increased venous diameter was also accompanied by increased distensibility and reduced elastin content, adaptations that would further enhance venous capacitance. Significant uterine venous enlargement during pregnancy was also reported in the mouse (34).

Mechanisms Underlying Gestational Uterine Vascular Remodeling

The role of local vs. systemic influences on uterine vascular remodeling

As pointed out by Poston and colleagues (55), vascular remodeling is an active process that is dependent on at least four cellular processes: growth, death, migration, and production/degradation of extracellular matrix. Each of these processes is subject to multiple regulatory and interactive influences, including physical forces such as pressure/stretch and shear stress, humoral/endocrine influences such as VEGF and the sex steroids, and local factors generated within the vascular wall, e.g., endothelial NO, angiotensin, and endothelin. Multiple linkages between cellular inputs and outputs further complicate understanding the precise mechanisms involved. For example, VEGF, estrogen, and shear stress all stimulate NO production, and both estrogen and NO regulate matrix metalloproteinase activity and, hence, passive mechanical properties and matrix composition (27,43,103).

In addressing the question of how uterine vascular remodeling is accomplished during pregnancy, one might first consider whether this process is due to the influence of local factors (e.g., those associated with implantation, placentation, and fetal growth), to systemic changes in hormonal levels, particularly those of estrogen and progesterone, or a combination thereof.

Three different experimental approaches have been utilized to investigate this concept: induction of pseudopregnancy (in animals such as the rat and rabbit that are induced ovulators), hormone replacement in oophorectomized animals to mimic the steroidal milieu of the pregnant state, and surgical ligation of (and, thereby, restriction of implantation to) one uterine horn in animals that have duplex uteri, such as the rat and rabbit.

Pseudopregnancy—In animals that are induced ovulators, sexual stimulation induces endocrine changes similar to those of pregnancy (pseudopregnancy), at least for the first half of gestation. Circulating estrogen and progesterone levels increase progressively during this time, allowing the evaluation of their influence without the complicating presence of fetoplacental units.

Peeters' group from Maastricht explored this question in mice (134) and reported that pseudopregnancy resulted in a 25% increase in main uterine artery lumen radius and a 71% increase in cross-sectional area on *day 11* of pseudopregnancy. Structural changes were accompanied by reduced smoothelin expression and an increased proliferation of uterine artery medial smooth muscle cells. The effects of pseudopregnancy paralleled changes in control pregnant animals at 5 days and were about half as large as those measured in pregnant animals on *day 11*.

These data support the idea that placental and ovarian steroids may initiate the process of circumferential remodeling, which then becomes more dependent on local (fetoplacental) factors as pregnancy progresses. Interestingly, despite early arterial enlargement, increases in UPBF are quite modest, or absent, until much later in normal pregnancy in rodents (26), perhaps because changes in length (not documented in the Peeters' study) counteract those of increasing diameter. In addition, there may be an increased level of tone that maintains resistance and prevents an increase in blood flow; neither of these possibilities has been investigated to date.

Hormone replacement—Uterine arteries contain receptors for both estrogen and progesterone (66–68,72,73) and respond to these steroids via genomic and nongenomic mechanisms. Most attention has been directed at estrogen, especially in sheep (by Rosenfeld and Magness, e.g., Refs. 72,73,111,112), since its injection results in both acute and chronic increases in uterine blood flow related to vasodilation and increased cardiac output (19), and its *in vitro* application to isolated vessels induces vasorelaxation, albeit in concentrations that are several orders of magnitude higher than those present *in vivo*. Daily injections of estradiol benzoate (7–11 µg) for 3 wk induced a greater than twofold increase in the internal diameter of segmental mesometrial arteries of ovariectomized guinea pigs (85). Exogenous estradiol treatment in guinea pigs also stimulated DNA synthesis within the uterine artery (74) and increased cellular responsiveness to growth factors such as platelet-derived growth factor (PDGF) by a PKC-dependent mechanism (57). This latter study used isolated vascular smooth muscle cells from the guinea pig uterine artery and reported that cells from the uterine artery, but not the aorta of pregnant animals, showed enhanced spontaneous DNA synthesis and growth. Changes in pseudopregnancy (134), already discussed, are presumably due to an altered endocrine profile and, although more modest than those that occur during pregnancy, are nonetheless measurable and significant.

Exactly how estrogen increases arterial size is not known. Its actions may be indirect, via augmentation of endothelial NO synthesis (58,59) since nitric oxide (NO) has been implicated

in outward circumferential remodeling (96,132) and by the induction of uterine artery vasodilation (123). Along with progesterone, it is also an important determinant of growth factor secretion, e.g., VEGF (50), and is known to affect matrix metalloproteinase (MMP) activity and arterial biomechanical properties (138,139).

It thus appears that steroids promote uterine vascular remodeling through their modulation of numerous cellular processes associated with vascular contractility, growth, and matrix deposition. The reader is referred to a recent review by Chang and Zhang, which provides an excellent summary of steroidal influences on uterine vascular adaptation to pregnancy (19).

Unilateral horn ligation—In animals with duplex uteri, especially rodents, the two horns are physically separate so that one horn can be made to remain barren either by ligating the cervical end (which prevents the passage of sperm) or by tying off the ovarian end (which prevents eggs from entering the uterine lumen), and both experimental approaches have been used.

The uterine horn ligation model has been used to explore a variety of pregnancy-related questions in several species, including the cow (35), pig (36,60), rabbit (137), guinea pig (128), rat (136), and mouse (34), and the available evidence points to local rather than systemic factors playing a primary role in gestational uterine vascular remodeling and increases in blood flow.

In a rat model, this can be readily seen in the photographs shown in FIGURE 4, where the increase in arterial length is clearly apparent (also in diameter, although this parameter is more difficult to evaluate visually without higher magnification). Vessel dimensions in the nonpregnant horn were similar to those of a nonpregnant animal in terms of the length and diameter of both large and small arteries. Consistent with this, uterine vein size was found to be increased only in the pregnant horn of single horn ligated mice, with values from the nonpregnant horn being similar to that of nonpregnant controls (34).

Physiological mechanisms of uterine vascular remodeling during pregnancy

Shear stress and NO—Studies by Langille, DeMey, and others (5,13,16,30,63,64,69, 117,131) have firmly established that an increase in shear stress stimulates expansive remodeling of both large and small arteries in a number of vascular beds. Interestingly, despite the number of citations, the species diversity is limited since virtually all of the data in these studies have been obtained using rodent models.

This mechanism is appealing from a physiological standpoint, since an increase in vessel caliber allows the maintenance of increased flow, while at the same time normalizing endothelial shear stress due to a decrease in blood flow velocity. A good discussion of the biophysical and theoretical aspects of shear stress in vascular tissues can be found in several recent papers (102,107), and the inquisitive reader is also referred to a classic 1975 paper by Rodbard (110) for a more historic perspective of our understanding of the relationship between vascular caliber and flow.

Elevation in shear stress is a plausible mechanism for uterine arterial enlargement during gestation in view of its well established nature as a physiological mechanism and the fact that the reduction in downstream resistance that occurs secondary to placentation would be an effective stimulus for increasing the velocity of flow (and, therefore, shear stress) in upstream arteries. Vasodilation and/or vessel growth would allow the augmented flow to continue, but with a slower velocity, thereby normalizing shear stress in the process.

Yet, the velocity of blood in the uterine artery of women in *week 36* of pregnancy was nearly eight times faster than in the nonpregnant state (averaging 61.4 vs. 8.4 cm/s), whereas arterial diameter increased approximately twofold (99). These data do not support normalization of shear stress, although, unfortunately, it is not possible to calculate vascular wall shear without knowing the actual velocity and viscosity of blood at the endothelial surface—a difficult proposition for a non-Newtonian fluid delivered in a pulsatile and sometimes non-laminar fashion.

The molecular mechanism(s) by which shear stress leads to circumferential vessel growth is not known, although several recent studies have indicated that endothelial NO may be an important mediator of this process (114,130–132). This observation is pertinent to uterine gestational remodeling in view of the well established upregulation of eNOS and NO signaling in the pregnant state and is supported by several lines of evidence. First, both message and protein levels of NOS-3 (or eNOS), the enzyme responsible for NO production by the endothelium, are increased by shear stress, as is the release of NO (131). Accordingly, NO synthase activity and expression (91) and relaxation to acetylcholine (90) are augmented in uterine arteries from pregnant women. Second, mice lacking the gene for NOS-3 show a reduction in uterine vascular remodeling and decreased uterine blood flow (132). And third, chemical inhibition of NO production by NO inhibitors such as L-NAME virtually abolished the expansive remodeling of the main uterine artery and of smaller radial arteries in the rat, although, notably, axial growth was completely unaffected (96).

There is also some evidence against a role for NO in flow-induced remodeling (16), although the focus of this paper was not on gestation, and observations were limited to the mesenteric circulation, which does not undergo expansive remodeling during pregnancy and may develop different, and sometimes even opposite, patterns of reactivity (23,24). Differences in regional vascular responses have also been documented in other studies (23,100). In a study that utilized NOS-3 knockout mice (132), the growth of the main uterine artery was less than that of wild-type animals but still significantly greater than that of nonpregnant controls, suggesting that mechanisms other than those linked to shear stress-induced endothelial NO release play a role. NO production by guinea pig extravillous trophoblast has also been described (88), raising the possibility that other cell types such as macrophages or trophoblast may migrate into the periarterial space and stimulate vasodilation and remodeling of both the arterial and venous wall through mechanisms linked to NO.

A number of other signaling pathways have been implicated in flow-induced arterial remodeling (although not in the setting of gestational remodeling) including MMP activation (130), adrenergic influences (30), toll-like receptors (51), cytoskeleton (69), including vimentin (117), and membrane-associated tyrosine kinases such as PYK2 (125).

Matrix metalloproteinases in gestational vascular remodeling—Matrix metalloproteinases (MMPs) play a vital role in matrix turnover and are, in turn, subject to regulation by both sex steroids and NO. For example, both estrogen and progesterone, alone and in combination, have been shown to modulate smooth muscle cell MMP gene and protein expression, matrix turnover, and deposition (43,89). The activity of several MMPs is elevated in vascular tissues during pregnancy (55) and MMP levels, particularly those of MMP-2 and MMP-9, are altered in preeclamptic women (87,104). This combination of regulation by gestational signals and a broad spectrum of actions on the structure of the vascular wall (e.g., matrix reorganization and composition, and smooth muscle cell migration) has established a central role for MMPs in vascular remodeling (77). The linkage between estrogen and MMP activity was already discussed above, and several reviews highlight our understanding of these processes in greater depth and consider the role they may play in pregnancy under physiological and pathological conditions (55,104).

Other mechanisms (e.g., the renin-angiotensin system and altered VEGF/PIGF signaling) in uterine vascular remodeling—If elevated shear stress secondary to decreased downstream vascular resistance (due to placentation) were the only stimulus for remodeling, one might not expect to see any remodeling in vessels that deliver blood to the myometrium. Yet, although the extent of remodeling of premyometrial radial arteries is less than that of preplacental radial vessels, at least in rats (41,47), these arteries are still clearly larger and longer than those of age-matched nonpregnant animals. Thus it is likely that other mechanisms are involved in the remodeling process.

For example, the renin-angiotensin system may play a role in view of the association between angiotensin II and smooth muscle cellular hypertrophy (39), and one current theory of preeclampsia posits that, at least in some women, this disease may have an automimmune component related to the production of agonistic antibodies directed at the AT-1 receptor (25,108). Exposure to auto-antibodies extracted from the serum of preeclamptic women resulted in the development of a preeclampsia-like syndrome in pregnant mice, and this syndrome could be prevented by co-administration of losartan, an AT-1 receptor blocker, or by an antibody-neutralizing seven-amino acid epitope peptide (140). Although uterine vascular remodeling was not measured in this study, placental abnormalities and small fetus size were reported, and uterine blood flow is a well established determinant of fetal growth (62). Presumably, this is a humoral mechanism that would therefore have a similar effect on both preplacental and premyometrial vessels.

A related observation was made in a study by St-Louis et al. (119), who showed that administering a low-sodium diet, which is associated with over-activation of the renin-angiotensin system (113), to pregnant rats during the last third of gestation attenuated the normal increase in arcuate artery diameter by ~50%.

In contrast, although a recent study (135) describing a transgenic rat model with an overactivated reninangiotensin system reported fetal growth restriction, impaired uterine artery endothelial relaxation, and hypersensitivity to the constrictor effects of phenylephrine, the Doppler resistance index decreased during pregnancy compared with control pregnant animals, suggesting unimpaired uteroplacental flow in the uterine artery. Unfortunately, arterial remodeling was not evaluated directly, and the differentiation of large vs. small, or premyometrial vs. preplacental artery adaptations as contributing factors was not evaluated.

Changes in vasodilator influences may also occur. For example, a number of studies (12,53, 61,70,121,126) have implicated abnormal VEGF/PIGF (placental growth factor) signaling in preeclamptic women due to overexpression of a soluble receptor for VEGF/PIGF (called sFlt-1 or sVEGFR-1). Rats injected with an adenovirus that overexpresses sFlt-1 develop several features associated with preeclampsia (76), although direct evaluation of uterine blood flow or vascular remodeling, either in women or animals, has not been carried out to date. Growth factors of the VEGF family are appealing candidates for mediating this process since they induce vasodilation, stimulate endothelial mitosis, and are associated with hypervascularization and enlargement of existing vessels (at least in the mouse ear; Ref. 93), but their role in pregnancy-induced uterine vascular remodeling and of effects on different segments of the vasculature (e.g., mesometrial vs. intraplacental, large vs. small, and preplacental vs. premyometrial arteries, or in arteries vs. veins) has not been defined.

Venoarterial signaling as a pathway for arterial remodeling—That the molecular signal or signals that induce remodeling originate from the uterus or placenta is an intriguing idea, especially since the latter is a fetal organ known to be a rich source of vasoactive and growth-promoting molecular signals. Involvement of periplacental maternal tissues, e.g., the decidua, is also possible and has yet to be investigated. The principal question, however, is:

How can molecules secreted into the uterine or placental venous effluent affect the structure (and/or tone) of the afferent (arterial) circulation?

One mechanism that deserves mention is venoarterial transfer. Illustrated diagrammatically in FIGURE 5, it posits that secreted placental and/or myometrial signals (such as growth factors) pass across the uterine venous wall and thereby influence the structure of adjacent uterine arteries. Since growth factors such as VEGF and PlGF are also potent vasodilators, this mechanism could also provide a short loop pathway for regulating placental perfusion by inducing changes in arterial tone.

An anatomical basis for venoarterial transfer does exist in the uterine circulation, since arteries and veins are arranged in close apposition in a number of species (17,18,40) and in humans (122). Moreover, physiological plausibility derives from studies that have established this as a mechanism of luteolysis (40). Prostaglandins secreted by the uterus and carried in its veins are transferred to the ovarian artery, where they travel to the corpus luteum and induce a number of processes leading to luteal demise, with ischemic vasoconstriction among them.

Thus venoarterial exchange is plausible from both an anatomical and a physiological standpoint. The venous permeability of isolated rat uterine veins is considerable, even to large molecular weight signals (70 kDa), significantly enhanced during pregnancy, and regulated by physical forces such as wall tension and molecular signals such as VEGF (17,18).

At the same time, evidence that conclusively demonstrates (or refutes) the importance of venoarterial transfer as an *in vivo* mechanism for uterine vascular gestational remodeling is currently lacking, and the determination of whether this mechanism is physiologically important awaits further research.

Summary

Coordinated and sufficient uterine vascular remodeling must occur to facilitate the many-fold increase in uteroplacental blood flow that is required for normal pregnancy outcome. This physiological process is unique in the adult vascular system and likely occurs via a combination of mechanisms. Table 1 and FIGURE 6 are intended to provide a summary of the pathways, signals, and physiological processes that may be involved in uterine vascular remodeling during gestation. This table and figure also provide a basis for the consideration of future research, since the evidence implicating the involvement of these pathways and signals is largely indirect and inferential, i.e., based on other studies conducted in settings outside of the subject of this review.

Further investigations whose experimental design is clever enough to provide more specific, definitive information would help us bridge the theoretical with the real and provide deeper understanding of a process that is, at this point in time, still largely defined in descriptive rather than mechanistic terms.

Acknowledgments

The authors gratefully acknowledge the research support of the National Heart, Lung, and Blood Institute (NIH RO1 HL-079253 and HL-073895).

References

1. Adamson SL, Lu Y, Whiteley KJ, Holmyard D, Hemberger M, Pfarrer C, Cross JC. Interactions between trophoblast cells and the maternal and fetal circulation in the mouse placenta. *Dev Biol* 2002;250:358–373. [PubMed: 12376109]

2. Annibale DJ, Rosenfeld CR, Stull JT, Kamm KE. Protein content and myosin light chain phosphorylation in uterine arteries during pregnancy. *Am J Physiol Cell Physiol* 1990;259:C484–C489.
3. Assali NS, Douglass RA Jr, Baird WW, Nicholson DB, Suyemoto R. Measurement of uterine blood flow and uterine metabolism. IV. Results in normal pregnancy. *Am J Obstet Gynecol* 1953;66:248–253. [PubMed: 13065329]
4. Assali NS, Rauramo L, Peltonen T. Measurement of uterine blood flow and uterine metabolism VIII. Uterine and fetal blood flow and oxygen consumption in early human pregnancy. *Am J Obstet Gynecol* 1960;79:86–98. [PubMed: 13794853]
5. Ben Driss A, Benessiano J, Poitevin P, Levy BI, Michel JB. Arterial expansive remodeling induced by high flow rates. *Am J Physiol Heart Circ Physiol* 1997;272:H851–H858.
6. Bjellin L, Sjoquist PO, Carter AM. Uterine, maternal placental and ovarian blood flow throughout pregnancy in the guinea pig. *Zeitschrift fur Geburtshilfe und Perinatologie* 1975;179:179–187. [PubMed: 1189513]
7. Blechner JN, Stenger VG, Prystowsky H. Uterine blood flow in women at term. *Am J Obstet Gynecol* 1974;120:633–640. [PubMed: 4421293]
8. Brosens I, Robertson WB, Dixon HG. The physiological response of the vessels of the placental bed to normal pregnancy. *J Pathol Bacteriol* 1967;93:569–579. [PubMed: 6054057]
9. Brosens JJ, Pijnenborg R, Brosens IA. The myometrial junctional zone spiral arteries in normal and abnormal pregnancies: a review of the literature. *Am J Obstet Gynecol* 2002;187:1416–1423. [PubMed: 12439541]
10. Bruce NW. The distribution of blood flow to the reproductive organs of rats near term. *J Reprod Fertil* 1976;46:359–362. [PubMed: 1255563]
11. Buhimschi CS, Buhimschi IA, Malinow AM, Weiner CP. Myometrial thickness during human labor and immediately post partum. *Am J Obstet Gynecol* 2003;188:553–559. [PubMed: 12592271]
12. Bujold E, Romero R, Chaiworapongsa T, Kim YM, Kim GJ, Kim MR, Espinoza J, Goncalves LF, Edwin S, Mazor M. Evidence supporting that the excess of the sVEGFR-1 concentration in maternal plasma in preeclampsia has a uterine origin. *J Matern Fetal Neonatal Med* 2005;18:9–16. [PubMed: 16105786]
13. Buus CL, Pourageaud F, Fazzi GE, Janssen G, Mulvany MJ, De Mey JG. Smooth muscle cell changes during flow-related remodeling of rat mesenteric resistance arteries. *Circ Res* 2001;89:180–186. [PubMed: 11463726]
14. Caluwaerts S, Vercruysse L, Luyten C, Pijnenborg R. Endovascular trophoblast invasion and associated structural changes in uterine spiral arteries of the pregnant rat. *Placenta* 2005;26:574–584. [PubMed: 15993707]
15. Caton D, Kalra PS. Endogenous hormones and regulation of uterine blood flow during pregnancy. *Am J Physiol Regul Integr Comp Physiol* 1986;250:R365–R369.
16. Ceiler DL, De Mey JG. Chronic N^G-nitro-L-arginine methyl ester treatment does not prevent flow-induced remodeling in mesenteric feed arteries and arcading arterioles. *Arterioscler Thromb Vase Biol* 2000;20:2057–2063.
17. Celia G, Osol G. Uterine venous permeability in the rat is altered in response to pregnancy, vascular endothelial growth factor, and venous constriction. *Endothelium* 2005;12:81–88. [PubMed: 16036319]
18. Celia G, Osol G. Venoarterial communication as a mechanism for localized signaling in the rat uterine circulation. *Am J Obstet Gynecol* 2002;187:1653–1659. [PubMed: 12501079]
19. Chang K, Lubo Z. Review article: steroid hormones and uterine vascular adaptation to pregnancy. *Reprod Sci* 2008;15:336–348. [PubMed: 18497342]
20. Cipolla M, Osol G. Hypertrophic and hyperplastic effects of pregnancy on the rat uterine arterial wall. *Am J Obstet Gynecol* 1994;171:805–811. [PubMed: 8092233]
21. Cipolla MJ, Binder ND, Osol G. Myoendometrial versus placental uterine arteries: structural, mechanical, and functional differences in late-pregnant rabbits. *Am J Obstet Gynecol* 1997;177:215–221. [PubMed: 9240609]
22. Craven CM, Morgan T, Ward K. Decidual spiral artery remodelling begins before cellular interaction with cytotrophoblasts. *Placenta* 1998;19:241–252. [PubMed: 9639319]

23. D'Angelo G, Osol G. Modulation of uterine resistance artery lumen diameter by calcium and G protein activation during pregnancy. *Am J Physiol Heart Circ Physiol* 267:H952–H961.
24. D'Angelo G, Osol G. Regional variation in resistance artery diameter responses to alpha-adrenergic stimulation during pregnancy. *Am J Physiol Heart Circ Physiol* 1993;264:H78–H85.
25. Dechend R, Muller DN, Wallukat G, Homuth V, Krause M, Dudenhausen J, Luft FC. AT1 receptor agonistic antibodies, hypertension, and preeclampsia. *Seminars Nephrol* 2004;24:571–579.
26. Dowell RT, Kauer CD. Maternal hemodynamics and uteroplacental blood flow throughout gestation in conscious rats. *Meth Find Exp Ciin Pharmacol* 1997;19:613–625.
27. Dumont O, Loufrani L, Henrion D. Key role of the NO-pathway and matrix metalloprotease-9 in high blood flow-induced remodeling of rat resistance arteries. *Arterioscler Thromb Vase Biol* 2007;27:317–324.
28. Eastabrook G, Hu Y, von Dadelszen P. The role of decidual natural killer cells in normal placentation and in the pathogenesis of preeclampsia. *J Obstet Gynaecol Can* 2008;30:467–476. [PubMed: 18611298]
29. Edman CD, Toofanian A, MacDonald PC, Gant NF. Placental clearance rate of maternal plasma androstenedione through placental estradiol formation: an indirect method of assessing uteroplacental blood flow. *Am J Obstet Gynecol* 1981;141:1029–1037. [PubMed: 7315914]
30. Erami C, Zhang H, Tanoue A, Tsujimoto G, Thomas SA, Faber JE. Adrenergic catecholamine trophic activity contributes to flow-mediated arterial remodeling. *Am J Physiol Heart Circ Physiol* 2005;289:H744–H753. [PubMed: 15849236]
31. Espinoza J, Romero R, Mee Kim Y, Kusanovic JP, Hassan S, Erez O, Gotsch F, Than NG, Papp Z, Jai Kim C. Normal and abnormal transformation of the spiral arteries during pregnancy. *J Perinat Med* 2006;34:447–458. [PubMed: 17140293]
32. Farrer-Brown G, Beilby JO, Tarbit MH. The blood supply of the uterus, 1. Arterial vasculature. *J Obst Gynaecol Br Commonwealth* 1970;77:673–681.
33. Farrer-Brown G, Beilby JO, Tarbit MH. The blood supply of the uterus, 2. Venous pattern. *J Obst Gynaecol Br Commonwealth* 1970;77:682–689.
34. Forbes TR, Taku E. Vein size in intact and hysterectomized mice during the estrous cycle and pregnancy. *Anat Rec* 1975;182:61–65. [PubMed: 1155791]
35. Ford SP, Chenault JR, Echternkamp SE. Uterine blood flow of cows during the oestrous cycle and early pregnancy: effect of the conceptus on the uterine blood supply. *J Reprod Fertil* 1979;56:53–62. [PubMed: 469858]
36. Ford SP, Christenson RK. Blood flow to uteri of sows during the estrous cycle and early pregnancy: local effect of the conceptus on the uterine blood supply. *Biol Reprod* 1979;21:617–624. [PubMed: 497319]
37. Ford SP, Reynolds LP, Ferrell CL. Blood flow, steroid secretion and nutrient uptake of the gravid uterus during the periparturient period in sows. *J Anim Sci* 1984;59:1085–1091. [PubMed: 6511680]
38. Ford SP, Stice SL. Effects of the ovary and conceptus on uterine blood flow in the pig. *J Reprod Fertil Suppl* 1985;33:83–90. [PubMed: 2418203]
39. Geisterfer AA, Peach MJ, Owens GK. Angiotensin II induces hypertrophy, not hyperplasia, of cultured rat aortic smooth muscle cells. *Circ Res* 1988;62:749–756. [PubMed: 3280155]
40. Ginther OJ. Internal regulation of physiological processes through local venoarterial pathways: a review. *J Anim Sci* 1974;39:550–564. [PubMed: 4213194]
41. Gokina NI, Mandala M, Osol G. Induction of localized differences in rat uterine radial artery behavior and structure during gestation. *Am J Obstet Gynecol* 2003;189:1489–1493. [PubMed: 14634590]
42. Goldman-Wohl DS, Yagel S. Examination of distinct fetal and maternal molecular pathways suggests a mechanism for the development of preeclampsia. *J Reprod Immunol* 2007;76:54–60. [PubMed: 17482678]
43. Grandas OH, Mountain DH, Kirkpatrick SS, Cassada DC, Stevens SL, Freeman MB, Goldman MH. Regulation of vascular smooth muscle cell expression and function of matrix metalloproteinases is mediated by estrogen and progesterone exposure. *J Vase Surg*. In press
44. Greiss FC Jr, Anderson SG. Uterine blood flow during early ovine pregnancy. *Am J Obstet Gynecol* 1970;106:30–38. [PubMed: 5460660]

45. Griendling KK, Fuller EO, Cox RH. Pregnancy-induced changes in sheep uterine and carotid arteries. *Am J Physiol Heart Circ Physiol* 1985;248:H658–H665.
46. Guenther AE, Conley AJ, Van Orden DE, Farley DB, Ford SP. Structural and mechanical changes of uterine arteries during pregnancy in the pig. *J Anim Sci* 1988;66:3144–3152. [PubMed: 2852664]
47. Hammer ES, Cipolla MJ. Arterial wall hyperplasia is increased in placental compared with myoendometrial radial uterine arteries from late-pregnant rats. *Am J Obstet Gynecol* 2005;192:302–308. [PubMed: 15672040]
48. Hart MV, Hosenpud JD, Hohimer AR, Morton MJ. Hemodynamics during pregnancy and sex steroid administration in guinea pigs. *Am J Physiol Regul Integr Comp Physiol* 1985;249:R179–R185.
49. Hees H, Moll W, Wrobel KH, Hees I. Pregnancy-induced structural changes and trophoblastic invasion in the segmental mesometrial arteries of the guinea pig (*Cavia porcellus* L.). *Placenta* 1987;8:609–627. [PubMed: 3438257]
50. Herve MA, Meduri G, Petit FG, Domet TS, Lazennec G, Mourah S, Perrot-Applanat M. Regulation of the vascular endothelial growth factor (VEGF) receptor Flk-1/KDR by estradiol through VEGF in uterus. *J Endocrinol* 2006;188:91–99. [PubMed: 16394178]
51. Hollestelle SC, De Vries MR, Van Keulen JK, Schoneveld AH, Vink A, Strijder CF, Van Middelaar BJ, Pasterkamp G, Quax PH, De Kleijn DP. Toll-like receptor 4 is involved in outward arterial remodeling. *Circulation* 2004;109:393–398. [PubMed: 14699006]
52. Jeyabalan A, Conrad KP. Renal function during normal pregnancy and preeclampsia. *Front Biosci* 2007;12:2425–2437. [PubMed: 17127252]
53. Karumanchi SA, Lindheimer MD. Advances in the understanding of eclampsia. *Curr Hyperten Rep* 2008;10:305–312.
54. Kaufmann P, Black S, Huppertz B. Endovascular trophoblast invasion: implications for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod* 2003;69:1–7. [PubMed: 12620937]
55. Kelly B, Stone S, Poston L. Cardiovascular adaptation to pregnancy: the role of altered vascular structure. *Fetal Matern Med Rev* 1999;11:105–116.
56. Keyes LE, Majack R, Dempsey EC, Moore LG. Pregnancy stimulation of DNA synthesis and uterine blood flow in the guinea pig. *Pediatr Res* 1997;41:708–715. [PubMed: 9128295]
57. Keyes LE, Moore LG, Walchak SJ, Dempsey EC. Pregnancy-stimulated growth of vascular smooth muscle cells: importance of protein kinase C-dependent synergy between estrogen and platelet-derived growth factor. *J Cell Physiol* 1996;166:22–32.
58. Kim KH, Bender JR. Rapid, estrogen receptor-mediated signaling: why is the endothelium so special? *Sci STKE* 2005;2005:pe28. [PubMed: 15956360]
59. Kim KH, Moriarty K, Bender JR. Vascular cell signaling by membrane estrogen receptors. *Steroids* 2008;73:864–869. [PubMed: 18325557]
60. Knight JW, Bazer FW, Thatcher WW, Franke DE, Wallace HD. Conceptus development in intact and unilaterally hysterectomized-ovariectomized gilts: interrelations among hormonal status, placental development, fetal fluids and fetal growth. *J Anim Sci* 1977;44:620–637. [PubMed: 853021]
61. Koga K, Osuga Y, Yoshino O, Hirota Y, Ruimeng X, Hirata T, Takeda S, Yano T, Tsutsumi O, Taketani Y. Elevated serum soluble vascular endothelial growth factor receptor 1 (sVEGFR-1) levels in women with preeclampsia. *J Clin Endocrinol Metab* 2003;88:2348–2351. [PubMed: 12727995]
62. Lang U, Baker RS, Braems G, Zygmunt M, Kunzel W, Clark KE. Uterine blood flow: a determinant of fetal growth. *Eur J Obstet Gynecol Reprod Biol* 2003;110:S55–S61. [PubMed: 12965091]
63. Langille BL. Arterial remodeling: relation to hemodynamics. *Can J Physiol Pharmacol* 1996;74:834–841. [PubMed: 8946070]
64. Langille BL, O'Donnell F. Reductions in arterial diameter produced by chronic decreases in blood flow are endothelium-dependent. *Science* 1986;231:405–407. [PubMed: 3941904]
65. Lees MH, Hill JD, Ochsner AJ 3rd, Thomas CL, Novy MJ. Maternal placental and myometrial blood flow of the rhesus monkey during uterine contractions. *Am J Obstet Gynecol* 1971;110:68–81. [PubMed: 4995593]

66. Leiberman JR, van Vroonhoven CC, Beckmann I, van der Kwast TH, Wallenburg HC. Uterine artery estrogen receptors in the nonpregnant and pregnant guinea pig. *Am J Obstet Gynecol* 1990;163:1685–1688. [PubMed: 2240124]
67. Leiberman JR, Wiznitzer A, Glezerman M, Feldman B, Levy J, Sharoni Y. Estrogen and progesterone receptors in the uterine artery of rats during and after pregnancy. *Eur J Obstet Gynecol Reprod Biol* 1993;51:35–40. [PubMed: 8282141]
68. Liao WX, Magness RR, Chen DB. Expression of estrogen receptors-alpha and -beta in the pregnant ovine uterine artery endothelial cells in vivo and in vitro. *Biol Reprod* 2005;72:530–537. [PubMed: 15564597]
69. Loufrani L, Henrion D. Role of the cytoskeleton in flow (shear stress)-induced dilation and remodeling in resistance arteries. *Med Biol Eng Comput* 2008;46:451–460. [PubMed: 18246377]
70. Lu F, Longo M, Tamayo E, Maner W, Al-Hendy A, Anderson GD, Hankins GD, Saade GR. The effect of over-expression of sFlt-1 on blood pressure and the occurrence of other manifestations of preeclampsia in unrestrained conscious pregnant mice. *Am J Obstet Gynecol* 2007;196(396):e391–e397.
71. Lyall F. Priming and remodelling of human placental bed spiral arteries during pregnancy: a review. *Placenta* 2005;26:S31–S36. [PubMed: 15837064]
72. Magness RR, Parker CR Jr, Rosenfeld CR. Systemic and uterine responses to chronic infusion of estradiol-17 beta. *Am J Physiol Endocrinol Metab* 1993;265:E690–E698.
73. Magness RR, Rosenfeld CR. Local and systemic estradiol-17 beta: effects on uterine and systemic vasodilation. *Am J Physiol Endocrinol Metab* 1989;256:E536–E542.
74. Makinoda S, Moll W. Deoxyribonucleic acid synthesis in mesometrial arteries of guinea pigs during oestrous cycle, pregnancy and treatment with oestradiol benzoate. *Placenta* 1986;7:189–198. [PubMed: 3737574]
75. Mateev SN, Mouser R, Young DA, Mecham RP, Moore LG. Chronic hypoxia augments uterine artery distensibility and alters the circumferential wall stress-strain relationship during pregnancy. *J Appl Physiol* 2006;100:1842–1850. [PubMed: 16714414]
76. Maynard SE, Venkatesha S, Thadhani R, Karumanchi SA. Soluble Fms-like tyrosine kinase 1 and endothelial dysfunction in the pathogenesis of preeclampsia. *Pediatr Res* 2005;57:1R–7R. [PubMed: 15557110]
77. Merchant SJ, Davidge ST. The role of matrix metalloproteinases in vascular function: implications for normal pregnancy and pre-eclampsia. *Bjog* 2004;111:931–939. [PubMed: 15327607]
78. Metcalfe J, Romney SL, Ramsey LH, Reid DE, Burwell CS. Estimation of uterine blood flow in normal human pregnancy at term. *J Clin Invest* 1955;34:1632–1638. [PubMed: 13271546]
79. Moll W. Structure adaptation and blood flow control in the uterine arterial system after hemochorial placentation. *Eur J Obstet Gynecol Reprod Biol* 2003;110:S19–S27. [PubMed: 12965087]
80. Moll W, Espach A, Wrobel KH. Growth of mesometrial arteries in guinea pigs during pregnancy. *Placenta* 1983;4:111–123. [PubMed: 6878181]
81. Moll W, Kunzel W. The blood pressure in arteries entering the placentae of guinea pigs, rats, rabbits, and sheep. *Pflugers Arch* 1973;338:125–131. [PubMed: 4734441]
82. Moll W, Kunzel W. Blood pressures in the uterine vascular system of anaesthetized pregnant guinea pigs. *Pflugers Arch* 1971;330:310–322. [PubMed: 5169236]
83. Moll W, Kunzel W, Herberger J. Hemodynamic implications of hemochorial placentation. *Eur J Obstet Gynecol Reprod Biol* 1975;5:67–74. [PubMed: 1053578]
84. Moll W, Kunzel W, Stolte A, Kleinhout J, de Jong P, Veth A. The blood pressure in the decidual part of the uteroplacental arteries (spiral arteries) of the Rhesus monkey. *Pflugers Arch* 1973;346:291–297.
85. Moll W, Nienartowicz A, Hees H, Wrobel KH, Lenz A. Blood flow regulation in the uteroplacental arteries. *Troph Res* 1988;3:83–96.
86. Mulvany MJ, Baumbach GL, Aalkjaer C, Heagerty AM, Korsgaard N, Schiffrin EL, Heistad DD. Vascular remodeling. *Hypertension* 1996;28:505–506. [PubMed: 8794840]
87. Myers JE, Merchant SJ, Macleod M, Mires GJ, Baker PN, Davidge ST. MMP-2 levels are elevated in the plasma of women who subsequently develop preeclampsia. *Hypertens Pregnancy* 2005;24:103–115. [PubMed: 16036395]

88. Nanaev A, Chwalisz K, Frank HG, Kohnen G, Hegele-Hartung C, Kaufmann P. Physiological dilation of uteroplacental arteries in the guinea pig depends on nitric oxide synthase activity of extravillous trophoblast. *Cell Tissue Res* 1995;282:407–421. [PubMed: 8581935]
89. Natoli AK, Medley TL, Ahimastos AA, Drew BG, Thearle DJ, Dilley RJ, Kingwell BA. Sex steroids modulate human aortic smooth muscle cell matrix protein deposition and matrix metalloproteinase expression. *Hypertension* 2005;46:1129–1134. [PubMed: 16230520]
90. Nelson SH, Steinsland OS, Suresh MS, Lee NM. Pregnancy augments nitric oxide-dependent dilator response to acetylcholine in the human uterine artery. *Hum Reprod* 1998;13:1361–1367. [PubMed: 9647573]
91. Nelson SH, Steinsland OS, Wang Y, Yallampalli C, Dong YL, Sanchez JM. Increased nitric oxide synthase activity and expression in the human uterine artery during pregnancy. *Circ Res* 2000;87:406–411. [PubMed: 10969039]
92. Nienartowicz A, Link S, Moll W. Adaptation of the uterine arcade in rats to pregnancy. *J Dev Physiol* 1989;12:101–108. [PubMed: 2621336]
93. Odorisio T, Schietroma C, Zaccaria ML, Cianfarani F, Tiverson C, Tatangelo L, Failla CM, Zambruno G. Mice overexpressing placenta growth factor exhibit increased vascularization and vessel permeability. *J Cell Sci* 2002;115:2559–2567. [PubMed: 12045226]
94. Ong SS, Baker PN, Mayhew TM, Dunn WR. Remodeling of myometrial radial arteries in preeclampsia. *Am J Obstet Gynecol* 2005;192:572–579. [PubMed: 15696005]
95. Oosterhof H, Wichers G, Fidler V, Aarnoudse JG. Blood viscosity and uterine artery flow velocity waveforms in pregnancy: a longitudinal study. *Placenta* 1993;14:555–561. [PubMed: 8290495]
96. Osol G, Barron C, Gokina N, Mandala M. Inhibition of nitric oxide synthases abrogates pregnancy-induced uterine vascular expansive remodeling. *J Vase Res*. In press
97. Osol G, Cipolla M. Pregnancy-induced changes in the three-dimensional mechanical properties of pressurized rat uteroplacental (radial) arteries. *Am J Obstet Gynecol* 1993;168:268–274. [PubMed: 8420338]
98. Page KL, Celia G, Leddy G, Taatjes DJ, Osol G. Structural remodeling of rat uterine veins in pregnancy. *Am J Obstet Gynecol* 2002;187:1647–1652. [PubMed: 12501078]
99. Palmer SK, Zamudio S, Coffin C, Parker S, Stamm E, Moore LG. Quantitative estimation of human uterine artery blood flow and pelvic blood flow redistribution in pregnancy. *Obstet Gynecol* 1992;80:1000–1006. [PubMed: 1448242]
100. Pasterkamp G, Galis ZS, de Kleijn DP. Expansive arterial remodeling: location, location, location. *Arterioscler Thromb Vase Biol* 2004;24:650–657.
101. Pijnenborg R, Robertson WB, Brosens I, Dixon G. Review article: trophoblast invasion and the establishment of haemochorial placentation in man and laboratory animals. *Placenta* 1981;2:71–91. [PubMed: 7010344]
102. Pries AR, Reglin B, Secomb TW. Remodeling of blood vessels: responses of diameter and wall thickness to hemodynamic and metabolic stimuli. *Hypertension* 2005;46:725–731. [PubMed: 16172421]
103. Pustovrh MC, Jawerbaum A, White V, Capobianco E, Higa R, Martinez N, Lopez-Costa JJ, Gonzalez E. The role of nitric oxide on matrix metalloproteinase 2 (MMP2) and MMP9 in placenta and fetus from diabetic rats. *Reproduction* 2007;134:605–613. [PubMed: 17890296]
104. Raffetto JD, Khalil RA. Matrix metalloproteinases and their inhibitors in vascular remodeling and vascular disease. *Biochem Pharmacol* 2008;75:346–359. [PubMed: 17678629]
105. Ramsey EM, Houston ML, Harris JW. Interactions of the trophoblast and maternal tissues in three closely related primate species. *Am J Obstet Gynecol* 1976;124:647–652. [PubMed: 816200]
106. Rekonen A, Luotola H, Pitkanen M, Kuikka J, Pyorala T. Measurement of intervillous and myometrial blood flow by an intravenous ¹³³Xe method. *Br J Obstet Gynaecol* 1976;83:723–728. [PubMed: 974049]
107. Reneman RS, Arts T, Hoeks AP. Wall shear stress: an important determinant of endothelial cell function and structure—in the arterial system in vivo. Discrepancies with theory. *J Vase Res* 2006;43:251–269.
108. Roberts JM. Angiotensin-1 receptor autoantibodies: a role in the pathogenesis of preeclampsia? *Circulation* 2000;101:2335–2337. [PubMed: 10821805]

109. Robertson WB, Manning PJ. Elastic tissue in uterine blood vessels. *J Pathol* 1973;109:P3. [PubMed: 4719740]
110. Rodbard S. Vascular caliber. *Cardiology* 1975;60:4–49. [PubMed: 126799]
111. Rosenfeld CR. Mechanisms regulating angiotensin II responsiveness by the uteroplacental circulation. *Am J Physiol Regul Integr Comp Physiol* 2001;281:R1025–R1040. [PubMed: 11557608]
112. Rosenfeld CR, Morriss FH Jr, Makowski EL, Meschia G, Battaglia FC. Circulatory changes in the reproductive tissues of ewes during pregnancy. *Gynecol Invest* 1974;5:252–268. [PubMed: 4461407]
113. Roy-Clavel E, Picard S, St-Louis J, Brochu M. Induction of intrauterine growth restriction with a low-sodium diet fed to pregnant rats. *Am J Obstet Gynecol* 1999;180:608–613. [PubMed: 10076136]
114. Rudic RD, Shesely EG, Maeda N, Smithies O, Segal SS, Sessa WC. Direct evidence for the importance of endothelium-derived nitric oxide in vascular remodeling. *J Clin Invest* 1998;101:731–736. [PubMed: 9466966]
115. Santoni A, Carlino C, Gismondi A. Uterine NK cell development, migration and function. *Reprod Biomed Online* 2008;16:202–210. [PubMed: 18284874]
116. Sargent IL, Borzychowski AM, Redman CW. Immunoregulation in normal pregnancy and preeclampsia: an overview. *Reprod Biomed Online* 2006;13:680–686. [PubMed: 17169180]
117. Schiffers PM, Henrion D, Boulanger CM, Colucci-Guyon E, Langa-Vuves F, van Essen H, Fazzi GE, Levy BI, De Mey JG. Altered flow-induced arterial remodeling in vimentin-deficient mice. *Arterioscler Thromb Vase Biol* 2000;20:611–616.
118. Sladek SM, Magness RR, Conrad KP. Nitric oxide and pregnancy. *Am J Physiol Regul Integr Comp Physiol* 1997;272:R441–R463.
119. St-Louis J, Sicotte B, Beausejour A, Brochu M. Remodeling and angiotensin II responses of the uterine arcuate arteries of pregnant rats are altered by low- and high-sodium intake. *Reproduction* 2006;131:331–339. [PubMed: 16452726]
120. St-Louis J, Sicotte B, Bedard S, Brochu M. Blockade of angiotensin receptor subtypes in arcuate uterine artery of pregnant and postpartum rats. *Hypertension* 2001;38:1017–1023. [PubMed: 11711491]
121. Staff AC, Braekke K, Harsem NK, Lyberg T, Holthe MR. Circulating concentrations of sFlt1 (soluble fms-like tyrosine kinase 1) in fetal and maternal serum during pre-eclampsia. *Eur J Obstet Gynecol Reprod Biol* 2005;122:33–39. [PubMed: 15935542]
122. Stewart EA, Nowak RA. New concepts in the treatment of uterine leiomyomas. *Obstet Gynecol* 1998;92:624–627. [PubMed: 9764641]
123. Storment JM, Meyer M, Osol G. Estrogen augments the vasodilatory effects of vascular endothelial growth factor in the uterine circulation of the rat. *Am J Obstet Gynecol* 2000;183:449–453. [PubMed: 10942485]
124. Tabata M, Negishi H, Yamaguchi T, Makinoda S, Fujimoto S, Moll W. The effects of the proliferation of the radial arteries of the placenta on oxygen transport to the fetal guinea pig. *Adv Exp Med Biol* 1988;222:675–681. [PubMed: 3364293]
125. Tai LK, Okuda M, Abe J, Yan C, Berk BC. Fluid shear stress activates proline-rich tyrosine kinase via reactive oxygen species-dependent pathway. *Arterioscler Thromb Vase Biol* 2002;22:1790–1796.
126. Taylor RN, Grimwood J, Taylor RS, McMaster MT, Fisher SJ, North RA. Longitudinal serum concentrations of placental growth factor: evidence for abnormal placental angiogenesis in pathologic pregnancies. *Am J Obstet Gynecol* 2003;188:177–182. [PubMed: 12548214]
127. Thaler I, Manor D, Itskovitz J, Rottem S, Levit N, Timor-Tritsch I, Brandes JM. Changes in uterine blood flow during human pregnancy. *Am J Obstet Gynecol* 1990;162:121–125. [PubMed: 2301480]
128. Thorbert G, Aim P, Owman C, Sjoberg NO, Sporrang B. Regional changes in structural and functional integrity of myometrial adrenergic nerves in pregnant guinea-pig, and their relationship to the localization of the conceptus. *Acta Physiol Scand* 1978;103:120–131. [PubMed: 676764]
129. Thornburg KL, Jacobson SL, Giraud GD, Morton MJ. Hemodynamic changes in pregnancy. *Semin Perinatol* 2000;24:11–14. [PubMed: 10709851]

130. Tronc F, Mallat Z, Lehoux S, Wassef M, Esposito B, Tedgui A. Role of matrix metalloproteinases in blood flow-induced arterial enlargement: interaction with NO. *Arterioscler Thromb Vase Biol* 2000;20:E120–E126.
131. Tuttle JL, Nachreiner RD, Bhuller AS, Condict KW, Connors BA, Herring BP, Dalsing MC, Unthank JL. Shear level influences resistance artery remodeling: wall dimensions, cell density, and eNOS expression. *Am J Physiol Heart Circ Physiol* 2001;281:H1380–H1389. [PubMed: 11514310]
132. van der Heijden OW, Essers YP, Fazzi G, Peeters LL, De Mey JG, van Eys GJ. Uterine artery remodeling and reproductive performance are impaired in endothelial nitric oxide synthase-deficient mice. *Biol Reprod* 2005;72:1161–1168. [PubMed: 15659709]
133. van der Heijden OW, Essers YP, Simkens LH, Teunissen QG, Peeters LL, De Mey JG, van Eys GJ. Aging blunts remodeling of the uterine artery during murine pregnancy. *J Soc Gynecol Invest* 2004;11:304–310.
134. van der Heijden OW, Essers YP, Spaanderman ME, De Mey JG, van Eys GJ, Peeters LL. Uterine artery remodeling in pseudopregnancy is comparable to that in early pregnancy. *Biol Reprod* 2005;73:1289–1293. [PubMed: 16120827]
135. Verlohren S, Niehoff M, Hering L, Geusens N, Herse F, Tintu AN, Plagemann A, LeNoble F, Pijnenborg R, Muller DN, Luft FC, Dudenhausen JW, Gollasch M, Dechend R. Uterine vascular function in a transgenic preeclampsia rat model. *Hypertension* 2008;51:547–553. [PubMed: 18195162]
136. Waddell BJ, Burton PJ. Full induction of rat myometrial 11beta-hydroxysteroid dehydrogenase type 1 in late pregnancy is dependent on intrauterine occupancy. *Biol Reprod* 2009;62:1005–1009. [PubMed: 10727270]
137. Whitney EA, Ducsay CA, Valenzuela GJ. Is uterine blood flow controlled locally or systemically in the pregnant rabbit? *Am J Obstet Gynecol* 1993;169:1507–1509. [PubMed: 8267054]
138. Zhang Y, Stewart KG, Davidge ST. Endogenous estrogen mediates vascular reactivity and distensibility in pregnant rat mesenteric arteries. *Am J Physiol Heart Circ Physiol* 2001;280:H956–H961. [PubMed: 11179035]
139. Zhang Y, Stewart KG, Davidge ST. Estrogen replacement reduces age-associated remodeling in rat mesenteric arteries. *Hypertension* 2000;36:970–974. [PubMed: 11116109]
140. Zhou CC, Zhang YY, Irani RA, Zhang H, Mi T, Popek EJ, Hicks MJ, Ramin SM, Kellems RE, Xia Y. Angiotensin receptor agonistic autoantibodies induce pre-eclampsia in pregnant mice. *Nat Med* 2008;14:855–862. [PubMed: 18660815]

FIGURE 1. Patterns of vascular remodeling

This drawing relates changes in lumen diameter and cross-sectional area to illustrate the various two-dimensional patterns of arterial remodeling (adapted, with permission, from R. H. Hilgers). The remodeling of a vessel to a larger lumen with the same wall thickness, such as occurs in the uterine circulation during pregnancy (white shaded box), is termed “outward hypertrophic,” since cross-sectional area is increased. Conversely, vessel narrowing with increased wall thickness occurs in chronic hypertension and may be inward eutrophic (smaller lumen with a somewhat thicker wall, but the same cross-sectional area, characteristic of smaller resistance arteries) or inward hypertrophic (smaller lumen with sufficient wall thickening to increase cross-sectional area, characteristic of larger conduit vessels). A common assumption is that changes in cross-sectional area indicate changes in wall mass (as implied by the terms “hypotrophic” or “hypertrophic”). This, of course, is only correct if vessel length is not altered. Although changes in venous or arterial length rarely occur in the adult and are therefore rarely measured, the uterine circulation during pregnancy is one notable and pertinent exception since existing vessels do undergo considerable elongation, thereby increasing wall mass further.

Thus changes in both cross-sectional area and axial length must be considered to evaluate true changes in mass.

FIGURE 2. Comparative anatomy of the uterine circulation in humans, rodents, and ungulates
 In most mammals, including humans (A), blood is delivered to the uterus bidirectionally via a dual arterial anastomotic loop in which one end (the two ovarian arteries) originates from the aorta and the other (uterine arteries) from the internal iliacs. Unlike the linear, branching pattern of many vascular networks in which blockage leads to downstream ischemia, this bilateral anatomical arrangement provides the uterus with a dual source of blood and considerable redundancy in case of occlusion. Perpendicular vessels arise from the main utero-ovarian arteries and pass into the body of the uterus to form the arcuate arteries, which encircle the organ by coursing within the myometrium just beneath its serosal (outer) surface; vessels from each side anastomose along the uterine midline. Smaller radial arteries (see *inset*) emanate

from the arcuates and penetrate the myometrium centripetally before ramifying into either straight (basal) or coiled (spiral) arteries at the myoendometrial border. The basal vessels spread to form a network along the myoendometrial border, while the spiral arteries penetrate further into the endometrium and terminate close to the uterine lumen in capillaries that are, in turn, drained by venules that coalesce into larger veins that eventually enter the inferior vena cava (not shown). Rodents (*B*) have a duplex uterus, with the main utero-ovarian (or parametrial) arteries and veins running parallel to, but well outside of, the uterine wall within a planar sheet of connective tissue called the mesometrium (shown by tan shading). The vessels of the mesometrium are perfused by arterial blood coming from either the uterine or the ovarian end, i.e., with bidirectional flow. Secondary vessels analogous to the arcuate arteries in humans may form redundant loops with the main artery, and tertiary radial arteries connect the arcuate loops with the uterine wall. These radial (also called mesometrial or segmental) arteries can be further categorized based on their destination as being either premyometrial or preplacental. The premyometrial radial arteries enter the uterine wall between placentation sites and ramify into an intrauterine arterial plexus that supplies the myometrium, whereas preplacental radial vessels widen before entering the placenta through a process of endovascular trophoblast invasion. In addition to displaying distinctive patterns of remodeling during gestation, premyometrial vs. preplacental arteries have also been shown to have different patterns of reactivity (see text). In ungulates such as the sheep or pig, the main (middle) uterine artery originates from the umbilical branch of the internal iliac artery and divides into four primary branches that anastomose with contralateral vessels along the lesser curvature of the uterine horn. These vessels give rise to coiled, branching vessels that run along the ventral and dorsal surface of the uterus to form the arcuate arteries, with smaller branches (radial arteries) that penetrate the myometrium and terminate in arterioles within the endometrium. Most notably, whereas humans and rodents exhibit a hemochorial type of placentation (having low resistance), the placenta of sheep and pigs is epitheliochorial and therefore more analogous to a true microcirculatory bed. In humans, as well as other species (e.g., sheep, pig, rat, guinea pig), the uterus is drained by a venous system that parallels the arterial tree, with closely apposed arteries and veins.

FIGURE 3. Preplacental vs. premyometrial vessels in the rat

Photograph of a segment of the uterine mesometrium from a 20-day pregnant rat. A proximal radial artery (RA) divides into two branches, both of which approach the uterine wall (not visible). The lower vessel perfuses the placenta [preplacental artery (PPL)] while the upper courses to and penetrates the myometrium between implantation sites [premyometrial (PMM)]. A segment of a vein draining the placenta is visible at *top right*. Note consistent shape of the PMM vs. significant (2–3×) widening of the PPL. The area of widening likely denotes the extent of endovascular trophoblast invasion. Once the vessel has widened, it loses the ability to contract or dilate due to ablation and/or de-differentiation of vascular smooth muscle in the arterial media.

FIGURE 4. Local mechanisms predominate in uterine vascular remodeling in the rodent

A: photograph of a 20-day pregnant rat uterus in which one uterine horn was ligated at the ovarian end, preventing oocyte descent and fertilization. Thirteen fetoplacental units are present in the pregnant uterine horn (PH); the nonpregnant horn is at the bottom of the picture (NPH). CERV, cervix. **B:** the arterial vasculature of the specimen in A was infused with a latex casting compound and allowed to harden. Uterine and fetoplacental tissues were digested in KOH, leaving a cast of the arterial circulation of each horn. The main uterine arteries (MUA) and the mesometrial arteries (MESOM; arcuate, radial) are labeled, as are two small tree-like structures that represent the maternal intraplacental compartment typical of rodent hemochorial placentation (PL).

FIGURE 5. Venoarterial exchange

Signals secreted from the placenta, decidua, or myometrium (due to stretch) pass into the venous effluent, where their concentrations would be highest before dilution into the systemic circulation. Signals may be molecules that are vasoactive, mitogenic, or hypertrophic. These, in turn, pass across the venous wall to the adjacent artery to alter arterial tone and/or structure. This pathway could provide a mechanism for fetoplacental regulation of maternal blood flow. Although venoarterial exchange has been documented in the uterine circulation of a number of species as a mechanism for luteolysis (see text), its role in maternal uterine vascular remodeling during pregnancy is still hypothetical.

FIGURE 6. Four physiological mechanisms that may play a role in uterine arterial and venous widening and elongation during pregnancy

*Note: This generalized drawing of the uteroplacental circulation with hemochorial placentation is not meant to be anatomically accurate; rather, it summarizes the four principal mechanisms likely to be involved in gestational uterine vascular remodeling, as discussed in the text. 1) **Placentation/endovascular trophoblast invasion.** Hemochorial placentation provides a low-resistance pathway for maternal blood flow by eliminating the intramyometrial microcirculation and creating an intervillous space (IVS). Flow resistance is further decreased by arterial widening secondary to endovascular trophoblast invasion (green cells) of the preplacental (in humans, spiral) arteries. Invasion of the veins has also been documented, at*

least in rodents, although to a lesser depth. The depth of invasion is exaggerated for purposes of illustration. 2) **Increased shear stress.** Decreased resistance due to placentation and endovascular remodeling of distal (preplacental) arteries results in an acceleration of blood flow (Q) in proximal vessels, elevating shear stress at the endothelial surface (2, black arrows). Shear stimulates the release of endothelial NO and initiates outward circumferential (expansive) arterial remodeling. As vessels enlarge, higher flow can be maintained at a slower velocity, thereby normalizing shear stress at the endothelial surface. Venous shear stress must increase as well (since inflow must equal outflow), and venous circumferential enlargement and elongation clearly occur, although a role for shear stress in venous expansive remodeling has not yet been demonstrated experimentally. 3) **Venoarterial exchange.** 3a: placentally derived signals. The fetoplacental unit secretes a variety of signals (blue circles) into the IVS. These pass into the venous outflow, where they are most concentrated before systemic dilution and may 1) induce changes in venous structure and 2) enhance venous permeability through molecular mechanisms (e.g., VEGF, PlGF) and altered physical forces secondary to growth (e.g., increased wall tension). Both mechanisms have been described in the uterine circulation of rodents (see text), where arteries and veins are often in close apposition and would affect vessels within the area shown by the tan shaded box and indicated by the placement of symbols between the arteries and veins. Placental signals may include molecules that reduce arterial tone and stimulate cell division and/or hypertrophy, leading to vessel widening. 3b: myometrial signals. Mechanisms involved in the axial remodeling (elongation) of arteries and veins have not been identified to date. Here, we postulate that increasing volume of the conceptus (white arrows) leads to myometrial stretch. This, in turn, induces the release of mitogenic signals (black triangles) into the venous outflow, stimulating hyperplasia within the venous wall (and in the adjacent arterial wall via venoarterial exchange) and leading to structural elongation. Human arteries must also elongate during pregnancy, since the volume of the uterus increases approximately 1,000-fold, although it is not clear whether lengthening is due to arterial growth, elastic stretch, straightening of tortuous segments, or any combination thereof. In rodents, much of the vasculature is external to the myometrium and located in a planar mesometrium; in humans, arcuate and radial arteries are contained within the myometrium and may therefore be subject to direct stretch. 4) **Humoral factors.** The endocrine milieu of pregnancy results in altered systemic concentrations of hormones and growth factors (red circles). The source of hormones may be placental, although this is species dependent (e.g., in rodents, the ovary, not the placenta, produces estrogen and progesterone). Increased hormone concentrations may induce vasodilation, alter permeability, and stimulate both cellular and matrix remodeling.

Table 1
Processes, pathways, signals, and events implicated in maternal uterine vascular remodeling during pregnancy

Processes

Implantation/placentation
Endovascular trophoblast invasion
Myometrial stretch
Altered endocrine regulatory mechanisms

Pathways

Local
Physical (e.g., increased shear stress on endothelium)
Molecular (e.g., venoarterial exchange)
Systemic (humoral/endocrine)

Signals

Sex steroids
Nitric oxide
Prostaglandins
Growth factors and angiogenic molecules
Angiotensin
Other (?)

Events occurring within the vascular wall

Vasodilation
Cellular hypertrophy
Cellular hyperplasia
Matrix remodeling
Recruitment of periadventitial cells (?)